


# Microphysical Cloud Modeling of Substellar Atmospheres: Preliminary Findings

Wolf Cukier, Diana Powell  
University of Chicago




## Motivation





Clouds, being coupled to atmospheric processes, are highly complex and critical to a complete understanding of substellar atmospheres. Clouds are thought to cause color trends in the M-L-T-Y sequence of brown dwarfs and the abnormally red color of directly imaged giant planets. CARMA is a well-validated model that calculates the formation of clouds from first principles through modeling fundamental microphysical processes. It has been used to model clouds on every atmospheric solar system body as well as hot Jupiters, sub-Neptunes and Y-Dwarfs

## CARMA's Microphysics





\*Other species also included

## Preliminary Results



Size Distribution Profiles for  
Teff = 1400 K, log g = 3.5, Z = +0.0

- We have calculated a grid of cloud particle size distributions for:
  - Teff in (900K, 1000K, ... 2400 K)
  - log g in (3.5, 4.5, ... 5.5)
  - Metallicity in (-0.5, 0.0, +0.5)
- Above ~1000 K  $\text{Mg}_2\text{SiO}_4$  is dominant  
Below ~1000 K KCl is dominant  
(by total atmosphere cross-section)
- Hotter objects tend to have larger cloud particles. Cooler objects tend to have smaller cloud particles
- Clouds are variable on time scales of a few days



Size Distributions for  
log g = 3.5, Z = +0.0

## Next Steps

- Release particle size grid to the community
- Compare distributions with log-normal, bimodal, etc.
- Create emission spectra across the grid
- Release carmapy: a CARMA python wrapper

## Contact + About



Wolf Cukier (He/Him)  
1st year Graduate Student

Email: [wcukier@uchicago.edu](mailto:wcukier@uchicago.edu)  
Website: [wolfcukier.com](http://wolfcukier.com)

