

1 MEASUREMENT OF ELECTROWEAK PRODUCTION OF
2 SAME-SIGN W BOSON PAIRS WITH ATLAS (WORKING
3 TITLE)

4 William Kennedy DiClemente

8 Presented to the Faculties of The University of Pennsylvania
9 in Partial Fulfillment of the Requirements for the Degree of Doctor of Philosophy
10 2019 Last compiled: February 2, 2019

¹¹
¹² I. Joseph Kroll, Professor, Physics
¹³ Supervisor of Dissertation

23 MEASUREMENT OF ELECTROWEAK PRODUCTION OF SAME-SIGN W BOSON PAIRS
24 WITH ATLAS (WORKING TITLE)

25 COPYRIGHT
26 2019
27 William Kennedy DiClemente

28 All rights reserved.

Acknowledgements

³⁰ I'd like to thanks the Ghosts of Penn Students Past for providing me with such an amazing thesis
³¹ template.

32

ABSTRACT

33 MEASUREMENT OF ELECTROWEAK PRODUCTION OF SAME-SIGN W BOSON PAIRS
34 WITH ATLAS (WORKING TITLE)

35

William Kennedy DiClemente

36

J. Kroll

37

This is the abstract text.

Contents

39	Acknowledgements	iii
40	Abstract	iv
41	Contents	v
42	List of Tables	ix
43	List of Figures	xii
44	Preface	xviii
45	Preface	xviii
46	1 Introduction	1
47	2 Theoretical Framework	2
48	2.1 Introduction to the Standard Model	2
49	2.2 Electroweak symmetry breaking and the Higgs boson	4
50	2.3 Diboson physics	7
51	3 LHC and the ATLAS Detector	9
52	3.1 The Large Hadron Collider	9
53	3.2 The ATLAS Detector	9
54	3.2.1 The Inner Detector	9
55	3.2.1.1 Pixel Detector	9

56	3.2.1.2	Semiconductor Tracker	9
57	3.2.1.3	Transition Radiation Tracker	9
58	3.2.2	The Calorimeters	10
59	3.2.2.1	Liquid Argon Calorimeters	10
60	3.2.2.2	Tile Calorimeters	10
61	3.2.3	The Muon Spectrometer	10
62	3.2.4	Particle reconstruction	11
63	3.2.4.1	Track reconstruction	11
64	3.2.4.2	Muon reconstruction	11
65	3.2.4.3	Electron reconstruction	11
66	3.2.4.4	Jet reconstruction	11
67	4 Alignment of the ATLAS Inner Detector		12
68	4.1	The alignment method	13
69	4.1.1	Alignment levels	16
70	4.1.2	Alignment coordinate systems	17
71	4.2	Early 2015 alignment of the ATLAS detector	19
72	4.2.1	June alignment procedure	20
73	4.2.2	Alignment results	21
74	4.2.2.1	Residual distributions from collisions	22
75	4.2.2.2	Track parameter resolution from cosmic rays	24
76	4.2.3	Error scaling	25
77	4.3	Level 2 alignment of the TRT	28
78	4.4	Momentum bias from sagitta deformations	29
79	4.4.1	Sagitta bias monitoring with electron E/p	31
80	4.4.1.1	Measuring $\langle E/p \rangle$	32
81	4.4.1.2	Results in 13 TeV data	34
82	5 Measurement of same-sign WW production at $\sqrt{s} = 13$ TeV with ATLAS		37
83	5.0.2	Theoretical overview of vector boson scattering	37
84	5.0.3	Same-sign $W^\pm W^\pm$ scattering	39
85	5.0.4	Overview of backgrounds	41
86	5.1	Data and Monte Carlo samples	43

87	5.1.1	Monte Carlo samples	43
88	5.2	Object and event selection	44
89	5.2.1	Object selection	44
90	5.2.1.1	Muon candidate selection	45
91	5.2.1.2	Electron candidate selection	47
92	5.2.1.3	Jet candidate selection	47
93	5.2.1.4	Treatment of overlapping objects	47
94	5.2.2	Signal event selection	49
95	5.3	Background estimations	51
96	5.3.1	Estimation of the WZ background	51
97	5.3.2	Estimation of the $V\gamma$ background	52
98	5.3.3	Estimation of backgrounds from charge misidentification	54
99	5.3.3.1	Validation of the charge misidentification estimate	57
100	5.3.4	Estimation of non-prompt backgrounds with the fake factor method	57
101	5.3.4.1	Overview of the default fake factor method	58
102	5.3.4.2	The fake factor with p_T^{cone}	59
103	5.3.4.3	Application of the fake factor	61
104	5.3.4.4	Systematic uncertainties	65
105	5.3.4.5	Results of the fake factor	66
106	5.3.4.6	Validation of the fake factor	69
107	5.3.5	Reduction of WZ background using custom overlap removal	70
108	5.4	Cross section measurement	74
109	5.4.1	Maximum likelihood fit	74
110	5.4.2	Definition of the fiducial volume	76
111	5.4.3	Cross section extraction	78
112	5.5	Summary of uncertainties	79
113	5.5.1	Experimental uncertainties	79
114	5.5.2	Theoretical uncertainties	81
115	5.5.2.1	Uncertainties from EWK-QCD interference	81
116	5.6	Results	82
117	6	Prospects for same-sign WW at the High Luminosity LHC	86
118	6.0.1	Analysis Overview	87

119	6.1 Theoretical motivation	87
120	6.1.1 Experimental sensitivity to longitudinal polarization	88
121	6.2 Monte Carlo samples	88
122	6.3 Background estimations	90
123	6.3.1 Truth-based isolation	91
124	6.4 Object and event selection	92
125	6.4.1 Object selection	92
126	6.4.2 Event selection	92
127	6.5 Selection optimization	93
128	6.5.1 Random grid search algorithm	93
129	6.5.2 Inputs to the optimization	95
130	6.5.3 Results of the optimization	96
131	6.6 Results	97
132	6.6.1 Event yields	97
133	6.6.2 Uncertainties	103
134	6.6.3 Cross section measurement	103
135	6.6.4 Longitudinal scattering significance	104
136	7 Conclusion	108
137	A Additional material on truth isolation	109
138	B Additional material on $W^\pm W^\pm jj$ measurement at $\sqrt{s} = 13\text{TeV}$	110
139	B.1 Impact of experimental uncertainty on MC background estimations	110
140	Bibliography	112

List of Tables

142 4.1	The four alignment levels for each of the detector subsystems. The total number of alignable structures and degrees of freedom (DoF) to be aligned are given for each level.	18
144 4.2	Summary of the highest level of alignment applied to each ID subsystem when deriving the June alignment.	21
146 4.3	Estimated value of the error scaling term b for each subdetector component with the June alignment.	28
148 5.1	Predicted cross sections for EQK and QCD production of diboson processes relevant to VBS at $\sqrt{s} = 8\text{TeV}$ using the SHERPA MC generator. Loose generator level cuts are applied on lepton $p_T > 5\text{ GeV}$, dilepton invariant mass $m_{ll} > 4\text{ GeV}$, and at least two jets with $m_{jj} > 10\text{ GeV}$	40
152 5.2	Summary of MC samples used in the analysis.	45
153 5.3	Muon selection criteria. All muons are required to pass the preselection (top), and then either the signal (middle) or loose (bottom) criteria is applied to the preselected electrons.	46
155 5.4	Electron selection criteria. All electrons are required to pass the preselection (top), and then either the signal (middle) or loose (bottom) criteria is applied to the preselected electrons.	48
158 5.5	Jet selection criteria. All jets are required to pass the above selection in order to be used in the analysis.	48
160 5.6	Summary of the overlap removal procedure used in the analysis. If the criteria in the “check” column is met, in the “result” column, the object on the left of the arrow is removed in favor of the object on the right.	49
163 5.7	Summary of trigger requirements for electrons and muons for $\sqrt{s} = 13\text{TeV}$ data collected in 2015 and 2016. At least one of the triggers must be satisfied.	50
165 5.8	The signal event selection.	51
166 5.9	Event yields in the WZ control region before normalization. All lepton flavor channels are combined.	52
168 5.10	Selection criteria for the $V\gamma$ control region.	53
169 5.11	Event yields in the $V\gamma$ control region. The $V\gamma$ scale factor of 1.77 is calculated by scaling up the $Z\gamma$ and $Z+\text{jets}$ backgrounds to account for the difference between the data and predicted total background.	54
172 5.12	Selection criteria for the same-sign inclusive validation region.	57

173	5.13	Event selection for the dijet region used for calculating the fake factor. The selected lepton can pass either the nominal (signal) or loose selections. In the case of the nominal leptons, the $p_T > 27$ GeV requirement is replaced with $p_T > 15$ GeV.	63
174			
175	5.14	Values of the fake factor in each p_T bin and for each individual systematic source.	68
176			
177	5.15	Estimated yields for the fake lepton background. The estimated yield is shown in the first column together with the statistical uncertainty followed by the systematic uncertainties from variations of the the fake factors within their statistical (stat.) and systematic (syst.) uncertainties. The labels f_e and f_μ indicate the fake factors for electrons and muons, respectively.	69
178			
179	5.16	Selection criteria for the same-sign top fakes validation region.	69
180			
181	5.17	Custom OR definition. Leptons must pass this selection in order to be counted for the trilepton veto.	73
182			
183	5.18	Definition of the fiducial volume.	77
184			
185	5.19	Impact of various systematic effects on the fiducial cross section measurement. The impact of a given source of uncertainty is computed by performing the fit with the corresponding nuisance parameter varied up or down by one standard deviation from its nominal value.	79
186			
187	5.20	List of sources of experimental uncertainties on the reconstruction of physics objects.	80
188			
189	5.21	Impact of experimental uncertainties for the $W^\pm W^\pm jj$ EWK processes in all channels.	80
190			
191	5.22	Impact of experimental uncertainties for the WZ process in all channels.	81
192			
193	5.23	The set of generator level cuts used for generating the interference samples with <code>MadGraph</code>	82
194			
195	5.24	Cross sections for each different $W^\pm W^\pm jj$ production mode (inclusive, EWK only, QCD only, and interference only) generated using <code>MadGraph</code> . The cross sections are calculated using a minimal set of generator level cuts from events where the W decays to a muon.	82
196			
197	5.25	Table of the data and prediction event yields in the signal region before the fit. Numbers are shown for the six lepton flavor and charge channels and for all channels combined. Here the WZ background yields are normalized to the data in the WZ control region. The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.	83
198			
199			
200			
201			
202			
203			
204	5.26	Table of the data and prediction event yields in the signal region after the fit. Numbers are shown for the six lepton flavor and charge channels and for all channels combined. The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.	83
205			
206			
207			
208			
209			
210	6.1	Summary of MC samples used in the analysis.	90
211	6.2	Truth-based isolation requirements for electrons and muons.	91
212	6.3	Summary of the signal event selection.	93
213	6.4	Updates to the $W^\pm W^\pm jj$ event selection criteria after optimization. Cuts not listed remain unchanged from the default selection in Table 6.3.	97
214			
215	6.5	Signal and background event yields using the default event selection for an integrated luminosity of $\mathcal{L} = 3000 \text{ fb}^{-1}$. Events containing a fake or charge-flipped electron are removed from their respective sources and combined into a single entry each.	101
216			
217			
218	6.6	Signal and background event yields using the optimized event selection for an integrated luminosity of $\mathcal{L} = 3000 \text{ fb}^{-1}$. Events containing a fake or charge-flipped electron are removed from their respective sources and combined into a single entry each.	101
219			
220			

221	6.7 Summary of estimated experimental and rate uncertainties.	103
222	A.1 Event yields prior to applying any form of truth-based isolation criteria.	109
223	A.2 Event yields after applying a test version of the truth-based isolation.	109
224	B.1 Experimental systematics uncertainties for the $W^\pm W^\pm jj$ QCD processes in all channels.	110
225	B.2 Experimental systematics uncertainties for triboson process in all channels.	110
226	B.3 Experimental systematics uncertainties for $t\bar{t}V$ processes in all channels.	111
227	B.4 Experimental systematics uncertainties for the $W\gamma$ process in all channels.	111
228	B.5 Experimental systematics uncertainties for the $Z\gamma$ process in all channels.	111
229	B.6 Experimental systematics uncertainties for the ZZ process in all channels.	111

List of Figures

231 2.1	An illustration of the potential term $V(\phi)$ in the cases where $\mu^2 > 0$ (left) and $\mu^2 < 0$ (right). The right-hand plot shows the Higgs potential, or ‘‘Mexican hat potential’’, with the minimum at $ \phi = \sqrt{-\frac{\mu^2}{\lambda}}$ rather than at $ \phi = 0$ as in the left-hand plot.	5
234 2.2	Vector boson fusion involving triple gauge couplings (left) and vector boson scattering involving a quartic gauge coupling (right).	7
236 2.3	Diboson diagrams involving the s -channel (left) and t -channel (right) exchanges of a Higgs boson.	8
238 3.1	General cut-away view of the ATLAS detector.	10
239 4.1	Graphical representation of the effect of a misaligned detector element. The reconstructed particle track (dashed arrow) differs from the actual trajectory of the particle (solid arrow) due to the shift in one of the detector elements. The cyan lines represent the track-to-hit residuals.	13
243 4.2	Graphical representation of the ID alignment chain.	16
244 4.3	A schematic representation of the Inner Detector in the longitudinal plane with the global coordinate system overlaid on top. The Pixel detector and IBL are shown in blue, the SCT in green, and the TRT in red. The local coordinates for each subdetector module are inset on the right.	19
248 4.4	The d_0 distributions with respect to the reconstructed primary vertex using the $\sqrt{s} = 13\text{TeV}$ collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.	22
252 4.5	Local x (left) and local y (right) residual distributions of the IBL planar sensors using the $\sqrt{s} = 13\text{TeV}$ collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.	23
256 4.6	The mean of the local x (left) and local y (right) residual distributions as a function of the global z position of each IBL module using the $\sqrt{s} = 13\text{TeV}$ collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red).	24

260	4.7	Local x (top) and local y (bottom) residual distributions for the Pixel barrel (excluding 261 the IBL, left) and end-caps (right) using the $\sqrt{s} = 13\text{TeV}$ collision data sample recon- 262 structed using the June (black) and March (green) alignments. The data is compared to a 263 MC simulation using a perfect detector geometry (red). The distributions are normalized 264 to the same number of entries.	25
265	4.8	Local x residual distributions for the SCT barrel (left) and end-caps (right) using the 266 $\sqrt{s} = 13\text{TeV}$ collision data sample reconstructed using the June (black) and March 267 (green) alignments. The data is compared to a MC simulation using a perfect detector 268 geometry (red). The distributions are normalized to the same number of entries.	26
269	4.9	Residual distributions for the TRT barrel (left) and end-caps (right) using the $\sqrt{s} =$ 270 13TeV collision data sample reconstructed using the June (black) and March (green) 271 alignments. The data is compared to a MC simulation using a perfect detector geometry 272 (red). The distributions are normalized to the same number of entries.	26
273	4.10	Representation of splitting a single cosmic ray track passing through the entire detector 274 (left) into two separate tracks (right).	27
275	4.11	Distribution of the difference in the impact parameter Δd_0 (left) and charge over trans- 276 verse momentum $\Delta q/p_T$ (right) between the two cosmic ray split tracks. The June 277 (black) and March (green) alignments are compared. The distributions are normalized 278 to the same number of entries.	27
279	4.12	Pull distributions in local x (left) and y (right) for the IBL using the $\sqrt{s} = 13\text{TeV}$ collision 280 data sample after applying the error scaling.	28
281	4.13	Residual means by straw layer in two TRT ϕ -sectors affected by a tilt misalignment. The 282 tilts in each of the three modules are clearly visible in the red points representing the 283 reconstructed data prior to alignment. After four iterations of L2 alignment, the residual 284 means in the gray points are flat.	29
285	4.14	Two dimensional map of residuals in the first layer of the TRT barrel vs z and ϕ . Each 286 bin represents the mean of a Gaussian fit to the TRT residuals in that bin. The map on 287 the left is after the L3 (wire-by-wire) alignment of the TRT performed in 2010, and the 288 map on the right is after the L2 alignment at the end of 2015. The z -axis for both plots 289 use the same scale.	30
290	4.15	Representation of a curl distortion in the detector. The image shows a cutaway in the 291 transverse plane. The deformation is represented by the red arrows, and the impact on 292 the reconstructed positive (blue) and negative (green) tracks are shown. The dashed lines 293 represent the true particle trajectories, and the solid lines represent the reconstructed 294 trajectories.	31
295	4.16	Geometric definition of the sagitta s in relation to the length of the chord l and the radius 296 r of a circular arc.	31
297	4.17	E/p distributions of electrons and positrons in two different η - ϕ bins of the detector. The 298 left hand plot is taken from a region with no momentum bias where $\langle E/p \rangle^+ = \langle E/p \rangle^-$, 299 while the right hand plot shows an 8% disagreement in $\langle E/p \rangle$ between electrons and 300 positrons.	33
301	4.18	Sagitta bias in the $\sqrt{s} = 13\text{TeV}$ data collected by ATLAS in 2016 as a function of η and 302 ϕ for the prompt (left) and reprocessed (right) alignments using the E/p method.	35
303	4.19	Sagitta bias in the $\sqrt{s} = 13\text{TeV}$ data collected by ATLAS in 2016 projected along η for 304 the prompt (gray) and reprocessed (blue) alignments using the E/p method.	35
305	4.20	Sagitta bias in the $\sqrt{s} = 13\text{TeV}$ data collected by ATLAS in 2017 as a function of η 306 and ϕ in reconstructed electrons (left) and after two iterations of momentum corrections 307 (right) from the E/p method.	35

308	4.21 Sagitta bias in the $\sqrt{s} = 13\text{TeV}$ data collected by ATLAS in 2017 projected along η in 309 reconstructed electrons (gray) and after one (red) and two (blue) iterations of momentum 310 corrections from the E/p method.	36
311	5.1 Cross sections in nanobarns for five different scattering processes of longitudinally polarized 312 vector bosons as a function of center of mass energy \sqrt{s} . Without a SM Higgs boson 313 (left), the cross sections grow unbounded with \sqrt{s} ; however with a 120 GeV Higgs boson 314 (right), the cross sections no longer diverge.	38
315	5.2 Tree-level Feynman diagrams for VBS EWK $VVjj$ production including triple gauge 316 couplings involving W and/or Z bosons (top left and top middle), quartic gauge coupling 317 (top right), or the exchange of a Higgs boson (s -channel bottom left and t -channel bottom 318 right). The labels are quarks (q), fermions (f), and gauge bosons ($V = W, Z$).	39
319	5.3 Tree-level Feynman diagrams for non-VBS EWK $VVjj$ production. The labels are quarks 320 (q), fermions (f), and gauge bosons ($V = W, Z$).	39
321	5.4 Tree-level Feynman diagrams for QCD $VVjj$ production. The labels are quarks (q), 322 fermions (f), and gauge bosons ($V = W, Z$).	40
323	5.5 Feynman diagrams for VBS EWK production of $W^\pm W^\pm jj$ events. The leftmost diagram 324 contains a quartic gauge coupling vertex, and the rightmost diagram contains an exchange 325 of a Higgs boson. TODO: Make diagrams consistent with others	41
326	5.6 $W^\pm W^\pm jj$ VBS event topology containing two leptons (1 and 2) with the same elec- 327 tric charge, two neutrinos, and two forward tagging jets (3 and 4) with large rapidity 328 separation Δy	41
329	5.7 Generator level comparisons at $\sqrt{s} = 8\text{TeV}$ of dijet invariant mass (m_{jj} , left) and dijet 330 rapidity (Δy_{jj} , right) in EWK (red) and QCD (blue) $W^\pm W^\pm jj$ events. Both data sets 331 have been normalized to the same area.	42
332	5.8 ATLAS event display of a $pp \rightarrow W^+ W^+ \rightarrow \mu^+ \nu_\mu \mu^+ \nu_\mu jj$ event. The muons are rep- 333 resented by the red lines travelling from the ID through the MS, and the forward jets 334 are represented by the blue cones with yellow energy deposits in the calorimeters. The 335 direction of the E_T^{miss} in the transverse plane is indicated by the gray dashed line in the 336 inset image.	42
337	5.9 Leading lepton p_T (left) and η (right) distributions in the WZ control region before 338 normalization. All lepton channels are combined.	53
339	5.10 Trilepton invariant mass m_{lll} distribution in the WZ control region before normalization. 340 All lepton channels are combined.	53
341	5.11 Charge misidentification rates for electrons as a function of $ \eta $ and p_T . Rates are calcu- 342 lated from $Z \rightarrow e^+ e^-$ MC after applying scale factors to approximate the charge mis-ID 343 rates in data.	55
344	5.12 Dilepton invariant mass distribution m_{ll} for the ee channel in the same-sign inclusive VR.	57
345	5.13 p_T distributions for the leading (left) and subleading (right) electron for the ee channel 346 in the same-sign inclusive VR. In these plots, the cut requiring m_{ee} to fall within the Z 347 mass window has been inverted in order to test the modelling away from the Z peak. .	58
348	5.14 Δp_T distributions for nominal (blue) and loose (red) muons in simulated $t\bar{t}$ events. Each 349 muon has been matched to a truth-level jet. Both distributions are normalized to unit 350 area.	60
351	5.15 Δp_T distributions for muons (top) and electrons (bottom) in simulated $t\bar{t}$ events. Each 352 lepton has been matched to a truth-level jet, and that truth jet has had its p_T corrected 353 to include all truth muons within a cone of $\Delta R < 0.4$. The nominal leptons are in black. 354 Δp_T is calculated for the loose leptons using p_T (red) and $p_T + p_T^{\text{cone}}$ (blue).	62
355	5.16 Distributions of p_T^{cone}/p_T for nominal (black) and loose (red) muons in simulated $t\bar{t}$ events.	63

356	5.17	The measured fake factor as a function of muon $p_T + p_T^{\text{cone}}$. The error bars represent the statistical uncertainty only.	64
357			
358	5.18	The measured fake factor as a function of electron $p_T + p_T^{\text{cone}}$ in the central ($ \eta < 1.37$, blue) and forward ($ \eta > 1.37$, red) regions of the detector. The error bars represent the statistical uncertainty only.	64
359			
360			
361	5.19	Graphical representation of the fake factor application using $p_T + p_T^{\text{cone}}$. The value of $p_T + p_T^{\text{cone}}$ for the loose lepton is used to “look up” the fake factor weight which is then applied to the event. The loose lepton’s p_T becomes $p_T + p_T^{\text{cone}}$ for the purpose of the fake background estimation.	65
362			
363			
364			
365	5.20	Systematic variations in the fake factor as a function of muon $p_T + p_T^{\text{cone}}$. The individual fake factors obtained for each systematic variation are displayed with their statistical uncertainties.	66
366			
367			
368	5.21	Systematic variations in the fake factor as a function of electron $p_T + p_T^{\text{cone}}$ in the central ($ \eta < 1.37$, top) and forward ($ \eta > 1.37$, bottom) regions of the detector. The individual fake factors obtained for each systematic variation are displayed with their statistical uncertainties.	67
369			
370			
371			
372	5.22	Distributions of the subleading lepton p_T in the same-sign top fakes VR for $\mu^\pm\mu^\pm$ events (top right), $e^\pm e^\pm$ events (top left), $\mu^\pm e^\pm$ events (bottom left), and all events combined (bottom right). All errors are statistical only.	70
373			
374			
375	5.23	Pseudorapidity (η) distributions of truth muons (top) and electrons (bottom) for Sherpa $W^\pm W^\pm jj$ and WZ MC samples. The blue vertical lines represent the allowed η range for each lepton flavor. The numbers correspond to the number of raw MC events that fall within and outside of the allowed η range for each MC sample.	71
376			
377			
378			
379	5.24	Distributions of $p_{T,\text{ratio}}(\mu, j)$ for EWK and QCD $W^\pm W^\pm jj$ signal (black) and WZ background (teal) for truth-matched third muons in events that pass the trilepton veto. Both distributions are normalized to unit area. The associated efficiency curves are on the right where efficiency is defined as the percentage of total events that would pass a cut on $p_{T,\text{ratio}}(\mu, j)$ at a given value on the x -axis.	72
380			
381			
382			
383			
384	5.25	Distributions of $\Delta R(\mu, j)$ for EWK and QCD $W^\pm W^\pm jj$ signal (black) and WZ background (teal) for truth-matched third muons in events that pass the trilepton veto. Both distributions are normalized to unit area. The associated efficiency curves are on the right where efficiency is defined as the percentage of total events that would pass a cut on $\Delta R(\mu, j)$ at a given value on the x -axis.	72
385			
386			
387			
388			
389	5.26	Distributions of $p_{T,\text{ratio}}(e, j)$ for EWK and QCD $W^\pm W^\pm jj$ signal (black) and WZ background (teal) for truth-matched third electrons in events that pass the trilepton veto. Both distributions are normalized to unit area. The associated efficiency curves are on the right where efficiency is defined as the percentage of total events that would pass a cut on $p_{T,\text{ratio}}(e, j)$ at a given value on the x -axis.	73
390			
391			
392			
393			
394	5.27	Two-dimensional plots of $p_{T,\text{ratio}}(\mu, j)$ vs $\Delta R(\mu, j)$ for truth-matched third muons in events that pass the trilepton veto for EWK and QCD $W^\pm W^\pm jj$ signal (left) and WZ background (right). The blue overlay indicates the area in which the third leptons will pass the custom OR and result in the event failing the trilepton veto.	74
395			
396			
397			
398	5.28	Visual representation of the different kinematic regions relevant to the cross section measurement. The acceptance factor \mathcal{A} converts from the truth level total phase space to the truth level fiducial region, and the efficiency correction \mathcal{C} translates the fiducial region in to the reconstruction level signal region.	77
399			
400			
401			

402	5.29	The dijet invariant mass m_{jj} distributions for data and predicted signal and background in the signal region after the fit. The shaded band represents the statistical and systematic uncertainties added in quadrature. Note that the bins have been scaled such that they represent the number of events per 100 GeV in m_{jj} . The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.	84
410	5.30	The event yields for data and predicted signal and background in the WZ and low- m_{jj} control regions after the fit. The shaded band represents the statistical and systematic uncertainties added in quadrature. The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.	84
416	5.31	Comparison of the measured $W^\pm W^\pm jj$ EWK fiducial cross section with theoretical calculations from SHERPA v2.2.2 and POWHEG-BOX v2 . The light orange band represents the total experimental uncertainty on the measured value, and the dark orange hashed band is the statistical uncertainty. For the simulations, the light blue band represents the total theoretical uncertainty, and the dark blue hashed band are the scale uncertainties. The theory predictions do not include the interference between the EWK and QCD production.	85
423	6.1	Comparison of the leading (top) and subleading (bottom) lepton p_T distributions for purely longitudinal (LL, black) and mixed polarization (LT+TT, cyan) $W^\pm W^\pm jj$ events.	89
425	6.2	Comparison of the azimuthal dijet separation ($ \Delta\phi_{jj} $) for purely longitudinal (LL, black) and mixed polarization (LT+TT, cyan) $W^\pm W^\pm jj$ events.	90
427	6.3	A visual representation of a rectangular grid search algorithm. The signal events are the blue triangles, and the red circles are the background events. TODO: replace with own figure	94
430	6.4	A visual representation of a random grid search algorithm. The signal events are the blue triangles, and the red circles are the background events. TODO: replace with own figure	95
433	6.5	Leading lepton p_T distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram). TODO: Move to appendix or omit	98
437	6.6	Dilepton invariant mass distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram). TODO: Move to appendix or omit	98
441	6.7	Leading (top) and subleading (bottom) jet p_T distributions. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram).	99
445	6.8	Dijet invariant mass distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram). TODO: Move to appendix or omit	100

449	6.9	Lepton-jet centrality distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram).	100
450	6.10	p_T distributions for the leading jet using the default (left) and optimized (right) event selections for all channels combined.	102
451	6.11	p_T distributions for the subleading jet using the default (left) and optimized (right) event selections for all channels combined.	102
452	6.12	p_T distributions for lepton-jet centrality ζ using the default (left) and optimized (right) event selections for all channels combined.	102
453	6.13	Projections of the statistical (black), theoretical (blue), systematic (yellow), and total (red) uncertainties on the measured cross section as a function of integrated luminosity using the optimized event selection.	104
454	6.14	Dijet azimuthal separation ($ \Delta\phi_{jj} $) for the low m_{jj} region ($520 < m_{jj} < 1100$ GeV, top) and the high m_{jj} region ($m_{jj} > 1100$ GeV, bottom). The purely longitudinal (LL, gray) is plotted separately from the mixed and transverse (LT+TT, cyan) polarizations.	106
455	6.15	Projections of the expected longitudinal scattering significance as a function of integrated luminosity when considering all sources of uncertainties (black) or only statistical uncertainties (red).	107
456			
457			
458			
459			
460			
461			
462			
463			
464			
465			
466			

Preface

468 This thesis presents the major highlights of my work with the ATLAS experiment as a graduate
469 student at the University of Pennsylvania from Fall of 2013 until early Spring of 2019.

470 The first step of working on the experiment is to complete a *qualification task* in order to be
471 included on the author list of ATLAS publications. These tasks are an opportunity to contribute
472 to the experiment as a whole, such as maintaining detector hardware or monitoring physics perfor-
473 mance. For my qualification task, I worked with the Inner Detector Alignment group which works
474 to make sure we have accurate knowledge of the locations of each and every sensor in the detector.
475 My qualification task involved investigating a possible momentum bias in the Monte Carlo (MC)
476 simulated data. The MC is supposed to be reconstructed with a perfect detector geometry which
477 should in principle be free of any momentum biases. Ultimately I determined that the size of the
478 biases were small enough to be negligible compared to what is seen in the real data, and that they
479 could be corrected for if necessary.

480 My work with the alignment group would continue for the duration of my time here at Penn. In
481 early 2015, at the start of the LHC’s second data-taking run (Run 2), I worked assisted in validating
482 the first set of alignment constants using $\sqrt{s} = 8$ TeV proton-proton collision data. At this point
483 I took over the responsibility of alignment of the TRT subdetector. The TRT was aligned to high
484 accuracy in Run 1, and over the course of my time working on alignment, the TRT never required
485 a straw-by-straw alignment; however it did require a module-level alignment at the end of 2015.
486 My final responsibility in the alignment group was monitoring momentum biases using the energy-
487 momentum ratio (E/p) of electrons. For the large data reprocessing, the E/p method served as a
488 cross check to a similar method using Z boson events for monitoring and aligning out momentum
489 biases in the detector. The results from both methods were also used in the uncertainties for the
490 tracking measurements.

491 On the analysis side, I had previous experience in Standard Model (SM) electroweak physics
492 from my time as an undergraduate at Duke University, and it remained a point of interest for
493 me in graduate school. As such, I was happy to work with fellow Penn students on the cross
494 section measurement of SM WZ diboson production with the early $\sqrt{s} = 13$ TeV ATLAS data.
495 My contribution to the analysis was primarily on the software side, as I maintained and updated
496 the analysis framework. While the WZ measurement is not covered by this thesis, it provided me
497 with invaluable analysis experience in electroweak physics, as well as a detailed understanding of a
498 major background to many diboson processes. The results for this analysis can be found published
499 in Physics Letters B in 2016 [1].

500 The final two analyses I worked on involve the scattering of same-sign W bosons, and they make
501 up the majority of this thesis. The first analysis is a measurement of the $W^\pm W^\pm jj$ cross section
502 at $\sqrt{s} = 13$ TeV. This measurement along with that of the CMS collaboration represent the first
503 observation of the $W^\pm W^\pm jj$ scattering process. My primary contribution to the analysis is in the
504 estimation of the fake lepton background, where we implemented a brand new version of the fake
505 factor method using particle isolation variables. I also did a preliminary study of the interference
506 between electroweak and strong production of $W^\pm W^\pm jj$ events, assisted in the production of private
507 data samples for use with the analysis framework, and used my familiarity with the WZ process
508 to optimize the rejection of the background. Ultimately the majority of the WZ rejection was not
509 included in the final result; however, it is still covered in the thesis in the hopes that it will be useful
510 for similar analyses in the future. The formal publication for this measurement will likely be coming
511 out within the next few months.

512 The second $W^\pm W^\pm jj$ analysis is a study on the prospects for a measurement of the process at the
513 upgraded High-Luminosity LHC, scheduled to begin operation in 2026. Here my main contribution
514 was an optimization of the event selection using a Random Grid Search algorithm. Through the
515 optimization we expect to take advantage of the higher center of mass energy and greater volume of
516 data and tighten certain selection cuts to increase the strength of the $W^\pm W^\pm jj$ signal. In addition, I
517 once again maintained and updated the analysis framework and produced the group's data samples,
518 and I also developed a truth-based particle isolation criteria in order to reduce contributions from
519 backgrounds involving the top quark. The results of this prospects study will be published as a part
520 of the annual Yellow Report for the High-Luminosity LHC.

521

Will K. DiClemente
Philadelphia, February 2019

522

CHAPTER 1

523

Introduction

524 The Standard Model (SM)¹ has been remarkably successful...

¹Here's a footnote.

CHAPTER 2

Theoretical Framework

527 This chapter outlines the theoretical groundwork for the rest of the thesis. An overview of the Stan-
 528 dard Model of particle physics (SM) is given in Section 2.1, followed by the electroweak symmetry
 529 breaking mechanism involving the Higgs boson in Section 2.2. Finally, Section 2.3 will go into some
 530 detail on the interests of diboson physics specifically.

531 **2.1 Introduction to the Standard Model**

532 The Standard Model of particle physics serves as a mathematical description of the fundamental
 533 particles of the universe and their interactions. It has been developed over the course of the past
 534 century, incorporating both predictions from theory and results from experiments. All in all, the SM
 535 has proven remarkably successful in accurately describing particle interactions seen in experiments.

536 The SM is a quantum field theory (QFT) [2, 3] in which the fundamental particles are represented
 537 as excited states of their corresponding fields. The spin- $\frac{1}{2}$ fermionic fields give rise to the quarks
 538 and leptons comprising ordinary matter, the spin-1 fields correspond to the electroweak bosons and
 539 the gluon which mediate the electroweak and strong forces, respectively, and finally the scalar Higgs
 540 field is responsible for electroweak symmetry breaking. The excitations and interactions of the fields
 541 are governed by the SM Lagrangian, which is invariant under local transformations of the group
 542 $SU(3) \times SU(2) \times U(1)$. **TODO: need more detail/refinement here**

543 The first quantum field theory to be developed is quantum electrodynamics (QED) [4], which
 544 describes the electromagnetic interaction. The theory predicts the existence of a $U(1)$ gauge field
 545 that interacts with the electrically charged fermions. This field corresponds to the photon. A key
 546 aspect of QED is that it is perturbative. The coupling constant $\alpha = e^2/4\pi$ is small, where e is

547 electrical charge of the field, allowing for the use of perturbation theory in calculations. In this case,
 548 calculations can be written as a power series in α , where successive higher order terms contribute
 549 less to the final result. **TODO: renormalizability here?**

550 The strong interaction—the theory of quarks and gluons—has also been described using QFT
 551 as quantum chromodynamics (QCD). The symmetry group for QCD is $SU(3)$, and its eight gen-
 552 erators correspond to the eight differently charged, massless gluons [5]. Unlike in QED, which has
 553 positive and negative charges, the strong force has three “colors”. Color charge combined with the
 554 non-Abelian nature of $SU(3)$, which allows the gluons to interact with each other, result in the
 555 most well-known property of QCD: color confinement. In order to increase the separation between
 556 two color-charged quarks, the amount of energy required increases until it becomes energetically
 557 favorable to pair-produce a new quark-antiquark pair, which then bind to the original quarks. The
 558 end result of this is that only color-neutral objects exist in isolation. What this means for the strong
 559 coupling constant α_s is that its value at the low energies where confinement occurs is large, on
 560 the order of $\alpha_s \sim 1$. The consequence of this is that perturbation theory cannot be used to accu-
 561 rately approximate interactions. While this appears at first to be a critical problem for predictions,
 562 fortunately it turns out that α_s “runs”, or decreases in magnitude at higher energy [6, 7]. This so-
 563 called “asymptotic freedom” allows QCD to be calculated perturbatively [8] at energies accessible
 564 by collider experiments including the LHC.

565 The last gauge field corresponds to the weak interaction. Ultimately, the weak $SU(2)$ and the
 566 electromagnetic $U(1)$ mix to form the $SU(2) \times U(1)$ *electroweak* (EWK) interaction [9, 10]. A
 567 more detailed description of the mixing will be discussed in conjunction with electroweak symmetry
 568 breaking (EWSB) in Section 2.2; however, a summary of the resulting EWK interaction is presented
 569 here, at the risk of some repeated information to follow. There are three weak isospin bosons arising
 570 from the $SU(2)$ group (W_μ^1 , W_μ^2 , and W_μ^3) and one weak hypercharge boson from the $U(1)$ group
 571 (B_μ). The W_3 and B bosons mix according to the weak mixing angle θ_W to form the Z boson and
 572 the photon according to:

$$\begin{pmatrix} \gamma \\ Z \end{pmatrix} = \begin{pmatrix} \cos \theta_W & \sin \theta_W \\ -\sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^3 \end{pmatrix} \quad (2.1)$$

573 The value of θ_W is not predicted by the SM; it is one example of an experimental input to the
 574 theory, measured to be $\sin^2 \theta_W = 0.23153 \pm 0.00016$ [11]. The charged W^\pm bosons are a mixture of
 575 the remaining W_μ^1 and W_μ^2 bosons:

$$W^\pm = \frac{1}{\sqrt{2}}(W_\mu^1 \mp iW_\mu^2) \quad (2.2)$$

576 Unlike the photon (and the gluon of QCD), the W^\pm and Z bosons are massive. This means that
 577 even though SU(2) is non-Abelian, the range of interaction is short and confinement does not occur.
 578 Lastly, the EWK interaction is chiral, only coupling to the left-handed component of the fermion
 579 fields

580 One final field remains within the SM: the scalar Higgs field. It was originally proposed in the
 581 1960's to explain the masses of the W^\pm and Z bosons [12, 13, 14] and is the mechanism for the
 582 EWSB process. The particle associated with the field is a massive scalar boson, which was at last
 583 discovered by ATLAS and CMS in 2012 [15, 16] with a mass of 125 GeV.

584 2.2 Electroweak symmetry breaking and the Higgs boson

585 The results of electroweak mixing and the implications of the Higgs field have been introduced
 586 in the previous section. If the EWK theory were an unbroken symmetry, the associated W^\pm and
 587 Z bosons would be massless; however, when observed experimentally, they were found to be quite
 588 heavy [17, 18], at around 80 GeV and 91 GeV, respectively [19]. Here, a more detailed explanation
 589 of the Higgs mechanism and how it "spontaneously breaks" the EWK symmetry, resulting in the
 590 three massive bosons (W^\pm and Z) and one massless boson (photon), is presented.

591 To see how the Higgs mechanism results in the massive vector bosons and a massless photon,
 592 consider the following. Beginning with a complex scalar doublet ϕ defined as:

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \sqrt{\frac{1}{2}} \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix} \quad (2.3)$$

593 a Lagrangian \mathcal{L} can be written:

$$\mathcal{L} = (\mathcal{D}_\mu \phi)^\dagger (\mathcal{D}^\mu \phi) - \mu^2 \phi^\dagger \phi - \lambda(\phi^\dagger \phi)^2 \quad (2.4)$$

594 where $\lambda > 0$ and \mathcal{D}_μ is the covariant derivative. \mathcal{D}_μ is defined such that \mathcal{L} is invariant under a local
 595 SU(2) \times U(1) gauge transformation:

$$\mathcal{D}_\mu \phi = \left(\partial_\mu + \frac{ig}{2} \tau_a W_\mu^a + \frac{ig'}{2} B_\mu \right) \phi \quad (2.5)$$

596 where W_μ^a ($a = 1, 2, 3$) are the SU(2) fields with generators τ_a and coupling constant g , and B_μ is
 597 the U(1) field with coupling constant g' .

598 Isolating the potential term:

$$V(\phi) = \mu^2 \phi^\dagger \phi + \lambda(\phi^\dagger \phi)^2 \quad (2.6)$$

599 a choice must be made on the sign of μ^2 , and the case of interest is for $\mu^2 < 0$. This results in
600 the famous “mexican hat potential” shown in Figure 2.1, which is minimized along the collection of
601 points:

$$\phi^\dagger \phi = \frac{1}{2} (\phi_1^2 + \phi_2^2 + \phi_3^2 + \phi_4^2) = -\frac{\mu^2}{2\lambda} \quad (2.7)$$

602 This means that the minimum of the potential is not at $\phi = 0$ (as it would be in the case where
603 $\mu^2 > 0$), but rather at a value:

$$v \equiv \sqrt{-\frac{\mu^2}{\lambda}} \quad (2.8)$$

604 With no loss of generality due to the SU(2) symmetry, $\phi_1 = \phi_2 = \phi_4 = 0$ can be set in Equation 2.7
605 leaving $\phi_3^2 = v^2$. Finally, the *vacuum expectation value* (VEV) of the field can be written as:

$$\langle \phi \rangle = \sqrt{\frac{1}{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} \quad (2.9)$$




Figure 2.1: An illustration of the potential term $V(\phi)$ in the cases where $\mu^2 > 0$ (left) and $\mu^2 < 0$ (right). The right-hand plot shows the Higgs potential, or “Mexican hat potential”, with the minimum at $|\phi| = \sqrt{-\frac{\mu^2}{\lambda}}$ rather than at $|\phi| = 0$ as in the left-hand plot.

606 The VEV can be substituted back into the original Lagrangian in Equation 2.4, and, following
607 quite a bit of math, a collection of mass terms can be identified:

$$\mathcal{L} \subset \mathcal{L}_M \equiv \frac{1}{8} v^2 g^2 \left[(W_\mu^1)^2 + (W_\mu^2)^2 \right] + \frac{1}{8} v^2 \left[g^2 (W_\mu^3)^2 - 2gg' W_\mu^3 B^\mu + g'^2 (B_\mu)^2 \right] \quad (2.10)$$

608 Focusing on the first term for the moment, substituting in Equation 2.2 for the physical W^\pm bosons,
 609 the mass term can be seen clearly:

$$M_W^2 W^+ W^- = \left(\frac{1}{2}vg\right)^2 W^+ W^- \quad (2.11)$$

610

$$M_W = \frac{1}{2}vg \quad (2.12)$$

611 With a bit of clever forward-thinking, the second term of Equation 2.10 can be rewritten as:

$$\frac{1}{8}v^2 \left[gW_\mu^3 - g'B_\mu \right]^2 + 0 \left[g'W_\mu^3 - gB_\mu \right]^2 = \frac{1}{2}M_Z^2 Z_\mu^2 + \frac{1}{2}M_A^2 A_\mu^2 \quad (2.13)$$

612 where Z_μ^2 and A_μ^2 represent the physical Z boson and photon, respectively, and are defined as:

$$Z_\mu = \frac{gW_\mu^3 - g'B_\mu}{\sqrt{g^2 + g'^2}} \quad (2.14)$$

613

$$A_\mu = \frac{g'W_\mu^3 - gB_\mu}{\sqrt{g^2 + g'^2}} \quad (2.15)$$

614 From this, it can be seen that the photon is massless, and the mass of the Z boson is identified as:

$$M_Z = \frac{1}{2}v\sqrt{g^2 + g'^2} \quad (2.16)$$

615 Lastly, the Higgs field can couple directly to the fermions. Taking the electron as an example,
 616 an additional Lagrangian term can be written:

$$\mathcal{L}_e = -G_e [\bar{e}_L \phi e_R + \bar{e}_R \phi^\dagger e_L] \quad (2.17)$$

617 where e_L and e_R are the left-handed doublet and right-handed singlet, respectively, and ϕ is as in
 618 Equation 2.3. The symmetry can be spontaneously broken by a perturbation about the VEV:

$$\phi = \sqrt{\frac{1}{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix} \quad (2.18)$$

619 which, when substituted into \mathcal{L}_e gives:

$$\begin{aligned} \mathcal{L}_e &= -\frac{G_e}{\sqrt{2}}v(\bar{e}_L e_R + \bar{e}_R e_L) - \frac{G_e}{\sqrt{2}}(\bar{e}_L e_R + \bar{e}_R e_L)h \\ &= -m_e \bar{e}e - \frac{m_e}{v} \bar{e}eh \end{aligned} \quad (2.19)$$

620 for electron mass $m_e = \frac{G_e v}{\sqrt{2}}$. From the second term, it can be seen that the strength of the Higgs
 621 coupling to the electron is proportional to the mass of the electron. The rest of the fermion couplings
 622 follow from this example.

623 What is accomplished here is quite remarkable. The weak and electromagnetic interactions have
 624 been unified into a single $SU(2) \times U(1)$ interaction, and the physical bosons observed in nature arise
 625 as mixtures of the four gauge fields. Additionally, the non-zero VEV of the Higgs field results in
 626 masses for the W^\pm and Z bosons while the photon remains massless. Additionally, it is shown that
 627 the Higgs couples to fermions in proportion to their mass. From experimental measurements, the
 628 value of the VEV has been determined to be $v \approx 246$ GeV [19]. However, it should be noted that the
 629 theory does not predict the mass of the Higgs boson or of the fermions; these must all be determined
 630 from experiment.

631 2.3 Diboson physics

632 Processes involving two final state gauge bosons are of particular interest for testing the predictive
 633 power of the SM. Due to the non-Abelian nature of the EWK interaction, the corresponding gauge
 634 bosons are allowed to self-interact. This results in triple and quartic couplings of gauge bosons
 635 (TGCs and QGCs, respectively). The SM allowed TGCs are the $WW\gamma$ and WWZ vertices, and
 636 the QGCs predicted by the model include $WWZ\gamma$, $WW\gamma\gamma$, $WWZZ$, and $WWWW$. These ver-
 637 tices are accessible via a number of production modes at hadron colliders, including vector boson
 638 fusion and scattering (VBF and VBS, as in Figure 2.2). The LHC in particular has provided the
 639 first opportunity to observe some of these vertices, such as the $WW\gamma\gamma$ QGC via the $W\gamma\gamma$ final
 640 state [20], or the $WWWW$ QGC via same-sign $W^\pm W^\pm$ production [21]. Precise measurements of
 641 these processes can be compared to the theoretical predictions, and deviations can point to deficien-
 642 cies in the prediction, such as needing an additional order in α_s in the calculation, or even hints at
 643 new physics, like anomalous gauge couplings² [22].




Figure 2.2: Vector boson fusion involving triple gauge couplings (left) and vector boson scattering involving a quartic gauge coupling (right).

644 Diboson interactions also make up one of the most sensitive tests of EWSB. Aside from the

²In the SM, the TGCs and QGCs are fixed by the electromagnetic coupling constant α and the electroweak mixing angle θ_W . In many Beyond the Standard Model scenarios, these couplings are modified by additional contributions.

645 top quark, the EWK bosons have the strongest coupling to the Higgs mechanism, as there are
646 several production diagrams involving the exchange of a Higgs boson, including those in Figure 2.3.
647 In this instance, VBS processes are particularly important as the Higgs unitarizes the scattering
648 amplitude [23] (this particular aspect will be explored in more detail in the context of same-sign
649 $W^\pm W^\pm$ scattering in Section 5.0.2). Since the Higgs boson has only recently been discovered, it has
650 not yet been extensively tested, and there could still be deviations in the EWSB mechanism that
651 may manifest in the VBF and VBS measurements [24].




Figure 2.3: Diboson diagrams involving the s -channel (left) and t -channel (right) exchanges of a Higgs boson.

652

CHAPTER 3

653

LHC and the ATLAS Detector

654 **3.1 The Large Hadron Collider**

655 The Large Hadron Collider (LHC) [25] is...

656 **3.2 The ATLAS Detector**

657 ATLAS is a general-purpose particle detector...

658 **3.2.1 The Inner Detector**

659 The Inner Detector serves the primary purpose of measuring the trajectories of charged particles...

660 **3.2.1.1 Pixel Detector**

661 The Pixel detector consists of four cylindrical barrel layers and three disk-shaped endcap layers...

662 **3.2.1.2 Semiconductor Tracker**

663 The Semiconductor Tracker uses the same basic technology as the Pixels, but the fundamental unit
664 of silicon is a larger “strip”...

665 **3.2.1.3 Transition Radiation Tracker**

666 The Transition Radiation Tracker is the outermost component of the ID...




Figure 3.1: General cut-away view of the ATLAS detector [26].

667 3.2.2 The Calorimeters

668 ATLAS includes two types of calorimeter system for measuring electromagnetic and hadronic showers.
 669 These are the Liquid Argon (LAr) calorimeters and the Tile calorimeters. Together, these cover
 670 the region with $|\eta| < 4.9\dots$

671 3.2.2.1 Liquid Argon Calorimeters

672 The Liquid Argon system consists of...

673 3.2.2.2 Tile Calorimeters

674 The Tile calorimeter provides coverage for hadronic showers...

675 3.2.3 The Muon Spectrometer

676 Muon spectrometer stuff.

677 **3.2.4 Particle reconstruction**

678 Particle reconstruction algorithms

679 **3.2.4.1 Track reconstruction**

680 **3.2.4.2 Muon reconstruction**

681 **3.2.4.3 Electron reconstruction**

682 **3.2.4.4 Jet reconstruction**

CHAPTER 4

684 Alignment of the ATLAS Inner Detector

685 When a charged particle passes through the ATLAS ID, it leaves hits in the sensors along its path.
686 In order to accurately measure the track of the particle, it is necessary to know where these hits
687 occurred as precisely as possible, which in turn requires knowledge of the physical location of the
688 element that registered the hit. If one of these elements is *misaligned*, or displaced relative to its
689 position in the known detector geometry, the assumed location of the corresponding hit will not
690 match its actual location, resulting in an incorrect track fit. These misalignments can occur for
691 any number of reasons, including but not limited to elements shifting during maintenance periods
692 or cycles in ATLAS's magnetic field, or small movements during normal detector operations. The
693 effect of a misaligned detector element on the track reconstruction is shown in Figure 4.1.

694 In order to correct the misalignments, the ID alignment procedure is applied to accurately
695 determine the physical position and orientation of each detector element. The baseline accuracy of
696 the alignment is required to be such that the track parameter resolutions are not degraded by more
697 than 20% with respect to those derived from a perfect detector geometry³. This corresponds to a
698 precision of better than $10\mu\text{m}$ in the positioning of the elements of the silicon detectors [27].

699 This chapter outlines the ID alignment procedure, the alignment of the detector during the 2015
700 data taking period, and the steps taken to measure momentum biases in the alignment.

³The so-called *perfect geometry* refers to the description of the ATLAS detector in which every sensor precisely matches its design specifications. The perfect geometry contains no misalignments, and the position of each sensor is known exactly.




Figure 4.1: Graphical representation of the effect of a misaligned detector element. The reconstructed particle track (dashed arrow) differs from the actual trajectory of the particle (solid arrow) due to the shift in one of the detector elements. The cyan lines represent the track-to-hit residuals.

701 4.1 The alignment method

702 The alignment procedure uses track-based algorithm that updates the locations of detector elements
 703 in order to minimize the set of track-hit *residuals*. These residuals are defined as the distance between
 704 the fitted track position in a given detector element to the position of the hit recorded by the same
 705 element, and are shown by the cyan lines in Figure 4.1. Tracks in ATLAS are parameterized as
 706 five-dimensional vectors [28]:

$$\vec{\tau} = (d_0, z_0, \phi_0, \theta, q/p) \quad (4.1)$$

707 where d_0 and z_0 are the transverse and longitudinal impact parameters with respect to the origin,
 708 respectively, ϕ_0 is the azimuthal angle of the track at the point of closest approach to the origin, θ
 709 is the polar angle, and q/p is the charge of the track divided by its momentum. The residual for the
 710 i^{th} hit of a given track can then be written in terms of the track parameters $\vec{\tau}$ and a set of alignment
 711 parameters \vec{a} that describe the hit location [29]:

$$r_i(\vec{\tau}, \vec{a}) = (\vec{m}_i - \vec{e}_i(\vec{\tau}, \vec{a})) \cdot \hat{k} \quad (4.2)$$

712 where \vec{e}_i is the intersection point of the extrapolated track with the sensor, \vec{m}_i is the position of the
 713 associated hit within the sensor, and \hat{k} is the unit vector defining the direction of the measurement
 714 within the sensor. \vec{r} is then the vector of residuals for the given track.

715 A χ^2 function can be built from the residuals of all collected tracks:

$$\chi^2 = \sum_{\text{tracks}} \vec{r}^T V^{-1} \vec{r} \quad (4.3)$$

716 where V is the covariance matrix of the hit measurements. The χ^2 function is then minimized with
 717 respect to the alignment parameters \vec{a} , which contain all degrees of freedom being aligned. The
 718 minimization condition with respect to \vec{a} is:

$$\frac{d\chi^2}{d\vec{a}} = 0 \rightarrow 2 \sum_{\text{tracks}} \left(\frac{d\vec{r}}{d\vec{a}} \right)^T V^{-1} \vec{r} = 0 \quad (4.4)$$

719 This equation can be difficult to solve exactly, so the residual is rewritten as a first order Taylor
 720 expansion:

$$\vec{r} = \vec{r}_0 + \frac{d\vec{r}}{d\vec{a}} \delta\vec{a} \quad (4.5)$$

721 where \vec{r}_0 is dependent on an initial set of track and alignment parameters \vec{r}_0 and \vec{a}_0 , respectively;
 722 the track parameter dependence has also been folded into the total derivative $\frac{d\vec{r}}{d\vec{a}}$. Equation 4.5 can
 723 then be inserted into the minimization condition from Equation 4.4 to give:

$$\left[\sum_{\text{tracks}} \left(\frac{d\vec{r}}{d\vec{a}} \right)^T V^{-1} \left(\frac{d\vec{r}}{d\vec{a}} \right) \right] \delta\vec{a} + \sum_{\text{tracks}} \left(\frac{d\vec{r}}{d\vec{a}} \right)^T V^{-1} \vec{r}_0 = 0 \quad (4.6)$$

724 From this equation, the alignment matrix \mathcal{M}_a and alignment vector $\vec{\nu}_a$ can be defined:

$$\mathcal{M}_a = \sum_{\text{tracks}} \left(\frac{d\vec{r}}{d\vec{a}} \right)^T V^{-1} \left(\frac{d\vec{r}}{d\vec{a}} \right) \quad (4.7)$$

725

$$\vec{\nu}_a = \sum_{\text{tracks}} \left(\frac{d\vec{r}}{d\vec{a}} \right)^T V^{-1} \vec{r}_0 \quad (4.8)$$

726 Finally, the alignment corrections $\delta\vec{a}$ can be solved for by inverting the alignment matrix:

$$\delta\vec{a} = -\mathcal{M}_a^{-1} \vec{\nu}_a \quad (4.9)$$

727 which is a linear system of equations with a number of equations equal to the number of alignment
 728 degrees of freedom [30].

729 Inverting the matrix and solving this system of equations is referred to as *Global χ^2* align-
 730 ment [29]. This can be useful, as \mathcal{M}_a contains all the correlations between the alignable structures.
 731 However, inverting the matrix becomes difficult when the number of degrees of freedom becomes
 732 large, and as the number of alignable structures increases, so too does the size of the matrix \mathcal{M}_a .
 733 Eventually inverting the matrix becomes too computationally intensive to be practical.

734 This problem is solved by the *Local* χ^2 algorithm [31]. In this case, the alignment matrix is
 735 constructed to be block-diagonal, allowing for it to be inverted even for large numbers of degrees of
 736 freedom. This is achieved by replacing the full derivative in Equation 4.6 with the partial derivative
 737 $\frac{\partial \vec{r}}{\partial \vec{a}}$. The new alignment matrix \mathcal{M}'_a and alignment vector $\vec{\nu}'_a$ become:

$$\mathcal{M}_a = \sum_{\text{tracks}} \left(\frac{\partial \vec{r}}{\partial \vec{a}} \right)^T V^{-1} \left(\frac{\partial \vec{r}}{\partial \vec{a}} \right) \quad (4.10)$$

738

$$\vec{\nu}_a = \sum_{\text{tracks}} \left(\frac{\partial \vec{r}}{\partial \vec{a}} \right)^T V^{-1} \vec{r}_0 \quad (4.11)$$

739 Inverting \mathcal{M}'_a is considerably faster and less intensive even for large numbers of degrees of freedom;
 740 however, the correlations between the alignable structures is lost.

741 Due to the Taylor expansion used in Equation 4.6, several iterations of the alignment algorithm
 742 may be necessary to converge on a final set of alignment constants. The Local χ^2 alignment typically
 743 requires more iterations due to the loss of the correlation information [32]. In practice, the ATLAS
 744 reconstruction is run over a set of events, and the resulting tracks are fed to the alignment algorithm.
 745 The residuals are calculated, the alignment matrix is built and inverted, and a new set of alignment
 746 constants is obtained. The convergence of the alignment can be checked by:

747 1. Measure the $\Delta\chi^2$ with the previous iteration. If it is near zero, then the χ^2 is approaching its
 748 minimum.

749 2. Looking at the residual distributions for different alignable structures. A well aligned detector
 750 will have a mean residual of zero with a width approximating the intrinsic resolution of the
 751 detector.

752 If the above checks are satisfied, the process is finished and the final alignment constants are read
 753 out; if not, another iteration is performed. A visual representation of the alignment chain is shown
 754 in Figure 4.2.

755 Since a χ^2 minimization is used to align the detector, if there is a systematic misalignments in
 756 the detector that does not adversely affect the χ^2 , the algorithm will be insensitive to it. These
 757 misalignments are referred to as *weak modes*, and special care is taken to remove them [33]. One
 758 potential impact of weak modes is a bias in the track momentum of reconstructed particles. This
 759 particular effect is the subject of Section 4.4.

760 In practice, the detector is aligned both in “real-time” as data is collected, and during dedicated
 761 offline alignment campaigns. The real-time alignment is run in ATLAS’s so-called *calibration loop*,




Figure 4.2: Graphical representation of the ID alignment chain.

which comprises the first stage in the preparation of data for physics analysis. The calibration loop requires the alignment as well as various other detector calibrations to be available within 48 hours for initial data processing. A fast, coarse-grained alignment⁴ is run on a subset of the available data containing full tracking information, and the results are propagated to the reconstruction of that particular run [34]. Due to the time constraints of the calibration loop, a full sensor-by-sensor alignment is not possible.

The more thorough and finely tuned alignments are reserved for the dedicated alignment campaigns. These generally occur early in data taking campaigns, typically once a sufficient amount of data is collected after a detector shutdown, in order to obtain a good baseline alignment for use in the remainder of the data collection period. Once data taking is complete, another campaign determines an improved set of alignment constants (divided into several “blocks” to account for time-dependent misalignments), and the full data is reprocessed using the newly derived detector geometry. The initial offline alignment of the ATLAS detector at the beginning of Run 2 in 2015 is the subject of Section 4.2.

4.1.1 Alignment levels

The alignment of the detector is performed at several levels of increasing granularity. This adds flexibility in being able to align only as finely as needed, and it also allows for global, detector-level

⁴The calibration loop runs up to a Level 2 alignment in the silicon detectors, which involves treating each layer of sensors as a single object, defined in greater detail in Table 4.1.

779 misalignments to be corrected first before dealing with finer adjustments.

- 780 • Level 1 (L1) alignment involves moving entire subdetector components as a single unit, such
781 as the entire Pixel detector, or the SCT barrel. These often have the largest misalignments,
782 but they are easily corrected and do not require large volumes of data to do so.
- 783 • Level 2 (L2) alignment treats individual layers in the silicon detectors (modules in the TRT)
784 and end cap disks as individual alignable objects.
- 785 • Level 2.7 (L27) alignment was introduced with the addition of the IBL to the ID in Run 2. It
786 involves the stave-by-stave alignment of the IBL and Pixel barrel⁵.
- 787 • Level 3 (L3) alignment treats each sensor in the silicon detectors and each straw in the TRT
788 as an individual alignable object. It is the finest grained alignment available but also the most
789 computationally intensive due to the large number of degrees of freedom. The large number
790 of individual detector sensors being aligned also requires the largest amount of statistics.

791 The different alignment levels are listed in more detail in Table 4.1, including the number of alignable
792 structures and associated degrees of freedom for each detector component.

793 The implementation of the alignment algorithm in the software is flexible enough to allow each
794 subsystem to be aligned individually at a specified level. Each alignable structure has six degrees of
795 freedom: 3 translations (T_x, T_y, T_z) and 3 rotations (R_x, R_y, R_z)⁶; however individual degrees of
796 freedom may be turned on and off as required. In a typical alignment job, L1 and L2 contain few
797 enough degrees of freedom that the Global χ^2 algorithm can be used, but L3 alignments (which can
798 contain over 36,000 degrees of freedom in the silicon detectors alone) require the Local χ^2 algorithm.

799 4.1.2 Alignment coordinate systems

800 The global coordinate system (x, y, z) used by the ID alignment matches that of the ATLAS
801 detector in general, as detailed in Section 3.2 TODO: update with actual location of figure when
802 it's in.... The positions and orientations of individual detector modules of the ID are defined by
803 a right-handed local coordinate system (x', y', z') where the origin is defined as the geometrical
804 center of the module. The x' -axis for each silicon module is defined to point along the most sensitive

⁵For the purposes of this Chapter, the term “Pixel” will refer to the original three layers of the Pixel detector, and the IBL will be referenced separately.

⁶The TRT is an exception, as the subdetector does not have any resolution along the length of the straw. Therefore, for the barrel, T_z is omitted. Similarly for the straws themselves, only two parameters are defined: translation with respect to the radial direction (T_ϕ) and rotation with respect to the radial axis (R_r for the barrel and R_z for the end-caps) [35].

Level	Description of alignable structure	Structures	DoF
1	IBL detector	1	6
	Whole Pixel detector	1	6
	SCT barrel and 2 end-caps	3	18
	TRT barrel and 2 end-caps (T_z fixed)	3	17
Total:		8	47
2	IBL detector	1	6
	Pixel barrel layers	3	18
	Pixel end-cap disks	2×3	36
	SCT barrel layers	4	24
	SCT end-cap disks	2×9	108
	TRT barrel 32 modules (T_z fixed)	3×32	480
	TRT end-cap wheels	2×40	480
Total:		208	792
2.7	IBL staves	14	84
	Pixel barrel staves	$22+38+52$	672
	Pixel end-cap disks	2×3	18
	Total:		132
3	IBL modules	280	1,680
	Pixel modules	1,744	10,464
	SCT modules	4,088	24,528
	TRT barrel wires (T_ϕ, R_r only)	105,088	210,176
	TRT end-cap wires (T_ϕ, R_Z only)	245,760	491,520
	Total silicon sensors:		6,112
	Total TRT wires:		350,848
Total:		36,672	701,696

Table 4.1: The four alignment levels for each of the detector subsystems. The total number of alignable structures and degrees of freedom (DoF) to be aligned are given for each level.

805 direction of the module, the y' -axis is oriented along the long side of the module, and the z' -axis is
 806 orthogonal to the (x', y') plane. For the TRT straws, the x' -axis is perpendicular to both the wire
 807 and the radial direction, defined from the origin of the global frame to the straw center, the y' -axis
 808 points along the straw, and once again the z' -axis is orthogonal to the (x', y') plane. A depiction of
 809 the global and local coordinate systems for the ID is shown in Figure 4.3.

810 When considering the alignment degrees of freedom listed earlier in Section 4.1.1, grouped collec-
 811 tions of modules, layers, or entire subdetectors use the global coordinate system; individual modules
 812 use their respective local coordinate systems. The translations T_i are with respect to the origin of
 813 the given reference frame, and the rotations R_i are taken about the Cartesian axes.




Figure 4.3: A schematic representation of the Inner Detector in the longitudinal plane with the global coordinate system overlaid on top. The Pixel detector and IBL are shown in blue, the SCT in green, and the TRT in red. The local coordinates for each subdetector module are inset on the right. Image taken from [36].

814 4.2 Early 2015 alignment of the ATLAS detector

815 At the end of Run 1, the LHC was shut down for upgrades and maintenance. During this time,
 816 a number of upgrades were performed on the ATLAS detector, including the installation of a new
 817 innermost layer of the Pixel detector, the Insertable B-Layer (IBL) [37]. **TODO: This will certainly**
 818 **be defined in the detector description, so maybe the citation and abbreviation are not needed** These
 819 changes to the ID required some detector components to be removed temporarily, and many elements
 820 shifted relative to each other over the course of the maintenance process. In order to correct for these
 821 large detector movements prior to $\sqrt{s} = 13$ TeV collision data taking, an alignment was performed
 822 using cosmic ray data collected in early 2015 [36]. This alignment was able to correct for the majority
 823 of the large detector-wide misalignments as well as determine the global position of the IBL at the
 824 micron level.

825 In June of 2015, shortly after the data taking period began, the first track-based alignment
 826 of the refurbished ID was performed using $\mathcal{L} = 7.9 \text{ pb}^{-1}$ of $\sqrt{s} = 13$ TeV pp collision data [38].
 827 Starting from the initial geometry determined by the cosmic ray alignment, referred to hereafter
 828 as the *March alignment*, an improved set of alignment constants, called the *June alignment*, was
 829 derived from a data set of approximately 1.4 million selected tracks. For comparison, a MC sample
 830 containing approximately 2.7 million tracks was constructed from dijet events simulated using a
 831 perfect detector geometry; the MC events are reweighted to match the η and p_T distributions found
 832 in the data. Additional validation of the alignment results uses a set of cosmic ray data collected

833 by the detector during the LHC collisions. **TODO:** find some way of saying that this section follows
834 the alignment note [38]? or is it ok not to...

835 **4.2.1 June alignment procedure**

836 The data set used as the input for the alignment contains a subset of physics events used for prompt
837 reconstruction recorded at a rate of 10 Hz. To ensure that only high quality tracks are used for the
838 alignment, each track is required to have transverse momentum $p_T > 3$ GeV, contain at least one
839 hit in the Pixel detector, at least seven hits in the combined silicon detectors, and at least 25 hits
840 in the TRT.

841 A full L3 alignment of the IBL was included in the March alignment; however, a realignment
842 was necessary. Since the cosmic rays pass through the detector top-down, the staves on the sides of
843 the IBL could not be aligned as precisely as those on the top and bottom due to lower statistics.
844 Additionally, the IBL was operating at a temperature of -20°C during the cosmic data taking
845 and at -10°C for collision data taking. This proved to be significant, as it was observed that
846 the IBL staves experience a temperature-dependent, parabolic bowing in the local x' -direction of
847 approximately $-10\mu\text{m}/\text{K}$ [39]. As a result, a full L3 alignment of the IBL was essential in order to
848 correct for the bowing. Due to it being a brand new element of the detector as well as its importance
849 in vertexing and b jet tagging, aligning the IBL sensors with a high degree of precision was of great
850 importance.

851 The June alignment was performed in two stages, with the first pass focusing on relative move-
852 ments of the big structures and correcting for the bowing of the IBL. The March alignment corrected
853 for these larger movements as well; however, it was observed during Run 1 that these sort of mis-
854 alignments are introduced by changing conditions in the detector [33], such as in the cooling system
855 or magnet power cycling, which may have occurred between the early cosmic data taking and the
856 first $\sqrt{s} = 13$ TeV collisions. The silicon detectors were aligned at several different levels and the
857 IBL was aligned at the module level; the TRT detector was kept fixed to act as a global reference
858 frame. The full alignment chain for the first pass consisted of the following steps:

- 859 1. The IBL, Pixel, and SCT detectors were aligned at L1. The SCT barrel was not aligned in
860 T_z in order to constrain global displacements along the z -axis, as the TRT is not sensitive to
861 that degree of freedom.
- 862 2. The IBL, and Pixel barrel and end-caps, and SCT barrel were aligned at L2. The SCT end-caps
863 were aligned at L1.

864 3. The IBL and pixel barrel were aligned at L27, using all six degrees of freedom. The Pixel
 865 end-cap disks were only aligned in the plane (T_x , T_y , and R_z). The SCT was treated the same
 866 as in the previous step.

867 4. The IBL was aligned at L3 using all six degrees of freedom for each module.

868 The primary goal for the second pass was to remove a bias in the transverse impact parameter
 869 d_0 present in the March alignment. The resolution of d_0 was also poorer than expected. In order
 870 to correct for this, an additional constraint was passed to the alignment by adding an impact
 871 parameter with respect to the beam spot as a pseudo-measurement [40]. When the alignment
 872 algorithm minimizes the χ^2 , it will take care of the impact parameter minimization as well. Only
 873 the IBL and Pixel detectors were aligned in this step. The stages of the second pass are listed below,
 874 and the beam spot constraint was used in each:

875 1. The IBL and Pixel detectors were aligned at L2 with the SCT fixed.

876 2. The IBL was aligned at L27

877 3. The IBL and Pixel barrel and end-caps were aligned at L3.

878 The set of alignment constants obtained at the end of the second pass represents the June alignment.
 879 The highest level of alignment over the course of the two passes for each subdetector is listed in
 880 Table 4.2.

Detector	Highest level of alignment	
IBL	L3	
Pixel	Barrel	L3
	End-caps	L3 (T_x , T_y , and R_z only)
SCT	Barrel	L2 (except T_z)
	End-caps	L1
TRT	None	

Table 4.2: Summary of the highest level of alignment applied to each ID subsystem when deriving the June alignment.

881 4.2.2 Alignment results

882 The primary measure of alignment quality is assessed by looking at the track-hit residual distributions.
 883 If the detector is well aligned, the residuals will be Gaussian-distributed with a mean

of zero and a width approximating the detector's resolution. The residual distributions are constructed from the same selection of tracks that were used to perform the alignment, and are the focus of Section 4.2.2.1. A second check on the alignment involves observables sensitive to the track parameter resolution. In this case, cosmic rays are used, making use of a “split track” technique that takes advantage of the top-to-bottom cosmic ray trajectory (compared to the center-out trajectory of collision tracks). This method and the corresponding tests of the alignment are detailed in Section 4.2.2.2

Additionally, the effect of the beam spot constrained alignment on the impact parameter d_0 needs to be checked. The d_0 distributions for both the March and June alignments are compared to the MC simulation using a perfect geometry in Figure 4.4. In the March alignment, there is a bias of 18 μm in the mean of the distribution and the width is nearly twice that of the perfect geometry. After the second pass of the June alignment, the mean has shifted to 1 μm and the distribution has narrowed considerably. From this, it appears that the constrained alignment successfully removed the d_0 bias.




Figure 4.4: The d_0 distributions with respect to the reconstructed primary vertex using the $\sqrt{s} = 13$ TeV collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.

4.2.2.1 Residual distributions from collisions

As mentioned previously, the primary focus of the June alignment campaign was on the IBL and the Pixel detectors. The detectors are the closest to the beam line and have the finest resolutions of the

901 ID subdetectors. The residual distributions in local x and y of the IBL planar sensors⁷ are shown
 902 in Figure 4.5. These and subsequent figures in this section compare the June and March alignments
 903 to the perfectly-aligned MC simulation. Noticeable improvement in the distribution widths can be
 904 seen in both the local x - and y -directions, nearly matching the simulation in local x , which is the
 905 most sensitive direction.

906 Due to the temperature-dependent bowing of the IBL, it is also interesting to look at the means of
 907 the residual distributions for each ring of IBL sensors along the beam line, as shown in Figure 4.6. A
 908 deformation is clearly visible in the March alignment in both measurement directions, and the shape
 909 in the local x -direction is consistent with an average stave bowing due to the different operating
 910 temperature of the IBL during the March alignment and the 13 TeV collisions. This feature was
 911 nearly eliminated in both directions through the L3 alignment of the IBL sensors.




Figure 4.5: Local x (left) and local y (right) residual distributions of the IBL planar sensors using the $\sqrt{s} = 13$ TeV collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.

912 The local x and y residual distributions for the Pixel detector barrel and end-caps are shown in
 913 Figure 4.7. Even though the IBL is not included in the plots of the barrel, some of the noticeable
 914 improvement in the more sensitive local x direction is an effect of the improved IBL alignment.
 915 Similarly, the relatively broad local y residual distribution in the barrel likely indicates that further
 916 refinement of the IBL alignment was needed along that direction. Even so, the June alignment
 917 outperforms the March alignment and rivals the simulation in most of the plots.

⁷The IBL contains 12 planar sensors in the center of a stave, with four 3D sensors on either end. Only the planar sensors are shown here due to low statistics in the 3D sensors as well as poor MC modeling of these sensors.




Figure 4.6: The mean of the local x (left) and local y (right) residual distributions as a function of the global z position of each IBL module using the $\sqrt{s} = 13$ TeV collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red).

Similar distributions for the SCT and TRT barrel and end-caps are shown in Figures 4.8 and 4.9, respectively. Much like with the Pixel residuals, there is a reduction in the width of the TRT residuals between the March and June alignments due to the alignment of the other subdetectors improving the quality of the track fit. Even though neither subdetector was aligned at module-level, the residuals indicate that the previous L3 alignment performed in Run 1 has not degraded significantly during the upgrade and maintenance period.

4.2.2.2 Track parameter resolution from cosmic rays

Cosmic ray data is very useful as an independent check on the alignment in the barrel of the detector. While tracks from pp collisions originate within the detector and travel outwards, a cosmic ray that passes through the center of the detector leaves a track in both halves of the detector. If the cosmic ray is split in half, as in Figure 4.10, then it can be treated as two separate tracks each with nearly identical track parameters (some differences arise due to energy loss as the particle passes through the detector). The distribution of the difference in a given track parameter $\Delta\tau$ is approximately Gaussian with a variance $\sigma^2(\Delta\tau)$. Since both tracks come from the same particle, each track individually has a variance equal to $\sigma^2(\Delta\tau)/2$. The resolution of the track parameter is then given by the root mean square of the distribution divided by $\sqrt{2}$.

Cosmic rays whose split tracks each had transverse momentum $p_T > 2$ GeV and at least one, eight, and 25 hits in the barrels of the Pixel, SCT, and TRT detectors, respectively, were selected to




Figure 4.7: Local x (top) and local y (bottom) residual distributions for the Pixel barrel (excluding the IBL, left) and end-caps (right) using the $\sqrt{s} = 13$ TeV collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.

measure a collection of track parameters. Figure 4.11 shows the difference in the impact parameter Δd_0 and the charge divided by the transverse momentum $\Delta q/p_T$ of the selected split-track cosmic rays for both the March and June alignments. Both distributions show a reduction in width in the June alignment, corresponding to an improvement in the resolution of each track parameter. The Δd_0 plot shows a significant improvement in the June alignment, further validating the removal of the bias in the impact parameter.

4.2.3 Error scaling

The final step in preparing the new set of June alignment constants deals with the adjustment of the hit errors, or *error scaling*. Knowledge of the exact position of a hit measurement on a track is limited by the accuracy with which the sensors' positions are known. Let σ represent the hit




Figure 4.8: Local x residual distributions for the SCT barrel (left) and end-caps (right) using the $\sqrt{s} = 13 \text{ TeV}$ collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.




Figure 4.9: Residual distributions for the TRT barrel (left) and end-caps (right) using the $\sqrt{s} = 13 \text{ TeV}$ collision data sample reconstructed using the June (black) and March (green) alignments. The data is compared to a MC simulation using a perfect detector geometry (red). The distributions are normalized to the same number of entries.




Figure 4.10: Representation of splitting a single cosmic ray track passing through the entire detector (left) into two separate tracks (right).




Figure 4.11: Distribution of the difference in the impact parameter Δd_0 (left) and charge over transverse momentum $\Delta q/p_T$ (right) between the two cosmic ray split tracks. The June (black) and March (green) alignments are compared. The distributions are normalized to the same number of entries.

uncertainty used in track fitting, and σ_0 be the detector's intrinsic uncertainty. If $\sigma = \sigma_0$, the pull of the track-hit residual distributions should form a Gaussian distribution centered at zero with a standard deviation $\sigma = 1$ [30]. In the case of residual misalignment, the pull distributions' standard deviations will stray from unity. The hit uncertainty can be written as:

$$\sigma = a \cdot \sigma_0 \oplus b \quad (4.12)$$

where a is a scaling factor, and b is a constant term which can be interpreted as a measure of any remaining misalignment of the detector elements. In this alignment campaign, the value of a is fixed at $a = 1$ and b is evaluated from the residual pull distributions for each subdetector in its sensitive directions.

Once the value of b is determined, pull distributions derived from the new value of σ should have unit width. The error scaling values for each subdetector are listed in Table 4.3, and the pull distributions for the IBL after error scaling are shown in Figure 4.12.

Detector	Coordinate	$b(\mu\text{m})$
IBL	x	6.4
	y	43.6
Pixel	x	5.2
	y	28.6
SCT	x	7.5
	y	0
TRT	x	10.8
	$r\phi$	0
End-caps	x	8.6
	$r\phi$	0

Table 4.3: Estimated value of the error scaling term b for each subdetector component with the June alignment.




Figure 4.12: Pull distributions in local x (left) and y (right) for the IBL using the $\sqrt{s} = 13 \text{ TeV}$ collision data sample after applying the error scaling.

4.3 Level 2 alignment of the TRT

During validation of the final end-of-year reprocessing of the 2015 data, a misalignment was found in the barrel of the TRT detector, as several modules (triangular clusters of straws) showed rotations in the local y coordinate. The then-best available constants included a full L3 alignment of the silicon detectors and a separate L2 alignment of the TRT. However, not all degrees of freedom were enabled when the TRT was aligned. To correct for these tilts, an additional four iterations of L2 alignment

963 TRT was performed on the TRT enabling all available degrees of freedom (T_x , T_y , R_x , R_y , and R_z
 964 in the barrel, and T_x , T_y , and R_z for the endcaps). Plots of the residual means from barrel ϕ sectors
 965 containing modules affected by the tilt misalignment are shown in Figure 4.13 before and after the
 966 L2 alignment.




Figure 4.13: Residual means by straw layer in two TRT ϕ -sectors affected by a tilt misalignment. The tilts in each of the three modules are clearly visible in the red points representing the reconstructed data prior to alignment. After four iterations of L2 alignment, the residual means in the gray points are flat.

967 Following the L2 alignment, some additional time was taken to determine if a full wire-by-wire
 968 L3 alignment of the TRT was necessary. The TRT was last aligned at L3 during Run 1, but initial
 969 alignment campaigns in Run 2 did not show signs of misalignment, as can be seen from the residual
 970 distributions in Figure 4.9. In order to assess the alignment more carefully, two dimensional residual
 971 maps in ϕ and z were constructed for each layer in the TRT barrel and endcaps using the current
 972 alignment. These maps were compared to a similar set using the L3 alignment from 2010, from which
 973 it was determined that the straw-level alignment indeed hadn't degraded and a new L3 alignment
 974 was not needed. The maps for the first layer of the TRT barrel are shown in Figure 4.14 for both
 975 sets of alignment constants.

976 4.4 Momentum bias from sagitta deformations

977 A variety of weak mode deformations can exist in the detector even after alignment. As mentioned
 978 previously, these weak modes consist of misalignments which don't affect the χ^2 of the residuals
 979 and thus are not handled by the basic alignment algorithm. In the presence of a weak mode, the
 980 description of the detector geometry can still provide efficient and high quality track fits, but there




Figure 4.14: Two dimensional map of residuals in the first layer of the TRT barrel vs z and ϕ . Each bin represents the mean of a Gaussian fit to the TRT residuals in that bin. The map on the left is after the L3 (wire-by-wire) alignment of the TRT performed in 2010, and the map on the right is after the L2 alignment at the end of 2015. The z -axis for both plots use the same scale. Left figure taken from [30].

981 may also be systematic biases in one or more track parameters. Several weak modes, their impacts
 982 on the reconstruction, and the steps taken to eliminate them are detailed in [33, 41]. This section
 983 focuses specifically on sagitta distortions that result in a bias in the reconstructed track momentum.

984 These *sagitta* distortions consist of detector movements orthogonal to the trajectory of the outgoing
 985 particle. The effect on the reconstructed track curvature is different for positively and negatively
 986 charged particles, resulting in a charge-antisymmetric bias. This effect is illustrated in the curl
 987 deformation shown in Figure 4.15.

988 In the plane transverse to ATLAS’s magnetic field, outgoing particle tracks form circular arcs.
 989 The sagitta is defined as the distance from the center of this arc to the center of its base, as shown in
 990 Figure 4.16, and it represents the “amount of bending” in the track. In the case where the sagitta s
 991 is considerably smaller than the detector radius R_0 , which is a valid assumption when working with
 992 high momentum tracks, the transverse momentum of a particle of charge q can be written as [42]:

$$p_T \propto qB \frac{R_0^2}{8s} \quad (4.13)$$

993 where B is the strength of the detector’s magnetic field. If a sagitta bias is present, the track’s
 994 transverse momentum shifts by [41]:

$$q/p_T \rightarrow q/p_T + \delta_s \quad \text{or} \quad p_T \rightarrow p_T \cdot (1 + qp_T\delta_s)^{-1} \quad (4.14)$$

995 where δ_s is a universal bias parameter that uniquely defines the deformation. Finally, since the
 996 reconstructed polar angle does not change under a sagitta deformation, the longitudinal component

997 of the momentum scales along with the transverse component, and an equivalent equation can be
 998 written for the total momentum:

$$p \rightarrow p \cdot (1 + qp_T\delta_s)^{-1} \quad (4.15)$$




Figure 4.15: Representation of a curl distortion in the detector. The image shows a cutaway in the transverse plane. The deformation is represented by the red arrows, and the impact on the reconstructed positive (blue) and negative (green) tracks are shown. The dashed lines represent the true particle trajectories, and the solid lines represent the reconstructed trajectories.




Figure 4.16: Geometric definition of the sagitta s in relation to the length of the chord l and the radius r of a circular arc.

999 **4.4.1 Sagitta bias monitoring with electron E/p**

1000 Since a sagitta bias results in changes in the momentum of particles' tracks as measured by the ID,
 1001 they can be identified using independent measurements from other systems in the detector. One
 1002 such method involves using the energy-momentum ratio of electrons (E/p). Since the electron's
 1003 energy is measured in ATLAS's calorimeter systems, it is not sensitive to any sagitta bias that may
 1004 exist in the ID and the corresponding track momentum. Under the assumption that the calorimeter

1005 response is independent of the charge of incoming particles, a charge-dependent momentum bias in
 1006 the ID will manifest as a difference in the E/p ratio for electrons and positrons.

1007 In the presence of a sagitta bias, the momentum will change according to Equation 4.15 and the
 1008 average measured $\langle E/p \rangle$ can be written as:

$$\langle E/p \rangle^\pm \rightarrow \langle E/p \rangle^\pm \pm \langle E_T \rangle \delta_s \quad (4.16)$$

1009 where the approximation $p_T \approx E_T$ is used. Assuming that $\langle E/p \rangle^+ = \langle E/p \rangle^-$ in the absense of a
 1010 bias, the sagitta bias parameter can be written as:

$$\delta_s = \frac{\langle E/p \rangle^+ - \langle E/p \rangle^-}{2\langle E_T \rangle} \quad (4.17)$$

1011 If the kinematic selections for electrons and positrons are identical, the energy scale of the calorimeter
 1012 will not factor into the $\langle E/p \rangle$ difference; however, it will affect $\langle E_T \rangle$ which would scale the measured
 1013 δ_s . This is expected to be a small effect, as the energy scale for electrons has been measured at
 1014 $\sqrt{s} = 13$ TeV with uncertainties on the per-mil level across the entire detector [43].

1015 4.4.1.1 Measuring $\langle E/p \rangle$

1016 The E/p ratio is measured using electrons and positrons from $Z \rightarrow e^\pm e^\mp$ events in order to obtain
 1017 a high purity sample of electron candidates. They are required to pass a basic selection criteria to
 1018 ensure they are well measured by both the ID and the calorimeters:

- 1019 • $E_T > 25$ GeV
- 1020 • $|\eta| < 2.47$, excluding the calorimeter's barrel-to-endcap transition region in $1.37 < |\eta| < 1.52$
- 1021 • Pass MediumLH identification working point detailed in Section 3.2.4.3
- 1022 • Pass a selection of quality cuts, including a requirement that the electron be identified using
 1023 cluster information in the calorimeter
- 1024 • The associated track must have at least one hit in the IBL, three in the Pixel detector, and
 1025 five in the SCT detector.

1026 Events with exactly two opposite-charge electrons passing this selection with a dielectron invariant
 1027 mass within 30 GeV of the Z boson mass are then used for the E/p calculation.

1028 Since the size of the sagitta bias δ_s is not expected to be constant across the entire detector,
 1029 a two-dimensional rectangular grid binned in detector η and ϕ is constructed. From the selected

1030 events, separate distributions of E/p are made for electrons and positrons lying in each bin. Each
 1031 distribution is fit with Crystal Ball function⁸, and the peak of the distribution is taken as the value
 1032 of $\langle E/p \rangle$. If there is no bias on the track momentum in the bin, the peaks for electrons and positrons
 1033 should match. Example E/p distributions including the Crystal Ball fits are shown in Figure 4.17.

1034 It is important to emphasize that deviations from one in the *ratio* of $\langle E/p \rangle$ for electrons and
 1035 positrons indicates that a momentum bias may be present. The value of $\langle E/p \rangle$ itself is not expected
 1036 to equal one exactly, as the track momentum on average tends to be slightly lower than the energy
 1037 measurement in the calorimeter. This is due to the fact that if the electron were to radiate a photon,
 1038 its momentum would change slightly, while it is likely that both the electron and the emitted photon
 1039 would leave energy deposits near each other in the calorimeter and be reconstructed into the same
 1040 object.




Figure 4.17: E/p distributions of electrons and positrons in two different $\eta\phi$ bins of the detector. The left hand plot is taken from a region with no momentum bias where $\langle E/p \rangle^+ = \langle E/p \rangle^-$, while the right hand plot shows an 8% disagreement in $\langle E/p \rangle$ between electrons and positrons.

1041 Once the $\langle E/p \rangle^\pm$ distributions in each $\eta\phi$ bin have been extracted from the fits, a two dimen-
 1042 sional map of δ_s can be constructed using Equation 4.17. The map gives an overview of sagitta
 1043 biases that may be present in the detector, and can be used by the alignment algorithm to reduce
 1044 the bias in the next iteration. In this case, the tracks fed to the alignment have their momenta
 1045 corrected according to [41]:

$$q/p_{\text{corr}} = q/p_{\text{reco}}(1 - qp_T\delta_s) \quad (4.18)$$

1046 where p_{reco} is the reconstructed momentum of the track. The corrected momentum is then con-
 1047 strained in the alignment.

⁸The Crystal Ball function is a probability density function consisting of a Gaussian core and a power-law tail.

1048 **4.4.1.2 Results in 13 TeV data**

1049 The E/p method has been used to monitor sagitta biases in the detector several times over the
1050 course of Run 2. During this time, it has primarily served as an independent cross-check to a
1051 second method using $Z \rightarrow \mu^\pm \mu^\mp$ events [41]. The $Z \rightarrow \mu^\pm \mu^\mp$ method identifies individual track
1052 momentum biases through shifts in the reconstructed Z mass, which leaves it relatively insensitive
1053 to global sagitta biases. For this reason, the sagitta bias maps produced using this technique are
1054 normalized to those from the E/p method before being used to constrain the alignment.

1055 The results of two implementations of the E/p method are presented here.

- 1056 1. The first follows the end-of-year reprocessing of the entire ATLAS 2016 data set at $\sqrt{s} =$
1057 13 TeV. Two sets of alignment constants are compared: the *prompt* alignment, which was
1058 derived shortly after each run was recorded, and the *reprocessed* alignment. The maps of the
1059 sagitta bias comparing the two alignments calculated using the E/p method are shown in
1060 Figure 4.18, and the comparison of the η projection of each map is shown in Figure 4.19.
- 1061 2. The second uses the 2017 data after reprocessing, and compares the effects of multiple it-
1062 erations of the method. In each iteration, the momenta of the electrons and positrons are
1063 corrected based on Equation 4.15 using the value of δ_s computed in the previous iteration,
1064 and a new sagitta bias map is calculated. If the method is indeed characterizing the sagitta
1065 biases correctly, the corrections should converge quickly. The initial sagitta bias map is com-
1066 pared to the map after two such iterations in Figure 4.20, and the sagitta bias projected along
1067 η for each iteration is shown in Figure 4.21. Indeed, after just two iterations, δ_s is consistent
1068 with zero in nearly all bins.




Figure 4.18: Sagitta bias in the $\sqrt{s} = 13$ TeV data collected by ATLAS in 2016 as a function of η and ϕ for the prompt (left) and reprocessed (right) alignments using the E/p method.




Figure 4.19: Sagitta bias in the $\sqrt{s} = 13$ TeV data collected by ATLAS in 2016 projected along η for the prompt (gray) and reprocessed (blue) alignments using the E/p method.




Figure 4.20: Sagitta bias in the $\sqrt{s} = 13$ TeV data collected by ATLAS in 2017 as a function of η and ϕ in reconstructed electrons (left) and after two iterations of momentum corrections (right) from the E/p method.




Figure 4.21: Sagitta bias in the $\sqrt{s} = 13$ TeV data collected by ATLAS in 2017 projected along η in reconstructed electrons (gray) and after one (red) and two (blue) iterations of momentum corrections from the E/p method.

1069

CHAPTER 5

1070

1071

Measurement of same-sign WW production at $\sqrt{s} = 13$ TeV with ATLAS

1072 Production of same-sign W boson pairs is a particularly interesting SM process. When produced
 1073 via vector boson scattering (VBS), $W^\pm W^\pm jj$ is particularly sensitive to the electroweak symmetry
 1074 breaking (EWSB) mechanism as well as potential “beyond the Standard Model” (BSM) physics.
 1075 $W^\pm W^\pm jj$ events can be produced via electroweak-mediated (EWK) diagrams, of which VBS is a
 1076 subset, or QCD-mediated diagrams. The biggest advantage of same-sign $W^\pm W^\pm jj$ lies in its ratio
 1077 of electroweak (EWK) to QCD production cross sections. Despite the opposite-sign $W^\pm W^\mp$ having
 1078 a considerably larger total cross section, its EWK-mediated diagrams are considerably smaller than
 1079 its QCD-mediated diagrams, while for same-sign $W^\pm W^\pm$ the ratio is approximately one to one.
 1080 This makes $W^\pm W^\pm jj$ one of the best channels for studying VBS at the LHC.

1081 The first evidence of electroweak (EWK) $W^\pm W^\pm jj$ production was seen by the ATLAS and
 1082 CMS experiments at $\sqrt{s} = 8$ TeV with excesses of 3.6σ [21] and 2.0σ [44] over backgrounds, respec-
 1083 tively. More recently, ATLAS and CMS have both observed the EWK process at $\sqrt{s} = 13$ TeV
 1084 with significances of 6.9σ [45] and 5.5σ [46], respectively. The analysis presented in this chap-
 1085 ter is based off of the ATLAS $\sqrt{s} = 13$ TeV observation and cross section measurement of EWK
 1086 $W^\pm W^\pm jj$ production [45, 47].

1087 **5.0.2 Theoretical overview of vector boson scattering**

1088 VBS processes are very important to understand due to their sensitivity to the EWSB mechanism.
 1089 The scattering amplitude of longitudinally polarized vector bosons grows with center-of-mass energy
 1090 and ultimately violates unitarity above $\sqrt{s} = 1$ TeV in the absence of a light SM Higgs boson [48, 49].

1091 However, once the Higgs is introduced, the divergences cancel and the cross section no longer grows
 1092 unbounded, as can be seen in Figure 5.1, which consists of plots from [50].




Figure 5.1: Cross sections in nanobarns for five different scattering processes of longitudinally polarized vector bosons as a function of center of mass energy \sqrt{s} . Without a SM Higgs boson (left), the cross sections grow unbounded with \sqrt{s} ; however with a 120 GeV Higgs boson (right), the cross sections no longer diverge. Plots taken from [50].

1093 With the discovery of the Higgs boson in 2012 [15, 16], the EWSB mechanism can now be directly
 1094 studied. Due to the exchange of a Higgs in the s - and t -channel VBS diagrams ($W^\pm W^\pm jj$ itself only
 1095 contains the t -channel diagram), VBS processes are directly sensitive to properties of the Higgs. For
 1096 example, the high-mass tail in the VV scattering system allows an approximation of the effective
 1097 coupling strength of the Higgs to vector bosons that is independent of any assumptions on the Higgs
 1098 width [51]. Additionally, the center of mass energy dependence of the VV scattering can reveal
 1099 whether the Higgs boson unitarizes the longitudinal scattering amplitude fully or only partially [52].

1100 VBS events are characterized by two quarks from the colliding protons each radiating a massive
 1101 vector boson which then scatter and decay in the detector. The incoming quarks carry a large
 1102 amount of momentum and only deflect a small amount upon radiating the vector boson; as a result,
 1103 they often travel very close to the beam line. Ignoring the decay products of the bosons, these VBS
 1104 events result in a final state of two vector bosons (V) and two jets (j) at high pseudorapidities
 1105 (called *forward jets*) from the outgoing quarks. The shorthand $VVjj$ is used to represent this final
 1106 state.

1107 $VVjj$ events can be produced via two different physical processes. The first involves purely
 1108 electroweak interactions in the tree-level diagrams, with $\mathcal{O}(\alpha_{EWK}) = 6$ and will be referred to as
 1109 *EWK production*. This can be further broken down into VBS and non-VBS production. In the
 1110 VBS EWK production, the scattering occurs via triple or quartic gauge couplings, as well as the

s- or t -channel exchange of a Higgs boson. The non-VBS EWK production contains the same final state of two vector bosons and two outgoing quarks, but the bosons do not scatter. Due to gauge invariance, it is not possible to separate the VBS from the non-VBS productions [53]; therefore, both are included in the signal generation and are indistinguishable from one another. The second process involves a mix of the EWK and strong interactions, of order $\mathcal{O}(\alpha_s) = 2 \otimes \mathcal{O}(\alpha_{\text{EWK}}) = 4$ and will be referred to as *QCD production*. The tree-level Feynman diagrams for VBS EWK, non-VBS EWK, and QCD $VVjj$ production are found in Figures 5.2, 5.3, and 5.4, respectively.




Figure 5.2: Tree-level Feynman diagrams for VBS EWK $VVjj$ production including triple gauge couplings involving W and/or Z bosons (top left and top middle), quartic gauge coupling (top right), or the exchange of a Higgs boson (s -channel bottom left and t -channel bottom right). The labels are quarks (q), fermions (f), and gauge bosons ($V = W, Z$).




Figure 5.3: Tree-level Feynman diagrams for non-VBS EWK $VVjj$ production. The labels are quarks (q), fermions (f), and gauge bosons ($V = W, Z$).

5.0.3 Same-sign $W^\pm W^\pm$ scattering

Same-sign $W^\pm W^\pm jj$ scattering is considered to be one of the best channels for studying VBS at the LHC [51]. This is due primarily to the ratio of the EWK to the QCD production, which matters




Figure 5.4: Tree-level Feynman diagrams for QCD $VVjj$ production. The labels are quarks (q), fermions (f), and gauge bosons ($V = W, Z$).

a great deal due to the VBS events being a subset of the total EWK production. In an analysis the EWK production would be considered the signal and the QCD production a background, so a favorable ratio of the two helps greatly when comparing the size of the signal to the backgrounds. A study at $\sqrt{s} = 8$ TeV [54] was done using the **SHERPA** Monte Carlo (MC) generator to calculate EWK and QCD production cross sections at leading order for a variety of $VVjj$ processes decaying to leptons and can be found in Table 5.1. Despite its lower cross section compared to other $VVjj$ processes, the EWK to QCD ratio for $W^\pm W^\pm jj$ is approximately one-to-one, whereas for opposite-sign $W^\pm W^\mp jj$ the ratio is closer to 3%.

Process	Final state	σ_{EWK}	σ_{QCD}
$W^\pm W^\pm$	$l^\pm l^\pm \nu\nu jj$	19.5 fb	18.8 fb
$W^\pm W^\mp$	$l^\pm l^\mp \nu\nu jj$	91.3 fb	3030 fb
$W^\pm Z$	$l^\pm l^\pm l^\mp \nu jj$	30.2 fb	687 fb
ZZ	$l^+ l^- \nu\nu jj$	2.4 fb	162 fb
ZZ	$l^+ l^- l^+ l^- jj$	1.5 fb	106 fb

Table 5.1: Predicted cross sections for EQK and QCD production of diboson processes relevant to VBS at $\sqrt{s} = 8$ TeV using the **SHERPA** MC generator. Loose generator level cuts are applied on lepton $p_T > 5$ GeV, dilepton invariant mass $m_{ll} > 4$ GeV, and at least two jets with $m_{jj} > 10$ GeV. Numbers taken from [54].

This analysis studies $W^\pm W^\pm jj$ scattering where both W bosons decay leptonically to $e\nu$ or $\mu\nu$ ⁹. The $W^\pm W^\pm jj$ VBS final state consists of two leptons with the same electric charge, two neutrinos, and two high energy forward jets with a large invariant mass. Tree-level Feynman diagrams of VBS $W^\pm W^\pm jj$ production can be found in Figure 5.5 and a visual representation of the VBS topology can be found in Figure 5.6. The two forward jets also serve as a powerful tool to suppress the QCD production mode. In EWK events, the two jets tend to have much higher separation and a larger combined invariant mass than the two leading jets in a QCD event. The two plots shown in

⁹Throughout the rest of this chapter, l denotes either electrons (e) or muons (μ) unless stated otherwise. Additionally, e , μ , and ν (neutrino) with no charge or anti-particle designation refer interchangeably to either the particle or anti-particle.

1136 Figure 5.7 highlight the differences in these dijet quantities between the two production modes. An
 1137 ATLAS event display of a real $W^\pm W^\pm jj$ candidate event is shown in Figure 5.8.




Figure 5.5: Feynman diagrams for VBS EWK production of $W^\pm W^\pm jj$ events. The leftmost diagram contains a quartic gauge coupling vertex, and the rightmost diagram contains an exchange of a Higgs boson. **TODO: Make diagrams consistent with others**




Figure 5.6: $W^\pm W^\pm jj$ VBS event topology containing two leptons (1 and 2) with the same electric charge, two neutrinos, and two forward tagging jets (3 and 4) with large rapidity separation Δy .

1138 5.0.4 Overview of backgrounds

1139 In addition to QCD production of $W^\pm W^\pm jj$ events, there are several other processes that can end
 1140 up with a final state of two same-sign leptons, two neutrinos, and two jets. However, due to the ±2
 1141 final state charge, there is a considerable reduction in SM backgrounds (such as Z boson events)
 1142 when compared to an analysis like opposite-sign $W^\pm W^\mp jj$.

1143 One of the largest sources of background involves processes with prompt leptons¹⁰. These are
 1144 events that contain two leptons with the same electric charge and one or more additional leptons

¹⁰Prompt leptons are those that are produced in the primary collision and are a direct decay product of the process of interest. Non-prompt leptons originate from some secondary process, such as a b -hadron decay, or are jets that get mis-reconstructed as a lepton.




Figure 5.7: Generator level comparisons at $\sqrt{s} = 8$ TeV of dijet invariant mass (m_{jj} , left) and dijet rapidity (Δy_{jj} , right) in EWK (red) and QCD (blue) $W^\pm W^\pm jj$ events. Both data sets have been normalized to the same area. Plots taken from [54].




Figure 5.8: ATLAS event display of a $pp \rightarrow W^\pm W^\pm \rightarrow \mu^\pm \nu_\mu \mu^\pm \nu_\mu jj$ event. The muons are represented by the red lines travelling from the ID through the MS, and the forward jets are represented by the blue cones with yellow energy deposits in the calorimeters. The direction of the E_T^{miss} in the transverse plane is indicated by the gray dashed line in the inset image. Event display taken from [45].

that are “lost”, either by failing the selection criteria or falling outside of the detector’s acceptance. The number of processes that can contribute is limited by the requirement of same-sign leptons, and as a result this background is dominated by processes involving two or more vector bosons, with the largest contribution coming from WZ events and smaller contributions from ZZ and $t\bar{t}V$ events. Triboson events where one boson decays hadronically also contribute to this background; however, the jets are generally softer and more central than in a typical VBS event, and the cuts applied on the forward jets suppress these contributions.

The other dominant background comes from non-prompt, or “fake”, leptons. Here one or more leptons originate from the decay of another particle unrelated to the signal process, such as a heavy-flavor decay or photon conversion, or come from a jet that is misidentified as a lepton. This background is mostly made up of events from $t\bar{t}$ and $W+\text{jets}$ processes, with a much smaller contribution from $V\gamma$ events. **TODO: check whether $V\gamma$ really qualifies as non-prompt, we lump $Z\gamma$ in with the charge flip background in the paper...**

Finally, opposite-sign lepton pairs can enter the signal region if one of the leptons is reconstructed with the wrong charge (called *charge misidentification*¹¹). In practice, this only affects events with electrons, as the charge misidentification rate for muons is negligible [55]. This is a major background in events with two electrons, but is a much smaller contribution for events with one electron and one muon.

5.1 Data and Monte Carlo samples

This analysis uses 36.1 fb^{-1} of $\sqrt{s} = 13$ TeV proton-proton collision data recorded by ATLAS during 2015 and 2016. The uncertainty in the combined integrated luminosity is 2.1%. It is derived following a methodology similar to that detailed in [56] and using the LUCID-2 detector for the baseline luminosity measurements [57] from calibration of the luminosity scale using x - y beam-separation scans.

5.1.1 Monte Carlo samples

A number of Monte Carlo (MC) simulations are employed to model signal and background processes. In order to model the real collision data as closely as possible, each MC has been run through a full simulation of the ATLAS detector [58] in GEANT4 [59], and events have been reconstructed

¹¹Charge misidentification is also referred to interchangeably as *charge mis-ID* and *charge flip*.

1173 using the same algorithms as the data. The simulation reproduces as closely as possible the momentum
 1174 resolutions and calorimeter responses of the detector, and also includes the effects of pileup by
 1175 including soft QCD interactions using PYTHIA v8.1 [60]. The MC samples used in this analysis are
 1176 detailed in this section and summarized in Table 5.2.

1177 The $W^\pm W^\pm jj$ samples are modeled using SHERPA v2.2.2 [61, 62, 63] with the NNPDF3.0 PDF
 1178 set [64]. The EWK signal samples are generated by fixing the electroweak coupling constant to
 1179 $\mathcal{O}(\alpha_W) = 6$, and a QCD background sample was also generated with $\mathcal{O}(\alpha_W) = 4$. SHERPA includes
 1180 up to one parton at next-to-leading order (NLO) and up to three at leading order (LO) in the
 1181 strong coupling constant α_s . A second $W^\pm W^\pm jj$ EWK sample is generated using POWHEG-BOX
 1182 v2 [65] with the NNPDF3.0 PDF set and at NLO accuracy. This sample is only used for systematic
 1183 studies, as POWHEG-BOX does not include resonant triboson contributions in its matrix element, which
 1184 are non-negligible at NLO [66].

1185 Diboson processes (VV where $V = W, Z$) are simulated with SHERPA v2.2.2 for mixed hadronic
 1186 and leptonic decays and SHERPA v2.2.1 for fully leptonic decays of the bosons. Similarly, triboson
 1187 (VVV) and $V\gamma$ processes are simulated using SHERPA v2.1.1 with up to one parton at NLO and up
 1188 to three at LO. $W+jets$ processes are simulated with SHERPA2.2.1 with up to two partons at NLO
 1189 and four at LO. All the above SHERPA samples use the NNPDF3.0 PDF set and SHERPA's own parton
 1190 showering. The $Z+jets$ events are generated with Madgraph5_aMC@NLO [67] at LO and interfaced
 1191 with PYTHIA v8.1 for parton showering.

1192 $t\bar{t}$ events are generated using POWHEG-BOX v2 with the CT10 PDF set [68]. $t\bar{t}V$ samples are
 1193 generated at NLO with Madgraph5_aMC@NLO and the NNPDF3.0 PDF set interfaced with PYTHIA v8
 1194 for parton showering. Finally, single top events are generated with POWHEG-BOX v1 and the CT10f4
 1195 PDF set interfaced with PYTHIA v6 [69] for parton showering.

1196 5.2 Object and event selection

1197 This section details the selection criteria for objects used in the analysis as well as the selection for
 1198 signal events.

1199 5.2.1 Object selection

1200 Muons, electrons, and jets all must pass strict selection requirements to ensure that only high quality,
 1201 well measured objects are used. For leptons, a baseline selection is defined (called the *preselection*),
 1202 which all leptons must pass in order to be considered for the analysis. This preselection is an

Process	Generator	Comments
$W^\pm W^\pm jj$ (EWK)	SHERPA v2.2.2	Signal sample
$W^\pm W^\pm jj$ (EWK)	POWHEG-BOX v2	Systematics sample
$W^\pm W^\pm jj$ (QCD)	SHERPA v2.2.2	
Diboson	SHERPA v2.2.2	Both bosons decay leptonically ($llll$, $lll\nu$, $ll\nu\nu$)
Triboson	SHERPA v2.2.1	One boson decays leptonically, the other hadronically
$W + \text{jets}$	SHERPA v2.2.1	
$Z + \text{jets}$	Madgraph5_aMC@NLO	
$V\gamma$	SHERPA v2.1.1	
$V\gamma jj$ (EWK)	SHERPA v2.2.4	
$t\bar{t}V$	Madgraph5_aMC@NLO	
$t\bar{t}$	POWHEG-BOX v2	
Single top	POWHEG-BOX v1	EWK t -, s -, & Wt -channels

Table 5.2: Summary of MC samples used in the analysis.

1203 intentionally loose set of criteria in order to have high acceptance for rejecting backgrounds with
 1204 additional leptons (i.e. $WZ \rightarrow 3l\nu jj$). Signal leptons are then required to satisfy a much tighter
 1205 *signal selection* aimed at suppressing backgrounds from non-prompt or fake leptons. A third set of
 1206 lepton selection criteria, the *loose selection*, defines a sample enriched in non-prompt leptons, and
 1207 it is used in the fake factor method for estimating the non-prompt background, discussed in detail
 1208 in Section 5.3.4. Jets are only required to pass one set of selection criteria. These selections are
 1209 detailed in the following sections and summarized in Table 5.3 for muons, Table 5.4 for electrons,
 1210 and Table 5.5 for jets.

1211 5.2.1.1 Muon candidate selection

1212 Cuts on muon p_T serve to reject low momentum leptons from background processes and additional
 1213 collisions from pileup events. Preselected muons must have $p_T > 6$ GeV and signal muons $p_T >$
 1214 27 GeV. The p_T requirement for loose muons is lower than for signal muons, $p_T > 15$ GeV, for
 1215 reasons that are discussed in Section 5.3.4. **TODO:** reference proper subsection when it's done
 1216 Muons are required to fall within the detector's η acceptance: $|\eta| < 2.7$ for preselected muons,
 1217 which is tightened to $|\eta| < 2.5$ for the signal muons.

1218 Cuts on the transverse and longitudinal impact parameters are applied to ensure that the can-
 1219 didate muon originated from the primary particle interaction and not some other source, such as a
 1220 heavy flavor decay. The preselection and the loose selection both have looser requirements on the
 1221 transverse impact parameter significance (d_0/σ_{d_0}) than the signal selection; all three have the same
 1222 requirement on the transverse impact parameter ($|z_0 \times \sin \theta|$).

Finally, the muon candidates are required to pass a particle identification and an isolation criteria as defined in [70]. The methods used in constructing the identification and isolation workingpoints are described in more detail in Section 3.2.4.2. The muon identification serves to select prompt muons with high efficiency and well measured momenta. This analysis uses two different workingpoints, **Loose** for preselected muons and **Medium** for loose and signal muons, where **Medium** muons are a tighter subset of those that pass the **Loose** requirement. Muon isolation is a measurement of detector activity around the muon candidate, and it is measured with both track-based and calorimeter-based variables. The isolation workingpoint used for the signal muons, **Gradient**, is defined such that there is 90% or better background rejection efficiency for 25 GeV muons, and 99% efficiency at 60 GeV. There is no minimum isolation requirement for preselected or loose muons. Loose muons are additionally required to fail one or both of the signal transverse impact parameter cut and signal isolation requirement.

Muon preselection	
Momentum cut	$p_T > 6$ GeV
Angular acceptance	$ \eta < 2.7$
Longitudinal impact parameter	$ z_0 \times \sin \theta < 0.5$ mm
Transverse impact parameter	$d_0/\sigma_{d_0} < 10$
Particle identification	Loose

Muon signal selection	
Momentum cut	$p_T > 27$ GeV
Angular acceptance	$ \eta < 2.5$
Longitudinal impact parameter	$ z_0 \times \sin \theta < 0.5$ mm
Transverse impact parameter	$d_0/\sigma_{d_0} < 3$
Particle identification	Medium
Particle isolation	Gradient

Muon loose selection	
Momentum cut	$p_T > 15$ GeV
Angular acceptance	$ \eta < 2.5$
Longitudinal impact parameter	$ z_0 \times \sin \theta < 0.5$ mm
Transverse impact parameter	$d_0/\sigma_{d_0} < 10$
Particle identification	Medium
Fail signal transverse impact parameter and/or isolation cuts	

Table 5.3: Muon selection criteria. All muons are required to pass the preselection (top), and then either the signal (middle) or loose (bottom) criteria is applied to the preselected electrons.

1235 **5.2.1.2 Electron candidate selection**

1236 The electron candidate selections are very similar to those for muons. The p_T cut starts at $p_T >$
 1237 6 GeV for the preselection, increases to $p_T > 20$ GeV for loose electrons, and finally to $p_T > 27$ GeV
 1238 for signal electrons. The $|\eta|$ cut for electrons requires $|\eta| < 2.47$ for all electrons, with the region
 1239 $1.37 \leq |\eta| \leq 1.52$ removed from loose and signal electrons. This region is where the electromagnetic
 1240 calorimeter transitions from the barrel to the endcaps and is not fully instrumented. Both the
 1241 transverse and longitudinal impact parameter cuts are the same for all electron selections.

1242 The electron particle identification uses a multivariate likelihood technique (LH) [71] detailed
 1243 in Section 3.2.4.3. Preselected electrons must pass the loosest LH workingpoint `LooseLH` with
 1244 an additional requirement that there be a reconstructed track hit in the first layer of the pixel
 1245 detector (a so-called *B*-layer hit). The LH requirement for the loose and signal electrons the tightness
 1246 of the identification using `MediumLH` and `TightLH`, respectively. As for isolation, the `Gradient`
 1247 workingpoint is required for signal electrons only. The loose electrons must fail one or both of the
 1248 signal identification and isolation requirements.

1249 **5.2.1.3 Jet candidate selection**

1250 The final objects that need to pass selection are jets. Jets are clustered using the anti- k_t algo-
 1251 rithm [72] within a radius of $\Delta R = 0.4$. The jets are then calibrated using E_T - and η -dependent
 1252 correction factors that are trained using MC simulations [73]. These calibrated jets are then re-
 1253 quired to have $p_T > 30$ GeV if they lie in the forward regions of the detector ($2.4 < |\eta| < 4.5$) and
 1254 $p_T > 25$ GeV in the central region ($|\eta| \leq 2.4$). In order to suppress pileup jets, the so-called jet-
 1255 vertex-tagger (JVT) discriminant associates a jet with the primary interaction vertex [74]; central
 1256 jets with $p_T > 60$ GeV are required to pass the `Medium` JVT workingpoint, which corresponds to
 1257 an average efficiency of over 92%. Finally, the jets are required to be separated by selected prompt
 1258 leptons by at least $\Delta R(j, l) > 0.3$.

1259 **5.2.1.4 Treatment of overlapping objects**

1260 In the event that one or more objects are reconstructed very close to each other, there is the
 1261 possibility for double-counting if both originated from the same object. The procedure by which
 1262 this ambiguity is resolved is called *overlap removal* (OR). The standard ATLAS recommendation
 1263 for OR is implemented in this analysis [75, 76] and is summarized in Table 5.6.

Electron preselection	
Momentum cut	$p_T > 6$ GeV
Angular acceptance	$ \eta < 2.47$
Longitudinal impact parameter	$ z_0 \times \sin \theta < 0.5$ mm
Transverse impact parameter	$d_0/\sigma_{d_0} < 5$
Particle identification	LooseLH + B-layer hit

Electron signal selection	
Momentum cut	$p_T > 27$ GeV
Angular acceptance	$ \eta < 2.47$, excluding $1.37 \leq \eta \leq 1.52$
Longitudinal impact parameter	$ z_0 \times \sin \theta < 0.5$ mm
Transverse impact parameter	$d_0/\sigma_{d_0} < 5$
Particle identification	TightLH
Particle isolation	Gradient

Electron loose selection	
Momentum cut	$p_T > 20$ GeV
Angular acceptance	$ \eta < 2.47$, excluding $1.37 \leq \eta \leq 1.52$
Longitudinal impact parameter	$ z_0 \times \sin \theta < 0.5$ mm
Transverse impact parameter	$d_0/\sigma_{d_0} < 5$
Particle identification	MediumLH
Fail signal identification and/or isolation cuts	

Table 5.4: Electron selection criteria. All electrons are required to pass the preselection (top), and then either the signal (middle) or loose (bottom) criteria is applied to the preselected electrons.

Jet selection	
Momentum cut	$p_T > 30$ GeV for $2.4 < \eta < 4.5$ $p_T > 60$ GeV for $ \eta < 2.4$
JVT cut	Medium
Jet-lepton separation	$\Delta R(j, l) > 0.3$

Table 5.5: Jet selection criteria. All jets are required to pass the above selection in order to be used in the analysis.

1264 Since electrons leave a shower in the EM calorimeter, every electron has a jet associated with
 1265 it. Therefore, any jets close to an electron (within $\Delta R(e, j) < 0.2$) are rejected due to the high
 1266 probability that they are the same object. On the other hand, when jets and electrons overlap
 1267 within a large radius of $0.2 < \Delta R(e, j) < 0.4$, it is likely that the electron and jet both are part of
 1268 a heavy-flavor decay, and the electron is rejected.

1269 High energy muons can produce photons via bremsstrahlung radiation or collinear final state
 1270 radiation which results in a nearby energy deposit in the calorimeters. Non-prompt muons from
 1271 hadronic decays produce a similar signature; however, in this case the jet has a higher track multiplicity
 1272 in the ID. It is possible to address both cases by rejecting the jet when the ID track multiplicity
 1273 is less than three and otherwise rejecting the muon for jets and muons within $\Delta R(\mu, j) < 0.4$.

1274 In addition to the case above where muon bremsstrahlung results in a nearby reconstructed jet,
 1275 the ID track from the muon and the calorimeter energy deposit can lead to it being reconstructed
 1276 as an electron. In this case, if both a muon and an electron share a track in the ID, the muon is
 1277 kept and the electron is rejected, unless the muon is calorimeter-tagged¹², in which case the muon
 1278 is removed in favor of the electron.

Overlap	Check	Result (remove → keep)
Electron & Jet	$\Delta R(e, j) < 0.2$	Jet → electron
	$0.2 < \Delta R(e, j) < 0.4$	Electron → jet
Muon & Jet	$\Delta R(\mu, j) < 0.4$ and Jet N_{ID} tracks < 3	Jet → muon
	$\Delta R(\mu, j) < 0.4$ and Jet N_{ID} tracks ≥ 3	Muon → jet
Electron & Muon	Shared ID track	Electron → muon
	Shared ID track & muon is calo-tagged	Muon → electron

Table 5.6: Summary of the overlap removal procedure used in the analysis. If the criteria in the “check” column is met, in the “result” column, the object on the left of the arrow is removed in favor of the object on the right.

1279 5.2.2 Signal event selection

1280 After the objects have been selected, cuts are applied on a per-event level to select $W^\pm W^\pm jj$ signal
 1281 events. The event selection is summarized in Table 5.8 and is detailed in this section. It includes
 1282 the results of an optimization performed using a multidimensional grid scan.

1283 The initial event selection begins by choosing events that pass one or more of the trigger re-
 1284 quirements listed in Table 5.7. At least one signal lepton is “matched” to a passed trigger in order

¹²A calorimeter-tagged (CT) muon is a muon that is identified by matching an ID track to a calorimeter energy deposit. CT muons have relatively low reconstruction efficiency compared to those measured by the MS, but can be used to recover acceptance in regions of the detector where the MS does not have full coverage [70].

1285 to ensure that it was indeed a signal lepton that fired the trigger. A collection of *event cleaning*
 1286 cuts must also be passed in order to remove events collected during periods in which one or more
 1287 components of the detector was not operating optimally. Finally, the events are required to contain
 1288 at least one interaction vertex. An event can have multiple reconstructed vertices from additional
 1289 proton-proton collisions that occurred in the same bunch crossing. In this case, the *primary vertex*
 1290 is determined by choosing the vertex with the largest sum of the p_T^2 of its associated tracks.

	2015 data	2016 data
Electrons	$p_T > 24$ GeV and Medium ID	$p_T > 26$ GeV and Tight ID and Loose isolation
	$p_T > 60$ GeV and Medium ID	$p_T > 60$ GeV and Medium ID
	$p_T > 120$ GeV and Loose ID	$p_T > 140$ GeV and Loose ID
Muons	$p_T > 20$ GeV and Loose isolation $p_T > 50$ GeV	$p_T > 26$ GeV and Medium isolation $p_T > 50$ GeV

Table 5.7: Summary of trigger requirements for electrons and muons for $\sqrt{s} = 13$ TeV data collected in 2015 and 2016. At least one of the triggers must be satisfied.

1291 Events are then required to contain exactly two signal leptons with the same electric charge.
 1292 The dilepton pair must have a combined invariant mass of $m_{ll} \geq 20$ GeV in order to suppress low
 1293 mass Drell-Yan backgrounds. Two additional selections are applied to events in the ee -channel:
 1294 both electrons are required to have $|\eta| < 1.37$ with an invariant mass at least 15 GeV away from
 1295 the Z -boson mass to reduce events where one electron is reconstructed with the wrong charge (this
 1296 background will be discussed in more detail in Section 5.3 TODO: Replace with proper subsection
 1297 once it's written). To suppress backgrounds from events with more than two leptons, events with
 1298 more than two leptons passing the preselection are vetoed.

1299 Missing transverse energy (E_T^{miss}) represents any particles that escape the detector without
 1300 being measured, such as neutrinos, and is defined as the magnitude of the vector sum of transverse
 1301 momenta of all reconstructed objects. It can be difficult to calculate accurately, as it involves
 1302 measurements from all subsystems within the detector, and it is sensitive to any corrections that
 1303 may be applied to the reconstructed physics objects [77]. These corrections, including the momentum
 1304 smearing for muons, energy scale and smearing for electrons, and jet calibrations, are propagated
 1305 to the E_T^{miss} calculation. Events are required to contain $E_T^{\text{miss}} > 30$ GeV in order to account for the
 1306 two neutrinos from the W boson decays.

1307 At least two jets are required. The leading and subleading jets must have $p_T > 65$ GeV and
 1308 $p_T > 35$ GeV, respectively, and are referred to as the *tagging jets*. Events are vetoed if they contain
 1309 one or more jets that have been tagged as a b -jet to suppress backgrounds from heavy flavor decays

1310 (especially top quark events). The b -tagging algorithm used by ATLAS is a boosted decision tree
 1311 (BDT) called MV2c10, and this analysis uses a workingpoint with 85% efficiency [78].

1312 Finally, cuts are applied on the VBS signature outlined in Section 5.0.3. The tagging jets are
 1313 required to have a dijet invariant mass $m_{jj} > 200$ GeV and be separated in rapidity by $|\Delta y_{jj}| > 2.0$.
 1314 This preferentially selects the VBS EWK events over the QCD-produced $W^\pm W^\pm jj$ events.

Event selection	
Event preselection	Pass at least one trigger with a matched lepton Pass event cleaning At least one reconstructed vertex
Lepton selection	Exactly two leptons passing signal selection Both signal leptons with the same electric charge $ \eta < 1.37$ and $ M_{ee} - M_Z > 15$ GeV (ee -channel only) Veto events with more than two preselected leptons
Missing transverse energy	$E_T^{\text{miss}} \geq 30$ GeV
Jet selection	At least two jets Leading jet $p_T > 65$ GeV Subleading jet $p_T > 35$ GeV $m_{jj} > 200$ GeV $N_{b\text{-jet}} = 0$ $ \Delta y_{jj} > 2.0$

Table 5.8: The signal event selection.

1315 5.3 Background estimations

1316 The major sources of background events are summarized in Section 5.0.4, and the methods used to
 1317 estimate them are detailed in this section. Prompt backgrounds from ZZ and $t\bar{t}V$ are estimated
 1318 directly from MC simulations. The shape of the WZ and $V\gamma$ backgrounds are taken from MC, and
 1319 the predicted yeilds are normalized to the data predictions in dedicated control regions, as outlined
 1320 in Sections 5.3.1 and 5.3.2, respectively. Opposite sign events with a charge misidentified electron
 1321 are estimated by a data-driven background method which is summarized in Section 5.3.3. Finally, a
 1322 *fake factor* method is used to estimate the contributions from non-prompt backgrounds and is the
 1323 subject of Section 5.3.4.

1324 5.3.1 Estimation of the WZ background

1325 The dominant background involving prompt leptons comes from $WZ + \text{jets}$ events. The contribution
 1326 is estimated from MC simulation and normalized to data in a control region enriched in WZ events

defined by the same event selection as Table 5.8 for the signal region, with the following changes applied to increase the purity of the WZ process:

- The third lepton veto is inverted, requiring a third lepton with $p_T > 15$ GeV
- Two of the leptons must make a same-flavor opposite-sign pair. If more than one pair exists, the one with m_{ll} closest to the Z boson mass is chosen.
- The trilepton invariant mass is required to be $m_{lll} > 106$ GeV to reduce contributions from $Z\gamma$ and $Z+jets$

Once the event yields in the control region are calculated, they are propagated to the final signal region fit, detailed in Section 5.4.1, in a single bin combining all the lepton channels. The systematic uncertainties of the WZ background are also calculated at this time. The event yields for the WZ control region are listed in Table 5.9, and distributions of the leading lepton p_T and η as well as trilepton invariant mass m_{lll} are found in Figures 5.10 and 5.9, respectively.

Event yields in the WZ control region	
WZ	197.9 ± 1.4
ZZ	14.1 ± 0.3
Triboson	1.26 ± 0.1
top	10.8 ± 1.1
$Z\gamma$	3.1 ± 1.1
$Z+jets$	2.5 ± 1.4
Total prediction	229.7 ± 2.5
Data	201 ± 14.2

Table 5.9: Event yields in the WZ control region before normalization. All lepton flavor channels are combined.

5.3.2 Estimation of the $V\gamma$ background

Events from $V\gamma$ processes can pass selection if the photon converts into an e^+e^- pair and one of the electrons passes the selection criteria. The background is estimated from MC simulations which are then scaled by a normalization factor calculated from a control region enriched in $Z(\mu^+\mu^-)\gamma$ events. This control region selects two opposite-sign muons and an additional electron that is assumed to come from the photon conversion. The full event selection is detailed in Table 5.10.

The $Z\gamma$ MC samples available do not cover the full range of p_T^γ and $\Delta R(\gamma, l)$; thus, additional Drell-Yan samples ($Z+jets$) are used to fill out the phase space. Overlap between the two samples




Figure 5.9: Leading lepton p_T (left) and η (right) distributions in the WZ control region before normalization. All lepton channels are combined.




Figure 5.10: Trilepton invariant mass m_{lll} distribution in the WZ control region before normalization. All lepton channels are combined.

$V\gamma$ control region
Exactly two muons with $p_T > 27$ GeV and $p_T > 20$ GeV
Exactly one additional electron with $p_T > 15$ GeV
Remove overlap between $Z+jets$ and $Z\gamma$
Di-muon + photon invariant mass $75 < M_{\mu\mu\gamma} < 100$ GeV
$E_T^{\text{miss}} < 30$ GeV

Table 5.10: Selection criteria for the $V\gamma$ control region.

1347 are removed based to avoid double counting. Events with final state photons at truth level are
 1348 checked to ensure that the photon did not originate from a hadronic decay. Cuts on $p_T^\gamma > 10$ GeV
 1349 and $\Delta R(\gamma, l) > 0.1$ are then applied at generator level, and $Z\gamma$ events that fail and $Z+jets$ events
 1350 that pass this additional selection are removed.

1351 The normalization factor is calculated directly from the event yields in the $V\gamma$ control region
 1352 rather than in the signal fit, as is done for the WZ background. The event yields are listed in
 1353 Table 5.11, and the normalization factor is determined to be 1.77. No MC events from $Z\gamma$ processes
 1354 survive the full event selection; thus, the scaling is only applied to the $W\gamma$ background in the signal
 1355 region. A systematic uncertainty of 44% is assigned to the background based off of the uncertainties
 1356 in the calculation of the normalization factor.

Event yields in the $V\gamma$ control region	
$Z\gamma$	24.6 ± 3.3
$Z+jets$	3.0 ± 1.5
diboson + triboson	6.7 ± 0.3
top	1.5 ± 0.5
Total prediction	35.8 ± 3.7
Data	57 ± 7.6

Table 5.11: Event yields in the $V\gamma$ control region. The $V\gamma$ scale factor of 1.77 is calculated by scaling up the $Z\gamma$ and $Z+jets$ backgrounds to account for the difference between the data and predicted total background.

1357 5.3.3 Estimation of backgrounds from charge misidentification

1358 If an electron's charge is mis-reconstructed, it can lead to a real, opposite-sign lepton pair passing
 1359 the same-sign requirement in the event selection. There are two primary reasons this can occur:

- 1360 1. An electron emits a photon via bremsstrahlung which then converts into an electron-positron
 1361 pair, and the conversion track with the wrong electric charge is matched to the original electron.
 1362 This is the dominant process leading to charge flip, and it is highly dependent on the electron
 1363 η due to the different amount of detector material the electron passes through.
- 1364 2. The curvature of the electron's track is mismeasured, resulting in the wrong charge being
 1365 assigned. This process is dependent on the momentum of the electron, as its track becomes
 1366 more straight as the momentum of the electron increases.

1367 In order to estimate this background, the rate at which an electron's charge is misidentified is
 1368 calculated from $Z \rightarrow e^+e^-$ MC simulation. It is known that the MC does not perfectly model




Figure 5.11: Charge misidentification rates for electrons as a function of $|\eta|$ and p_{T} . Rates are calculated from $Z \rightarrow e^+e^-$ MC after applying scale factors to approximate the charge mis-ID rates in data.

the material effects leading to charge flip; as a result, scale factors are applied to the MC in order for it to better reflect the real performance. These scale factors are obtained from the ratio of charge mis-ID rates in data and uncorrected MC in [47] following the method outlined in [79]. Once the scale factors are applied, the charge misidentification rate ε can be extracted by comparing the electron's reconstructed charge with the charge of its truth particle:

$$\varepsilon(\eta, p_{\text{T}}) = \frac{N_{\text{wrong charge}}}{N_{\text{prompt electrons}}} \quad (5.1)$$

The charge mis-ID rate is calculated in bins of electron $|\eta|$ and p_{T} and varies from below 0.1% in the

central region of the detector up to 8% in the forward regions for high p_{T} (above 90 GeV) electrons.

A two-dimensional plot of ε can be found in Figure 5.11.

Given the charge flip rate $\varepsilon(\eta, p_{\text{T}})$, the rate at which an electron has its charge correctly reconstructed is $(1 - \varepsilon)$. Thus there are three possible combinations of charge identification, assuming a two-electron event:

- 1380 1. Both electrons are reconstructed correctly: $(1 - \varepsilon)^2$
- 1381 2. Both electrons are mis-reconstructed: ε^2
- 1382 3. Only one electron is mis-reconstructed: $2\varepsilon(1 - \varepsilon)$

1383 In order to estimate the size of the background from charge misidentification, opposite-sign events
 1384 are selected using the default event selection for a given signal or control region with the same-sign
 1385 requirement inverted. These events are then weighted by the probability for one of the electrons to
 1386 be reconstructed with the wrong charge:

$$\omega = \frac{\varepsilon_1(1 - \varepsilon_2) + \varepsilon_2(1 - \varepsilon_1)}{(1 - \varepsilon_1)(1 - \varepsilon_2) + \varepsilon_1\varepsilon_2} \quad (5.2)$$

1387 where the subscripts 1 and 2 refer to the leading and subleading electrons, respectively, and ε_i is a
 1388 function of the η and p_T of the i^{th} electron. In the case of an event with only one electron and one
 1389 muon, Equation 5.2 simplifies:

$$\omega = \frac{\varepsilon}{1 - \varepsilon} \quad (5.3)$$

1390 This method assumes that there is little contamination from fake electrons in the opposite-sign
 1391 sample, and this has been verified with MC simulation.

1392 Additionally, charge-flipped electrons tend to be reconstructed with lower energy when compared
 1393 to electrons with the correct charge. This is due to energy loss from the material interactions that
 1394 can cause the charge to be misidentified. A correction factor is calculated from MC simulations,
 1395 comparing the p_T of the truth electron to its reconstructed counterpart:

$$\alpha = \frac{\left(\frac{p_T^{\text{reco}}}{p_T^{\text{truth}}} - 1\right)_{\text{correct charge}}}{\left(\frac{p_T^{\text{reco}}}{p_T^{\text{truth}}} - 1\right)_{\text{wrong charge}}} \quad (5.4)$$

1396 The correction is then applied to the p_T of the charge-flipped electron via

$$p_T = p_T^0 / (1 + \alpha) + dE \quad (5.5)$$

1397 where p_T^0 is the uncorrected p_T of the electron and dE is a gaussian smearing factor centered at
 1398 zero with a width related to the energy resolution. Since which electron is misreconstructed is never
 1399 determined in this method, in the case of a two-electron event, the energy correction is applied
 1400 randomly to one of the two electrons based on the probabilities for them to be charge-flipped. This
 1401 also determines the overall sign of the event; the charge of the electron that does not receive the
 1402 correction is taken to be the charge for both.

1403 Systematic uncertainties on the charge mis-ID rates are calculated by generating two additional
 1404 sets of rates with the uncertainties on the scale factors varied up and down. The size of the esti-
 1405 mated charge flip background without the energy correction applied is also taken as a systematic
 1406 uncertainty. These systematic uncertainties are estimated to be approximately $\pm 15\%$.




Figure 5.12: Dilepton invariant mass distribution m_{ll} for the ee channel in the same-sign inclusive VR.

1407 5.3.3.1 Validation of the charge misidentification estimate

1408 The performance of the charge misidentification estimation is tested in the same-sign inclusive
1409 validation region (VR), defined in Table 5.12. For ee events, the mass of the dilepton pair is required
1410 to lie within 15 GeV of the Z boson mass to increase the purity of the charge flip background.
1411 $t\bar{t}$ production, which can contribute to both the charge mis-ID and fake lepton backgrounds, is
1412 suppressed by the b -jet veto. The di-electron invariant mass is shown in Figure 5.12, and distributions
1413 of the leading and subleading electron p_T in the ee -channel are shown in Figure 5.13 with the Z
1414 mass cut inverted. Agreement between data and prediction is seen within the total statistical and
1415 systematic uncertainties in the VR.

Same-sign inclusive VR
Exactly 2 same-sign signal leptons
$p_T > 27$ GeV for both leptons
$m_{ll} > 20$ GeV
$ m_{ee} - m_Z > 15$ GeV ($e^\pm e^\pm$ -channel only)
$N_{b\text{-jet}} = 0$

Table 5.12: Selection criteria for the same-sign inclusive validation region.

1416 5.3.4 Estimation of non-prompt backgrounds with the fake factor method

1417 Events with one prompt lepton produced in association with hadronic jets can pass the event selection
1418 if a jet is misidentified as a charged lepton or if a non-prompt lepton from the decay of a heavy
1419 flavor particle (such as b - and c -hadrons) passes the signal lepton criteria. These misidentified jets




Figure 5.13: p_T distributions for the leading (left) and subleading (right) electron for the ee channel in the same-sign inclusive VR. In these plots, the cut requiring m_{ee} to fall within the Z mass window has been inverted in order to test the modelling away from the Z peak.

and non-prompt leptons are collectively referred to as *fake leptons*, or simply *fakes*. The rate at which a fake lepton is misidentified is generally not modelled well enough by the MC to accurately estimate their contributions directly from simulation. Therefore, a data-driven technique called the *fake factor* is used to estimate the size and shape of background processes from fake leptons. In this analysis, a new modification to the fake factor is used involving the particle isolation variables; the method is outlined in the context of the *default* fake factor in Section 5.3.4.1, and the modified fake factor is outlined in Section 5.3.4.2.

5.3.4.1 Overview of the default fake factor method

The goal of the fake factor method is to measure the fake rate from real collision events in a region enriched in fake leptons and use it to estimate the size of the fake lepton background in a chosen signal or control region. This is done by creating two samples using different lepton definitions:

1. The *nominal* sample is made up of leptons passing the signal selection.
2. The *loose* sample is made up of leptons that fail the signal selection while still passing a loosened set of criteria. This sample is enriched in fake leptons and is orthogonal to the set of signal leptons.

Using the sets of nominal and loose leptons, a fake factor f can be calculated in a region enriched in processes that are prone to producing fake leptons:

$$f = \frac{N_{\text{nominal}}}{N_{\text{loose}}} \quad (5.6)$$

1437 Since the fake rate is not expected to be constant over the entire phase space, the fake factor can
 1438 be divided into bins:

$$f(b) = \frac{N_{\text{nominal}}(b)}{N_{\text{loose}}(b)} \quad (5.7)$$

1439 where b represents the bin number. In this analysis, the fake factor is binned in lepton p_T .

1440 In order to estimate the fake background contribution in a given signal or control region, the
 1441 fake factor is applied to a second control region with a selection identical to the region of interest
 1442 with one of the leptons required to satisfy the loose criteria. The region for which the background
 1443 is estimated contains two nominal leptons and is referred to as *nominal+nominal* (*NN*), and the
 1444 associated control region where the fake factor is applied contains one nominal and one loose lepton
 1445 and is referred to as *nominal+loose* (*NL*). The fake background in a *NN* region can then be
 1446 calculated as:

$$N_{NN}^{\text{fake bkg.}} = \sum_b f(b) N_{NL}(b) \quad (5.8)$$

1447 Backgrounds containing two prompt leptons can also enter the *NL* region if one of the leptons
 1448 passes the nominal selection and the other passes the loose selection. Since the fake factor method
 1449 estimates the fake background by scaling the amount of non-prompt events in the *NL* region, if these
 1450 prompt contributions are not removed, they will be included in the scaling and the background
 1451 will be overpredicted. The final estimate of the fake background becomes:

$$N_{NN}^{\text{fake bkg.}} = \sum_b f(b) (N_{NL}(b) - N_{NL}^{\text{prompt}}(b)) \quad (5.9)$$

1452 5.3.4.2 The fake factor with p_T^{cone}

1453 When a jet produces a non-prompt lepton, that lepton only carries a fraction of the underlying jet's
 1454 total momentum. Due to the isolation cut applied to the nominal leptons, they typically carry a
 1455 much larger percentage of the underlying jet momentum¹³ than the loose leptons (which are allowed
 1456 to fail this criteria).

1457 This discrepancy in the underlying jet momentum fraction can cause problems in the calculation
 1458 of the fake factor f . Consider the case where two separate events have jets of identical momentum,
 1459 but one produces a non-prompt lepton that passes the nominal selection, and the other produces a
 1460 non-prompt lepton that passes the loose selection. The loose lepton on average will have lower p_T
 1461 than the nominal lepton despite both originating from jets with the same momentum. This can be

¹³Since the isolation variables are a measure of detector activity around the lepton, if other nearby particles carried a significant portion of the jet's momentum, the lepton would likely fail this cut.

1462 seen explicitly when comparing the p_T of a muon to its associated truth jet:

$$\Delta p_T(\mu, j) = \frac{p_T(j) - p_T(\mu)}{p_T(j) + p_T(\mu)} \quad (5.10)$$

1463 Since muons are not included in the jet reconstruction algorithm, Δp_T approximates the momentum
1464 of the muon compared to the rest of the jet. For muons that carry more than 50% of the jet's
1465 momentum, Δp_T will be negative and vice-versa. The Δp_T distributions for nominal and loose
1466 muons in $t\bar{t}$ MC events is shown Figure 5.14, where a 50 GeV jet on average corresponds to a
1467 35 GeV nominal muon and a 20 GeV loose muon¹⁴.




Figure 5.14: Δp_T distributions for nominal (blue) and loose (red) muons in simulated $t\bar{t}$ events. Each muon has been matched to a truth-level jet. Both distributions are normalized to unit area.

1468 Since the default fake factor defined in Equation 5.7 is binned in lepton p_T , within a given bin,
1469 the underlying jet p_T spectrum can differ substantially between the numerator and the denominator.
1470 Additionally, these differences can vary depending on the process producing the non-prompt leptons
1471 or on the specific kinematic selections of the signal or control regions where the fake factor is applied.

1472 Fortunately, the majority of the jet momentum not carried by the non-prompt lepton (excluding
1473 neutrinos) can be recovered using isolation variables. A track-based isolation is chosen, referred to
1474 as p_T^{cone} , and it contains the sum of the p_T of all particle tracks originating from the primary vertex
1475 within a cone of $\Delta R < 0.3$ around the lepton. Thus, the sample of loose leptons in the denominator
1476 of the fake factor calculation is binned in $p_T + p_T^{\text{cone}}$ rather than simply lepton p_T . Adding the

¹⁴To better illustrate the point, here the muon is added back into the jet p_T , and the corresponding muon p_T is obtained via $\Delta p_T(\mu, j) = \frac{(p_T(j) - p_T\mu) - p_T(\mu)}{(p_T(j) - p_T\mu) + p_T(\mu)} = \frac{p_T(j) - 2p_T(\mu)}{p_T(j)}$.

isolation cone greatly reduces the difference in the fraction of the underlying jet momentum carried by the nominal and loose leptons. To check this, a new Δp_T is calculated between a lepton and its matched truth jet, where the truth jet p_T has been corrected to include all muons within a cone of $\Delta R < 0.4$:

$$p_T(j) = p_T(j_{\text{truth}}) + \sum_{\Delta R < 0.4} p_T(\mu_{\text{truth}}) \quad (5.11)$$

The Δp_T distributions comparing p_T and $p_T + p_T^{\text{cone}}$ for nominal and loose leptons using the corrected jet p_T are found in Figure 5.15, and better agreement is seen between the numerator (nominal) and denominator (loose with $p_T + p_T^{\text{cone}}$) distributions.

The numerator remains binned in lepton p_T , due to the fact that it is meant to mirror the signal region as closely as possible, and the signal lepton selection does not use $p_T + p_T^{\text{cone}}$. The impact of this is expected to be negligible due to the p_T^{cone} isolation being small for signal leptons, as shown for muons in Figure 5.16. Finally, the fake factor f becomes:

$$f(b) = \frac{N_{\text{nominal}}(b(p_T))}{N_{\text{loose}}(b(p_T + p_T^{\text{cone}}))} \quad (5.12)$$

5.3.4.3 Application of the fake factor

The fake factor itself is measured from a sample data events passing a dijet selection requiring exactly one lepton (either passing the nominal or loose selections) and at least one jet. The leading jet must also be b -tagged and approximately back-to-back with the lepton in order to enhance non-prompt lepton contributions while reducing contributions from processes involving W and Z bosons. W boson events are further suppressed by requiring the sum of the E_T^{miss} and the transverse mass of the lepton and E_T^{miss} to be less than 50 GeV. The full event selection for the dijet region is summarized in Table 5.13.

The numerator sample is constructed from dijet events in which the lepton passes the nominal (signal) selection and is binned in the lepton p_T . Similarly, the denominator sample is made up of the remaining dijet events where the lepton passes the loose selection and is binned in the lepton $p_T + p_T^{\text{cone}}$. The nominal and loose leptons pass the signal selection¹⁵ and loose selection, respectively, defined earlier in Table 5.3 for muons and Table 5.4 for electrons. Backgrounds from $W + \text{jets}$, $Z + \text{jets}$, $t\bar{t}$, and single top processes are estimated from MC simulations requiring one lepton to be prompt using the truth information; these contributions are subtracted from the dijet data. The fake factor

¹⁵The $p_T > 27$ GeV cut in the signal lepton selection is dropped in favor of the $p_T > 15$ GeV requirement in the dijet selection.




Figure 5.15: Δp_T distributions for muons (top) and electrons (bottom) in simulated $t\bar{t}$ events. Each lepton has been matched to a truth-level jet, and that truth jet has had its p_T corrected to include all truth muons within a cone of $\Delta R < 0.4$. The nominal leptons are in black. Δp_T is calculated for the loose leptons using p_T (red) and $p_T + p_T^{\text{cone}}$ (blue).




Figure 5.16: Distributions of p_T^{cone}/p_T for nominal (black) and loose (red) muons in simulated $t\bar{t}$ events.

Dijet event selection	
Event preselection	
Exactly one lepton with $p_T > 15$ GeV	
$N_{\text{jet}} > 0$	
Leading jet is b -tagged	
$p_T^{\text{lead. jet}} > 25$ GeV	
$p_T^{\text{lead. jet}} > 30$ GeV if $ \eta_j > 2.5$	
$ \Delta\phi(l, \text{lead. jet}) > 2.8$	
$m_T(l, E_T^{\text{miss}}) + E_T^{\text{miss}} < 50$ GeV	

Table 5.13: Event selection for the dijet region used for calculating the fake factor. The selected lepton can pass either the nominal (signal) or loose selections. In the case of the nominal leptons, the $p_T > 27$ GeV requirement is replaced with $p_T > 15$ GeV.

is then calculated using Equation 5.12 for muons and for central and forward electrons separately.
 The muon fake factor is shown in Figure 5.17, and the two electron fake factors are shown in Figure 5.18. The numerical values of the fake factors, including their systematic uncertainties which will be discussed in Section 5.3.4.4, are listed in Table 5.14.

In order to properly account for the denominator being binned in $p_T + p_T^{\text{cone}}$, special care needs to be taken when estimating the fake background from the NL regions. For the purposes of the fake factor calculation, it is perhaps more intuitive to consider a loose *object* with $p_T = p_T + p_T^{\text{cone}}$ instead of simply a loose *lepton*, as the lepton and the underlying jet are treated as a whole with this method. When the lepton p_T cuts required by a particular signal or control region are applied to nominal and loose leptons, the cut is applied to the p_T of the nominal lepton and to the $p_T + p_T^{\text{cone}}$ of the loose object. Similarly, when looking up the fake factor weight for a given NL event, the value taken from the bin corresponding to the $p_T + p_T^{\text{cone}}$ of the loose object. Finally, when applying




Figure 5.17: The measured fake factor as a function of muon $p_T + p_T^{\text{cone}}$. The error bars represent the statistical uncertainty only.




Figure 5.18: The measured fake factor as a function of electron $p_T + p_T^{\text{cone}}$ in the central ($|\eta| < 1.37$, blue) and forward ($|\eta| > 1.37$, red) regions of the detector. The error bars represent the statistical uncertainty only.

1515 the weight to the event, $p_T + p_T^{\text{cone}}$ is assigned as the p_T of the loose object. Figure 5.19 contains a
 1516 graphical representation of this procedure.




Figure 5.19: Graphical representation of the fake factor application using $p_T + p_T^{\text{cone}}$. The value of $p_T + p_T^{\text{cone}}$ for the loose lepton is used to “look up” the fake factor weight which is then applied to the event. The loose lepton’s p_T becomes $p_T + p_T^{\text{cone}}$ for the purpose of the fake background estimation.

1517 Finally, it should be noted that the addition of p_T^{cone} to the loose object may cause the loose
 1518 leptons in the denominator sample to migrate into higher bins. This results in an overall decrease in
 1519 the number of loose objects in the lower $p_T + p_T^{\text{cone}}$ bins due to there not being additional leptons at
 1520 lower p_T to replace them. Since the fake factor is a ratio of the number of events in a bin, this effect
 1521 causes the first few bins of the fake factor to increase, as can be seen clearly in Figure 5.17. However,
 1522 the signal and control regions (and their corresponding NL regions) contain a $p_T > 27$ GeV cut that
 1523 prevents these migrations from negatively impacting the fake estimation.

1524 5.3.4.4 Systematic uncertainties

1525 Four sources of systematic uncertainty are considered: the dijet event selection, the prompt back-
 1526 ground subtraction, the jet flavor composition, and residual dependence on the underlying jet p_T
 1527 spectrum. In order to measure the impact of these systematics, new fake factors are computed
 1528 with each of the systematic variations and the differences from the nominal values are taken as the
 1529 uncertainty.

- 1530 1. In order to estimate uncertainties due to the dijet selection, the cut on $M_T + E_T^{\text{miss}}$ is varied
 1531 by ± 5 GeV, $\Delta\phi(l, j)$ by ± 0.1 , and the jet p_T cut by $+5$ GeV.
- 1532 2. To estimate the systematic uncertainty on the prompt background subtraction, the MC pre-
 1533 diction in a $W+\text{jets}$ control region is compared to data. The discrepancy between data and
 1534 MC is found to be approximately 10% [47]. Therefore, the prompt background used for the
 1535 subtraction is scaled up and down by $\pm 10\%$.
- 1536 3. The difference in the jet flavor composition between the dijet events and the events in the
 1537 NL regions can affect the accuracy of the fake background estimation. The dijet sample is
 1538 dominated by light jets, while the NL regions tend to be dominated by heavy flavor from $t\bar{t}$.
 1539 To account for this, the fake factor is computed with a b -jet veto.
- 1540 4. To measure any residual dependence on the underlying jet p_T spectrum, the leading jet p_T
 1541 distribution is reweighted to match the p_T spectrum of truth jets that produce fake leptons
 1542 in MC simulations. This results in an increase in the number of nominal and loose leptons at
 1543 high momentum [47].




Figure 5.20: Systematic variations in the fake factor as a function of muon $p_T + p_T^{\text{cone}}$. The individual fake factors obtained for each systematic variation are displayed with their statistical uncertainties.

1544 **5.3.4.5 Results of the fake factor**

- 1545 The fake background contribution in the signal region is estimated by applying the fake factors
 1546 to the equivalent NL region using Equation 5.9, where the fake factor used corresponds to the




Figure 5.21: Systematic variations in the fake factor as a function of electron $p_T + p_T^{\text{cone}}$ in the central ($|\eta| < 1.37$, top) and forward ($|\eta| > 1.37$, bottom) regions of the detector. The individual fake factors obtained for each systematic variation are displayed with their statistical uncertainties.

fake factor	p_T [15, 20]	p_T [20, 27]	p_T [27, 35]	p_T [35, 45]	p_T [45, 55]	p_T [55, 65]	p_T [65, 75]	p_T [75, 200]
nominal	0.649 ± 0.007	0.083 ± 0.002	0.024 ± 0.002	0.021 ± 0.003	0.044 ± 0.007	0.067 ± 0.018	0.160 ± 0.055	0.481 ± 0.088
MT+MET	0.649 ± 0.007	0.082 ± 0.002	0.082 ± 0.002	0.020 ± 0.003	0.045 ± 0.007	0.068 ± 0.018	0.207 ± 0.062	0.523 ± 0.086
$\Delta\phi(\ell, j)$	0.645 ± 0.008	0.083 ± 0.003	0.024 ± 0.002	0.021 ± 0.004	0.045 ± 0.008	0.064 ± 0.021	0.064 ± 0.058	0.438 ± 0.092
Jet p_T	0.650 ± 0.007	0.083 ± 0.002	0.024 ± 0.002	0.021 ± 0.003	0.045 ± 0.007	0.069 ± 0.018	0.159 ± 0.018	0.481 ± 0.088
$N_{b\text{-jet}} = 0$	0.724 ± 0.003	0.094 ± 0.001	0.035 ± 0.001	0.025 ± 0.002	0.022 ± 0.004	0.060 ± 0.015	0.026 ± 0.053	0.044 ± 0.134
Bkg. subtraction	0.648 ± 0.007	0.083 ± 0.002	0.024 ± 0.002	0.019 ± 0.003	0.037 ± 0.007	0.044 ± 0.019	0.096 ± 0.062	0.370 ± 0.082
Jet p_T Reweighting	0.539 ± 0.077	0.093 ± 0.007	0.025 ± 0.004	0.043 ± 0.019	0.063 ± 0.014	0.085 ± 0.025	0.141 ± 0.110	1.962 ± 0.492

(a) Fake factor for muons.

fake factor	p_T [20, 27]	p_T [27, 35]	p_T [35, 45]	p_T [45, 55]	p_T [55, 65]	p_T [65, 75]	p_T [75, 200]
nominal	0.491 ± 0.031	0.140 ± 0.020	0.111 ± 0.023	0.256 ± 0.049	0.546 ± 0.091	0.460 ± 0.140	0.939 ± 0.125
MT+MET	0.493 ± 0.030	0.138 ± 0.019	0.115 ± 0.022	0.261 ± 0.045	0.559 ± 0.084	0.656 ± 0.091	0.802 ± 0.016
$\Delta\phi(\ell, j)$	0.488 ± 0.032	0.137 ± 0.020	0.110 ± 0.025	0.283 ± 0.053	0.503 ± 0.097	0.351 ± 0.149	1.117 ± 0.255
Jet p_T	0.489 ± 0.035	0.134 ± 0.021	0.105 ± 0.025	0.224 ± 0.048	0.593 ± 0.093	0.356 ± 0.144	0.928 ± 0.177
$N_{b\text{-jet}} = 0$	0.506 ± 0.029	0.140 ± 0.018	0.111 ± 0.022	0.260 ± 0.046	0.545 ± 0.084	0.546 ± 0.120	0.882 ± 0.103
Bkg. subtraction	0.493 ± 0.032	0.146 ± 0.021	0.115 ± 0.024	0.259 ± 0.049	0.550 ± 0.091	0.460 ± 0.140	0.939 ± 0.125
$N_{b\text{-jet}} = 0$	0.387 ± 0.009	0.130 ± 0.008	0.321 ± 0.012	0.473 ± 0.015	0.716 ± 0.180	0.716 ± 0.180	0.716 ± 0.180
Bkg. subtraction	0.488 ± 0.031	0.138 ± 0.020	0.106 ± 0.023	0.248 ± 0.049	0.529 ± 0.092	0.434 ± 0.143	0.888 ± 0.115
Jet p_T Reweighting	0.493 ± 0.031	0.142 ± 0.020	0.115 ± 0.023	0.264 ± 0.049	0.563 ± 0.090	0.485 ± 0.136	0.989 ± 0.132

(b) Fake factor for central electrons ($|\eta| < 1.37$).

fake factor	p_T [20, 27]	p_T [27, 35]	p_T [35, 45]	p_T [45, 55]	p_T [55, 65]	p_T [65, 75]	p_T [75, 200]
nominal	0.487 ± 0.046	0.148 ± 0.031	0.253 ± 0.046	0.412 ± 0.071	0.556 ± 0.117	0.691 ± 0.117	1.340 ± 0.340
MT+MET	0.483 ± 0.045	0.152 ± 0.031	0.241 ± 0.043	0.443 ± 0.070	0.565 ± 0.106	0.668 ± 0.117	1.075 ± 0.189
$\Delta\phi(\ell, j)$	0.495 ± 0.047	0.156 ± 0.033	0.271 ± 0.052	0.364 ± 0.074	0.664 ± 0.107	0.749 ± 0.056	0.885 ± 0.084
Jet p_T	0.471 ± 0.051	0.158 ± 0.035	0.247 ± 0.051	0.474 ± 0.085	0.283 ± 0.107	0.546 ± 0.149	1.189 ± 0.266
$N_{b\text{-jet}} = 0$	0.478 ± 0.042	0.170 ± 0.031	0.274 ± 0.046	0.389 ± 0.066	0.645 ± 0.104	0.757 ± 0.102	1.319 ± 0.326
Bkg. subtraction	0.523 ± 0.048	0.149 ± 0.033	0.235 ± 0.045	0.429 ± 0.073	0.555 ± 0.117	0.691 ± 0.117	1.340 ± 0.340
$N_{b\text{-jet}} = 0$	0.525 ± 0.011	0.234 ± 0.013	0.644 ± 0.016	0.710 ± 0.014	0.274 ± 0.316	0.274 ± 0.316	0.274 ± 0.316
Bkg. subtraction	0.484 ± 0.046	0.146 ± 0.031	0.248 ± 0.046	0.406 ± 0.071	0.545 ± 0.118	0.676 ± 0.118	1.317 ± 0.337
Jet p_T Reweighting	0.328 ± 0.068	0.124 ± 0.048	0.297 ± 0.100	0.234 ± 0.061	0.680 ± 0.092	0.452 ± 0.138	2.385 ± 1.729

(c) Fake factor for forward electrons ($1.37 < |\eta|$).Table 5.14: Values of the fake factor in each p_T bin and for each individual systematic source.

flavor of the loose lepton in the event. As usual, the prompt background is subtracted from the NL events using MC simulation. Charge misidentification is handled using the same method as in Section 5.3.3, with an additional set of charge flip rates calculated for loose leptons. The fake background yields in the signal region are listed in Table 5.15. An overall uncertainty of 50% is assigned to the fake background estimation in $\mu^\pm\mu^\pm$ events, and between 40% to 90% for $e^\pm e^\pm$ and $\mu^\pm e^\pm$ events, including both statistical and systematic effects.

	estimated yield	f_e stat. up	f_e stat. dn	f_e syst. up	f_e syst. dn	f_μ stat. up	f_μ stat. dn	f_μ syst. up	f_μ syst. dn
$e^\pm e^\pm$	11.42 ± 3.13	1.69	-1.69	1.67	-5.56	—	—	—	—
$\mu^\pm\mu^\pm$	4.82 ± 0.77	—	—	—	—	0.65	-0.65	3.64	-0.61
$\mu^\pm e^\pm$	37.08 ± 5.16	4.90	-4.90	5.59	-14.34	1.39	-1.39	16.10	-1.98

Table 5.15: Estimated yields for the fake lepton background. The estimated yield is shown in the first column together with the statistical uncertainty followed by the systematic uncertainties from variations of the the fake factors within their statistical (stat.) and systematic (syst.) uncertainties. The labels f_e and f_μ indicate the fake factors for electrons and muons, respectively.

5.3.4.6 Validation of the fake factor

The accuracy of the fake factor method is tested in several validation regions, the most sensitive of which is the same-sign top fakes VR (SS top VR), defined in Table 5.16. This region inverts the signal region's b -jet veto to accept events with exactly one b -jet. Due to this requirement, the dominant source of events comes from the $t\bar{t}$ process where a b -jet fakes an isolated lepton. The distribution of the subleading lepton p_T in this VR is shown in Figure 5.22 for all lepton flavor combinations. There is good agreement between the data and the prediction, even when only taking into account the statistical uncertainty and not the large systematic uncertainties assigned to the fake estimation.

Same-sign inclusive VR
Exactly 2 same-sign signal leptons
$p_T > 27$ GeV for both leptons
$m_{ll} > 20$ GeV
$ m_{ee} - m_Z > 15$ GeV ($e^\pm e^\pm$ -channel only)
$N_{b\text{-jet}} = 1$
$N_{\text{jet}} \geq 2$
Leading jet $p_T > 65$ GeV
Subleading jet $p_T > 35$ GeV

Table 5.16: Selection criteria for the same-sign top fakes validation region.




Figure 5.22: Distributions of the subleading lepton p_T in the same-sign top fakes VR for $\mu^\pm\mu^\pm$ events (top right), $e^\pm e^\pm$ events (top left), $\mu^\pm e^\pm$ events (bottom left), and all events combined (bottom right). All errors are statistical only.

1562 5.3.5 Reduction of WZ background using custom overlap removal

1563 The dominant source of prompt background in this analysis comes from WZ events where both
 1564 bosons decay leptonically. Traditionally, the background is dealt with by imposing a veto on any
 1565 event with a third lepton passing some loose identification criteria (the so-called *trilepton veto*). In
 1566 the case of this analysis, if one or more leptons (in addition to the two signal leptons) passed the
 1567 preselection criteria, the event would be rejected. However, WZ events can still enter the signal
 1568 region if one of the leptons fails the veto selection or falls outside of the detector's acceptance.

1569 In order to understand the sources of WZ events that are not removed by the trilepton veto,
 1570 a study was performed on truth-level leptons¹⁶ on $W^\pm W^\pm jj$ and WZ MC samples. Events with
 1571 three truth leptons were selected, and each was matched to its reconstruction-level partner by finding

¹⁶Truth particles are the particles produced directly by the MC generator before being passed through the full detector simulation, at which point they are considered *reconstruction-level* (or *reco-level*) particles.

the closest $\Delta R(\text{truth, reco})$ and $\Delta p_{\text{T,truth,reco}}$ match. For events surviving the trilepton veto, the two signal leptons were removed, and the remaining leptons represent real leptons that failed to be selected for the veto. Between 40-50% of these leptons fell outside of the eta acceptance of the analysis (see Figure 5.23) and were unrecoverable. The second largest source of leptons failing the preselection was the OR, defined in Section 5.2.1.4. The standard OF procedure appeared to be too aggressive in removing leptons in favor of jets, causing many three lepton events to “lose” their third lepton and pass the trilepton veto. Therefore a *Custom OR* was investigated which would replace the standard OR in the preselection and allow for better WZ rejection by removing fewer third leptons.

TODO: Mention how the extra leptons in the $W^{\pm}W^{\pm}jj$ are background leptons since there are only 2 from the main decay




Figure 5.23: Pseudorapidity (η) distributions of truth muons (top) and electrons (bottom) for Sherpa $W^{\pm}W^{\pm}jj$ and WZ MC samples. The blue vertical lines represent the allowed η range for each lepton flavor. The numbers correspond to the number of raw MC events that fall within and outside of the allowed η range for each MC sample.

In order to construct a “custom” OR, a new quantity is defined between a lepton (l) and a nearby jet (j)

$$p_{\text{T,ratio}}(l, j) = \frac{p_{\text{T},l}}{p_{\text{T},j}} \quad (5.13)$$

which, along with $\Delta R(l, j)$, will allow for more third leptons to pass the preselection. The idea behind including $p_{\text{T,ratio}}$ is to be able to preferentially remove background leptons originating from jets (i.e. those that carry a low percentage of the total jet momentum) instead of removing *any* lepton near to jet. The distributions of $p_{\text{T,ratio}}$ and the associated efficiency curves for muons and electrons can be found in Figures 5.24 and 5.26, respectively, and the distributions for $\Delta R(\mu, j)$ for

1590 muons can be found in Figure 5.25. Since all electrons have an associated jet in the calorimeters,
 1591 the $\Delta R(e, j)$ variable is not a good quantity to use for this custom OR.




Figure 5.24: Distributions of $p_{T,\text{ratio}}(\mu, j)$ for EWK and QCD $W^\pm W^\pm jj$ signal (black) and WZ background (teal) for truth-matched third muons in events that pass the trilepton veto. Both distributions are normalized to unit area. The associated efficiency curves are on the right where efficiency is defined as the percentage of total events that would pass a cut on $p_{T,\text{ratio}}(\mu, j)$ at a given value on the x -axis.




Figure 5.25: Distributions of $\Delta R(\mu, j)$ for EWK and QCD $W^\pm W^\pm jj$ signal (black) and WZ background (teal) for truth-matched third muons in events that pass the trilepton veto. Both distributions are normalized to unit area. The associated efficiency curves are on the right where efficiency is defined as the percentage of total events that would pass a cut on $\Delta R(\mu, j)$ at a given value on the x -axis.

1592 A workingpoint for the Custom OR was chosen by requiring 90% signal retention for muons
 1593 and 90% background rejection for electrons. The cut on electrons was allowed to be much tighter
 1594 because the number of signal events with a third electron is considerably smaller than for muons.
 1595 It should be re-emphasized the signal events that are present in Figures 5.24–5.26 do not represent




Figure 5.26: Distributions of $p_{T,\text{ratio}}(e,j)$ for EWK and QCD $W^\pm W^\pm jj$ signal (black) and WZ background (teal) for truth-matched third electrons in events that pass the trilepton veto. Both distributions are normalized to unit area. The associated efficiency curves are on the right where efficiency is defined as the percentage of total events that would pass a cut on $p_{T,\text{ratio}}(e,j)$ at a given value on the x -axis.

the full set of signal events, but only those with a real third lepton (which must come from some source other than the signal $W^\pm W^\pm jj$ process). For muons, an or of $p_{T,\text{ratio}}(\mu,j)$ and $\Delta R(\mu,j)$ is used to maximize the third lepton acceptance due to correlations between the quantities, as shown in Figure 5.27; for electrons, only a cut on $p_{T,\text{ratio}}(e,j)$ is used. The Custom OR workingpoint is outlined in Table 5.17.

Custom OR Definition	
Muons	$p_{T,\text{ratio}}(\mu,j) > 0.40$ or $\Delta R(\mu,j) > 0.15$
Electrons	$p_{T,\text{ratio}}(e,j) > 0.18$

Table 5.17: Custom OR definition. Leptons must pass this selection in order to be counted for the trilepton veto.

Tests of the performance of the Custom OR yield promising results, with approximately 20% reduction in WZ background compared to less than 2% signal loss in the signal region. Unfortunately, due to differences between the primary analysis framework and the one used for testing, in practice the gains in WZ rejection are not nearly as substantial, and ultimately the Custom OR is not included in the final analysis. However, it is still a potentially useful tool for improving background rejection via lepton number vetoes in analyses with overly aggressive OR procedures.




Figure 5.27: Two-dimensional plots of $p_{T,\text{ratio}}(\mu, j)$ vs $\Delta R(\mu, j)$ for truth-matched third muons in events that pass the trilepton veto for EWK and QCD $W^\pm W^\pm jj$ signal (left) and WZ background (right). The blue overlay indicates the area in which the third leptons will pass the custom OR and result in the event failing the trilepton veto.

1607 5.4 Cross section measurement

1608 The $W^\pm W^\pm jj$ EWK cross section is extracted from the signal region using a maximum-likelihood
 1609 fit applied simultaneously to four m_{jj} bins in the signal region as well as to the low- m_{jj} and WZ
 1610 control regions. For the fit and cross section extraction, the signal region is defined as in Table 5.8
 1611 with the dijet invariant mass requirement raised to $m_{jj} > 500$ GeV. The low- m_{jj} region is defined
 1612 to mirror the signal region exactly with the dijet invariant mass inverted to $200 < m_{jj} < 500$ GeV,
 1613 and the WZ control region is defined previously in Section 5.3.1.

1614 The signal and low- m_{jj} regions are split into six channels based on the flavor and charge of
 1615 the dilepton pair: $\mu^+\mu^+$, $\mu^-\mu^-$, μ^+e^+ , μ^-e^- , e^+e^+ , and e^-e^- . This split by charge increases the
 1616 sensitivity of the measurement due to the W^+/W^- charge asymmetry at hadron colliders favoring
 1617 the production of W^+ bosons [80]. Since the signal events contain two W bosons, the signal strength
 1618 compared to charge-symmetric backgrounds is much greater in the $++$ channels for both charges
 1619 combined. The WZ control region is included in the fit as a single bin ($l^\pm l^\pm l^\pm$).

1620 The maximum likelihood fit and cross section extractions are outlined in Sections 5.4.1 and
 1621 5.4.3, respectively. The results of the cross section measurement and of the analysis as a whole are
 1622 presented in Section 5.6.

1623 5.4.1 Maximum likelihood fit

1624 TODO: This section is very similar to what is written in the support note... May need to put

1625 some work into flushing it out so it's not so close to copy-paste The number of predicted signal
 1626 events in each channel c and m_{jj} bin b can be calculated from the SM predicted signal cross section
 1627 $\sigma_{\text{theo}}^{\text{tot}}$, the total integrated luminosity \mathcal{L} , the signal acceptance \mathcal{A} , and the efficiency corrections $\mathcal{C}(\theta)$,
 1628 where θ represents the set of nuisance parameters that parameterize the effects of each systematic
 1629 uncertainty on the signal and background expectations. The acceptance and efficiency corrections
 1630 will be covered in more detail in Section 5.4.2.

$$N_{cb}^{\text{sig}}(\theta) = \sigma_{\text{theo}}^{\text{tot}} \mathcal{A}_b \mathcal{C}_b(\theta) \mathcal{L} \quad (5.14)$$

1631 A signal strength parameter μ is defined as the ratio of the measured cross section to the SM
 1632 predicted cross section. The expected number of events in a given channel and bin can then be
 1633 expressed as the sum of the estimated background ($N_{cb}^{\text{bkg}}(\theta)$) and the number of predicted signal
 1634 events scaled by μ :

$$\begin{aligned} N_{cb}^{\text{exp}}(\theta) &= \mu N_{cb}^{\text{sig}}(\theta) + N_{cb}^{\text{bkg}}(\theta) \\ &= \mu \sigma_{\text{theo}}^{\text{tot}} \mathcal{A}_b \mathcal{C}_b(\theta) \mathcal{L} + N_{cb}^{\text{bkg}}(\theta) \end{aligned} \quad (5.15)$$

1635 The nuisance parameters are constrained by Gaussian probability distribution functions, and
 1636 the normalization of the WZ background mentioned in Section 5.3.1 is included in the fit as a free
 1637 parameter. The expected yields for signal and background processes are adjusted by the set of
 1638 nuisance parameters within the constraints of the systematic uncertainties. The yields after the fit
 1639 correspond to the value that best matches the observed data.

1640 The number of events per channel and bin after the fit can be written as a sum of the predicted
 1641 event yields for each sample s :

$$\nu_{cb}(\phi, \theta, \gamma_{cb}) = \gamma_{cb} \sum_s [\eta_{cs}(\theta) \phi_{cs}(\theta) \lambda] h_{cbs}(\theta) \quad (5.16)$$

1642 In this equation, the fitted number of events in a given channel and bin is obtained by weighting
 1643 the histogram of predicted yields h_{cbs} by the product of a given luminosity λ and any normalization
 1644 factors ϕ_{cs} that may be given for each channel and sample. The input histogram and the normal-
 1645 ization factors may depend on the nuisance parameters θ taking into account sources of systematic
 1646 uncertainty. Uncertainties on the normalization factors $\eta_{cs}(\theta)$ are also included. Finally, bin-by-bin
 1647 scale factors γ_{cb} are included to parameterize the statistical uncertainties of the MC predictions.

1648 The binned likelihood function is given by a product of Gaussian functions for the luminosity
 1649 and for the background uncertainties and a product of Poisson functions for the number of observed
 1650 events in each bin and channel:

$$L(\mu|\theta) = \mathcal{G}(\mathcal{L}|\theta_{\mathcal{L}}, \sigma_{\mathcal{L}}) \cdot \prod_c \prod_b \mathcal{P}(N_{cb}^{\text{meas.}} | \nu_{cb}(\mu)) \prod_p \mathcal{G}(\theta_p^0 | \theta_p) \quad (5.17)$$

where \mathcal{G} and \mathcal{P} are the Gaussian and Poisson functions, respectively. As before, \mathcal{L} represents the integrated luminosity with uncertainty $\sigma_{\mathcal{L}}$ and associated nuisance parameter $\theta_{\mathcal{L}}$. The number of measured events in a given bin and channel is represented by $N_{cb}^{\text{meas.}}$, and $\nu_{cb}(\mu)$ is the predicted number of events defined in Equation 5.16 expressed as a function of the signal strength μ . Finally, the set of nuisance parameters θ and any auxiliary measurements used to constrain them θ^0 are multiplied for each parameter p .

The profile likelihood ratio is defined as

$$q_{\mu} = -2 \ln \frac{L(\mu, \hat{\theta}_{\mu})}{L(\hat{\mu}, \hat{\theta})} \quad (5.18)$$

with $\hat{\mu}$ and $\hat{\theta}$ as the unconditional maximum likelihood estimates and $\hat{\theta}$ as the conditional maximum likelihood estimate for a given value of μ . The fitted signal strength $\hat{\mu}$ is obtained by maximizing the likelihood function with respect to all parameters. The compatibility of the observed data with the background-only hypothesis can then be calculated by setting $\mu = 0$. Observation of the $W^{\pm}W^{\pm}jj$ EWK process is claimed if the data is found to be inconsistent with the background-only hypothesis by more than 5σ .

5.4.2 Definition of the fiducial volume

Before extracting the cross section, it is necessary to define the fiducial volume, or the phase space of measureable events. It is a subset of the total phase space defined by selection requirements designed to mirror those applied in the analysis as closely as possible. The selection criteria for the fiducial volume are listed in Table 5.18.

In MC simulations, the total phase space is generated, providing the total theoretical cross section $\sigma_{\text{theo}}^{\text{tot}}$ and the total number of signal events $\mathcal{N}_{\text{sig}}^{\text{tot}}$ ¹⁷. After applying the fiducial selection at truth level, the total number of signal events in the fiducial region $\mathcal{N}_{\text{sig}}^{\text{fid}}$ is obtained. An acceptance factor \mathcal{A} is used to represent the efficiency of events falling in the fiducial region at truth level:

$$\mathcal{A} = \frac{\mathcal{N}_{\text{sig}}^{\text{fid}}}{\mathcal{N}_{\text{sig}}^{\text{tot}}} \quad (5.19)$$

A correction factor \mathcal{C} is also necessary to translate from the truth level fiducial volume to the reconstruction level signal region and is defined in terms of the number of reconstruction level MC

¹⁷For the purpose of clarity, the number of events at truth level is represented by a script \mathcal{N} , and the number of events at reconstruction level uses a regular N .


Fiducial region selection	
Lepton selection	Two prompt leptons (e, μ) $p_T > 27$ GeV and $ \eta < 2.5$ for both leptons Both leptons with the same electric charge Dilepton invariant mass $m_{ll} > 20$ GeV Dilepton separation $\Delta R(ll) > 0.3$
Missing transverse energy	Two neutrino system with $p_T^{\nu\nu} > 30$ GeV
Jet selection	At least two jets Leading jet $p_T > 65$ GeV Subleading jet $p_T > 35$ GeV Leading and subleading jet $ \eta < 4.5$ Jet-lepton separation $\Delta R(l, j) > 0.3$ Dijet invariant mass $m_{jj} > 500$ GeV Dijet separation $\Delta y_{jj} > 2.0$

Table 5.18: Definition of the fiducial volume.

1675 events in the signal region $N_{\text{sig}, \text{MC}}^{\text{SR}}$:

$$\mathcal{C} = \frac{N_{\text{sig}, \text{MC}}^{\text{SR}}}{N_{\text{sig}}^{\text{fid}}} \quad (5.20)$$

1676 Since the fit is binned in m_{jj} , the acceptance and efficiency correction factors need to be as well.
 1677 Therefore, \mathcal{A}_i and \mathcal{C}_{ij} are written in terms of truth m_{jj} bins i and reconstruction m_{jj} bins j . A
 1678 graphical representation of these regions and the use of the acceptance and correction factors can
 1679 be seen in Figure 5.28.

Figure 5.28: Visual representation of the different kinematic regions relevant to the cross section measurement. The acceptance factor \mathcal{A} converts from the truth level total phase space to the truth level fiducial region, and the efficiency correction \mathcal{C} translates the fiducial region into the reconstruction level signal region.

1680 **5.4.3 Cross section extraction**

1681 The $W^\pm W^\pm jj$ EWK fiducial cross section is measured using the signal strength parameter μ that is
 1682 determined by the maximum likelihood fit. This parameter is dependent on the nuisance parameters
 1683 θ and can be written explicitly in terms of the measured and theoretical cross sections as:

$$\mu(\theta) = \frac{\sigma_{\text{meas}}^{\text{SR}}}{\sigma_{\text{theo}}^{\text{SR}}} \quad (5.21)$$

1684 In the simple case with only one bin, the equation for the total number of expected events in the
 1685 signal region first introduced in Equation 5.15 can be written as:

$$N_{\text{exp}}^{\text{SR}}(\theta) = \mu(\theta) \cdot \sigma_{\text{theo}}^{\text{tot}} \cdot \mathcal{L} \cdot \mathcal{A} \cdot \mathcal{C}(\theta) + N_{\text{bkg}}^{\text{SR}}(\theta) \quad (5.22)$$

1686 with the non-binned versions of \mathcal{A} and \mathcal{C} defined in Equations 5.19 and 5.20, respectively.

1687 If the measured fiducial cross section is written as:

$$\sigma_{\text{meas}}^{\text{fid}} = \mu \cdot \mathcal{A} \cdot \sigma_{\text{theo}}^{\text{tot}} \quad (5.23)$$

1688 then Equation 5.22 can be rearranged to read:

$$\sigma_{\text{meas}}^{\text{fid}} = \frac{N_{\text{exp}}^{\text{SR}}(\theta) - N_{\text{bkg}}^{\text{SR}}(\theta)}{\mathcal{L} \cdot \mathcal{C}(\theta)} \quad (5.24)$$

1689 The measured fiducial cross section can finally be rewritten in terms of $\hat{\mu}$, which is the best estimator
 1690 of the signal strength as extracted from the fit:

$$\begin{aligned} \sigma_{\text{meas}}^{\text{fid}} &= \hat{\mu}(\theta) \cdot \sigma_{\text{theo}}^{\text{tot}} \cdot \mathcal{A} \\ &= \hat{\mu}(\theta) \cdot \sigma_{\text{theo}}^{\text{fid}} \end{aligned} \quad (5.25)$$

1691 In practice, however, the cross section is not extracted from a single bin, and Equation 5.22
 1692 becomes for a single channel in truth and reconstruction level m_{jj} bins i and j , respectively:

$$N_{\text{exp}}^{\text{SR}}(\theta) = \mu(\theta) \cdot \sigma_{\text{theo}}^{\text{tot}} \cdot \mathcal{L} \cdot \sum_i \mathcal{A}_i \cdot \sum_j \mathcal{C}_{ij} + \sum_j N_{\text{bkg},j}^{\text{SR}}(\theta) \quad (5.26)$$

1693 where now the binned versions of \mathcal{A}_i and \mathcal{C}_{ij} are used. This equation can be extended to include all
 1694 the analysis channels by increasing the number of bins i and j . Additionally, it can be shown that
 1695 Equation 5.25 holds for this more complex case as well [47], provided care is taken to ensure that
 1696 all the uncertainties are handled properly.

1697 **5.5 Summary of uncertainties**

1698 Systematic uncertainties enter the final fit as nuisance parameters which can impact the estimated
1699 signal and background yields and the shapes of the m_{jj} distributions. These uncertainties can arise
1700 from the experimental methods or from the theoretical calculations used in the analysis. This section
1701 summarizes the systematic uncertainties; the experimental uncertainties are detailed in Section 5.5.1,
1702 and the theoretical uncertainties are covered in Section 5.5.2. The impacts of the systematic uncer-
1703 tainties on the final cross section measurement are summarized in Table 5.19.

Source	Impact [%]
Reconstruction	± 4.0
Electrons	± 0.5
Muons	± 1.2
Jets and E_T^{miss}	± 2.8
b -tagging	± 2.0
Pileup	± 1.5
Background	± 5.0
Misid. leptons	± 3.9
Charge misrec.	± 0.3
WZ	± 1.3
$W^\pm W^\pm jj$ QCD	± 2.8
Other	± 0.8
Signal	± 3.6
Interference	± 1.0
EW Corrections	± 1.3
Shower, Scale, PDF & α_s	± 3.2
Total	± 7.4

Table 5.19: Impact of various systematic effects on the fiducial cross section measurement. The impact of a given source of uncertainty is computed by performing the fit with the corresponding nuisance parameter varied up or down by one standard deviation from its nominal value.

1704 **5.5.1 Experimental uncertainties**

1705 Experimental uncertainties include detector effects as well as uncertainties on the background es-
1706 timation methods. Sources of systematic uncertainty on the measurement of physics objects are
1707 listed in Table 5.20, grouped by the relevant object type. For backgrounds estimated from MC
1708 simulations, variations in these sources of uncertainty are propagated through the analysis to obtain
1709 the corresponding uncertainties on the event yields. Additional experimental uncertainties include
1710 the integrated luminosity, the photon conversion rate from Section 5.3.2, and the data driven charge

1711 misidentification and fake lepton background estimations from Sections 5.3.3 and 5.3.4.5, respectively.
1712

1713 The largest sources of experimental uncertainty on the MC estimations come from the jet-related
1714 uncertainties and the b -tagging efficiency, while the largest uncertainty on the background estimation
1715 comes from the fake factor. The effects of the uncertainties on the $W^\pm W^\pm jj$ EWK signal and the
1716 dominant MC estimated background, WZ , are listed in Tables 5.21 and 5.22, respectively. Since
1717 the overall contributions from other processes estimated with MC are small, the uncertainties on
1718 these backgrounds have a lesser impact on the final measurement; these tables can be found in
1719 Appendix B.1.

Experimental uncertainties	
Electrons	Energy resolution
	Energy scale
	Identification efficiency
	Isolation efficiency
	Reconstruction efficiency
	Trigger efficiency
Muons	Energy scale
	Identification efficiency
	Inner detector track resolution
	Muon spectrometer resolution
	Trigger efficiency
E_T^{miss}	Resolution
	Scale
Jets	Energy resolution
	Energy scale
	JVT cut efficiency
	b -tagging efficiency
	Jets from pileup

Table 5.20: List of sources of experimental uncertainties on the reconstruction of physics objects.

$W^\pm W^\pm jj$ EWK	$e^\pm e^\pm$ % Yield	$\mu^\pm e^\pm$ % Yield	$\mu^\pm \mu^\pm$ % Yield
Jet-related Uncertainties	2.28	2.22	2.28
b -tagging efficiency	1.81	1.76	1.74
Pile-up	0.48	0.97	2.42
Trigger efficiency	0.02	0.08	0.47
Lepton reconstruction/ID	1.45	1.14	1.83
MET reconstruction	0.26	0.17	0.21

Table 5.21: Impact of experimental uncertainties for the $W^\pm W^\pm jj$ EWK processes in all channels.

WZ	$e^\pm e^\pm$ % Yield	$\mu^\pm e^\pm$ % Yield	$\mu^\pm \mu^\pm$ % Yield
Jet-related Uncertainties	9.58	5.03	8.45
b-tagging efficiency	2.49	2.23	2.40
Pile-up	2.99	3.49	3.33
Trigger efficiency	0.03	0.09	0.43
Lepton reconstruction/ID	1.52	1.24	3.07
MET reconstruction	0.93	0.79	1.63

Table 5.22: Impact of experimental uncertainties for the WZ process in all channels.1720 **5.5.2 Theoretical uncertainties**

1721 It is also necessary to consider uncertainties on the theoretical predictions in the fiducial region. They
 1722 include the choice of PDF set, the value of the strong coupling constant α_s , the renormalization
 1723 scale μ_R , the factorization scale μ_F , and the parton showering. The size of these uncertainties are
 1724 measured by generating new samples with variations in a chosen parameters and comparing them
 1725 to samples using the nominal choice of the parameter. Internal variations on the PDF sets or using
 1726 a different set entirely results in a relative uncertainty of up to 2.25% on the nominal sample. The
 1727 impact from varying α_s is very small, on the order of < 0.01%. The factorization and renormalization
 1728 scales are independently varied between 0.5-2.0 from their nominal values of 1.0. This results in
 1729 relative uncertainties on the prediction of up to 15%. Finally, varying the parameters in the parton
 1730 showering results in up to 8% uncertainty.

1731 **5.5.2.1 Uncertainties from EWK-QCD interference**

1732 As mentioned in Section 5.0.2, $W^\pm W^\pm jj$ production consists of both EWK processes. The two
 1733 production modes cannot be naively separated due to cross terms in the matrix element calculation.
 1734 These cross terms are referred to as *interference* terms. Since the $W^\pm W^\pm jj$ EWK production is
 1735 the focus of the analysis, and the signal region is designed to preferentially select those events, it is
 1736 important to measure the size of the EWK-QCD interference contributions.

1737 The interference effects are estimated using the `MadGraph` MC generator, as it has a feature that
 1738 allows direct modelling of the interference term. This allows four samples to be generated:

- 1739 1. Inclusive: All available diagrams are used in the matrix element calculation
- 1740 2. EWK only: Only EWK diagrams ($\mathcal{O}(\alpha_{\text{EWK}}) = 4$) are used
- 1741 3. QCD only: Only QCD diagrams ($\mathcal{O}(\alpha_s) = 2 \otimes \mathcal{O}(\alpha_{\text{EWK}}) = 2$) are used

1742 4. Interference: Only the interference terms are used

1743 A minimal set of generator level cuts, listed in Table 5.23, is applied in order to avoid biasing the
 1744 sample towards either production mode. The cross sections for each of the four channels can be
 1745 found in Table 5.24. The size of the interference is found to be approximately 6% of the total cross
 1746 section and is taken as a systematic uncertainty.

Generator level cuts
$\Delta\eta_{jj} < 10$
Jet $p_T > 20$ GeV
$M_{jj} > 10$ GeV

Table 5.23: The set of generator level cuts used for generating the interference samples with **MadGraph**.

Sample	σ (fb)
Inclusive	3.646 ± 0.0012
EWK only	2.132 ± 0.0005
QCD only	1.371 ± 0.0008
Interference	0.227 ± 0.0002

Table 5.24: Cross sections for each different $W^\pm W^\pm jj$ production mode (inclusive, EWK only, QCD only, and interference only) generated using **MadGraph**. The cross sections are calculated using a minimal set of generator level cuts from events where the W decays to a muon.

1747 5.6 Results

1748 After running the full analysis chain, the event yields in the signal region, low- m_{jj} control region,
 1749 and WZ control region as well as associated nuisance parameters representing the uncertainties are
 1750 passed to the maximum likelihood fit. From this fit, the normalization factor for the WZ control
 1751 region μ_{WZ} and the signal strength parameter in the signal region μ_{obs} are determined, and the
 1752 predicted yields in each input bin have been shifted according to the process detailed in Section 5.4.1.

1753 The WZ normalization factor is measured to be:

$$\mu_{WZ} = 0.88^{+0.07}_{-0.07}(\text{stat})^{+0.31}_{-0.21}(\text{theory})^{+0.22}_{-0.11}(\text{sys}) \quad (5.27)$$

1754 and is constrained primarily by the number of data events in the WZ control region. The observed
 1755 signal strength of $W^\pm W^\pm jj$ EWK production, defined in Equation 5.21, is extracted from the fit
 1756 and measured with respect to the prediction of the **SHERPA v2.2.2** MC generator:

$$\mu_{\text{obs}} = 1.45^{+0.25}_{-0.24}(\text{stat})^{+0.06}_{-0.08}(\text{theory})^{+0.27}_{-0.22}(\text{sys}) \quad (5.28)$$

1757 This corresponds to a rejection of the background-only hypothesis with a significance of 6.9σ .

1758 The observed number of data events are compared to the predicted signal and background yields
 1759 in the signal region in Table 5.25 before applying the fit and in Table 5.26 after the fit. The m_{jj}
 1760 distributions for data and prediction are shown in Figure 5.29 after the fit, and the fitted event
 1761 yields in the low- m_{jj} and WZ control regions are shown in Figure 5.30. Additional distributions
 1762 can be found in Appendix B.

	e^+e^+	e^-e^-	μ^+e^+	μ^-e^-	$\mu^+\mu^+$	$\mu^-\mu^-$	combined
WZ	1.9 ± 0.6	1.3 ± 0.4	14 ± 4	8.9 ± 2.6	5.5 ± 1.6	3.6 ± 1.1	35 ± 10
Non-prompt	4.1 ± 2.3	2.3 ± 1.7	9 ± 5	6 ± 4	0.57 ± 0.15	0.67 ± 0.25	23 ± 10
e/γ conversions	1.74 ± 0.29	1.8 ± 0.4	6.1 ± 1.6	3.7 ± 0.8	—	—	13.4 ± 2.5
Other prompt	0.17 ± 0.05	0.14 ± 0.04	0.90 ± 0.19	0.60 ± 0.14	0.36 ± 0.10	0.19 ± 0.05	2.4 ± 0.5
$W^\pm W^\pm jj$ QCD	0.38 ± 0.13	0.16 ± 0.05	3.0 ± 1.0	1.2 ± 0.4	1.8 ± 0.6	0.76 ± 0.25	7.3 ± 2.5
Expected background	8.2 ± 2.4	5.7 ± 1.8	33 ± 7	21 ± 5	8.2 ± 1.8	5.3 ± 1.2	81 ± 14
$W^\pm W^\pm jj$ EWK	3.8 ± 0.6	1.49 ± 0.22	16.5 ± 2.5	6.5 ± 1.0	9.1 ± 1.4	3.5 ± 0.5	41 ± 6
Data	10	4	44	28	25	11	122

Table 5.25: Table of the data and prediction event yields in the signal region before the fit. Numbers are shown for the six lepton flavor and charge channels and for all channels combined. Here the WZ background yields are normalized to the data in the WZ control region. The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.

	e^+e^+	e^-e^-	μ^+e^+	μ^-e^-	$\mu^+\mu^+$	$\mu^-\mu^-$	combined
WZ	1.49 ± 0.30	1.10 ± 0.26	11.7 ± 1.7	8.0 ± 1.3	5.0 ± 0.6	3.5 ± 0.6	31 ± 4
Non-prompt	2.2 ± 1.3	1.2 ± 0.7	5.7 ± 2.8	4.5 ± 1.8	0.57 ± 0.06	0.65 ± 0.14	15 ± 6
e/γ conversions	1.6 ± 0.4	1.6 ± 0.5	6.3 ± 1.6	4.3 ± 1.1	—	—	13.8 ± 2.9
Other prompt	0.16 ± 0.04	0.14 ± 0.04	0.90 ± 0.19	0.63 ± 0.13	0.39 ± 0.09	0.22 ± 0.05	2.4 ± 0.5
$W^\pm W^\pm jj$ QCD	0.35 ± 0.13	0.15 ± 0.05	2.9 ± 1.0	1.2 ± 0.4	1.8 ± 0.6	0.76 ± 0.25	7.2 ± 2.4
Expected background	5.8 ± 1.5	4.1 ± 1.1	27 ± 4	18.7 ± 2.6	7.7 ± 0.8	5.1 ± 0.6	69 ± 7
$W^\pm W^\pm jj$ EWK	5.6 ± 1.0	2.2 ± 0.4	24 ± 5	9.4 ± 1.8	13.5 ± 2.5	5.2 ± 1.0	60 ± 11
Data	10	4	44	28	25	11	122

Table 5.26: Table of the data and prediction event yields in the signal region after the fit. Numbers are shown for the six lepton flavor and charge channels and for all channels combined. The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.

1763 The last ingredient necessary to measure the $W^\pm W^\pm jj$ EWK cross section is the theory predicted
 1764 cross section in the fiducial region defined in Table 5.18. SHERPA v2.2.2 is used for the calculation,
 1765 and the cross section in the total generator phase space is 40.81 ± 0.05 fb, and the fiducial cross section
 1766 is 2.01 ± 0.02 fb. This corresponds to an acceptance factor of $\mathcal{A} = 0.0493 \pm 0.0002$. Uncertainties on
 1767 the simulation are estimated using variations of the scale, parton shower, and PDF set. The final




Figure 5.29: The dijet invariant mass m_{jj} distributions for data and predicted signal and background in the signal region after the fit. The shaded band represents the statistical and systematic uncertainties added in quadrature. Note that the bins have been scaled such that they represent the number of events per 100 GeV in m_{jj} . The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.




Figure 5.30: The event yields for data and predicted signal and background in the WZ and low- m_{jj} control regions after the fit. The shaded band represents the statistical and systematic uncertainties added in quadrature. The background estimations from the fake factor are included in the “Non-prompt” category, and backgrounds from $V\gamma$ production and electron charge misidentification are combined in the “ e/γ conversions” category. Finally, ZZ , VVV , and $t\bar{t}V$ backgrounds are combined in the “Other prompt” category.

1768 prediction used in the cross section measurement including uncertainties from Section 5.5.2 is:

$$\sigma_{\text{SHERPA}}^{\text{fid}} = 2.01 \pm 0.02(\text{stat}) \pm^{+0.29}_{-0.23}(\text{scale}) \pm^{+0.16}_{-0.02}(\text{parton shower}) \pm^{+0.05}_{-0.03}(\text{PDF}) \text{ fb} \quad (5.29)$$

1769 Combining this **SHERPA** prediction with the measured signal strength μ_{obs} from Equation 5.28,
 1770 the measured fiducial cross section $\sigma_{\text{meas}}^{\text{fid}}$ can be calculated using Equation 5.25:

$$\sigma_{\text{meas}}^{\text{fid}} = 2.91 \pm^{+0.51}_{-0.47}(\text{stat}) \pm^{+0.12}_{-0.16}(\text{theory}) \pm^{+0.24}_{-0.23}(\text{sys}) \pm^{+0.08}_{-0.06}(\text{luminosity}) \text{ fb} \quad (5.30)$$

1771 A plot comparing the measured fiducial cross section to two theoretical calculations is shown in
 1772 Figure 5.31. The measured value is compared to the **SHERPA v2.2.2** prediction used to calculate
 1773 μ_{obs} as well as to **POWHEG-BOX v2**. As mentioned in Section 5.1.1, this **POWHEG** sample does not
 1774 include the resonant triboson diagrams and is only used here for a visual comparison.




Figure 5.31: Comparison of the measured $W^\pm W^\pm jj$ EWK fiducial cross section with theoretical calculations from **SHERPA v2.2.2** and **POWHEG-BOX v2**. The light orange band represents the total experimental uncertainty on the measured value, and the dark orange hashed band is the statistical uncertainty. For the simulations, the light blue band represents the total theoretical uncertainty, and the dark blue hashed band are the scale uncertainties. The theory predictions do not include the interference between the EWK and QCD production.

1775

CHAPTER 6

1776

Prospects for same-sign WW at the High Luminosity LHC

1777

1778 On December 3, 2018, Run 2 of the LHC officially ended, and the collider was shut down to begin
1779 the first of two scheduled extended maintenance periods [81]. During these two long shutdowns,
1780 the Phase-I and Phase-II upgrades of the LHC and ATLAS will occur in order to prepare for the
1781 High-Luminosity LHC (HL-LHC) which is scheduled to begin operation in 2026 [82].

1782 The HL-LHC is planned to run at a center-of-mass energy of $\sqrt{s} = 14$ TeV with an instantaneous
1783 luminosity of $\mathcal{L} = 5 \times 10^{34}$ cm $^{-2}$ s $^{-1}$ with up to 200 collisions per beam-crossing. Over the course
1784 of operation, the HL-LHC is expected to collect a total integrated luminosity of $\mathcal{L} = 3000$ fb $^{-1}$ by
1785 2035 [83]. **TODO: Compare to current LHC numbers?**

1786 These run conditions will be much harsher than what ATLAS has experienced so far, and there
1787 are several upgrades planned for the detector to operate in the high luminosity environment. Most
1788 notably, the entire ID will be replaced with an all-silicon tracker which will extend the coverage from
1789 $|\eta| \leq 2.7$ up to $|\eta| \leq 4.0$. This will allow for reconstruction of charged particle tracks which can
1790 in turn be matched to clusters in the calorimeters for electron identification or forward jet tagging
1791 [84].

1792 The upgraded detector, the higher beam energy, and the increased volume of data to be collected
1793 provides the opportunity to measure rarer processes with a much higher precision than what was
1794 possible in Run 1. Same-sign $W^\pm W^\pm jj$ production, is one such process. With greater statistics,
1795 the accuracy of the cross section measurement can be improved over the 13 TeV analysis detailed in
1796 Chapter 5, and it also will allow for more detailed physics studies, such as measuring the polarization
1797 of the W bosons. A measurement of the longitudinal polarization of the scattered W bosons has

1798 not yet been possible, but it remains of great interest due to its sensitivity to electroweak symmetry
 1799 breaking [85]. The analysis detailed in this chapter is based off of the 2018 ATLAS HL-LHC
 1800 $W^\pm W^\pm jj$ prospects study [86] which is itself an extension of the 2017 ATLAS study [87]. **TODO:**
 1801 mention CMS's study + yellow report?

1802 6.0.1 Analysis Overview

1803 The experimental signature of interest is identical to the 13 TeV analysis: two prompt leptons (either
 1804 electrons or muons) with the same charge, missing transverse energy, and two high energy, forward
 1805 jets. These jets are again required to have a large angular separation and a high combined invariant
 1806 mass to preferentially select EWK- over QCD-produced $W^\pm W^\pm jj$ events.

1807 Background processes are again similar to the 13 TeV analysis and are summarized again here.
 1808 The dominant source of prompt background from $WZ+jets$ events where both bosons decay lepton-
 1809 ically. If the lepton from the Z -decay with opposite charge from the W falls outside of the detector
 1810 acceptance or is not identified, the remainder could appear to be a $W^\pm W^\pm jj$ signal event. To a
 1811 lesser extent, $ZZ+jets$ events can enter the signal region in much the same way provided two lep-
 1812 tons are “lost”. Other prompt sources include $t\bar{t}+V$ and multiple parton interactions, however
 1813 these processes do not contribute much. These prompt backgrounds are expected to contribute
 1814 less than in Run 2 with the addition of forward tracking in the upgraded ATLAS detector. Jets
 1815 mis-reconstructed as leptons or leptons from hadronic decays (such as $t\bar{t}$ and $W+jets$ production)
 1816 comprise the non-prompt lepton background. Lastly, events with two prompt, opposite-charge elec-
 1817 trons can appear as a same-sign event provided one of the electrons is mis-reconstructed as the
 1818 wrong charge.

1819 In this analysis, the EWK production of $W^\pm W^\pm jj$ is studied in the context of the planned
 1820 HL-LHC run conditions and upgraded ATLAS detector. An optimized event selection (referred to
 1821 as the *optimized selection*) is also explored in an effort to gain increased signal significance over
 1822 the *default selection*. The cross section of the inclusive EWK production is measured for both the
 1823 default and optimized selections, and the extraction of the longitudinal scattering significance is
 1824 measured with the optimized selection.

1825 6.1 Theoretical motivation

1826 The theoretical motivation for studying the ssWW process—and VBS in general—is detailed in Sec-
 1827 tion 5.0.2. Since it is specifically the scattering of *longitudinally polarized* vector bosons that violates

1828 unitarity without a SM Higgs boson, a direct measurement of this cross section will be very useful
 1829 for understanding how the Higgs unitarizes the process [85].

1830 6.1.1 Experimental sensitivity to longitudinal polarization

1831 **TODO:** mention that since there are so many polarization possibilities, a large integrated luminosity
 1832 is needed to measure just one of them individually There are three possible polarization states for
 1833 a massive vector boson: two transverse (+ or -) and one longitudinal (0). Therefore, in a system
 1834 with two W bosons, the overall polarization can be purely longitudinal (00), purely transverse (++,
 1835 --, and +-), or mixed (+0 and -0). The three combinations will be referred to as LL , TT , and
 1836 LT respectively.

1837 In order extract the longitudinal scattering component, it is necessary to find variables that
 1838 distinguish the LL from the TT and LT . Several were studied, and those with the best discriminating
 1839 power between the polarizations are the leading and subleading lepton p_T as well as the azimuthal
 1840 separation ($|\Delta\phi_{jj}|$) of the two VBS jets. The LL events prefer lower p_T for both signal leptons
 1841 (see Figure 6.1), which motivates keeping cuts on these quantities as low as possible in the event
 1842 selection. In the case of $|\Delta\phi_{jj}|$, the LL events generally had a larger dijet separation (see Figure 6.2),
 1843 and this variable is used in a binned likelihood fit to extract the longitudinal scattering significance.

1844 6.2 Monte Carlo samples

1845 As no real HL-LHC data will be available for many years, all signal and background processes
 1846 are modeled using MC simulations generated at $\sqrt{s} = 14$ TeV, with the event yields scaled to the
 1847 anticipated HL-LHC integrated luminosity of $\mathcal{L} = 3000 \text{ fb}^{-1}$. The MC samples used in the analysis
 1848 are generated at particle-level and have not been run through the typical full simulation of the
 1849 ATLAS detector. Instead, smearing functions derived from a **GEANT4** simulation of the upgraded
 1850 ATLAS detector are used to estimate detector effects such as momentum resolution. In addition,
 1851 pileup events are fully simulated. The MC samples used in this analysis are summarized in Table 6.1.

1852 The signal sample consists of both VBS and non-VBS electroweak (EWK) $W^\pm W^\pm jj$ production,
 1853 and it is simulated with the **Madgraph5_aMC@NLO** generator using the NNPDF3.0 PDF set and in-
 1854 terfaced with **PYTHIA v8** [88] for hadronization and parton showering. To study the longitudinal
 1855 polarization more directly, two additional **Madgraph5_aMC@NLO** $W^\pm W^\pm jj$ samples are used: one
 1856 containing only the longitudinal contribution (LL) and a second containing the transverse (TT) and
 1857 mixed (LT) contributions.




Figure 6.1: Comparison of the leading (top) and subleading (bottom) lepton p_T distributions for purely longitudinal (LL, black) and mixed polarization (LT+TT, cyan) $W^\pm W^\pm jj$ events.

1858 There are many other processes that can produce the same final state as the $W^\pm W^\pm jj$ and
 1859 must also be accounted for using MC simulations. WZ events are generated using **SHERPA v2.2.0**,
 1860 which includes up to one parton at NLO in the strong coupling constant and up to three addi-
 1861 tional partons at LO. Both EWK and QCD production are included in these samples. ZZ and
 1862 triboson VVV ($V = W, Z$) events are generated using **SHERPA v2.2.2** with up to two additional
 1863 partons in the final state. For the triboson backgrounds, the bosons can decay leptonically or
 1864 hadronically. $W+jets$ backgrounds are generated for electron, muon, and tau final states at LO
 1865 with **Madgraph5_aMC@NLO** and the **NNPDF3.0** set with showering from **PYTHIA v8**. $Z+jets$ events are
 1866 produced using **POWHEG-BOX v2** and the **CT10** PDF set interfaced with **PYTHIA v8**. Finally, $t\bar{t}$ and
 1867 single-top events are generated using **POWHEG-BOX** with showering from **PYTHIA v6**.




Figure 6.2: Comparison of the azimuthal dijet separation ($|\Delta\phi_{jj}|$) for purely longitudinal (LL, black) and mixed polarization (LT+TT, cyan) $W^{\pm}W^{\pm}jj$ events.

Process	Generator	Comments
$W^{\pm}W^{\pm}jj$ (EWK)	Madgraph5_aMC@NLO	Signal sample
$W^{\pm}W^{\pm}jj$ (QCD)	Madgraph5_aMC@NLO	
$W^{\pm}W^{\pm}jj$ (LL)	Madgraph5_aMC@NLO	Pure longitudinal polarization sample
$W^{\pm}W^{\pm}jj$ (TT+LT)	Madgraph5_aMC@NLO	Mixed and transverse polarization sample
Diboson	SHERPA v2.2.0	WZ events
	SHERPA v2.2.2	ZZ events
Triboson	SHERPA v2.2.2	
$W+jets$	Madgraph5_aMC@NLO	
$Z+jets$	POWHEG-BOX v2	
$t\bar{t}$	POWHEG-BOX	
Single top	POWHEG-BOS	

Table 6.1: Summary of MC samples used in the analysis.

1868 6.3 Background estimations

1869 In this analysis, all background contributions are estimated using MC simulations. Backgrounds such
 1870 as electron charge misidentification and fake electrons from jets (which are traditionally estimated
 1871 using data-driven techniques) are estimated using a set of parameterization functions applied to the
 1872 MC. These functions calculate the probability that an electron is assigned the wrong charge or a
 1873 jet is mis-reconstructed as an electron parameterized by the p_T and η of the electron or jet. The
 1874 probabilities are derived from studies on expected electron performance with the upgraded ATLAS
 1875 detector [89].

1876 Processes involving two W and Z bosons are grouped together as *diboson* backgrounds, with the

exception of $W^\pm W^\pm jj$ events produced via QCD interactions, which are kept separate. Similarly, all backgrounds with three vector bosons are combined and labeled as *triboson*. Any $W+jets$ or top events that pass selection and do not contain a fake electron, as well as any $Z+jets$ events without an electron identified as having its charge misidentified are combined as *other non-prompt* backgrounds.

6.3.1 Truth-based isolation

To properly calculate particle isolation, it requires information from several detector subsystems including tracking and calorimeter responses. Since the MC samples used in this analysis have not been run through a full detector simulation, it is not possible to construct the canonical isolation variables used in analyses. At truth-level, this is generally not a serious concern as p_T signal leptons tend to be well isolated to begin with. However, isolation is one of the most powerful tools for rejecting leptons from non-prompt sources such as top events, which are produced in association with additional nearby particles from b and c quark decays. In this analysis, with the absence of any sort of isolation requirement, contributions from top backgrounds (including single top, $t\bar{t}$ and $t\bar{t} + V$) are more than an order of magnitude higher than expected.

As a result, it is necessary to find one or more quantities that are comparable to the isolation information that is available in fully-simulated samples. Analogues to track- and calorimeter-based isolation variables are constructed by summing the momentum and energy, respectively, of stable truth particles with $p_T > 1$ GeV within a specified radius of each signal lepton. For the track-based isolation, only charged truth particles are used; both charged and neutral particles (excluding neutrinos) are included for the calorimeter-based isolation. Ultimately, a set of isolation cuts are chosen that are similar to those recommended by ATLAS for Run 2 analyses. The truth-based isolation requirements are listed in Table 6.2.

	Electron Isolation	Muon Isolation
Track-based isolation cone size	$\Delta R < 0.2$	$\Delta R < 0.3$
Track-based isolation requirement	$\sum p_T/p_T^e < 0.06$	$\sum p_T/p_T^\mu < 0.04$
Calorimeter-based isolation cone size	$\Delta R < 0.2$	$\Delta R < 0.2$
Calorimeter-based isolation requirement	$\sum E_T/p_T^e < 0.06$	$\sum E_T/p_T^\mu < 0.15$

Table 6.2: Truth-based isolation requirements for electrons and muons.

The truth-based isolation requirement reduces the top background by over 99%, and the percentage of the total background consisting of top events is reduced from 83% to 2%. Additional details on the truth-based isolation studies are presented in Appendix A.

¹⁹⁰² **6.4 Object and event selection**

¹⁹⁰³ **6.4.1 Object selection**

¹⁹⁰⁴ Electrons and muons are preselected to have $p_T > 7$ and 6 GeV, respectively, and $|\eta| \leq 4.0$. The
¹⁹⁰⁵ likelihood of a given lepton to pass the trigger and identification requirements is estimated by
¹⁹⁰⁶ calculating an efficiency dependent on the p_T and η of the lepton. The leptons are also required to
¹⁹⁰⁷ pass the isolation criteria detailed in Table 6.2. Jets that have been tagged as a fake electron by the
¹⁹⁰⁸ functions described in Section 6.3 are treated as electrons for the purpose of the object selection and
¹⁹⁰⁹ are subject to the same criteria. In order to be considered a signal lepton, an additional requirement
¹⁹¹⁰ of $p_T > 25$ GeV is applied on top of the preselection. The two highest p_T leptons passing this
¹⁹¹¹ selection are chosen to be the leading and subleading signal leptons.

¹⁹¹² Jets are clustered using the anti- k_t algorithm [72] from final-state particles within a radius of
¹⁹¹³ $\Delta R = 0.4$ (excluding muons and neutrinos). Jets are required to have $p_T > 30$ GeV and lie within
¹⁹¹⁴ $|\eta| < 4.5$, with an additional cut of $p_T > 70$ GeV for jets above $|\eta| \geq 3.8$ in order to suppress jets
¹⁹¹⁵ from pileup interactions. Jets overlapping with a preselected electron within $\Delta R(e, j) < 0.05$ are
¹⁹¹⁶ removed in order to prevent double counting. The two highest p_T jets are defined as the leading
¹⁹¹⁷ and subleading *tag jets*.

¹⁹¹⁸ **6.4.2 Event selection**

¹⁹¹⁹ The default event selection is summarized in Table 6.3 and described here. Exactly two signal
¹⁹²⁰ leptons are required with the same electric charge and separated from each other by $\Delta R(l l) > 0.3$.
¹⁹²¹ In order to suppress contributions from Drell-Yan backgrounds, the two signal leptons must have
¹⁹²² an invariant mass m_{ll} greater than 20 GeV. Additionally, if both signal leptons are electrons, their
¹⁹²³ mass must be at least 10 GeV from the Z -boson mass in order to reduce background from Z -boson
¹⁹²⁴ decays¹⁸. The event is required to have at least 40 GeV of missing transverse energy (E_T^{miss}) to
¹⁹²⁵ account for the two neutrinos from the W decays. Events with additional preselected leptons are
¹⁹²⁶ vetoed, which greatly reduces WZ and ZZ backgrounds.

¹⁹²⁷ Each event must have at least two jets, and both tag jets are required to not overlap with the
¹⁹²⁸ signal leptons, and there is a veto on events with one or more b -jets. In order to preferentially select
¹⁹²⁹ EWK production, the tag jets are also required to have a large separation between them and a large

¹⁸The electron charge misidentification rate in the upgraded ATLAS detector is estimated to be high enough that contributions from $Z \rightarrow ee$ backgrounds are non-negligible.

¹⁹³⁰ invariant mass. Finally, a cut on the lepton centrality¹⁹, ζ , defined in Equation 6.1 enhances the
¹⁹³¹ EWK $W^\pm W^\pm jj$ signal.

$$\zeta = \min[\min(\eta_{\ell 1}, \eta_{\ell 2}) - \min(\eta_{j 1}, \eta_{j 2}), \max(\eta_{j 1}, \eta_{j 2}) - \max(\eta_{\ell 1}, \eta_{\ell 2})] \quad (6.1)$$

Selection requirement	Selection value
Lepton kinematics	$p_T > 25 \text{ GeV}$ $ \eta \leq 4.0$
Jet kinematics	$p_T > 30 \text{ GeV}$ for $ \eta \leq 4.5$ $p_T > 70 \text{ GeV}$ for $ \eta > 3.8$
Dilepton charge	Exactly two signal leptons with same charge
Dilepton separation	$\Delta R_{l,l} \geq 0.3$
Dilepton mass	$m_{ll} > 20 \text{ GeV}$
Z boson veto	$ m_{ee} - m_Z > 10 \text{ GeV}$ (ee -channel only)
E_T^{miss}	$E_T^{\text{miss}} > 40 \text{ GeV}$
Jet selection	At least two jets with $\Delta R_{l,j} > 0.3$
b jet veto	$N_{b\text{-jet}} = 0$
Dijet separation	$\Delta \eta_{jj} > 2.5$
Trilepton veto	No additional preselected leptons
Dijet mass	$m_{jj} > 500 \text{ GeV}$
Lepton-jet centrality	$\zeta > 0$

Table 6.3: Summary of the signal event selection.

¹⁹³² 6.5 Selection optimization

¹⁹³³ An upgraded detector along with an increase in center of mass energy and integrated luminosity
¹⁹³⁴ provides an opportunity to study whether the event selection can be optimized to improve the signal
¹⁹³⁵ to background ratio.

¹⁹³⁶ 6.5.1 Random grid search algorithm

¹⁹³⁷ The chosen method for optimizing the event selection is a cut-based algorithm known as the Random
¹⁹³⁸ Grid Search (RGS) [90]. Consider a simple case of two variables x and y chosen to differentiate signal
¹⁹³⁹ from background. In order to be considered a signal event, a given event would be required to pass
¹⁹⁴⁰ a set of selection criteria, called a *cut point*: $c = \{x > x_c, y > y_c\}$. A simple method to choose the
¹⁹⁴¹ optimal cut point (i.e. the “best” values of the cuts x_c and y_c) would be to construct an $n \times m$

¹⁹ ζ is a measurement of whether the two signal leptons lie between the two tagging jets in η , as is preferred by the VBS topology.

1942 rectangular grid in x and y consisting of points $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_m)$, as in Figure 6.3. One
 1943 can then choose a cut point $c_k = \{x > x_i, y > y_j\}$ that maximizes the signal significance as measured
 1944 by a chosen metric. This would be considered a *rectangular grid search*.

1945 While effective in principle, a rectangular grid search comes with two major drawbacks:

- 1946 1. The algorithm scales exponentially as the number of variables to be optimized increases, as
 1947 this is effectively increasing the dimensionality of the grid. In the simple case of a square grid
 1948 with N bins per variable v , the number of cut points to be evaluated grows as N^v .
- 1949 2. Signal and background samples are rarely evenly distributed over the entire grid, resulting
 1950 in many cut points being sub-optimal and evaluating them would be a waste of computing
 1951 resources.

1952 To combat these limitations, the RGS algorithm constructs a grid of cut points directly from
 1953 the signal sample itself. In the two-dimensional example, this means that the variables x_i and y_j
 1954 making up the cut point $c_k = \{x > x_i, y > y_j\}$ take their values directly from a given signal event.
 1955 This has the benefit of creating a *random grid* of cut points that is biased towards regions of high
 1956 signal concentration by construction. This reduces the need for exponentially increasing numbers of
 1957 cut points while ensuring that computing resources are not wasted in regions with few to no signal
 1958 events. An example of the the two-dimensional random grid is shown in Figure 6.4.




Figure 6.3: A visual representation of a rectangular grid search algorithm. The signal events are the blue triangles, and the red circles are the background events. **TODO: replace with own figure**

1959 Once the random grid of cut points is constructed, the optimal cut point can be chosen using any
 1960 number of metrics, such as signal to background ratio. For the purpose of the $W^\pm W^\pm jj$ upgrade




Figure 6.4: A visual representation of a random grid search algorithm. The signal events are the blue triangles, and the red circles are the background events. **TODO: replace with own figure**

1961 study, the optimal cut point is chosen to be the one that maximizes the signal significance Z as
 1962 defined in Equation 6.2 [91].

$$Z = \sqrt{2 \left[(s + b) \ln \left(\frac{s + b}{b_0} \right) + b_0 - s - b \right] + \frac{(b - b_0)^2}{\sigma_b^2}} \quad (6.2)$$

1963 where s and b are the number of signal and background events, respectively, σ_b is the total uncertainty
 1964 on the background, and b_0 is defined as:

$$b_0 = \frac{1}{2} \left(b - \sigma_b^2 + \sqrt{(b - \sigma_b^2)^2 + 4(s + b)\sigma_b^2} \right) \quad (6.3)$$

1965 In the case where the background is known precisely (i.e. $\sigma_b = 0$), Equation 6.2 simplifies to

$$Z = \sqrt{2 \left(b \left[(1 + s/b) \ln(1 + s/b) - s/b \right] \right)} \quad (6.4)$$

1966 which further reduces to the familiar $Z = s/\sqrt{b}$ for the case when $s \ll b$.

1967 6.5.2 Inputs to the optimization

1968 In order to train the RGS, signal and background samples are prepared from events passing the
 1969 event selection outlined in Table 6.3 up through the b -jet veto. The signal sample is chosen to be
 1970 the longitudinally polarized $W^\pm W^\pm jj$ EWK events, and the transverse and mixed polarizations
 1971 are treated as background along with $W^\pm W^\pm jj$ events from QCD interactions and the traditional
 1972 backgrounds listed in Section 6.3. Splitting the inclusive $W^\pm W^\pm jj$ EWK events by polarization

1973 allows the optimization to favor the longitudinally polarized events as much as possible, even though
 1974 they both contribute to the EWK signal.

1975 The following variables are chosen for optimization:

- 1976 • Leading lepton p_T
- 1977 • Dilepton invariant mass (m_{ll})
- 1978 • Leading and subleading jet p_T
- 1979 • Dijet invariant mass (m_{jj})
- 1980 • Lepton-jet centrality (ζ)

1981 Subleading lepton p_T is omitted as it is desirable to keep the cut value as low as possible due to
 1982 its sensitivity to the longitudinal polarization (as discussed in Section 6.1.1). Additionally, the dijet
 1983 separation $\Delta\eta_{jj}$ was included in the optimization originally, however it was dropped from the list due
 1984 to the cut value being motivated by differences between EWK and QCD produced $W^\pm W^\pm jj$ events.

1985 Two additional constraints were imposed when selecting the optimal cut point:

- 1986 1. At least 1000 signal events must survive in order to prevent the optimization from being too
 1987 aggressive and unnecessarily reducing signal statistics.
- 1988 2. The dijet invariant mass may only vary within a 50 GeV range of the default value (from
 1989 450 – 550 GeV) due to the cut being physically motivated by the VBS event topology (see
 1990 Section 5.0.3).

1991 Lastly, the signal significance is calculated without taking into account the uncertainty of the
 1992 background using Equation 6.4. This is due to the fact that the statistical uncertainties of the fake
 1993 electron and charge misidentification backgrounds are quite large, owing to poor MC statistics in a
 1994 few of the samples. If Equation 6.2 were used instead, the optimization will cut unreasonably hard
 1995 against these backgrounds. Since Monte Carlo statistics is not expected to be a limiting factor when
 1996 this analysis is performed at the HL-LHC, it is more realistic to simply ignore these large statistical
 1997 uncertainties for the purpose of the optimization.

1998 6.5.3 Results of the optimization

1999 Ultimately, the random grid is constructed from over 38,000 LL-polarized $W^\pm W^\pm jj$ events in the
 2000 six variables listed above. After applying the constraints, the optimal cut point reduces the total

background from 9900 to 2310 while reducing the signal from 3489 to 2958. This corresponds to an increase in signal significance from $Z = 33.26$ to $Z = 52.63$ as calculated by Equation 6.4. The updates to the event selection are listed in Table 6.4.

The large reduction in the background is primarily a result of the increase in the leading and subleading jet p_T from 30 GeV to 90 GeV and 45 GeV, respectively. As can be seen in Figure 6.7, this increase removes a significant portion of the backgrounds from jets faking electrons and charge mis-ID. Additionally, the loosening of the lepton-jet centrality cut ζ allows more signal events to survive the event selection (see Figure 6.9). Other changes to the event selection are minor and do not individually have a large impact on the signal or background yields.

The full event yields after optimization as well as the cross section measurement are detailed alongside those using the default selection in Section 6.6.

TODO: It's a bit awkward to reference the results of the default/optimized before they're properly presented. Maybe move the sections around? not sure...

Selection requirement	Selection value
Lepton kinematics	$p_T > 28$ GeV (leading lepton only)
Jet kinematics	$p_T > 90$ GeV (leading jet) $p_T > 45$ GeV (subleading jet)
Dilepton mass	$m_{ll} > 28$ GeV
Dijet mass	$m_{jj} > 520$ GeV
Lepton-jet centrality	$\zeta > -0.5$

Table 6.4: Updates to the $W^\pm W^\pm jj$ event selection criteria after optimization. Cuts not listed remain unchanged from the default selection in Table 6.3.

6.6 Results

6.6.1 Event yields

After applying the full event selection, the analysis is broken down into four channels based off of the flavor of the signal leptons: $\mu\mu$, ee , μe , and $e\mu$. The full signal and background event yields are shown in Table 6.5 for each channel separately and combined using the default event selection. 3489 EWK $W^\pm W^\pm jj$ events are expected compared to 9900 background events. The dominant sources of background are jets faking electrons followed by charge misidentification and diboson processes. Triboson events, QCD $W^\pm W^\pm jj$, and other non-prompt sources make up approximately 5% of the total background combined.




Figure 6.5: Leading lepton p_T distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram). **TODO: Move to appendix or omit**




Figure 6.6: Dilepton invariant mass distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram). **TODO: Move to appendix or omit**




Figure 6.7: Leading (top) and subleading (bottom) jet p_T distributions. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram).




Figure 6.8: Dijet invariant mass distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram). **TODO:** Move to appendix or omit




Figure 6.9: Lepton-jet centrality distribution. The default and optimized cuts are represented by the red and green dashed lines, respectively. The $W^\pm W^\pm jj$ EWK signal (black points) is normalized to the same area as the sum of the backgrounds (colored histogram).

	All channels	$\mu\mu$	ee	μe	$e\mu$
$W^\pm W^\pm jj$ (QCD)	206.4	91.1	22.8	38.4	54.1
Charge Misidentification	2300	0.0	2100	90	160
Jets faking electrons	5000	0.0	3400	1200	340
$WZ + ZZ$	2040	500	438	423	680
Tribosons	115	47	15.4	21.6	31.2
Other non-prompt	210	110	20	60	27
Total Background	9900	750	6000	1900	1290
Signal $W^\pm W^\pm jj$ (EWK)	3489	1435	432	679	944

Table 6.5: Signal and background event yields using the default event selection for an integrated luminosity of $\mathcal{L} = 3000 \text{ fb}^{-1}$. Events containing a fake or charge-flipped electron are removed from their respective sources and combined into a single entry each.

2023 The event yields for the optimized selection detailed in Section 6.5.3 are listed in Table 6.6. After
 2024 optimization, 2958 signal events and just 2310 background events are expected. Diboson events are
 2025 now the primary source of background, as the optimization greatly reduces the fake and charge
 2026 misidentification backgrounds. As discussed earlier, the increase in the leading and subleading jet
 2027 p_T cuts as well as the loosening of the centrality cut are most responsible for the changes in the
 2028 signal and background yields; distributions of these quantities using the default and the optimized
 2029 event selections can be found in Figures 6.10, 6.11, and 6.12, respectively.

	All channels	$\mu\mu$	ee	μe	$e\mu$
$W^\pm W^\pm jj$ (QCD)	168.7	74.6	19.7	32.2	42.2
Charge Misidentification	200	0.0	11	30	160
Jets faking electrons	460	0.0	130	260	70
$WZ + ZZ$	1286	322	289	271	404
Tribosons	76	30.1	9.6	15.1	21.6
Other non-prompt	120	29	16.6	50	19
Total Background	2310	455	480	660	710
Signal $W^\pm W^\pm jj$ (EWK)	2958	1228	380	589	761

Table 6.6: Signal and background event yields using the optimized event selection for an integrated luminosity of $\mathcal{L} = 3000 \text{ fb}^{-1}$. Events containing a fake or charge-flipped electron are removed from their respective sources and combined into a single entry each.

2030 It is important to note, however, that the MC sample used to estimate $Z + \text{jets}$ events suffers from
 2031 poor statistics which results in large per-event weights once scaled to $\mathcal{L} = 3000 \text{ fb}^{-1}$. This sample
 2032 contributes heavily to the fake and charge misidentification backgrounds, and a handful of these
 2033 events being cut out by the optimization contributes has a large effect on the dramatic reduction
 2034 of these backgrounds. As a result, these particular optimized results are likely overly optimistic.
 2035 However, given proper MC statistics, it is still expected that the optimization will outperform the




Figure 6.10: p_T distributions for the leading jet using the default (left) and optimized (right) event selections for all channels combined.




Figure 6.11: p_T distributions for the subleading jet using the default (left) and optimized (right) event selections for all channels combined.




Figure 6.12: p_T distributions for lepton-jet centrality ζ using the default (left) and optimized (right) event selections for all channels combined.

2036 default selection.

2037 6.6.2 Uncertainties

2038 TODO: Ask for details on how some of these uncertainties were calculated – specifically the fakes and
 2039 charge mis-ID The uncertainties considered for the analysis are summarized in Table 6.7. Values for
 2040 experimental systematics on the trigger efficiency, lepton and jet reconstruction, and flavor tagging
 2041 are taken directly from the 13 TeV analysis [45]. The rate uncertainties for the background processes
 2042 are halved from the 13 TeV values.

Source	Uncertainty (%)
$W^\pm W^\pm jj$ (EWK)	3
Luminosity	1
Trigger efficiency	0.5
Lepton reconstruction and identification	1.8
Jets	2.3
Flavor tagging	1.8
Jets faking electrons	20
Charge misidentification	25
$W^\pm W^\pm jj$ (QCD)	20
Top	15
Diboson	10
Triboson	15

Table 6.7: Summary of estimated experimental and rate uncertainties.

2043 6.6.3 Cross section measurement

2044 The cross section is calculated using the same method as in the 13 TeV analysis, detailed in Chap-
 2045 ter 5. TODO: update from chapter reference to subsection reference (once it's written)... Once
 2046 again, each of the four lepton flavor channels is further split by charge (i.e. $\mu\mu \rightarrow \mu^+\mu^+ + \mu^-\mu^-$),
 2047 as this increases the sensitivity of the analysis. Each channel's m_{jj} distribution is combined in a
 2048 profile likelihood fit to extract the EWK $W^\pm W^\pm jj$ production cross section. The expected cross
 2049 section calculated using the default event selection is:

$$\sigma_{W^\pm W^\pm jj}^{\text{expected}} = 16.89 \pm 0.36 \text{ (stat)} \pm 0.53 \text{ (theory)} \pm 0.84 \text{ (syst)} \text{ fb} \quad (6.5)$$

2050 The expected cross section calculated using the optimized event selection is:

$$\sigma_{W^\pm W^\pm jj}^{\text{expected}} = 16.94 \pm 0.36 \text{ (stat)} \pm 0.53 \text{ (theory)} \pm 0.78 \text{ (syst)} \text{ fb} \quad (6.6)$$

2051 The optimized selection should not change the measured value of the cross section, and indeed both
 2052 are consistent with within uncertainties. The systematic uncertainty is reduced by approximately 7%
 2053 with the optimized selection. Projections of the total uncertainty on the cross section as a function
 2054 of integrated luminosity made by [TODO: how was this made?](#) is shown in Figure 6.13. As the
 2055 integrated luminosity increases past $\mathcal{L} > 3000 \text{ fb}^{-1}$, the statistical uncertainty reduces faster than
 2056 the systematic uncertainties. However, the total uncertainty is expected to reduce by less than a
 2057 percent with increased luminosity past the planned 3000 fb^{-1} .




Figure 6.13: Projections of the statistical (black), theoretical (blue), systematic (yellow), and total (red) uncertainties on the measured cross section as a function of integrated luminosity using the optimized event selection.

2058 6.6.4 Longitudinal scattering significance

2059 [TODO: get some details on how this was all done...](#) The longitudinal scattering significance is
 2060 extracted from the $|\Delta\phi_{jj}|$ distribution using a simultaneous binned likelihood fit. In order to increase
 2061 sensitivity, the $|\Delta\phi_{jj}|$ distribution was split into two bins in m_{jj} , and an additional cut on the
 2062 pseudorapidity of the subleading lepton was applied ($|\eta| < 2.5$) to reduce background from fake and
 2063 charge misidentification. The $|\Delta\phi_{jj}|$ distributions used in the fit are shown in Figure 6.14. Due to
 2064 limited statistics, the four lepton flavor channels were not split by charge. The expected significance

2065 of the $W_L^\pm W_L^\pm jj$ process is 1.8σ with a precision of 47% on the measurement. Projections of the
2066 expected significance as a function of integrated luminosity is shown in Figure 6.15.




Figure 6.14: Dijet azimuthal separation ($|\Delta\phi_{jj}|$) for the low m_{jj} region ($520 < m_{jj} < 1100 \text{ GeV}$, top) and the high m_{jj} region ($m_{jj} > 1100 \text{ GeV}$, bottom). The purely longitudinal (LL, gray) is plotted separately from the mixed and transverse (LT+TT, cyan) polarizations.




Figure 6.15: Projections of the expected longitudinal scattering significance as a function of integrated luminosity when considering all sources of uncertainties (black) or only statistical uncertainties (red).

2067

CHAPTER 7

2068

Conclusion

2069 Here's where you wrap it up.

2070 **Looking Ahead**

2071

2072 Here's an example of how to have an "informal subsection".

APPENDIX A

Additional material on truth isolation

yields by type	all channels	$\mu\mu$	ee	μe	$e\mu$
signal	4011	1583.2	531.7	793.1	1103.1
ww qcd	252.6	105.8	30.4	48	68.4
charge flip	2528.4	0.0	2075.4	255.1	197.8
fakes	7135.4	0.0	4675.1	1904.3	555.9
diboson	2370.4	581.2	491.8	517.9	779.6
triboson	125.5	49.1	17.8	24.6	34.1
top	90150.5	26618	15301.6	25277.9	22953.1
z+jets	241.2	0.0	0.0	0.0	241.2
w+jets	31.4	3.9	7.6	13.2	6.7
total bkg	102803.9	27354	22592	28027.8	24830.1
signal	4011	1583.2	531.7	793.1	1103.1

Table A.1: Event yields prior to applying any form of truth-based isolation criteria.

yields by type	all channels	$\mu\mu$	ee	μe	$e\mu$
signal	3470.5	1427.3	428.8	675.8	938.7
ww qcd	205.8	90.8	22.7	38.3	54
charge flip	2398.3	0.0	2104.6	95.8	197.9
fakes	4309.7	0.0	3390.6	750.8	168.3
diboson	1552.4	311.3	355.6	346.8	538.7
triboson	115	46.8	15.4	21.6	31.2
top	156.9	42.3	14.8	76.6	23.3
z+jets	0.0	0.0	0.0	0.0	0.0
w+jets	0.3	0.0	0.0	0.3	0.0
total bkg	8738.1	491.3	5903.7	1329.8	1013.4
signal	3470.5	1427.3	428.8	675.8	938.7

Table A.2: Event yields after applying a test version of the truth-based isolation.

TODO: Add tables for tight vs loose working point, information on the necessity of TRUTH1++

APPENDIX B

Additional material on $W^\pm W^\pm jj$ measurement at $\sqrt{s} = 13$ TeV

2079 B.1 Impact of experimental uncertainty on MC background estimations

$W^\pm W^\pm jj$ QCD	ee % Yield	$e\mu$ % Yield	$\mu\mu$ % Yield
Jet-related Uncertainties	3.41	3.04	2.85
b-tagging efficiency	2.56	2.48	2.48
Pile-up	4.99	0.45	0.33
Trigger efficiency	0.02	0.08	0.41
Lepton reconstruction/ID	1.62	1.19	1.89
MET reconstruction	0.41	0.22	0.34

Table B.1: Experimental systematics uncertainties for the $W^\pm W^\pm jj$ QCD processes in all channels.

Triboson	ee % Yield	$e\mu$ % Yield	$\mu\mu$ % Yield
Jet-related Uncertainties	13.09	13.39	16.85
b-tagging efficiency	2.96	3.77	4.95
Pile-up	19.37	24.66	6.87
Trigger efficiency	0.02	0.07	0.47
Lepton reconstruction/ID	1.66	1.27	2.48
MET reconstruction	0.00	0.46	0.00

Table B.2: Experimental systematics uncertainties for triboson process in all channels.

$t\bar{t}V$	ee % Yield	$e\mu$ % Yield	$\mu\mu$ % Yield
Jet-related Uncertainties	17.65	11.97	14.27
b-tagging efficiency	15.02	9.04	13.83
Pile-up	8.73	10.69	4.18
Trigger efficiency	0.03	0.08	0.39
Lepton reconstruction/ID	2.57	3.27	2.66
MET reconstruction	1.75	4.16	1.62

Table B.3: Experimental systematics uncertainties for $t\bar{t}V$ processes in all channels.

$W\gamma$	ee % Yield	$e\mu$ % Yield	$\mu\mu$ % Yield
Jet-related Uncertainties	7.05	33.36	—
b-tagging efficiency	1.97	2.94	—
Pile-up	4.11	14.17	—
Trigger efficiency	0.01	0.14	—
Lepton reconstruction/ID	1.40	1.13	—
MET reconstruction	0.00	0.00	—

Table B.4: Experimental systematics uncertainties for the $W\gamma$ process in all channels.

$Z\gamma$	ee % Yield	$e\mu$ % Yield	$\mu\mu$ % Yield
Jet-related Uncertainties	16.22	370.44	—
b-tagging efficiency	1.08	3.10	—
Pile-up	12.57	11.51	—
Trigger efficiency	0.02	0.07	—
Lepton reconstruction/ID	1.26	22.01	—
MET reconstruction	0.00	0.00	—

Table B.5: Experimental systematics uncertainties for the $Z\gamma$ process in all channels.

ZZ	ee % Yield	$e\mu$ % Yield	$\mu\mu$ % Yield
Jet-related Uncertainties	15.71	15.76	35.18
b-tagging efficiency	2.23	2.35	2.89
Pile-up	1.22	3.20	4.58
Trigger efficiency	0.03	0.10	0.36
Lepton reconstruction/ID	3.59	3.10	5.70
MET reconstruction	4.84	3.26	3.24

Table B.6: Experimental systematics uncertainties for the ZZ process in all channels.

Bibliography

- 2082 [1] ATLAS Collaboration, M. Aaboud et al., *Measurement of the $W^\pm Z$ boson pair-production*
 2083 *cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector*, *Phys. Lett.* **B762**
 2084 (2016) 1–22, [arXiv:1606.04017 \[hep-ex\]](#). (document)
- 2085 [2] S. Weinberg, *The Quantum Theory of Fields, Volume 1: Foundations*. Cambridge University
 2086 Press, 1995. [2.1](#)
- 2087 [3] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*.
 2088 Addison-Wesley, 1995. [2.1](#)
- 2089 [4] R. P. Feynman, *Mathematical formulation of the quantum theory of electromagnetic*
 2090 *interaction*, *Phys. Rev.* **80** (1950) 440–457. [2.1](#)
- 2091 [5] M. Y. Han and Y. Nambu, *Three-triplet model with double SU(3) symmetry*, *Phys. Rev.* **139**
 2092 (1965) B1006–B1010. [2.1](#)
- 2093 [6] D. J. Gross and F. Wilczek, *Asymptotically Free Gauge Theories - I*, *Phys. Rev.* **D8** (1973)
 2094 3633–3652. [2.1](#)
- 2095 [7] D. J. Gross and F. Wilczek, *ASYMPTOTICALLY FREE GAUGE THEORIES. 2.*, *Phys.*
 2096 *Rev.* **D9** (1974) 980–993. [2.1](#)
- 2097 [8] H. D. Politzer, *Reliable Perturbative Results for Strong Interactions?*, *Phys. Rev. Lett.* **30**
 2098 (1973) 1346–1349. [,274(1973)]. [2.1](#)
- 2099 [9] A. Salam and J. C. Ward, *Weak and Electromagnetic Interactions*, *Nuovo Cimento* **11** (1959)
 2100 568–577. [2.1](#)
- 2101 [10] S. L. Glashow, *The Renormalizability of Vector Meson Interactions*, *Nucl. Phys.* **10** (1959)
 2102 107–117. [2.1](#)
- 2103 [11] ALEPH, DELPHI, L3, OPAL, SLD, LEP Electroweak Working Group, SLD Electroweak
 2104 Group, SLD Heavy Flavour Group Collaboration, S. Schael et al., *Precision electroweak*
 2105 *measurements on the Z resonance*, *Phys. Rept.* **427** (2006) 257–454, [arXiv:hep-ex/0509008](#)
 2106 [hep-ex]. [2.1](#)
- 2107 [12] F. Englert and R. Brout, *Broken symmetry and the mass of gauge vector mesons*, *Phys. Rev.*
 2108 *Lett.* **13** (1964) 321–323. [2.1](#)

- 2109 [13] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Physics Letters* **12**
2110 (1964) 132–133. 2.1
- 2111 [14] P. W. Higgs, *Broken symmetries and the masses of gauge bosons*, *Phys. Rev. Lett.* **13** (1964)
2112 508–509. 2.1
- 2113 [15] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model*
2114 *Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1,
2115 arXiv:1207.7214 [hep-ex]. 2.1, 5.0.2
- 2116 [16] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS*
2117 *experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30, arXiv:1207.7235 [hep-ex]. 2.1, 5.0.2
- 2118 [17] UA1 Collaboration, G. Arnison et al., *Experimental Observation of Isolated Large Transverse*
2119 *Energy Electrons with Associated Missing Energy at $s^{**}(1/2) = 540\text{-GeV}$* , *Phys. Lett. B* **122**
2120 (1983) 103–116. [,611(1983)]. 2.2
- 2121 [18] UA1 Collaboration, G. Arnison et al., *Experimental Observation of Lepton Pairs of Invariant*
2122 *Mass Around $95\text{-GeV}/c^{**2}$ at the CERN SPS Collider*, *Phys. Lett. B* **126** (1983) 398–410.
2123 [,7.55(1983)]. 2.2
- 2124 [19] Particle Data Group Collaboration, K. A. Olive et al., *Review of Particle Physics*, *Chin.*
2125 *Phys. C* **38** (2014) 090001. 2.2, 2.2
- 2126 [20] ATLAS Collaboration, G. Aad et al., *Evidence of $W\gamma\gamma$ Production in pp Collisions at*
2127 *$\sqrt{s} = 8\text{ TeV}$ and Limits on Anomalous Quartic Gauge Couplings with the ATLAS Detector*,
2128 *Phys. Rev. Lett.* **115** (2015) no. 3, 031802, arXiv:1503.03243 [hep-ex]. 2.3
- 2129 [21] ATLAS Collaboration, ATLAS Collaboration, *Evidence for Electroweak Production*
2130 *of $W^\pm W^\pm jj$ in pp Collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS Detector*, *Phys. Rev. Lett.* **113**
2131 (2014) no. 14, 141803, arXiv:1405.6241 [hep-ex]. 2.3, 5
- 2132 [22] A. Falkowski, M. Gonzalez-Alonso, A. Greljo, D. Marzocca, and M. Son, *Anomalous Triple*
2133 *Gauge Couplings in the Effective Field Theory Approach at the LHC*, *JHEP* **02** (2017) 115,
2134 arXiv:1609.06312 [hep-ph]. 2.3
- 2135 [23] B. W. Lee, C. Quigg, and H. B. Thacker, *Weak Interactions at Very High-Energies: The Role*
2136 *of the Higgs Boson Mass*, *Phys. Rev. D* **16** (1977) 1519. 2.3
- 2137 [24] D. R. Green, P. Meade, and M.-A. Pleier, *Multiboson interactions at the LHC*, *Rev. Mod.*
2138 *Phys.* **89** (2017) no. 3, 035008, arXiv:1610.07572 [hep-ex]. 2.3
- 2139 [25] L. R. Evans and P. Bryant, *LHC Machine*, JINST **3** (2008) S08001.
2140 <https://cds.cern.ch/record/1129806>. This report is an abridged version of the LHC
2141 Design Report (CERN-2004-003). 3.1
- 2142 [26] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST
2143 **3** (2008) S08003. 3.1
- 2144 [27] ATLAS Collaboration, *ATLAS inner detector: Technical Design Report*, Vol. 1. CERN,
2145 Geneva, 1997. <https://cds.cern.ch/record/331063>. 4

- 2146 [28] P. F. Åkesson, T. Atkinson, M. J. Costa, M. Elsing, S. Fleischmann, A. N. Gaponenko,
2147 W. Liebig, E. Moyse, A. Salzburger, and M. Siebel, *ATLAS Tracking Event Data Model*,
2148 Tech. Rep. ATL-SOFT-PUB-2006-004. ATL-COM-SOFT-2006-005.
2149 CERN-ATL-COM-SOFT-2006-005, CERN, Geneva, Jul, 2006.
2150 <https://cds.cern.ch/record/973401>. 4.1
- 2151 [29] P. Brückman, A. Hicheur, and S. J. Haywood, *Global chi2 approach to the Alignment of the*
2152 *ATLAS Silicon Tracking Detectors*, Tech. Rep. ATL-INDET-PUB-2005-002.
2153 ATL-COM-INDET-2005-004. CERN-ATL-INDET-PUB-2005-002, CERN, Geneva, 2005.
2154 <https://cds.cern.ch/record/835270>. 4.1, 4.1
- 2155 [30] ATLAS Collaboration Collaboration, *Alignment of the ATLAS Inner Detector Tracking*
2156 *System with 2010 LHC proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$* , Tech. Rep.
2157 ATLAS-CONF-2011-012, CERN, Geneva, Mar, 2011.
2158 <https://cds.cern.ch/record/1334582>. 4.1, 4.2.3, 4.14
- 2159 [31] R. Härtel and S. Bethke, *Iterative local Chi2 alignment approach for the ATLAS SCT*
2160 *detector*, Nov, 2005. <http://cds.cern.ch/record/1511044>. Presented 28 Nov 2005. 4.1
- 2161 [32] ATLAS Collaboration Collaboration, P. Brückman de Renstrom, *Alignment strategy for the*
2162 *Inner Detector of ATLAS*, . <https://cds.cern.ch/record/1047114>. 4.1
- 2163 [33] *Alignment of the ATLAS Inner Detector and its Performance in 2012*, Tech. Rep.
2164 ATLAS-CONF-2014-047, CERN, Geneva, Jul, 2014.
2165 <https://cds.cern.ch/record/1741021>. 4.1, 4.2.1, 4.4
- 2166 [34] J. J. Peña, *Alignment of the ATLAS Inner Detector Upgraded for the LHC Run II*, *J. Phys.*
2167 *Conf. Ser.* **664** (2015) no. 7, 072025. 4.1
- 2168 [35] J. Alison and I. J. Kroll, *The Road to Discovery: Detector Alignment, Electron Identification,*
2169 *Particle Misidentification, WW Physics, and the Discovery of the Higgs Boson*, Dec, 2015.
2170 <https://cds.cern.ch/record/1536507>. Presented 08 Nov 2012. 6
- 2171 [36] *ATLAS Inner Detector Alignment Performance with February 2015 Cosmic Rays Data*, Tech.
2172 Rep. ATL-PHYS-PUB-2015-009, CERN, Geneva, Apr, 2015.
2173 <https://cds.cern.ch/record/2008724>. 4.3, 4.2
- 2174 [37] ATLAS Collaboration Collaboration, M. Capeans, G. Darbo, K. Einsweiller, M. Elsing,
2175 T. Flick, M. Garcia-Sciveres, C. Gemme, H. Pernegger, O. Rohne, and R. Vuillermet, *ATLAS*
2176 *Insertable B-Layer Technical Design Report*, Tech. Rep. CERN-LHCC-2010-013.
2177 ATLAS-TDR-19, Sep, 2010. <https://cds.cern.ch/record/1291633>. 4.2
- 2178 [38] *Alignment of the ATLAS Inner Detector with the initial LHC data at $\sqrt{s} = 13 \text{ TeV}$* , Tech.
2179 Rep. ATL-PHYS-PUB-2015-031, CERN, Geneva, Jul, 2015.
2180 <https://cds.cern.ch/record/2038139>. 4.2
- 2181 [39] *Study of the mechanical stability of the ATLAS Insertable B-Layer*, Tech. Rep.
2182 ATL-INDET-PUB-2015-001, CERN, Geneva, Jun, 2015.
2183 <https://cds.cern.ch/record/2022587>. 4.2.1
- 2184 [40] *Common Framework Implementation for the Track-Based Alignment of the ATLAS Detector*,
2185 Tech. Rep. ATL-SOFT-PUB-2014-003, CERN, Geneva, Mar, 2014.
2186 <https://cds.cern.ch/record/1670354>. 4.2.1

- 2187 [41] ATLAS Collaboration Collaboration, *Study of alignment-related systematic effects on the*
 2188 *ATLAS Inner Detector track reconstruction*, Tech. Rep. ATLAS-CONF-2012-141, CERN,
 2189 Geneva, Oct, 2012. <https://cds.cern.ch/record/1483518>. 4.4, 4.4, 4.4.1.1, 4.4.1.2
- 2190 [42] S. Marti-Garcia, S. Camarda, T. Golling, M. Danninger, N. A. Styles, J. Wollrath, W. K.
 2191 Di Clemente, S. Henkelmann, J. Jimenez Peña, O. Estrada Pastor, A. K. Morley, H. Oide,
 2192 A. T. Vermeulen, P. Butti, E. Ricci, F. M. Follega, and R. Iuppa, *Studies of radial distortions*
 2193 *of the ATLAS Inner Detector*, Tech. Rep. ATL-COM-INDET-2018-010, CERN, Geneva, Mar,
 2194 2018. <https://cds.cern.ch/record/2306724>. 4.4
- 2195 [43] ATLAS Collaboration Collaboration, *Electron and photon energy calibration with the ATLAS*
 2196 *detector using data collected in 2015 at $\sqrt{s} = 13$ TeV*, Tech. Rep. ATL-PHYS-PUB-2016-015,
 2197 CERN, Geneva, Aug, 2016. <https://cds.cern.ch/record/2203514>. 4.4.1
- 2198 [44] CMS Collaboration, V. Khachatryan et al., *Study of vector boson scattering and search for*
 2199 *new physics in events with two same-sign leptons and two jets*, *Phys. Rev. Lett.* **114** (2015)
 2200 no. 5, 051801, [arXiv:1410.6315 \[hep-ex\]](https://arxiv.org/abs/1410.6315). 5
- 2201 [45] ATLAS Collaboration Collaboration, *Observation of electroweak production of a same-sign W*
 2202 *boson pair in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS*
 2203 *detector*, Tech. Rep. ATLAS-CONF-2018-030, CERN, Geneva, Jul, 2018.
 2204 <https://cds.cern.ch/record/2629411>. 5, 5.8, 6.6.2
- 2205 [46] CMS Collaboration, CMS Collaboration, *Observation of electroweak production of same-sign*
 2206 *W boson pairs in the two jet and two same-sign lepton final state in proton-proton collisions*
 2207 *at $\sqrt{s} = 13$ TeV*, [arXiv:1709.05822 \[hep-ex\]](https://arxiv.org/abs/1709.05822). 5
- 2208 [47] C. Bittrich, W. K. Di Clemente, et al., *Support note for measurement of electroweak*
 2209 *$W^\pm W^\pm jj$ production at $\sqrt{s} = 13$ TeV*, Tech. Rep. ATL-COM-PHYS-2018-252, CERN,
 2210 Geneva, Mar, 2018. <https://cds.cern.ch/record/2309552>. 5, 5.3.3, 2, 4, 5.4.3
- 2211 [48] B. W. Lee, C. Quigg, and H. B. Thacker, *The Strength of Weak Interactions at Very*
 2212 *High-Energies and the Higgs Boson Mass*, *Phys. Rev. Lett.* **38** (1977) 883–885. 5.0.2
- 2213 [49] S. D. Rindani, *Strong gauge boson scattering at the LHC*, in *Physics at the Large Hadron*
 2214 *Collider*, A. Datta, B. Mukhopadhyaya, A. Raychaudhuri, A. K. Gupta, C. L. Khetrapal,
 2215 T. Padmanabhan, and M. Vijayan, eds., pp. 145–155. 2009. [arXiv:0910.5068 \[hep-ph\]](https://arxiv.org/abs/0910.5068). 5.0.2
- 2216 [50] A. Alboteanu, W. Kilian, and J. Reuter, *Resonances and Unitarity in Weak Boson Scattering*
 2217 *at the LHC*, *JHEP* **11** (2008) 010, [arXiv:0806.4145 \[hep-ph\]](https://arxiv.org/abs/0806.4145). 5.0.2, 5.1
- 2218 [51] J. M. Campbell and R. K. Ellis, *Higgs Constraints from Vector Boson Fusion and Scattering*,
 2219 *JHEP* **04** (2015) 030, [arXiv:1502.02990 \[hep-ph\]](https://arxiv.org/abs/1502.02990). 5.0.2, 5.0.3
- 2220 [52] M. Szleper, *The Higgs boson and the physics of WW scattering before and after Higgs*
 2221 *discovery*, [arXiv:1412.8367 \[hep-ph\]](https://arxiv.org/abs/1412.8367). 5.0.2
- 2222 [53] E. Accomando, A. Ballestrero, A. Belhouari, and E. Maina, *Isolating Vector Boson Scattering*
 2223 *at the LHC: Gauge cancellations and the Equivalent Vector Boson Approximation vs complete*
 2224 *calculations*, *Phys. Rev.* **D74** (2006) 073010, [arXiv:hep-ph/0608019 \[hep-ph\]](https://arxiv.org/abs/hep-ph/0608019). 5.0.2

- 2225 [54] P. Anger, B. Axen, et al., *Same Sign $W^\pm W^\pm$ Production and Limits on Anomalous Quartic*
 2226 *Gauge Couplings*, Tech. Rep. ATL-COM-PHYS-2013-990, CERN, Geneva, Jul, 2013.
 2227 <https://cds.cern.ch/record/1561731>. Internal note for approved paper **STDM-2013-06**.
 2228 **5.0.3, 5.1, 5.7**
- 2229 [55] J. Almond, L. Bianchini, et al., *Search for heavy neutrino, W_R and Z_R gauge bosons in events*
 2230 *with two high- P_T leptons and jets with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV*,
 2231 Tech. Rep. ATL-COM-PHYS-2013-810, CERN, Geneva, Jun, 2013.
 2232 <https://cds.cern.ch/record/1555805>. Supporting note for EXOT-2012-24, published as
 2233 JHEP07 (2015) 162. **5.0.4**
- 2234 [56] ATLAS Collaboration, M. Aaboud et al., *Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC*, Eur. Phys. J. **C76** (2016) no. 12, 653,
 2235 [arXiv:1608.03953 \[hep-ex\]](https://arxiv.org/abs/1608.03953). **5.1**
- 2237 [57] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in*
 2238 *ATLAS*, JINST **13** (2018) no. 07, P07017. **5.1**
- 2239 [58] ATLAS Collaboration, G. Aad et al., *The ATLAS Simulation Infrastructure*, Eur. Phys. J.
 2240 **C70** (2010) 823–874, [arXiv:1005.4568 \[physics.ins-det\]](https://arxiv.org/abs/1005.4568). **5.1.1**
- 2241 [59] S. Agostinelli et al., *GEANT4 - a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003)
 2242 **250–303**. **5.1.1**
- 2243 [60] T. Sjostrand, S. Mrenna, and P. Skands, *A Brief Introduction to PYTHIA 8.1*, Comput.
 2244 Phys. Commun. **178** (2008) 852–867, [arXiv:0710.3820 \[hep-ph\]](https://arxiv.org/abs/0710.3820). **5.1.1**
- 2245 [61] T. Gleisberg et al., *Event generation with SHERPA 1.1*, JHEP **02** (2009) 007,
 2246 [arXiv:0811.4622 \[hep-ph\]](https://arxiv.org/abs/0811.4622). **5.1.1**
- 2247 [62] S. Schumann and F. Krauss, *A parton shower algorithm based on Catani-Seymour dipole*
 2248 *factorization*, JHEP **03** (2008) 038, [arXiv:0709.1027 \[hep-ph\]](https://arxiv.org/abs/0709.1027). **5.1.1**
- 2249 [63] S. Höche, F. Krauss, S. Schumann, and F. Siegert, *QCD matrix elements and truncated*
 2250 *showers*, JHEP **05** (2009) 053, [arXiv:0903.1219 \[hep-ph\]](https://arxiv.org/abs/0903.1219). **5.1.1**
- 2251 [64] R. D. Ball et al., *Parton distributions for the LHC Run II*, JHEP **04** (2015) 040,
 2252 [arXiv:1410.8849 \[hep-ph\]](https://arxiv.org/abs/1410.8849). **5.1.1**
- 2253 [65] S. Alioli, P. Nason, C. Oleari, and E. Re, *A general framework for implementing NLO*
 2254 *calculations in shower Monte Carlo programs: the POWHEG BOX*, JHEP **06** (2010) 043,
 2255 [arXiv:1002.2581 \[hep-ph\]](https://arxiv.org/abs/1002.2581). **5.1.1**
- 2256 [66] A. Ballestrero et al., *Precise predictions for same-sign W -boson scattering at the LHC*, Eur.
 2257 Phys. J. **C78** (2018) no. 8, 671, [arXiv:1803.07943 \[hep-ph\]](https://arxiv.org/abs/1803.07943). **5.1.1**
- 2258 [67] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao,
 2259 T. Stelzer, P. Torrielli, and M. Zaro, *The automated computation of tree-level and*
 2260 *next-to-leading order differential cross sections, and their matching to parton shower*
 2261 *simulations*, JHEP **07** (2014) 079, [arXiv:1405.0301 \[hep-ph\]](https://arxiv.org/abs/1405.0301). **5.1.1**
- 2262 [68] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin, and C. P. Yuan, *New*
 2263 *parton distributions for collider physics*, Phys. Rev. D **82** (2010) 074024, [arXiv:1007.2241](https://arxiv.org/abs/1007.2241)
 2264 [\[hep-ph\]](https://arxiv.org/abs/hep-ph). **5.1.1**

- 2265 [69] T. Sjostrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006)
 2266 026, [arXiv:0603175 \[hep-ph\]](https://arxiv.org/abs/0603175). 5.1.1
- 2267 [70] ATLAS Collaboration, G. Aad et al., *Muon reconstruction performance of the ATLAS*
 2268 *detector in protonproton collision data at $\sqrt{s} = 13$ TeV*, Eur. Phys. J. **C76** (2016) no. 5, 292,
 2269 [arXiv:1603.05598 \[hep-ex\]](https://arxiv.org/abs/1603.05598). 5.2.1.1, 12
- 2270 [71] ATLAS Collaboration Collaboration, *Electron efficiency measurements with the ATLAS*
 2271 *detector using the 2015 LHC proton-proton collision data*, Tech. Rep.
 2272 ATLAS-CONF-2016-024, CERN, Geneva, Jun, 2016.
 2273 <https://cds.cern.ch/record/2157687>. 5.2.1.2
- 2274 [72] M. Cacciari, G. P. Salam, G. Soyez, *The anti- k_t jet clustering algorithm*, JHEP **04** (2008) 063,
 2275 [arXiv:0802.1189 \[hep-ph\]](https://arxiv.org/abs/0802.1189). 5.2.1.3, 6.4.1
- 2276 [73] ATLAS Collaboration, M. Aaboud et al., *Jet energy scale measurements and their systematic*
 2277 *uncertainties in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Phys.
 2278 Rev. **D96** (2017) no. 7, 072002, [arXiv:1703.09665 \[hep-ex\]](https://arxiv.org/abs/1703.09665). 5.2.1.3
- 2279 [74] *Tagging and suppression of pileup jets with the ATLAS detector*, Tech. Rep.
 2280 ATLAS-CONF-2014-018, CERN, Geneva, May, 2014.
 2281 [http://cds.cern.ch/record/1700870](https://cds.cern.ch/record/1700870). 5.2.1.3
- 2282 [75] D. Adams, C. Anastopoulos, et al., *Recommendations of the Physics Objects and Analysis*
 2283 *Harmonisation Study Groups 2014*, Tech. Rep. ATL-PHYS-INT-2014-018, CERN, Geneva,
 2284 Jul, 2014. <https://cds.cern.ch/record/1743654>. 5.2.1.4
- 2285 [76] ATLAS Collaboration, M. Aaboud et al., *Measurement of the cross-section for producing a W*
 2286 *boson in association with a single top quark in pp collisions at $\sqrt{s} = 13$ TeV with ATLAS*,
 2287 JHEP **01** (2018) 063, [arXiv:1612.07231 \[hep-ex\]](https://arxiv.org/abs/1612.07231). 5.2.1.4
- 2288 [77] ATLAS Collaboration, M. Aaboud et al., *Performance of missing transverse momentum*
 2289 *reconstruction with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV*, Eur.
 2290 Phys. J. **C78** (2018) no. 11, 903, [arXiv:1802.08168 \[hep-ex\]](https://arxiv.org/abs/1802.08168). 5.2.2
- 2291 [78] ATLAS Collaboration, M. Aaboud et al., *Measurements of b-jet tagging efficiency with the*
 2292 *ATLAS detector using $t\bar{t}$ events at $\sqrt{s} = 13$ TeV*, JHEP **08** (2018) 089, [arXiv:1805.01845](https://arxiv.org/abs/1805.01845)
 2293 [\[hep-ex\]](https://arxiv.org/abs/1805.01845). 5.2.2
- 2294 [79] J.-F. Arguin, J. Claude, G. Gonella, B. P. Kersevan, K. Koneke, K. Lohwasser, K. Mochizuki,
 2295 M. Muskinja, C. Piaulet, and P. Sommer, *Support Note for Electron ID: charge*
 2296 *mis-identification*, Tech. Rep. ATL-COM-PHYS-2017-1014, CERN, Geneva, Jul, 2017.
 2297 <https://cds.cern.ch/record/2273365>. 5.3.3
- 2298 [80] C.-H. Kom and W. J. Stirling, *Charge asymmetry in W + jets production at the LHC*, Eur.
 2299 Phys. J. **C69** (2010) 67–73, [arXiv:1004.3404 \[hep-ph\]](https://arxiv.org/abs/1004.3404). 5.4
- 2300 [81] R. Steerenberg, *LHC Report: Another run is over and LS2 has just begun...*,
 2301 [https://home.cern/news/news/accelerators/](https://home.cern/news/news/accelerators/lhc-report-another-run-over-and-ls2-has-just-begun)
 2302 [lhc-report-another-run-over-and-ls2-has-just-begun](https://home.cern/news/news/accelerators/lhc-report-another-run-over-and-ls2-has-just-begun), 2018. Accessed: 2018-12-14. 6
- 2303 [82] *Letter of Intent for the Phase-I Upgrade of the ATLAS Experiment*, Tech. Rep.
 2304 CERN-LHCC-2011-012. LHCC-I-020, CERN, Geneva, Nov, 2011.
 2305 [http://cds.cern.ch/record/1402470](https://cds.cern.ch/record/1402470). 6

- 2306 [83] G. Apollinari, I. Bjar Alonso, O. Brning, M. Lamont, and L. Rossi, *High-Luminosity Large*
2307 *Hadron Collider (HL-LHC): Preliminary Design Report*. CERN Yellow Reports: Monographs.
2308 CERN, Geneva, 2015. <https://cds.cern.ch/record/2116337>. 6
- 2309 [84] ATLAS Collaboration Collaboration, ATLAS Collaboration, *ATLAS Phase-II Upgrade*
2310 *Scoping Document*, Cern-lhcc-2015-020, Geneva, Sep, 2015.
2311 <http://cds.cern.ch/record/2055248>. 6
- 2312 [85] D. Espriu and B. Yencho, *Longitudinal WW scattering in light of the “Higgs boson”*
2313 *discovery*, Phys. Rev. D **87** (2013) 055017, arXiv:1212.4158 [hep-ph]. 6, 6.1
- 2314 [86] ATLAS Collaboration Collaboration, *Prospects for the measurement of the $W^\pm W^\pm$ scattering*
2315 *cross section and extraction of the longitudinal scattering component in pp collisions at the*
2316 *High-Luminosity LHC with the ATLAS experiment*, Tech. Rep. ATL-PHYS-PUB-2018-052,
2317 CERN, Geneva, Dec, 2018. <http://cds.cern.ch/record/2652447>. 6
- 2318 [87] ATLAS Collaboration Collaboration, *Studies on the impact of an extended Inner Detector*
2319 *tracker and a forward muon tagger on $W^\pm W^\pm$ scattering in pp collisions at the*
2320 *High-Luminosity LHC with the ATLAS experiment*, Tech. Rep. ATL-PHYS-PUB-2017-023,
2321 CERN, Geneva, Dec, 2017. <https://cds.cern.ch/record/2298958>. 6
- 2322 [88] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel,
2323 C. O. Rasmussen, and P. Z. Skands, *An Introduction to PYTHIA 8.2*, Comput. Phys.
2324 Commun. **191** (2015) 159–177, arXiv:1410.3012 [hep-ph]. 6.2
- 2325 [89] ATLAS Collaboration Collaboration, *Expected performance for an upgraded ATLAS detector*
2326 *at High-Luminosity LHC*, Tech. Rep. ATL-PHYS-PUB-2016-026, CERN, Geneva, Oct, 2016.
2327 <http://cds.cern.ch/record/2223839>. 6.3
- 2328 [90] P. C. Bhat, H. B. Prosper, S. Sekmen, and C. Stewart, *Optimizing Event Selection with the*
2329 *Random Grid Search*, Comput. Phys. Commun. **228** (2018) 245–257, arXiv:1706.09907
2330 [hep-ph]. 6.5.1
- 2331 [91] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, *Asymptotic formulae for likelihood-based*
2332 *tests of new physics*, Eur. Phys. J. **C71** (2011) 1554, arXiv:1007.1727 [physics.data-an].
2333 [Erratum: Eur. Phys. J.C73,2501(2013)]. 6.5.1