

Lecture Notes for

Algebraic and Hermitian K -Theory

Lecturer
Fabian Hebestreit

Notes typed by
Ferdinand Wagner

Winter Term 2020/21
University of Bonn

This text consists of unofficial notes on the lecture *Advanced Topics in Algebra – Algebraic and Hermitian K-Theory*, taught at the University of Bonn by Fabian Hebestreit in the winter term (Wintersemester) 2020/21.

Some additions have been made by the author. To distinguish them from the lecture's actual contents, they are labelled with an asterisk. So any *Proof** or *Lemma** etc. that the reader might encounter are wholly the author's responsibility.

Please report errors, typos etc. through the *Issues* feature of GitHub, or just tell me before or after the lecture.

Contents

List of Lectures	iv
0. Introduction	1
Organisational Stuff	1
A Fairytale	1
Outline of the Course	5
I. Recollections and Preliminaries	6
∞ -Categories and Anima	6
Cocartesian/Left Fibrations and Lurie’s Straightening Equivalence	13
Yoneda’s Lemma, Limits and Colimits in ∞ -Categories	20
Yoneda’s Lemma and Adjunctions	20
First Properties and Examples of (Co)Limits	22
Colimits and the Yoneda embedding	30
Kan Extensions	34
Bousfield Localisations	43
The Rezk Nerve and Complete Segal Spaces	49
II. Symmetric Monoidal and Stable ∞-Categories	59
\mathbb{E}_1 -Monoids and \mathbb{E}_1 -Groups	59
\mathbb{E}_∞ -Monoids and \mathbb{E}_∞ -Groups	74
Spectra and Stable ∞ -Categories	87
An Interlude on ∞ -Operads	102
Towards the Tensor Product of Spectra	107
Some Brave New Algebra	124
\mathbb{E}_∞ -Ring Spectra	125
∞ -Categories of Modules	129
Tensor Products over Arbitrary Bases	137
Miscellanea	141
III. The Group Completion Theorem and the K-Theory of Finite Fields	144
Localisations of \mathbb{E}_∞ -Ring Spectra	144
The Group Completion Theorem and $K_1(R)$	149
Towards the K -Theory of Finite Fields	153
Rational K -Theory	153
The Cyclic Invariance Condition and Quillen’s Plus Construction	156
Topological K -Theory	164
A Sketch of Quillen’s Computation	171
IV. The K-Theory of Stable ∞-Categories	184
Bibliography	186

List of Lectures

Lecture 0 (27th October, 2020)	1
From Riemann–Roch to Quillen’s definition of K -theory. Outline of the course.	
Lecture 1 (29th October, 2020)	6
Fabian gives a history lesson. A brief recollection of ∞ -categories, nerve functors, anima, Joyal’s lifting theorem, and Hom spaces in ∞ -categories.	
Lecture 2 (3rd November, 2020)	10
Examples of ∞ -categories. Dold–Kan correspondence. Straightening/unstraightening equivalence and an invariant definition of cocartesian/left fibrations.	
Lecture 3 (5th November, 2020)	16
The Hom functor and the twisted arrow category. Yoneda’s lemma, adjunctions, (co)limits. Some computations of (co)limits.	
Lecture 4 (10th November, 2020)	24
Localisations. More properties of (co)limits. Left Kan extensions for presheaf categories. Hom anima in functor categories.	
Lecture 5 (12th November, 2020)	34
Kan extensions and a version of Theorem I.2 for ∞ -categories. Lots of examples to show how algebraic topology becomes a triviality after enough ∞ -category theory. — “I’m so happy he [Peter Scholze] uses ‘An’ [to denote anima]. I forced this on him. He wanted to call them ‘Ani’. That’s the name of Darth Vader!”	
Lecture 6 (17th November, 2020)	43
Bousfield localisations, limits and colimits inside them. Applications to derived categories of rings. — “Look up a paper on Bousfield colocalisations. And then cry yourself to sleep.”	
Lecture 7 (19th November, 2020)	49
The Rezk nerve and (complete) Segal spaces/anima. — “Today is brainf*ck time. There’s no deep content, except for the things I’m not saying.”	
Lecture 8 (24th November, 2020)	56
A sleep-deprived Fabian proves that $Q(\mathcal{C})$ is a Segal ∞ -category and introduces cartesian monoids.	

LIST OF LECTURES

Lecture 9 (26th November, 2020)	63
More on \mathbb{E}_1 -monoids and \mathbb{E}_1 -groups. Group completion. — “The fact that you can write down a group or a monoid in semester 2 is sort of an accident.”	
Lecture 10 (1st December, 2020)	68
A pushout computation. Monoidal 1-categories give monoidal ∞ -categories. Definition of algebraic and hermitian K -theory.	
Lecture 11 (3rd December, 2020)	71
Topological K -theory and its connection to deep theorems in geometric topology. Cartesian commutative monoids and groups.	
Lecture 12 (8th December, 2020)	76
Semi-additive and additive ∞ -categories, monoids and groups in these.	
Lecture 13 (10th December, 2020)	86
Spectra! The recognition principle for infinite loop spaces. Stable ∞ -categories. — “Peter May hates this proof. I love it.”	
Lecture 14 (15th December, 2020)	93
Spectra and prespectra, spectrification. Σ^∞ and the Baratt–Priddy–Quillen theorem. Another model for spectra. Towards a symmetric monoidal structure on Sp .	
Lecture 15 (17th December, 2020)	102
Operads. Symmetric monoidal ∞ -category as operads, strict vs. lax monoidal functors. Some examples. A technical lemma about presheaves.	
Lecture 16 (22nd December, 2020)	109
Cocartesian and cartesian monoidal structures. Algebras over ∞ -operads. Day convolution. Operadic limits and colimits. The tensor product on spectra. — “I didn’t feel like presenting this as a gift to you. It’s more like a punishment for Christmas.”	
Lecture 17 (7th January, 2021)	124
Some tensor product calculations. \mathbb{E}_∞ -ring spectra. The ∞ -operads Assoc and LMod , categories of modules over \mathbb{E}_∞ -ring spectra.	
Lecture 18 (12th January, 2021)	131
A user’s guide to Lurie’s treatment of brave new algebra. Localisations of \mathbb{E}_∞ -ring spectra.	
Lecture 19 (14th January, 2021)	148
The group completion theorem. Computation of $K_1(R)$. $K_1(R)$ as a generalised determinant. — “Aaahhh, we’re not gonna discuss more about this thing!”	
Lecture 20 (19th January, 2021)	152
Primitive elements in coalgebra and Hopf algebra structures on rational homology. Rational K -theory. The cyclic invariance criterion.	

LIST OF LECTURES

- Lecture 21 (21st January, 2021)** **159**
The Quillen plus construction. Topological K -theory, Bott periodicity, Adams operations.
—“*My mum is calling me ... [Rejects call] Now my phone has decided to tell me that my mum has called me.*”
- Lecture 22 (26th January, 2021)** **169**
More on Adams operations. A sketch of Quillen’s computation of the K -theory of finite fields.
- Lecture 23 (28th January, 2021)** **177**
Completions of spectra. Suslin rigidity and further results in the K -theory of fields. K_0 of stable ∞ -categories.

CHAPTER 0.

Introduction

0

Organisational Stuff

Under no circumstances you should call Fabian by his last name! Also there will be oral exams in the weeks from 15th–19th February and 22nd–26th March.

Disclaimer ⚠. — These are not official lecture notes. Instead, Fabian uploads his own handwritten notes [\[A&HK\]](#) (please suggest a better shorthand) to the lecture’s [website](#). These are an excellent resource and please the eye with their purple colour scheme! So why should I bother typing my own notes? This is because of two main reasons. First, I like having my notes as one single document with clickable hyperlinks and a somewhat reasonable file size (I once attempted to open Fabian’s straightening/unstraightening notes on my phone with slow mobile internet—don’t try this at home). The second reason is that typing notes and polishing them after the lecture really forces me to think through all the technical details, which can be a lot of work, but it’s the best way for me to learn that stuff.

I might also take the opportunity to try and work out some skipped details or omitted proofs myself (given the Fabian’s ambitious goals for this lecture, there can be no doubt that some cuts will have to be made). *My own additions are marked with an asterisk!* So whenever you encounter a *Proof** which is not just digging up some references, or a *Lemma**, be extra careful for mistakes. If you happen to spot some, do not hesitate to tell me via [GitHub](#) or in person.

Differences in Numbering ⚠. — Unfortunately, my numbers in Chapter [I](#) tend to diverge from Fabian’s notes, mainly because some examples discussed there haven’t been part of the lecture (that’s another reason to check out Fabian’s notes!). I’m trying to keep the numbers aligned where possible and to minimize the offset where not.

A Fairytale

LECTURE 0
27th Oct, 2020

We start with an overview of the mathematical developments that eventually led to the invention of K -theory. Don’t worry if you’re not familiar with the stuff on the next few pages, it is neither a prerequisite for the lecture, nor will it play any prominent role in it.

Let’s begin in the 1850’s: Consider a compact Riemann surface Σ and let $D \in \mathbb{Z}[\Sigma]$ be a divisor on Σ . That is, $D = \sum D_s \{s\}$ is a formal sum of points $s \in \Sigma$ with coefficients $D_s \in \mathbb{Z}$, all but finitely many of which are zero. For example, if $f: \Sigma \rightarrow \mathbb{C}$ is a meromorphic function, one could consider the *principal divisor* $D(f)$ given by

$$D(f)_s = \begin{cases} a & \text{if } f \text{ has a zero of order } a \text{ at } s \\ -a & \text{if } f \text{ has a pole of order } a \text{ at } s \\ 0 & \text{else} \end{cases}.$$

For some divisor $D = \sum D_s \{s\}$, put $\deg D = \sum D_s$ and consider the \mathbb{C} -vector space

$$M(D) = \{f: \Sigma \rightarrow \mathbb{C} \text{ meromorphic} \mid D(f)_s \geq -D_s \text{ for all } s \in \Sigma\}.$$

An important problem in the theory of Riemann surfaces is to determine the dimension $\dim M(D)$. Riemann proved the inequality $\dim M(D) \geq \deg D + 1 - g(\Sigma)$, which was soon improved upon by his student Roch, who obtained what is famously known today as the *Riemann–Roch theorem*.

0.1. Theorem (Riemann–Roch). — *Let Σ be a compact Riemann surface of genus $g(\Sigma)$ and D be a divisor on Σ . Put $D^\vee = K_\Sigma - D$, where K_Σ is the divisor of any 1-form on Σ . Then*

$$\dim M(D) - \dim M(D^\vee) = \deg D + 1 - g(\Sigma).$$

Note that K_Σ , and thus D^\vee , are only defined up to a principal divisor—in other words, only their *divisor class* is well-defined—but that’s all we need, since both $\dim M(D)$ and $\deg D$ don’t change if D is replaced by $D + D(f)$ for some meromorphic $f: \Sigma \rightarrow \mathbb{C}$.

0.2. Example. — Consider $\Sigma = \mathbb{CP}^1 = \mathbb{C} \cup \{\infty\}$ and choose $D = n\{\infty\}$. Then $M(D)$ is the space of all meromorphic functions $f: \mathbb{CP}^1 \rightarrow \mathbb{C}$ whose pole at ∞ has order at most n . In other words,

$$M(D) = \left\{ f: \mathbb{C} \rightarrow \mathbb{C} \text{ holomorphic} \mid \begin{array}{l} |f(z)| \text{ is bounded by } C|z|^n \text{ for some} \\ \text{suitable constant } C \geq 0 \text{ as } |z| \rightarrow \infty \end{array} \right\}.$$

To get $K_{\mathbb{CP}^1}$ we can choose the meromorphic 1-form dz , which is holomorphic on \mathbb{C} and has a pole of order 2 at ∞ (because $d(z^{-1}) = -z^{-2}dz$ has a pole of order 2 at 0). Thus the divisor class of $K_{\mathbb{CP}^1}$ is that of $-2\{\infty\}$. Plugging in Theorem 0.1 gives

$$\dim M(n\{\infty\}) - \dim M(-(2+n)\{\infty\}) = n + 1.$$

For $n = 0$ we obtain $M(0) \cong \mathbb{C}$, since all bounded holomorphic functions on \mathbb{C} are constant by Liouville’s theorem. By the same reason, $M(n\{\infty\}) = 0$ for $n < 0$. Hence

$$\dim M(n\{\infty\}) = n + 1 \quad \text{for all } n \geq 0,$$

which makes a lot of sense since we would expect (and have just proved) that $M(n\{\infty\})$ is precisely the space of polynomials of degree $\leq n$ in that case.

Now let’s try to restate the Riemann–Roch theorem in more modern terms. The first step is to replace divisors by line bundles, which can be done by means of the bijection

$$\{\text{divisors on } \Sigma\} / \{\text{principal divisors}\} \xrightarrow{\sim} \{\text{isomorphism classes of line bundles on } \Sigma\},$$

which is, in fact, an isomorphism of abelian groups between the *divisor class group* $\text{Cl}(\Sigma)$, whose group structure is inherited from $\mathbb{Z}[\Sigma]$, and the *Picard group* $\text{Pic}(\Sigma)$, whose group structure is given by the tensor product of line bundles. If a line bundle L corresponds to a divisor D under this isomorphism, then the space $M(D)$ corresponds to the \mathbb{C} -vector space $\Gamma(\Sigma, L)$ of holomorphic sections of L . Thus, Theorem 0.1 can be restated as

$$\dim \Gamma(\Sigma, L) - \dim \Gamma(\Sigma, T^*\Sigma \otimes L^{-1}) = \deg L + 1 - g(\Sigma)$$

Observe that $\Gamma(\Sigma, L) = H_{\text{sheaf}}^0(\Sigma, L)$ and $\Gamma(\Sigma, T^*\Sigma \otimes L^{-1}) = H_{\text{sheaf}}^1(\Sigma, L)$ by Serre duality. So the term on the left-hand side can be interpreted as the “Euler characteristic” $\chi(\Sigma, L)$. This was the starting point for a generalisation to arbitrary dimensions found by Hirzebruch, who was not only the founding father of all mathematics in Bonn after the war, but also incredibly good at guessing the correct generalisations.

0.3. Theorem (Hirzebruch–Riemann–Roch). — *Let $E \rightarrow X$ be a holomorphic vector bundle over a d -dimensional compact complex manifold X . Then*

$$\chi(X, E) = \sum_{i=0}^d (c_i(E) \cup \text{Td}_{d-i}(TX)).$$


Here $c_i(E)$ is the i^{th} Chern class of E and $\text{Td}_{d-i}(TX)$ is the $(d-i)^{\text{th}}$ Todd class of TX , so that the right-hand side lives in the cohomology group $H^{2d}(X, \mathbb{Z}) = \mathbb{Z}$ and the above equation makes sense.

Still no sign of K -theory though. This is when Grothendieck, the master of them all, entered the stage. Recall that $H^i(X, E) = R^i\Gamma(X, E)$. Consider the canonical map $f: X \rightarrow *$, so that the global sections functor is canonically isomorphic to the pushforward along f . In formulas, $f_* = \Gamma(X, -)$. Grothendieck’s idea was to generalize Theorem 0.3 to arbitrary proper morphisms $f: X \rightarrow Y$ of complex manifolds by replacing the $H^i(X, E)$ (appearing in the definition of $\chi(X, E)$) by $R^i f_* E$. This raises an immediate question though: What is $\sum (-1)^i R^i f_* E$ supposed to be? The summands are coherent sheaves on Y after all, which we can add (using the direct sum), but surely not subtract one from another. And that brings us straight to K -theory!

0.4. Definition. — Let X be a complex manifold. We define the 0^{th} K -groups $K_0(X)$ and $K^0(X)$ as follows:

- (a) $K_0(X)$ is the group completion of the monoid of isomorphism classes vector bundles on X (the monoid structure is given by taking direct sums), modulo the relation $[E] = [E'] + [E'']$ for every short exact sequence $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$ (that’s the condition that was missing in the lecture).
- (b) $K^0(X)$ is defined in the same way, with vector bundles replaced by arbitrary coherent sheaves on X .

0.5. Theorem (Grothendieck–Riemann–Roch). — *Let $f: X \rightarrow Y$ be a proper morphism of complex manifolds. It induces a morphism $f_! = \sum (-1)^i R^i f_*: K^0(X) \rightarrow K^0(Y)$ on K -groups which fits into the following hellish commutative diagram:*

$$\begin{array}{ccc} K^0(X) & \xrightarrow{f_!} & K^0(Y) \\ \text{Td}(X) \text{ ch}(-) \downarrow & & \downarrow \text{ch}(-) \text{Td}(Y) \\ H^*(X, \mathbb{Z}) & \xrightarrow{f_*} & H^*(Y, \mathbb{Z}) \end{array}$$


The bottom line is the usual pushforward in cohomology (use Poincaré duality on X , then the usual pushforward $f_*: H_*(X, \mathbb{Z}) \rightarrow H_*(Y, \mathbb{Z})$ on homology, and finally use Poincaré duality on Y to get back to cohomology).

The theory of $K_0(X)$ (and its higher versions $K_i(X)$) is called *topological K-theory* and was developed by Atiyah and Hirzebruch soon after Grothendieck had presented his result at the Arbeitstagung in Bonn. But $K_0(X)$ also has an algebraic analogue.

0.6. Definition. — Let R be a ring. The 0^{th} K -group $K_0(R)$ is the group completion of the monoid of finite projective R -modules (the monoid structure is given by taking direct sums, as usual).

Since every short exact sequence of projective R -modules splits, we don't need to divide out the relation from Definition 0.4. The group $K_0(R)$ is an interesting invariant of rings: It is the universal recipient for a “dimension function” for finite projective R -modules.

- 0.7. Example.** — (a) If $R = k$ is a field (or more generally a PID), then the usual dimension induces an isomorphism $K_0(k) \xrightarrow{\sim} \mathbb{Z}$.
- (b) In general, the map $\mathbb{Z} \rightarrow K_0(R)$ induced by $n \mapsto [R^{\oplus n}]$ for $n \geq 0$ is injective iff R has the *invariant basis number property*.
- (c) If $R = \mathbb{Q}[G]$ for some finite group G , then $K_0(R) = \bigoplus_{V \in \text{Irr}(G)} \mathbb{Z}$, where the indexing set $\text{Irr}(G)$ is the set of isomorphism classes of irreducible G -representations. Indeed, in this case $\mathbb{Q}[G] = \prod_{V \in \text{Irr}(G)} \text{Mat}_{n_V}(\text{End}(V))$ for some integers $n_V \geq 0$ holds by the Artin–Wedderburn theorem, which easily gives the above characterisation.

We can go one step further and give an ad hoc definition of $K_1(R)$.

0.8. Definition. — The 1^{st} K -group $K_1(R) = \text{GL}_{\infty}(R)^{\text{ab}}$ is the abelianisation of the infinite general linear group $\text{GL}_{\infty}(R) = \text{colim}_{n \geq 0} \text{GL}_n(R)$.

Moreover, if $I \subseteq R$ is an ideal and $S \subseteq R$ is a multiplicative subset, then one can define K -groups $K_0(I)$ and $K_0(R, S)$ fitting into exact sequences

$$\begin{aligned} K_1(R) &\longrightarrow K_1(R/I) \xrightarrow{\partial} K_0(I) \longrightarrow K_0(R) \longrightarrow K_0(R/I) \\ K_1(R) &\longrightarrow K_1(R[S^{-1}]) \xrightarrow{\partial} K_0(R, S) \longrightarrow K_0(R) \longrightarrow K_0(R[S^{-1}]) , \end{aligned}$$

which look a bit too much like long exact cohomology sequences to be a coincidence. People actually managed to produce an ad hoc definition of $K_2(R)$ fitting into the sequences above, but that was where consensus ended: People suggested several non-equivalent definitions of higher K -groups and it was not at all clear how to continue ...

... until Quillen came! He realized that $K_0(R)$ and $K_1(R)$ could be written as homotopy groups of a certain simplicial group: Let $\text{Proj}(R)$ denote the symmetric monoidal groupoid of finite projective R -modules. Now consider the inclusion

$$\{\text{Picard groupoids}\} \subseteq \{\text{symmetric monoidal groupoids}\} .$$

Here a symmetric monoidal groupoid (G, \otimes) is a *Picard groupoid* if for all $x \in G$ there exists a $y \in G$ such that $x \otimes y \simeq 1$. The above inclusion has a left adjoint $(-)^{\text{grp}}$, and using this we can write

$$K_i(R) = \pi_i N(\text{Proj}(R)^{\text{grp}}) \quad \text{for } i = 1, 2 .$$

Here N denotes the nerve construction.

OUTLINE OF THE COURSE

Quillen’s suggestion, although he didn’t have the words to say that yet, was to do the same, but in the setting of ∞ -categories. That is, we consider

$$\begin{array}{ccc}
 \{\text{Picard groupoids}\} & \xleftarrow{(-)^{\text{grp}}} \subseteq & \{\text{symmetric monoidal groupoids}\} \\
 \text{!} \cap & & \text{!} \cap \\
 \{\text{Picard } \infty\text{-groupoids}\} & \xleftarrow{(-)^{\infty\text{-grp}}} \subseteq & \{\text{symmetric monoidal } \infty\text{-groupoids}\}
 \end{array}$$

(the objects on the bottom line are also known as *grouplike* \mathbb{E}_∞ -spaces and \mathbb{E}_∞ -spaces respectively). In this framework, we can finally define the higher K -groups!

0.9. Definition. — The i^{th} K -group of a ring R is defined as $K_i(R) = \pi_i(\text{Proj}(R)^{\infty\text{-grp}})$ (these are abelian groups). More importantly, we define $K(R) = \text{Proj}(R)^{\infty\text{-grp}}$, which is an object of the ∞ -category of anima.

0.10. Warning $\triangle!$. — The functor

$$(-)^{\infty\text{-grp}}: \{\text{symmetric monoidal } \infty\text{-groupoids}\} \longrightarrow \{\text{Picard } \infty\text{-groupoids}\}$$

is bloody complicated. For example, one has $\{\text{Finite sets}, \sqcup\}^{\infty\text{-grp}} = \Omega^\infty \mathbb{S} = \text{colim}_{n \geq 0} \Omega^n \mathbb{S}^n$, so even in the simplest case—finite sets and disjoint union—the homotopy groups of what comes out are terrifying: they are the stable homotopy groups of spheres.

Outline of the Course

The course is planned to consist of three parts, but let’s see how far we will actually get.

Part 1. — *wherein we lay the required ∞ -categorical foundations.* We will discuss symmetric monoidal structures on ∞ -categories and anima. In particular, we will analyse \mathbb{E}_∞ -spaces/groups and develop the theory of spectra and stable ∞ -categories. Important examples of stable ∞ -categories are the ∞ -category Sp of spectra and the derived ∞ -category $\mathcal{D}(R)$ of R -modules. We will also prove that $\{\text{Picard } \infty\text{-groupoids}\} \simeq \{\text{connective spectra}\}$, which is a result due to Boardman–Vogt and May.

Part 2. — *wherein we finally define K -theory.* Apart from that, we will do some basic computations, including Quillens computation of $K_*(\mathbb{F}_q)$. There might also be some guest lectures to get an overview of the landscape.

Part 3. — *wherein we do some “modern” K -theory.* In particular, we will discuss K -theory as a functor $K: \text{Cat}_\infty^{\text{st}} \rightarrow \text{Sp}$ from the ∞ -category of stable ∞ -categories to the ∞ -category of spectra, and prove the basic results of “localisation, resolution, and dévissage”. For this we will follow [LT19] as well as the series of papers [Fab+20_I; Fab+20_{II}; Fab+20_{III}] that Fabian co-authored. These papers are actually concerned with *hermitian* K -theory, which arises if we replace $\text{Proj}(R)$ by

$$\text{Unimod}(R) = \{(P, q) \mid P \text{ is finite projective, } q \text{ is a unimodular form on } R\}.$$

Fabian hopes to develop the algebraic and the hermitian theory simultaneously. If time permits, we will also talk about K -theory being the “universal additive invariant” in the sense of [BGT13].

CHAPTER I.

Recollections and Preliminaries

I

∞ -Categories and Anima

LECTURE 1
29th Oct, 2020

The plan for today is to give a rapid review of Fabians lectures from the last two semesters. Naturally we won't really prove anything, but give references to Fabians notes [HC_I; HC_{II}] instead.

I.1. Definition. — The *simplex category* Δ is the category of finite totally ordered sets and order preserving maps. For \mathcal{C} a category we put $s\mathcal{C} = \text{Fun}(\Delta^{\text{op}}, \mathcal{C})$ and $c\mathcal{C} = \text{Fun}(\Delta, \mathcal{C})$ and call this the categories of *simplicial* and *cosimplicial objects* in \mathcal{C} .

I.2. Theorem. — Let A be a small category and let \mathcal{C} be a cocomplete category. Then the Yoneda embedding $Y: A \rightarrow \text{Fun}(A^{\text{op}}, \text{Set})$ induces an equivalence of categories

$$Y^*: \text{Fun}^L(\text{Fun}(A^{\text{op}}, \text{Set}), \mathcal{C}) \xrightarrow{\sim} \text{Fun}(A, \mathcal{C}).$$

Here Fun^L denotes the full subcategory of functors who are left adjoints (and thus admit right adjoints).

*Proof**. In [HC_I, Theorem I.41] we proved this statement with Fun^L replaced by $\text{Fun}^{\text{colim}}$, the full subcategory of colimit-preserving functors. But every such functor automatically admits a right adjoint by [HC_I, Proposition II.18], so $\text{Fun}^{\text{colim}} \subseteq \text{Fun}^L$. As all left adjoints preserve colimits, this inclusion is actually an equality. \square

In particular, we can apply Theorem I.2 to $A = \Delta$ and obtain $\text{Fun}^L(s\text{Set}, \mathcal{C}) \simeq \text{Fun}(\Delta, \mathcal{C})$ (and of course the right-hand side is the category $c\mathcal{C}$ of cosimplicial objects of \mathcal{C}). Given a functor $F: \Delta \rightarrow \mathcal{C}$, we denote the corresponding adjoint pair by

$$|_F: s\text{Set} \rightleftarrows \mathcal{C} : \text{Sing}_F.$$

In concrete terms, $|_F$ can be described as the left Kan extension

$$\begin{array}{ccc} \Delta & \xrightarrow{F} & \mathcal{C} \\ Y \downarrow & \nearrow \text{Lan}_Y F = |_F & \\ s\text{Set} & & \end{array}$$

of F along the Yoneda embedding $Y: \Delta \rightarrow s\text{Set}$. Moreover, as Sing_F is supposed to be right-adjoint to $|_F$, it is necessarily given by

$$\text{Sing}_F(X)_n = \text{Hom}_{\mathcal{C}}(|\Delta^n|_F, X) = \text{Hom}_{\mathcal{C}}(F([n]), X).$$

I.3. Example. — The following adjunctions arise in the way explained above.

- (a) $\pi_0: \mathbf{sSet} \rightleftarrows \mathbf{Set} : \text{const}$, where as usual π_0 denotes the set of connected components.
- (b) $\pi: \mathbf{sSet} \rightleftarrows \mathbf{Cat} : \mathbf{N}$. Here πX denotes the *homotopy category* of a simplicial set X , and $\mathbf{N}(\mathcal{C})$ the *nerve* of a category \mathcal{C} .
- (c) $| \cdot |: \mathbf{sSet} \rightleftarrows \mathbf{Top} : \text{Sing}$, the adjunction that motivates the above notation.
- (d) $X \times -: \mathbf{sSet} \rightleftarrows \mathbf{sSet} : F(X, -)$ for any simplicial set X . In particular, $F(X, Y)$ is our notation for the simplicial set of maps between X and Y , explicitly given by $F(X, Y)_n = \text{Hom}_{\mathbf{sSet}}(X \times \Delta^n, Y)$.
- (e) $\mathfrak{C}[-]: \mathbf{sSet} \rightleftarrows \mathbf{Cat}^{\mathbf{sSet}} : \mathbf{N}^c$, taking a simplicial set X to the simplicially enriched category $\mathfrak{C}[X]$. The right adjoint \mathbf{N}^c is called the *coherent nerve* functor.


I.4. Definition (“Horn filling conditions”). — A *quasicategory* or ∞ -category \mathcal{C} is a simplicial set such that for all $0 < i < n$ and all solid diagrams

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & \mathcal{C} \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^n & & \end{array}$$

there exists a dashed arrow as indicated rendering it commutative. A simplicial set X is a *Kan complex* if it is an ∞ -category and the above condition holds for $i = 0, n$ as well. The full subcategories of \mathbf{sSet} spanned by quasicategories and Kan complexes are denoted \mathbf{qCat} and \mathbf{Kan} .

If \mathcal{C} and \mathcal{D} are ∞ -categories, a *functor* $F: \mathcal{C} \rightarrow \mathcal{D}$ is a map of simplicial sets, or equivalently a 0-simplex in the simplicial set $F(\mathcal{C}, \mathcal{D})$, which we usually denote $\text{Fun}(\mathcal{C}, \mathcal{D})$ in this case. Similarly, a *natural transformation* $\eta: F \Rightarrow G$ between functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$ is a 1-simplex η in $\text{Fun}(\mathcal{C}, \mathcal{D})$ connecting the 0-simplices F and G .

The simplicial set Λ_i^n in Definition I.4 is called the i^{th} n -horn and is given by removing the interior of Δ^n as well as the face opposing its i^{th} vertex. The 2-horns look as follows:



The connection between ∞ -categories as defined in Definition I.4 and ordinary categories (“1-categories”) is given by the following theorem.

I.5. Theorem (“Unique horn filling”). — *The nerve functor $N: \mathbf{Cat} \rightarrow \mathbf{sSet}$ is fully faithful with essential image given by those simplicial sets X such that the dashed arrow in Definition I.4 exists uniquely. In fact, for a simplicial set X the following are equivalent:*

- (a) $X \cong \mathbf{N}(\mathcal{C})$ for some category \mathcal{C} .
- (b) For all $0 < i < n$, the restriction $F(\Delta^n, X) \xrightarrow{\sim} F(\Lambda_i^n, X)$ is an isomorphism.

- (c) For all n , the restriction $F(\Delta^n, X) \rightarrow F(I^n, X)$ is an isomorphism. Here I^n denotes the spine of Δ^n , i.e., the union of all edges connecting consecutive vertices.

Moreover, an ordinary category \mathcal{C} is a groupoid iff $N(\mathcal{C})$ is a Kan complex.

*Proof**. In [HC_I, Theorem II.25] we proved the equivalence of (a), (b), and (c), but with $\text{Hom}_{\text{sSet}}(-, X)$ instead of $F(-, X)$. However, the weaker version already implies the stronger one: We have

$$F(-, X)_n = \text{Hom}_{\text{sSet}}(- \times \Delta^n, X) = \text{Hom}_{\text{sSet}}(-, F(\Delta^n, X)),$$

and if $X \cong N(\mathcal{C})$ is the nerve of a category, then

$$F(\Delta^n, X) = F(N([n]), N(\mathcal{C})) = N\text{Fun}([n], \mathcal{C}),$$

so we are mapping into the nerve of a category again. The right equality follows from a straightforward calculation, using that $N: \text{Cat} \rightarrow \text{sSet}$ is fully faithful. The additional assertion about groupoids and Kan complexes is addressed in Theorem I.13 \square

I.6. Example. — “My first Kan complexes”:

- (a) For all topological spaces $X \in \text{Top}$, the simplicial set $\text{Sing } X$ is a Kan complex, which is pretty easy to check.
- (b) Any simplicial group is a Kan complex. That’s not entirely obvious, but not hard. See [Stacks, Tag 08NZ] for example.
- (c) If \mathcal{C} and \mathcal{D} are 1-categories, then $N(\text{Fun}(\mathcal{C}, \mathcal{D})) \cong \text{Fun}(N(\mathcal{C}), N(\mathcal{D}))$.

I.7. Theorem (Kan/Joyal). — *Let \mathcal{C} be a simplicial set.*

- (a) *If \mathcal{C} is a Kan complex or an ∞ -category, then the same holds for $F(X, \mathcal{C})$ for all simplicial sets X .*
- (b) *\mathcal{C} is an ∞ -category iff $F(\Delta^n, \mathcal{C}) \rightarrow F(\Lambda_i^n, \mathcal{C})$ is a trivial fibration for all $0 < i < n$, which is again equivalent to $F(\Delta^n, \mathcal{C}) \rightarrow F(I^n, \mathcal{C})$ being a trivial fibration for all n .*
- (c) *\mathcal{C} is a Kan complex iff the conditions from (b) hold and in addition $F(\Delta^n, \mathcal{C}) \rightarrow F(\Lambda_i^n, \mathcal{C})$ are trivial fibrations for $i = 0, n$.*

Actually, in (b) and (c) it suffices to have the respective conditions only for $n = 2$.

*Proof**. The “if” parts of (b) and (c) follow from the fact that trivial fibrations are surjective. For the rest of (a), (b), and (c), see [HC_I, Corollary V.2.23 and Corollary VI.2.4]. The fact that \mathcal{C} is already an ∞ -category if only $F(\Delta^2, \mathcal{C}) \rightarrow F(\Lambda_1^2, \mathcal{C})$ is a trivial fibration was proved in [HC_I, Corollary VI.2.5]. This easily implies the corresponding assertion for Kan complexes: If $F(\Delta^2, \mathcal{C}) \rightarrow F(\Lambda_i^2, \mathcal{C})$ is surjective for $i = 0, 2$, then every morphism in \mathcal{C} has a left- and right-inverse, hence \mathcal{C} is a Kan complex by Theorem I.13. \square

I.8. Warning $\triangle!$. — In view of Theorem I.7(b) it seems tempting to define ∞ -categories as simplicial sets that have lifting against $I^n \subseteq \Delta^n$. But that’s *wrong*! The class of simplicial sets obtained in that way, called *composers*, is larger than the class of ∞ -categories. If you want to replace Λ_i^n by I^n , you really need the stronger condition that $F(\Delta^n, \mathcal{C}) \rightarrow F(I^n, \mathcal{C})$ is a trivial fibration rather than just surjective on 0-simplices.

I.9. Definition. — Let \mathcal{C} be an ∞ -category. For 0-simplices $a, b \in \mathcal{C}_0$, define their *hom space/mapping anima/any combination of these* as the pullback

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(a, b) & \longrightarrow & \mathrm{Fun}(\Delta^1, \mathcal{C}) \\ \downarrow & \lrcorner & \downarrow (s, t) \\ \Delta^0 & \xrightarrow{(a, b)} & \mathcal{C} \times \mathcal{C} \end{array}$$

Here $s, t: \mathrm{Fun}(\Delta^1, \mathcal{C})$ send a 1-simplex in \mathcal{C} to its source and target respectively. The ∞ -category $\mathrm{Fun}(\Delta^1, \mathcal{C})$ is also called the *arrow category* of \mathcal{C} and denoted $\mathrm{Ar}(\mathcal{C})$.

- I.10. Example.** — (a) For any 1-category \mathcal{D} one has $\mathrm{Hom}_{N(\mathcal{D})}(a, b) \cong \mathrm{const} \mathrm{Hom}_{\mathcal{D}}(a, b)$. In other words, $\mathrm{Hom}_{N(\mathcal{D})}(a, b)$ is a discrete simplicial sets and it corresponds to the Hom set in the original category.
- (b) If \mathcal{C} is an ∞ -category, then the unit adjunction $\mathcal{C} \rightarrow N(\pi\mathcal{C})$ of the (π, N) adjunction induces isomorphisms

$$\mathrm{Hom}_{N(\pi\mathcal{C})}(a, b) \cong \mathrm{const} \mathrm{Hom}_{\pi\mathcal{C}}(a, b) \cong \mathrm{const} \pi_0 \mathrm{Hom}_{\mathcal{C}}(a, b).$$

In other words, $\pi\mathcal{C}$ is the 1-category having \mathcal{C}_0 as objects and homotopy classes of \mathcal{C}_1 as morphisms.

I.11. Definition. — A ∞ -category \mathcal{C} is called an *∞ -groupoid*—or, in Scholze’s fancy new terminology, an *anima*—if $\pi\mathcal{C}$ is a groupoid.

- I.12. Example.** — (a) For a 1-category \mathcal{D} let $\mathrm{core} \mathcal{D}$ denote the (usually non-full) subcategory spanned by the isomorphisms in \mathcal{D} ; in particular, $\mathrm{core} \mathcal{D}$ is a groupoid. If \mathcal{C} is an ∞ -category, we define $\mathrm{core} \mathcal{C}$ by the pullback

$$\begin{array}{ccc} \mathrm{core} \mathcal{C} & \longrightarrow & \mathcal{C} \\ \downarrow & \lrcorner & \downarrow \\ N(\mathrm{core} \pi\mathcal{C}) & \longrightarrow & N(\pi\mathcal{C}) \end{array}$$

Then $\mathrm{core} \mathcal{C}$ is an anima. In fact, one can check that it is the largest anima contained in the ∞ -category \mathcal{C} .

- (b) If \mathcal{C} is an ∞ -category, then $\mathrm{Hom}_{\mathcal{C}}(a, b)$ is an anima for all $a, b \in \mathcal{C}$ (that’s not entirely obvious though).

It’s pretty easy to check that every Kan complex is an anima. Surprisingly, and that was one of the first hard theorems in ∞ -category theory, the converse holds as well!

I.13. Theorem (Joyal). — *Every ∞ -groupoid is in fact a Kan complex.*

*Proof**. This follows from Joyal’s lifting theorem, see [HC_I, Theorem VI.3.20]. \square

By now, we haven’t been able to produce any examples of ∞ -categories yet, safe for nerves of 1-categories (these guys don’t count). The first real source of interesting examples is given by coherent nerves of Kan enriched categories.

I.14. Theorem (Cordier–Porter). — *If $\mathcal{C} \in \text{Cat}^{\text{Kan}}$ is a category enriched in Kan complexes, then its coherent nerve $N^c(\mathcal{C})$ is an ∞ -category. Moreover, for all $a, b \in \mathcal{C}$ we have canonical homotopy equivalences*

$$\text{Hom}_{N^c(\mathcal{C})}(a, b) \simeq F_{\mathcal{C}}(a, b).$$

Here $F_{\mathcal{C}}$ denotes Kan complex of morphisms in the enriched category \mathcal{C} .

*Proof**. See [HC1, Theorem VII.19] (but be warned that stuff gets *technical*). \square

I.15. Example. — “My first ∞ -categories”:

- (a) The full subcategory $\text{Kan} \subseteq \text{sSet}$ is enriched over itself via $F_{\text{Kan}}(X, Y) = F(X, Y)$. Its coherent nerve $N^c(\text{Kan}) = \text{An}$ is called the *∞ -category of anima*.
- (b) The full subcategory $\text{qCat} \subseteq \text{sSet}$ is enriched over Kan via $F_{\text{qCat}}(\mathcal{C}, \mathcal{D}) = \text{core } F(\mathcal{C}, \mathcal{D})$. Its coherent nerve $N^c(\text{qCat}) = \text{Cat}_{\infty}$ is the *∞ -category of ∞ -categories*. Note that Fabian will usually write Cat instead of Cat_{∞} to not drag the index ∞ around all the time.
- (c) The category Cat_1 of all small 1-categories is canonically enriched in groupoids via $F_{\text{Cat}_1}(\mathcal{C}, \mathcal{D}) = \text{core } \text{Fun}(\mathcal{C}, \mathcal{D})$. This enrichment can be upgraded to a Kan enrichment: Namely, taking the nerves of all hom groupoids (which produces Kan complexes by Theorem I.13) induces a functor $N_*: \text{Cat}^{\text{Grpd}} \rightarrow \text{Cat}^{\text{Kan}}$. We thus obtain an ∞ -category $\text{Cat}_1^{(2)} = N^c(N_* \text{Cat}_1)$, called the *2-category of 1-categories*.¹ Similarly one can define a 2-category $\text{Grpd}_1^{(2)}$. We end up with a chain of inclusions

$$\begin{array}{ccccc} & & \text{An} & & \\ & \subseteq & & \subseteq & \\ N(\text{Set}) \subseteq \text{Grpd}_1^{(2)} & & & & \text{Cat}_{\infty} \\ & \subseteq & & \subseteq & \\ & & \text{Cat}_1^{(2)} & & \end{array}$$

All of these induce homotopy equivalences on mapping anima. In other words, they are *fully faithful*.

- (d) The category Top is Kan enriched via $F_{\text{Top}}(X, Y)_n = \text{Hom}_{\text{Top}}(X \times |\Delta^n|, Y)$, which provides the *∞ -category of topological spaces* after taking coherent nerves. However, be warned that $N^c(\text{Top}) \not\simeq \text{An}!!$
- (e) Let R be a ring and $\text{Ch}(R)$ the category of chain complexes over R . It is enriched over simplicial R -modules, and thus Kan-enriched by Example I.6, via

$$F_{\text{Ch}(R)}(C, D)_n = \text{Hom}_{\text{Ch}(R)}(C \otimes_{\mathbb{Z}} C_{\bullet}^{\text{simp}}(\Delta^n), D).$$

Here $C_{\bullet}^{\text{simp}}(\Delta^n)$ is a chain complex of abelian groups given as follows: In degree m we put $\mathbb{Z}[\{\text{non-degenerate } m\text{-simplices of } \Delta^n\}]$ and the differentials are given by $\partial_m = \sum_{i=0}^m (-1)^i d_i$, where d_i is induced by the corresponding face map $d_i: (\Delta^n)_m \rightarrow (\Delta^n)_{m-1}$, which preserves non-degenerate simplices (that doesn't hold for arbitrary simplicial sets

¹Here “2-category” means that $\pi_i \text{Hom}_{\text{Cat}_1}(\mathcal{C}, \mathcal{D}) = 0$ for $i \geq 2$. To prove this, use Theorem I.14 and check that nerves of groupoids have vanishing higher homotopy groups.

though). You probably recognize $C_{\bullet}^{\text{simp}}(X)$ as the complex that computes the simplicial homology groups $H_i^{\text{simp}}(X)$.

We let $\mathcal{K}(R) = \mathbf{N}^c(\text{Ch}(R))$ be the ∞ -category of chain complexes (this “ \mathcal{K} ” has nothing to do with “ K -theory”) and let $\mathcal{K}_{\geq 0}(R), \mathcal{K}_{\leq 0}(R) \subseteq \mathcal{K}(R)$ the full sub- ∞ -categories spanned by chain complexes concentrated in nonnegative/nonpositive degrees. Inside $\mathcal{K}_{\geq 0}(R)$, there is the full sub- ∞ -category $\mathcal{D}_{\geq 0}(R) \subseteq \mathcal{K}_{\geq 0}(R)$ spanned by the degreewise projective chain complexes in concentrated nonnegative degrees. This is called the *bounded below derived category* of R . Similarly, there is the *bounded above derived category* $\mathcal{D}_{\leq 0}(R) \subseteq \mathcal{K}_{\leq 0}(R)$ spanned by degreewise injective chain complexes concentrated in nonpositive degrees.

Finally, there is $\mathcal{D}^{\text{perf}}(R) \subseteq \mathcal{K}(R)$ which is the full sub- ∞ -category spanned by bounded complexes of degreewise finite projective modules. Once we’ve defined K -theory, we’ll eventually see that $K(R) = \text{Proj}(R)^{\infty\text{-gfp}} \simeq K(\mathcal{D}^{\text{perf}}(R))$ (and yes, this time, “ K ” is really the “ K ” from “ K -theory”).

I.15a. Exercise. — For complexes $C, D \in \text{Ch}(R)$ let $\underline{\text{Hom}}_R(C, D)$ denote the internal Hom in $\text{Ch}(R)$, so that $\underline{\text{Hom}}_R(C, -)$ is right-adjoint to $C \otimes_R -$ (see the [nLab article](#) for an explicit construction). Show that the Kan enrichment in Example I.15(e) satisfies

$$\pi_i \text{F}_{\text{Ch}(R)}(C, D) \cong H_i(\underline{\text{Hom}}(C, D)) \quad \text{for all } i \geq 0.$$

In particular, $\pi_0 \text{F}_{\text{Ch}(R)}(C, D)$ is the set of chain homotopy classes of maps $C \rightarrow D$.

I.16. Theorem (Dold–Kan). — *By a variant of Theorem I.2, the cosimplicial object C_{\bullet} in $\text{Ch}_{\geq 0}(\mathbb{Z})$ given by $[n] \mapsto C_{\bullet}^{\text{simp}}(\Delta^n)$ gives rise to mutually inverse equivalences of (ordinary!) categories*

$$| \quad |_{C_{\bullet}} : \text{sAb} \xrightleftharpoons{\sim} \text{Ch}_{\geq 0}(\mathbb{Z}) : \text{Sing}_{C_{\bullet}}.$$

These equivalences translates homotopy groups into chain homology groups. Moreover, we have $\text{F}_{\text{Ch}(\mathbb{Z})}(C, D) = \text{Sing}_{C_{\bullet}}(\underline{\text{Hom}}_{\mathbb{Z}}(C, D))$ for all $C, D \in \text{Ch}_{\geq 0}(\mathbb{Z})$.

I.17. Philosophical Nonsense I. — We obtain a chain of adjunctions

$$\text{Top} \xleftarrow[\text{Sing}]{| \quad |} \text{sSet} \xleftarrow[\text{forget}]{\mathbb{Z}[-]} \text{sAb} \xleftarrow[\text{Sing}_{C_{\bullet}}]{| \quad |_{C_{\bullet}}} \text{Ch}_{\geq 0}(R)$$

(all bottom arrows are right-adjoints, that’s why the direction of the arrows on the left is swapped). Going from Top to $\text{Ch}_{\geq 0}(\mathbb{Z})$ sends a topological space X to its singular chain complex $C_{\bullet}^{\text{sing}}(X)$. Going in the reverse direction sends a chain complex C to the corresponding generalized Eilenberg–MacLane space. In particular, if $C = A[n]$ is given by a single abelian group A sitting in degree n , then it is sent to $K(A, n)$.

I.18. Philosophical Nonsense II. — To describe what an unbounded chain complex is, it suffices to understand bounded below chain complexes. Namely, a chain complex $C \in \text{Ch}(R)$ can be described by the sequence of truncations $\tau_{\geq i}C \in \text{Ch}_{\geq i}(R)$. These are given by

$$\tau_{\geq i}C = \left(\dots \xrightarrow{\partial_{i+3}} C_{i+2} \xrightarrow{\partial_{i+2}} C_{i+1} \xrightarrow{\partial_{i+1}} \ker \partial_i \longrightarrow 0 \longrightarrow 0 \longrightarrow \dots \right)$$

(this is chosen in such a way that the canonical map $\tau_{\geq i}C \rightarrow C$ induces isomorphisms on homology in degrees $\geq i$). One easily checks the following relation with way to many brackets:

$$(\tau_{\geq i}C)[-i] \cong \tau_{\geq 0}\left(\left((\tau_{\geq i-1}C)[- (i-1)]\right)[-1]\right).$$

Note that under the Dold–Kan correspondence (Theorem I.16), the operation $C \mapsto \tau_{\geq 0}(C[-1])$ on $\text{Ch}_{\geq 0}(\mathbb{Z})$ corresponds to $X \mapsto \Omega_0 X$ on sAb . Here Ω_0 denotes the simplicial loop space (or at least some model for it) with base point $0 \in X_0$, i.e., the neutral element of the simplicial abelian group X . So if we write $X_i = |(\tau_{\geq i}C)[-i]|_{C_\bullet}$, then we obtain canonical isomorphisms $X_i \cong \Omega_0 X_{i-1}$. I wonder where I’ve seen that before ...

... so let’s define an *abelian spectrum* to be a sequence of simplicial abelian groups X_i together with isomorphisms $X_i \cong \Omega_0 X_{i-1}$. In that way, Theorem I.16 extends to an equivalence $\{\text{abelian spectra}\} \simeq \text{Ch}(\mathbb{Z})$. But that’s only the beginning of a long story ...

I.19. Theorem (Joyal). — *Let \mathcal{C} and \mathcal{D} be ∞ -categories.*

- (a) *A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ of ∞ -categories is an equivalence (i.e. there exists $G: \mathcal{D} \rightarrow \mathcal{C}$ and natural equivalences $F \circ G \simeq \text{id}_{\mathcal{D}}$ and $G \circ F \simeq \text{id}_{\mathcal{C}}$) if and only if it is essentially surjective and fully faithful (i.e. the induced map $F_*: \pi_0 \text{core } \mathcal{C} \rightarrow \pi_0 \text{core } \mathcal{D}$ is surjective and $F_*: \text{Hom}_{\mathcal{C}}(x, y) \rightarrow \text{Hom}_{\mathcal{D}}(F(x), F(y))$ is a homotopy equivalence of anima for all $x, y \in \mathcal{C}$.)*
- (b) *A natural transformation $\eta: F \Rightarrow G$ of functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$ is a natural equivalence (i.e. η is an isomorphism in the homotopy category $\pi \text{Fun}(\mathcal{C}, \mathcal{D})$) if and only if induces equivalences on 0-simplices, i.e. $\eta_x: F(x) \rightarrow G(x)$ is an equivalence for all $x \in \mathcal{C}$.*

*Proof**. See [HC_I, Theorem VII.1, Theorem VII.8]. □

In 1-category theory, Theorem I.19 is a triviality (up to axiom of choice business). But in ∞ -land it is *hard*. An ∞ -category contains infinitely more data than just a bunch of objects and morphism spaces. All the more surprising, that these actually suffice to characterize equivalences.

As an application, we get a precise formulation of Grothendieck’s conjectural connection between (∞) -groupoids and spaces that became famously known as the *homotopy hypothesis* (at that time, Grothendieck was already way beyond scribbling devils around his diagrams, living somewhere deep in the Pyrenees).

I.20. Corollary (Grothendieck’s homotopy hypothesis). — *Let CW denote the category of CW -complexes with its Kan enrichment given by $F_{\text{CW}}(X, Y) = \text{Sing Map}(X, Y)$. The functors*

$$| \cdot | : \text{An} \rightleftarrows \text{N}^c(\text{CW}) : \text{Sing}$$

are inverse equivalences.

*Proof**. See [HC_I, Corollary VII.6]. □

This finishes our recollection of Fabian’s Higher Categories I lecture and we start reviewing the material from Higher Categories II.

Cocartesian/Left Fibrations and Lurie’s Straightening Equivalence

The main goal of Fabian’s Higher Categories II lecture last semester was to construct a functor $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{An}$ and prove the ∞ -analogue of Yoneda’s lemma. We know what $\mathrm{Hom}_{\mathcal{C}}$ ought to do on 0-simplices of \mathcal{C} by Definition I.9, but making it into a functor is incredibly complicated—as is any attempt at defining functors into \mathbf{An} or \mathbf{Cat}_{∞} . The key to define such functors is Lurie’s straightening/unstraightening equivalence.

I.21. Theorem (Lurie). — *Let $*/\mathbf{An}$ denote (any model for) the slice category of anima under a point. To each functor $F: \mathcal{C} \rightarrow \mathbf{An}$ of ∞ -categories associate an ∞ -category $\mathrm{Un}(F)$ via the pullback*

$$\begin{array}{ccc} \mathrm{Un}(F) & \longrightarrow & */\mathbf{An} \\ \downarrow & \lrcorner & \downarrow \\ \mathcal{C} & \xrightarrow{F} & \mathbf{An} \end{array}$$

This association upgrades to a fully faithful functor $\mathrm{Un}: \mathrm{Fun}(\mathcal{C}, \mathbf{An}) \rightarrow \mathbf{Cat}_{\infty}/\mathcal{C}$ (the “unstraightening” functor), whose essential image is the full subcategory $\mathrm{Left}(\mathcal{C})$ spanned by the left fibrations over \mathcal{C} (to be defined in Definition I.24). We let $\mathrm{St}: \mathrm{Left}(\mathcal{C}) \rightarrow \mathrm{Fun}(\mathcal{C}, \mathbf{An})$ (“straightening”) denote an inverse.

Proof.* This follows from the combined effort of all we did in Higher Categories II, so let me randomly choose [HCII, Remark X.57(iii)] as a reference. \square

I.22. Example. — For any $x \in \mathcal{C}$, the straightening of the left fibration $x/\mathcal{C} \rightarrow \mathcal{C}$ is defined to be $\mathrm{Hom}_{\mathcal{C}}(x, -): \mathcal{C} \rightarrow \mathbf{An}$. That this makes sense can be seen as follows. Observe that the fibre of $\mathrm{Un}(F) \rightarrow \mathcal{C}$ over some $y \in \mathcal{C}$ is just $F(y)$, since the fibre of $*/\mathbf{An} \rightarrow \mathbf{An}$ over some anima X is just X itself. Conversely, $\mathrm{St}(p: \mathcal{E} \rightarrow \mathcal{C})(y) \simeq (\text{fibre of } G \text{ over } y)$ holds for all left fibrations $p: \mathcal{E} \rightarrow \mathcal{C}$. In particular, evaluating $\mathrm{Hom}_{\mathcal{C}}(x, -)$ at y gives the fibre of x/\mathcal{C} over y , which is indeed $\mathrm{Hom}_{\mathcal{C}}(x, y)$.

Theorem I.21 generalizes as follows.

I.23. Theorem (Lurie again, of course). — *For any ∞ -category \mathcal{C} , the equivalences from Theorem I.21 extend to inverse equivalences*

$$\mathrm{St}: \mathrm{Cocart}(\mathcal{C}) \xrightleftharpoons[\sim]{\sim} \mathrm{Fun}(\mathcal{C}, \mathbf{Cat}_{\infty}) : \mathrm{Un},$$

where $\mathrm{Cocart}(\mathcal{C})$ is the (non-full!) subcategory of $\mathbf{Cat}_{\infty}/\mathcal{C}$ spanned by cocartesian fibrations and maps preserving cocartesian edges (to be defined in Definition I.24).

Proof.* Again, I should really cite the entirety of Fabian’s previous lecture. The proof of the statement in question finishes in [HCII, Chapter X p. 113]. \square

There is also a dual version of Theorem I.23, which asserts that there is an equivalence $\mathrm{Cart}(\mathcal{C}) \simeq \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{\infty})$, induced by the *cartesian* straightening and unstraightening functors. Whenever there is room for confusion, we’ll use $\mathrm{St}^{\mathrm{cocart}}$, $\mathrm{Un}^{\mathrm{cocart}}$ for the cocartesian versions, and $\mathrm{St}^{\mathrm{cart}}$, $\mathrm{Un}^{\mathrm{cart}}$ for the cartesian versions.

With that out of the way, let’s define what this “cocartesian” thing actually is!

I.24. Definition. — Let $p: \mathcal{E} \rightarrow \mathcal{C}$ be a functor of ∞ categories.

- (a) Informally, an edge $f: x \rightarrow y$ in \mathcal{E} is called *p-cocartesian* if the following condition holds: For all edges $g: x \rightarrow z$ in \mathcal{E} and all fillers $\sigma: \Delta^2 \rightarrow \mathcal{C}$ as depicted in the right diagram below,

$$\begin{array}{ccc} x & \xrightarrow{g} & z \\ f \downarrow & \dashrightarrow & \downarrow \exists(!) \\ y & & \end{array} \quad \xrightarrow{p} \quad \begin{array}{ccc} p(x) & \xrightarrow{p(g)} & p(z) \\ p(f) \downarrow & \searrow \sigma & \downarrow \\ p(y) & & \end{array}$$

there exists a unique (up to contractible choice) filler $\tau: \Delta^2 \rightarrow \mathcal{E}$ satisfying $p(\tau) = \sigma$. In other words, the space of lifts of σ is contractible.

Somewhat more formally, $f: x \rightarrow y$ is *p-cocartesian* if for all $z \in \mathcal{E}$ the diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{E}}(y, z) & \xrightarrow{f^*} & \mathrm{Hom}_{\mathcal{E}}(x, z) \\ p_* \downarrow & & \downarrow p_* \\ \mathrm{Hom}_{\mathcal{C}}(p(y), p(z)) & \xrightarrow{p(f)^*} & \mathrm{Hom}_{\mathcal{C}}(p(x), p(z)) \end{array}$$

is a *homotopy pullback* diagram of anima. However, that doesn't mean the above is a pullback diagram taken inside the 1-category \mathbf{Kan} (or \mathbf{sSet})—in fact, it likely doesn't even commute on the nose, but only up to homotopy. And worse: The “precomposition” maps f^* and $p(f)^*$ are not even canonically defined. To construct f^* , one has to choose a “composition law” in \mathcal{E} , i.e. a section of the trivial fibration $\mathrm{Fun}(\Delta^2, \mathcal{E}) \rightarrow \mathrm{Fun}(\Lambda_1^2, \mathcal{E})$, but there is no canonical one.

Despite all these difficulties, there is a way to give a well-defined definition of homotopy pullbacks using model categories (see [HCII, Example VIII.49(vi)]). However, once we have the ∞ -categorical machinery available, there is a better definition: A diagram of anima is a *homotopy pullback* if it is a pullback diagram taken inside the ∞ -category \mathbf{An} ! The connection to the model-theoretic definition will be made precise in Theorem I.34 below.

- (b) Let $p\text{-Cocart} \subseteq \mathrm{Ar}(\mathcal{E})$ denote the full subcategory spanned by the cocartesian edges. Then p is called a *cocartesian fibration* if the dashed arrow q in the diagram below is an equivalence of ∞ -categories.

$$\begin{array}{ccccc} p\text{-Cocart} & & & & p \\ & \searrow q & & & \downarrow \\ & P & \xrightarrow{\quad} & \mathrm{Ar}(\mathcal{C}) & \\ & \downarrow & \lrcorner & \downarrow s & \\ & \mathcal{E} & \xrightarrow{p} & \mathcal{C} & \end{array}$$

Informally, the way to think about this is that all edges of \mathcal{C} admit a lift with given starting point. Also $p\text{-Cocart} \rightarrow P$ is automatically fully faithful, as we'll prove below.

- (c) We call p a *left fibration* if it is a cocartesian fibration and satisfies the following equivalent conditions.
- (i) $\mathrm{St}(p): \mathcal{C} \rightarrow \mathbf{Cat}_{\infty}$ factors through \mathbf{An} .

- (ii) The *derived fibres* of p are anima. By that we mean the fibres of $\mathcal{E}' \rightarrow \mathcal{C}$, where $\mathcal{E} \xrightarrow{\sim} \mathcal{E}' \twoheadrightarrow \mathcal{C}$ is any factorisation of p into an Joyal equivalence followed by fibration in the Joyal model structure. Or just take the fibres as usual, but with pullbacks inside the ∞ -category Cat_∞ .
- (iii) p is *conservative*, i.e. for all edges f in \mathcal{E} we have that f is an equivalence iff $p(f)$ is an equivalence in \mathcal{C} .
- (iv) All edges in \mathcal{E} are p -cocartesian.

There is also a dual notion of *cartesian edges/fibrations* and *right fibrations*. An edge of \mathcal{E} is p -cartesian if its opposite is a cocartesian edge with respect to $p^{\text{op}}: \mathcal{E}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$ (in other words, we have to reverse all the arrows in the informal part of (a)). Similarly, p is a cartesian/right fibration if p^{op} is a cocartesian/left fibration.

In the lecture we had a brief discussion about how Definition I.24 relates to the definition that the participants of Fabian's previous lectures would have expected. I'll give an expanded version of that discussion below. To be safe, everything is labelled with an asterisk to indicate it is an addition to the lecture.

I.24a. Warning* $\triangle!$. — Definition I.24 does not coincide with the way we defined cocartesian and left fibrations in Fabian's previous lectures. The main difference is that we don't require p to be an inner fibration, resulting in the advantage that Definition I.24 is invariant under Joyal equivalence (Fabian promised that this will save us some headache). Up to replacing $p: \mathcal{E} \rightarrow \mathcal{C}$ by an isofibration, both definitions agree, as the following Lemma* I.24b shows.

Also note that it doesn't suffice to replace p by an inner fibration. In fact, the inclusion $\{0\} \hookrightarrow J$ of a point into the free-living isomorphism is an inner fibration (as the nerve of a map of 1-categories) and a Joyal equivalence, hence it satisfies Definition I.24(b), but it is not an isofibration and thus not cocartesian in the old sense by [HCII, Proposition IX.2].

I.24b. Lemma*. — *The dashed arrow q in Definition I.24(b) is automatically fully faithful. In particular, for a functor $p: \mathcal{E} \rightarrow \mathcal{C}$ the following conditions are equivalent:*

- (a) p is a cocartesian/left fibration in the new sense.
- (b) For all factorisations $\mathcal{E} \xrightarrow{\sim} \mathcal{E}' \twoheadrightarrow \mathcal{C}$ into a Joyal equivalence followed by a fibration in the Joyal model structure, $\mathcal{E}' \rightarrow \mathcal{C}$ is a cocartesian/left fibration in the old sense.
- (c) The above condition holds for some factorisation $\mathcal{E} \xrightarrow{\sim} \mathcal{E}' \twoheadrightarrow \mathcal{C}$ as above.

Moreover, the four conditions from Definition I.24(c) are indeed equivalent.

*Proof**. Let \times^R denote derived pullbacks (or homotopy pullbacks, that's the same). Let $f: x \rightarrow y$ and $g: x' \rightarrow y'$ be cocartesian edges. Since $p\text{-Cocart} \subseteq \text{Ar}(\mathcal{E})$ is a full subcategory, we can use the calculation of Hom anima in arrow categories from [HCII, Proposition VIII.5] to obtain

$$\text{Hom}_{p\text{-Cocart}}(f, g) \simeq \text{Hom}_{\mathcal{E}}(x, x') \times_{\text{Hom}_{\mathcal{E}}(x, y')}^R \text{Hom}_{\mathcal{E}}(y, y').$$

Our definition of Hom anima in Definition I.9 commutes with pullbacks of simplicial sets (as long as the pullback still gives an ∞ -category), so we get

$$\text{Hom}_P(q(f), q(g)) = \text{Hom}_{\mathcal{E}}(x, x') \times_{\text{Hom}_{\mathcal{C}}(p(x), p(x'))} \text{Hom}_{\text{Ar}(\mathcal{C})}(p(f), p(g)).$$

This is a pullback on the nose, but also a homotopy pullback because $s: \text{Ar}(\mathcal{C}) \rightarrow \mathcal{C}$ is an inner fibration (even a cartesian fibration by the dual of Example I.25; this also implies

that P is an ∞ -category), hence $s_*: \text{Hom}_{\text{Ar}(\mathcal{C})}(p(f), p(g)) \rightarrow \text{Hom}_{\mathcal{C}}(p(x), p(x'))$ is a Kan fibration. Now replacing $\text{Hom}_{\mathcal{E}}(y, y')$ by the homotopy pullback from Definition I.24(a), $\text{Hom}_{\text{Ar}(\mathcal{C})}(p(f), p(g))$ by its description from [HCII, Proposition VIII.5], and simplifying the resulting homotopy pullbacks shows that $q_*: \text{Hom}_{p\text{-Cocart}}(f, g) \rightarrow \text{Hom}_P(q(f), q(g))$ is indeed an equivalence, hence q is fully faithful.

To prove equivalence of (a), (b), (c), note that Definition I.24(b) is invariant under Joyal equivalences, so we may assume $p: \mathcal{E} \rightarrow \mathcal{C}$ is a fibration in the Joyal model structure, or equivalently an isofibration since both are ∞ -categories. If p is cocartesian in the old sense, then $q: p\text{-Cocart} \rightarrow P$ is surjective on the nose, hence a Joyal equivalence by what we just proved. This is enough to prove (b) \Rightarrow (c) \Rightarrow (a). For (a) \Rightarrow (b), we may again assume p is an isofibration and we must show that q is surjective on the nose rather than just essentially surjective. We will even show that q is a trivial fibration. Since q is a Joyal equivalence, it suffices to show that it is an isofibration. Observe that $\text{Ar}(\mathcal{E}) \rightarrow P$ is an isofibration by [HC1, Corollary VII.11]. Hence its restriction q to the full subcategory $p\text{-Cocart} \subseteq \text{Ar}(\mathcal{E})$ is an inner fibration. But $p\text{-Cocart}$ is closed under equivalences in $\text{Ar}(\mathcal{E})$ (which is evident from the homotopy pullback condition in Definition I.24(a)), so an easy argument shows that q also inherits lifting against $\{0\} \hookrightarrow J$ from the isofibration $\text{Ar}(\mathcal{E}) \rightarrow P$.

Now that we know equivalence of (a), (b), (c) (except for the assertions about left fibrations, but these will follow immediately), it's easy to show that the conditions in Definition I.24 are indeed equivalent: Note that the values of $\text{St}(p)$ are given by the (derived) fibres of p , hence (i) \Leftrightarrow (ii), and (ii) \Leftrightarrow (iii) \Leftrightarrow (iv) follows from [HCII, Proposition IX.3] \square

I.25. Example. — “My first cocartesian edges/fibrations”:

- (a) Any equivalence in \mathcal{E} is p -cocartesian by Joyal's lifting theorem.
- (b) The “target morphism” $t: \text{Ar}(\mathcal{C}) \rightarrow \mathcal{C}$ (given by restriction along $\{1\} \hookrightarrow \Delta^1$) is a cocartesian fibration (in both the old and new sense since it is an isofibration). Let $f: x \rightarrow x'$ and $g: y \rightarrow y'$ be objects in $\text{Ar}(\mathcal{C})$. Then a morphism $\sigma: f \rightarrow g$ in $\text{Ar}(\mathcal{C})$, i.e. a commutative square

$$\begin{array}{ccc} x & \longrightarrow & y \\ f \downarrow & \sigma & \downarrow g \\ x' & \longrightarrow & y' \end{array}$$

in \mathcal{C} , is a t -cocartesian edge if and only if the induced morphism $\sigma_0: x \rightarrow y$ is an equivalence in \mathcal{C} . We will prove this in Lemma* I.25a below (Fabian gave the idea in the lecture, but some more details can't hurt).

This immediately implies that t is a cocartesian fibration, because for every object $f: x \rightarrow x'$ of $\text{Ar}(\mathcal{C})$ and all morphisms $x' \rightarrow y'$ in \mathcal{C} , we can take

$$\begin{array}{ccc} x & \xlongequal{\quad} & x \\ f \downarrow & & \downarrow \\ x' & \longrightarrow & y' \end{array}$$

as a t -cocartesian lift. However, t is usually not a left fibration. The straightening $\text{St}(t)$ is the slice category functor $\mathcal{C}/-: \mathcal{C} \rightarrow \text{Cat}_\infty$, which not necessarily factors over An .

As a fun fact, let us mention that $t: \text{Ar}(\mathcal{C}) \rightarrow \mathcal{C}$ is also a cartesian fibration, provided \mathcal{C} has pullbacks (we'll talk more about limits in ∞ -categories soon), which we leave as Exercise I.25b. Spoiler warning: I've also included my attempt at a solution there.

- (c) For all $x \in \mathcal{C}$, the slice category projection $\mathcal{C}/x \rightarrow \mathcal{C}$ is a cartesian fibration, and in fact even a right fibration, since its (derived or non-derived) fibre over y is the anima $\mathrm{Hom}_{\mathcal{C}}(y, x)$.

I.25a. Lemma*. — *With notation as in Example I.25(b), σ is t -cocartesian iff the induced morphism $\sigma_0: x \rightarrow y$ is an equivalence in \mathcal{C} .*

Proof.* Indeed, if $x \rightarrow y$ is an equivalence, then an easy calculation involving the characterisation of Hom anima in arrow categories ([HCII, Proposition VIII.5]) shows that σ satisfies the homotopy pullback condition from Definition I.24(a). Conversely, if σ is cocartesian, then any diagram

$$\begin{array}{ccccc}
 & & z & & \\
 & \nearrow & \downarrow h & \nwarrow & \\
 x & \xrightarrow{\quad} & y & & \\
 \downarrow f & \swarrow & \downarrow g & \searrow & \\
 & z' & & & \\
 & \nwarrow & \swarrow & & \\
 x' & \xrightarrow{\quad} & y' & &
 \end{array}$$

in which the front face is σ , and only the top and right back face are missing, admits a filler (at least up to equivalence of such diagrams). Choose the left back face λ as follows: Put $z = x$ and $z' = y'$, also choose $\lambda_0: x \rightarrow x$ as the identity on x , and $\lambda_1: x' \rightarrow y'$ as $\sigma_1: x' \rightarrow y'$, and $h: x \rightarrow y'$ as a composition of f and σ_1 in \mathcal{C} . Then a filler of the above diagram provides a left inverse of $x \rightarrow y$. Moreover, the morphism $\rho: g \rightarrow h$ in $\mathrm{Ar}(\mathcal{C})$ induced by the right back face ρ is again t -cocartesian. This is because λ is a composition of ρ and σ , and both σ and λ are t -cocartesian (for λ this follows from $x \rightarrow z$ being the identity on x , hence an equivalence), so [HCII, Proposition IX.5] can be applied. Therefore, we may repeat the all the arguments for ρ instead of σ , which shows that the left inverse of $\sigma_0: x \rightarrow y$ has a left inverse itself, whence σ_0 must indeed be an equivalence. \square

I.25b. Exercise. — Show that \mathcal{C} admits pullbacks (in the ∞ -categorical sense defined in Example I.33(b) below) iff the cocartesian fibration $t: \mathrm{Ar}(\mathcal{C}) \rightarrow \mathcal{C}$ is also a cartesian fibration. More generally, the t -cartesian edges are given precisely by the pullback squares.

Proof sketch.* If the claimed characterisation of cartesian edges is true, then the lifting condition for t to be a cartesian fibration is satisfied if and only if \mathcal{C} has all pullbacks. So it suffices to show the second assertion.

We'll freely use the yoga of limits in ∞ -categories which is developed from Example I.33(b) onward. By Corollary I.50, $\mathrm{Hom}_{\mathcal{C}}(z, -): \mathcal{C} \rightarrow \mathbf{An}$ preserves all limits for every $z \in \mathcal{C}$. Also limits in \mathbf{An} can be computed by homotopy limits thanks to Theorem I.34 below.

Now let $f: x \rightarrow x'$ and $g: y \rightarrow y'$ be objects of $\mathrm{Ar}(\mathcal{C})$ and let $\sigma: f \rightarrow g$ be an edge in $\mathrm{Ar}(\mathcal{C})$. Then σ is cartesian iff for all $h: z \rightarrow z'$ we can compute $\mathrm{Hom}_{\mathrm{Ar}(\mathcal{C})}(h, f)$ by a certain homotopy pullback dual to that in Definition I.24(a). Writing that condition out explicitly and simplifying using the description of Hom anima in arrow categories given in [HCII, Proposition VIII.5], we obtain the condition

$$\mathrm{Hom}_{\mathcal{C}}(z, x) \times_{\mathrm{Hom}_{\mathcal{C}}(z, x')}^R \mathrm{Hom}_{\mathcal{C}}(z', x') \simeq \mathrm{Hom}_{\mathcal{C}}(z, y) \times_{\mathrm{Hom}_{\mathcal{C}}(z, y')}^R \mathrm{Hom}_{\mathcal{C}}(z', y').$$

If $z = z'$ and $h = \mathrm{id}_z$, then this condition precisely says that x is a pullback of $y \rightarrow y' \leftarrow x'$ by our remark on $\mathrm{Hom}_{\mathcal{C}}(z, -)$ in the previous paragraph. We should perhaps explain why

it suffices to check that on objects $z \in \mathcal{C}$. This is because one has, in fact, a natural transformation $\mathrm{Hom}_{\mathcal{C}}(-, x) \Rightarrow \mathrm{Hom}_{\mathcal{C}}(-, y) \times_{\mathrm{Hom}_{\mathcal{C}}(-, y')}^R \mathrm{Hom}_{\mathcal{C}}(-, x')$ of functors $\mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{An}$ (to construct this one has to dive into the straightening construction), and whether this is an equivalence can indeed be checked object-wise by Theorem I.19. Conversely, if x is a pullback of $y \rightarrow y' \leftarrow x'$, one easily checks that the above equivalence holds. \square

I.26. Constructing the functor $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{An}$. — There are at least four equivalent ways to do so.

- (a) Recall $\mathbf{An} = \mathbf{N}^c(\mathbf{Kan})$, so $\mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{An}$ corresponds to a simplicially enriched functor $\mathfrak{C}[\mathcal{C}^{\mathrm{op}} \times \mathcal{C}] \rightarrow \mathbf{Kan}$. This is provided by the commutative square

$$\begin{array}{ccc} \mathfrak{C}[\mathcal{C}^{\mathrm{op}} \times \mathcal{C}] & \longrightarrow & \mathbf{Kan} \\ \downarrow & & \uparrow \mathrm{Ex}^{\infty} \\ \mathfrak{C}[\mathcal{C}]^{\mathrm{op}} \times \mathfrak{C}[\mathcal{C}] & \xrightarrow{F_{\mathfrak{C}[\mathcal{C}]}} & \mathbf{sSet} \end{array}$$

where as usual $F_{\mathfrak{C}[\mathcal{C}]}: \mathfrak{C}[\mathcal{C}]^{\mathrm{op}} \times \mathfrak{C}[\mathcal{C}] \rightarrow \mathbf{sSet}$ denotes the simplicial set of morphisms in the simplicially enriched category $\mathfrak{C}[\mathcal{C}]$. Instead of Ex^{∞} we could use any functorial weak equivalence into a Kan complexes; for example, Lurie uses the functor $X \mapsto \mathrm{Sing}|X|$ in [HTT, §5.1.3].

- (b) We construct the corresponding left fibration to $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$ explicitly: It is the *twisted arrow category* $\mathrm{TwAr}(\mathcal{C})$ defined by $\mathrm{TwAr}(\mathcal{C})_n = \mathrm{Hom}_{\mathbf{sSet}}((\Delta^n)^{\mathrm{op}} \star \Delta^n, \mathcal{C})$. The functorial inclusions $(\Delta^n)^{\mathrm{op}} \sqcup \Delta^n \subseteq (\Delta^n)^{\mathrm{op}} \star \Delta^n$ induce a projection

$$\mathrm{TwAr}(\mathcal{C}) \longrightarrow \mathcal{C}^{\mathrm{op}} \times \mathcal{C},$$

which turns out to be a left fibration with the correct fibres. See [HA, §5.2.1] for proofs. The twisted arrow category construction of $\mathrm{Hom}_{\mathcal{C}}$ is the one that is most commonly used today.

- (c) Fabian's favourite construction considers the commutative diagram

$$\begin{array}{ccc} \mathrm{Ar}(\mathcal{C}) & \xrightarrow{(s,t)} & \mathcal{C} \times \mathcal{C} \\ & \searrow t & \swarrow \mathrm{pr}_2 \\ & \mathcal{C} & \end{array}$$

which is a morphism in $\mathrm{Cocart}(\mathcal{C})$ (in particular, (s, t) preserves cocartesian edges) by Example I.25(b) and the following simple fact: For all ∞ -categories \mathcal{C} and \mathcal{D} , the projection $\mathrm{pr}_2: \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{D}$ is cartesian and cocartesian, and the pr_2 -cartesian/cocartesian edges are those of the form (equivalence in \mathcal{C} , arbitrary morphism in \mathcal{D}).

Applying the cocartesian straightening construction gives a natural transformation $\mathrm{St}^{\mathrm{cocart}}(t) \Rightarrow \mathrm{const} \mathcal{C}$ of functors $\mathcal{C} \rightarrow \mathbf{Cat}_{\infty}$ by Theorem I.23. This natural transformation defines a map $\Delta^1 \rightarrow \mathrm{Fun}(\mathcal{C}, \mathbf{Cat}_{\infty})$, hence a functor $S: \mathcal{C} \rightarrow \mathrm{Ar}(\mathbf{Cat}_{\infty})$ by adjoining. Informally, S takes $x \in \mathcal{C}$ to the object $\mathcal{C}/x \rightarrow \mathcal{C}$ in the arrow category of \mathbf{Cat}_{∞} . In particular, these are right fibrations by Example I.25(c) and the target is always \mathcal{C} , whence S factors through $\mathrm{Right}(\mathcal{C}) \rightarrow \mathrm{Ar}(\mathbf{Cat}_{\infty})$. We denote the resulting functor $S: \mathcal{C} \rightarrow \mathrm{Right}(\mathcal{C})$ as well. Composing S with $\mathrm{St}^{\mathrm{cart}}: \mathrm{Right}(\mathcal{C}) \xrightarrow{\sim} \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{An})$, the cartesian straightening equivalence, finally provides a functor

$$Y: \mathcal{C} \longrightarrow \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{An})$$

(the *Yoneda embedding*), which gives the desired $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{An}$ after “currying” (which is how Fabian sometimes calls the adjunction $\mathrm{Fun}(\mathcal{C}, \mathrm{Fun}(\mathcal{D}, \mathcal{E})) \simeq \mathrm{Fun}(\mathcal{C} \times \mathcal{D}, \mathcal{E})$).

(d) You can also dualize all steps in (c) and start with

$$\begin{array}{ccc} \mathrm{Ar}(\mathcal{C}) & \xrightarrow{(s,t)} & \mathcal{C} \times \mathcal{C} \\ & \searrow s \quad \swarrow \mathrm{pr}_1 & \\ & \mathcal{C} & \end{array}$$

I.27. Theorem. — *All constructions in I.26 give the same functor $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{An}$ up to equivalence.*

We won’t prove Theorem I.27. Lurie shows that I.26(a) and (b) are equivalent in [HA, Proposition 5.2.1.11], and for equivalence of (c) and (d) you probably shouldn’t go for a direct proof, but show that either one is equivalent to (b). Fabian believes that if one works with any of these constructions long enough, it will become obvious that the others are equivalent.

I.28. More on the twisted arrow category. — By I.26(b) and Theorem I.21, the twisted arrow category of \mathcal{C} sits inside a pullback (up to equivalence, or take the pullback in Cat_{∞})

$$\begin{array}{ccc} \mathrm{TwAr}(\mathcal{C}) & \longrightarrow & */\mathbf{An} \\ \downarrow & \lrcorner & \downarrow \\ \mathcal{C}^{\mathrm{op}} \times \mathcal{C} & \xrightarrow{\mathrm{Hom}_{\mathcal{C}}} & \mathbf{An} \end{array}$$

(if you don’t want to use I.26(b) to construct $\mathrm{Hom}_{\mathcal{C}}$, this pullback square can also be taken as a definition of $\mathrm{TwAr}(\mathcal{C})$).

Unwinding, this shows that the objects of $\mathrm{TwAr}(\mathcal{C})$ are given by pairs $(x, y) \in \mathcal{C}^{\mathrm{op}} \times \mathcal{C}$ together with a map $f: \Delta^0 \rightarrow \mathrm{Hom}_{\mathcal{C}}(x, y)$. In other words, objects are arrows $f: x \rightarrow y$ in \mathcal{C} . Informally, an edge $f \rightarrow g$ in $\mathrm{TwAr}(\mathcal{C})$ between $f: x \rightarrow y$ and $g: x' \rightarrow y'$ is given by a “twisted square” $(\Delta^1)^{\mathrm{op}} \star \Delta^1 \rightarrow \mathcal{C}$ and a 2-simplex $\Delta^2 \rightarrow \mathbf{An}$ as follows:


$$\begin{array}{ccc} x \xleftarrow{m_s} x' & & \mathrm{Hom}_{\mathcal{C}}(x, y) \xrightarrow{m_s^* \circ m_t, *} \mathrm{Hom}_{\mathcal{C}}(x', y') \\ f \downarrow & \text{and} & \swarrow f \quad \searrow g \\ y \xrightarrow{m_t} y' & & \Delta^0 \end{array}$$

More formally, since $\mathrm{TwAr}(\mathcal{C})$ sits in the pullback above, $\mathrm{Hom}_{\mathrm{TwAr}(\mathcal{C})}(f, g)$ can be computed as the pullback of the corresponding Hom anima in $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$, \mathbf{An} , and $*/\mathbf{An}$. This pullback is also a homotopy pullback, as $*/\mathbf{An} \rightarrow \mathbf{An}$ is an inner fibration (even a left fibration in the old sense), hence induces Kan fibrations on Hom anima. Plugging in the calculation of Hom anima in slice categories from [HCII, Corollary VIII.6] thus provides a homotopy pullback


$$\begin{array}{ccc} \mathrm{Hom}_{\mathrm{TwAr}(\mathcal{C})}(f, g) & \longrightarrow & \{g\} \\ \downarrow & \lrcorner_h & \downarrow \\ \mathrm{Hom}_{\mathcal{C}}(x', x) \times \mathrm{Hom}_{\mathcal{C}}(y, y') & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(x', y') \end{array}$$

where the bottom arrow sends $(m_s, m_t) \mapsto m_t \circ f \circ m_s$ for some choice of composition.

The twisted arrow category will be important to us since it has to do with *spans* in \mathcal{C} . This becomes apparent when we look at $\mathrm{TwAr}(\Delta^n)$. Its objects are arrows in Δ^n . We organize these arrows in an upwards pointing triangle with rows numbered $0, 1, \dots, n$ from the bottom to the top: In the i^{th} row, we put the morphisms that “go up by i ”, i.e., $(0 \rightarrow i), (1 \rightarrow i+1), \dots, (n-i \rightarrow n)$. In particular, the bottom row contains the identities and the top row contains only the morphism $(0 \rightarrow n)$. If we draw in some of the edges in $\mathrm{TwAr}(\mathcal{C})$ (every other edge can be obtained as a composite of these), we obtain a picture as follows:



And voilà, plenty of spans! The corresponding picture for $\mathrm{Ar}(\Delta^n)$ is instead the following:



As a bit of foreshadowing, Fabian mentioned that the first picture is related to the Quillen Q -construction (which will come up later, see Example I.68(d)), whereas the second is related to the Waldhausen S -construction (which we probably won't discuss in the course).

I.28a. Exercise. — Show that $\mathrm{core} \mathrm{TwAr}(\mathcal{C}) \simeq \mathrm{core} \mathrm{Ar}(\mathcal{C})$ holds for all ∞ -categories \mathcal{C} .

Proof sketch. See [A&HK, Proposition I.31] for a rough sketch. \square

Yoneda's Lemma, Limits and Colimits in ∞ -Categories

Yoneda's Lemma and Adjunctions

Now that we have a sufficient supply of mutually equivalent definitions of the Hom functor, the next step is to prove Yoneda's lemma.

I.29. Theorem (Yoneda's lemma). — *Let \mathcal{C} be an ∞ -category. Given a functor $F: \mathcal{C} \rightarrow \mathrm{An}$ and an object $x \in \mathcal{C}$, the evaluation map*

$$\mathrm{ev}_{\mathrm{id}_x}: \mathrm{Nat}(\mathrm{Hom}_{\mathcal{C}}(x, -), F) \xrightarrow{\sim} F(x)$$

is an equivalence (we use Nat as a shortcut for $\mathrm{Hom}_{\mathrm{Fun}(\mathcal{C}, \mathrm{An})}$). Moreover, adjoining the functor $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathrm{Fun}(\mathcal{C}, \mathrm{An})$ gives fully faithful functors (“Yoneda embeddings”)

$$Y^{\mathcal{C}}: \mathcal{C} \longrightarrow \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{An}) \quad \text{and} \quad Y_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \longrightarrow \mathrm{Fun}(\mathcal{C}, \mathrm{An}).$$

*Proof**. See [HCII, Corollaries XI.2 and XI.4]. \square

I.30. Definition. — For an ∞ -category \mathcal{C} we denote by $\mathcal{P}(\mathcal{C}) = \text{Fun}(\mathcal{C}^{\text{op}}, \text{An})$ the ∞ -category of presheaves on \mathcal{C} .

Onwards to adjunctions!

I.31. Definition. — Let $R: \mathcal{D} \rightarrow \mathcal{C}$ be a functor of ∞ -categories.

- (a) Given objects $y \in \mathcal{C}$, $x \in \mathcal{D}$, and a morphism $\eta: y \rightarrow Rx$ in \mathcal{C} , we say η *witnesses* x as a *left-adjoint object to y under R* if the composite

$$\text{Hom}_{\mathcal{D}}(x, -) \xrightarrow{R} \text{Hom}_{\mathcal{C}}(Rx, R-) \xrightarrow{\eta^*} \text{Hom}_{\mathcal{C}}(y, R-)$$

is an equivalence of functors $\mathcal{D} \rightarrow \text{An}$ (which may be tested on objects by Theorem I.19(b)).

- (b) An *adjunction* between R and some functor $L: \mathcal{C} \rightarrow \mathcal{D}$ is an equivalence

$$\text{Hom}_{\mathcal{D}}(L-, -) \simeq \text{Hom}_{\mathcal{C}}(-, R-)$$

as functors $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \text{An}$.

A simple but still somewhat surprising and incredibly useful consequence of Yoneda's lemma is that to define left-adjoint functors, it suffices to do so on objects (something that is wildly false for arbitrary functors)!

I.32. Corollary. — A functor $R: \mathcal{D} \rightarrow \mathcal{C}$ of ∞ -categories admits a left adjoint if and only if every $y \in \mathcal{C}$ admits a left-adjoint object under R . More generally, if $\mathcal{C}_R \subseteq \mathcal{C}$ is the full subcategory spanned by those objects $y \in \mathcal{C}$ which admit a left-adjoint object, extracting these left-adjoint objects defines a functor

$$L: \mathcal{C}_R \longrightarrow \mathcal{D}.$$

Proof. We repeat the proof given in [HCII, Lemma XI.6]. Via currying, the functor $\text{Hom}_{\mathcal{C}}(-, R-): \mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \text{An}$ corresponds to a functor $H: \mathcal{C}^{\text{op}} \rightarrow \text{Fun}(\mathcal{D}, \text{An})$. When restricted to $\mathcal{C}_R^{\text{op}}$, it lands in the representables, i.e., in the essential image of the Yoneda embedding $Y_{\mathcal{D}}: \mathcal{D}^{\text{op}} \rightarrow \text{Fun}(\mathcal{D}, \text{An})$. By Theorem I.29, $Y_{\mathcal{D}}$ is fully faithful, hence an equivalence onto its essential image by Theorem I.19(a). Composing $H|_{\mathcal{C}_R^{\text{op}}}$ with an inverse of this equivalence gives a functor $\mathcal{C}_R^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$. Taking $(-)^{\text{op}}$ we get $L: \mathcal{C}_R \rightarrow \mathcal{D}$ as required.

By construction, $Y_{\mathcal{D}} \circ L^{\text{op}}$ is equivalent to $H|_{\mathcal{C}_R^{\text{op}}}$ in $\text{Fun}(\mathcal{C}_R^{\text{op}}, \text{Fun}(\mathcal{D}, \text{An}))$. But this means that $\text{Hom}_{\mathcal{D}}(L-, -)$ and $\text{Hom}_{\mathcal{C}}(-, R-)$ are equivalent in $\text{Fun}(\mathcal{C}_R^{\text{op}} \times \mathcal{D}, \text{An})$. If $\mathcal{C}_R = \mathcal{C}$, this proves that L is indeed a left adjoint of R . \square

I.33. Example. — “My first adjoint pairs of functors”:

- (a) The inclusion $\text{An} \subseteq \text{Cat}_{\infty}$ has both a left adjoint, the functor $| \cdot |: \text{Cat}_{\infty} \rightarrow \text{An}$ (which sends an ∞ -category to its localisation at all its morphisms), and a right adjoint, given by $\text{core}: \text{Cat}_{\infty} \rightarrow \text{An}$. We proved this in [HCII, Example XI.8]. Moreover, the inclusion $\text{Set} \rightarrow \text{An}$ has $\pi_0: \text{An} \rightarrow \text{Set}$ as a left adjoint. In pictures,

$$\begin{array}{ccccc} & & \pi_0 & & | \cdot | \\ & \swarrow & \text{---} & \nwarrow & \text{---} \\ \text{Set} & \subseteq & \text{An} & \subseteq & \text{Cat}_{\infty} \\ & & \text{core} & & \end{array}$$

Of course we should actually write $N(\text{Set})$ instead of Set , but from now on we'll frequently abuse notation that way since writing N all the time will inevitably get on our ... *nerves* (sorry).

- (b) Let \mathcal{I} be any simplicial set (which we think of as a “diagram shape”) and consider the functor $\text{const}: \mathcal{C} \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$. Given a map $F: \mathcal{I} \rightarrow \mathcal{C}$, a left-adjoint object to F under const is called a *colimit* of F and denoted $\text{colim}_{\mathcal{I}} F$. Similarly, a right-adjoint object to F is called a *limit* and denoted $\text{lim}_{\mathcal{I}} F$. In particular, one has

$$\text{Hom}_{\mathcal{C}} \left(\text{colim}_{\mathcal{I}} F, x \right) \simeq \text{Nat}(F, \text{const } x)$$

for all objects $x \in \mathcal{C}$, and similarly for $\text{lim}_{\mathcal{I}} F$. We will discuss limits and colimits in detail in the next subsection.

First Properties and Examples of (Co)Limits

First we'll discuss the main existence theorem for limits and colimits in ∞ -categories. For that matter, recall the notion of *Kan simplicial model categories* from [HCII, Digression IV Definition G.2]. For a (not necessarily Kan simplicial) model category A , we denote by A^{cf} its full subcategory of bifibrant objects (Lurie uses the notation A° instead). Also $(-)^c: A \rightarrow A$ and $(-)^f: A \rightarrow A$ denote any choice of cofibrant/fibrant replacement functors.

I.34. Theorem. — *If A is a Kan simplicial model category, then $N^c(A^{\text{cf}})$ has all limits and colimits. For a diagram $F: \mathcal{I} \rightarrow N^c(A^{\text{cf}})$, these are given by*

$$\text{colim}_{\mathcal{I}} F \simeq \left(\text{hocolim}_{\mathfrak{C}[\mathcal{I}]} \tilde{F} \right)^f \quad \text{and} \quad \text{lim}_{\mathcal{I}} F \simeq \left(\text{holim}_{\mathfrak{C}[\mathcal{I}]} \tilde{F} \right)^c,$$

where $\tilde{F}: \mathfrak{C}[\mathcal{I}] \rightarrow A^{\text{cf}}$ is adjoint to $F: \mathcal{I} \rightarrow N^c(A^{\text{cf}})$.

*Proof**. See [HCII, Theorem XI.21]. □

I.34a. Remark*. — We'll soon talk about cofinal and final maps (see Definition I.44 or [HTT, §4.1]), but let me already mention that the theory of these allows us to assume that all diagram shapes \mathcal{I} are ∞ -categories. This hardly ever matters, but we formulated Theorem I.23 for ∞ -categories only, so I sleep better knowing that \mathcal{I} can be chosen that way.

I.35. Example. — Applying Theorem I.34 to $A = \text{sSet}$ with its Kan–Quillen model structure or $A = \text{sSet}^+$ (the category of marked simplicial sets) with its marked Joyal model structure, which are Kan simplicial model categories by [HCII, Digression IV Examples G.3], shows that An and Cat_{∞} are complete and cocomplete (i.e. have all limits and colimits).

Another example of a complete and cocomplete ∞ -category is $\mathcal{K}(R)$ for any ring R , as defined in Example I.15(e). In this case though, we can't apply Theorem I.34, since $\mathcal{K}(R)$ is (probably) not given by a Kan simplicial model structure on $\text{Ch}(R)$, at least not for the projective or injective model structure (these have quasi-isomorphisms as weak equivalences, whereas $\mathcal{K}(R)$ wants homotopy equivalences). Instead, one has to use some more properties of limits and colimits, which we will develop until Exercise I.50b.

I.35a. Example*. — Per Bastiaan's suggestion, we give two more examples. Recall that a pullback on the nose in a model category is also a homotopy pullback if all objects are fibrant and one of its legs is a fibration (this follows from Reedy's lemma for example, see [HCII, Corollary VIII.52]).

- (a) Combining this observation with Theorem I.34 shows that the pullback defining $\mathrm{Hom}_{\mathcal{C}}(a, b)$ from Definition I.9 is also a pullback in Cat_{∞} , because its vertical leg $\mathrm{Fun}(\Delta^1, \mathcal{C}) \rightarrow \mathrm{Fun}(\partial\Delta^1, \mathcal{C}) \simeq \mathcal{C} \times \mathcal{C}$ is an isofibration (by [HC_I, Corollary VII.11]), hence a fibration in the Joyal model structure on \mathbf{sSet} .
- (b) Likewise, the pullback defining $\mathrm{Un}(F)$ in Theorem I.21 is a pullback in Cat_{∞} , because $\ast/\mathrm{An} \rightarrow \mathrm{An}$ is a left fibration, hence an isofibration.

For a cocartesian fibration $p: E \rightarrow S$, we denote by $\Gamma(p)$ its ∞ -category of sections defined by the pullback

$$\begin{array}{ccc} \Gamma(p) & \longrightarrow & \mathrm{Fun}(S, E) \\ \downarrow & \lrcorner & \downarrow p_{\ast} \\ \{\mathrm{id}_S\} & \longrightarrow & \mathrm{Fun}(S, S) \end{array}$$

If we use the new notion from Definition I.24(b) and p is not necessarily an isofibration, we need to take this pullback in Cat_{∞} , otherwise it can also be taken in \mathbf{sSet} by the arguments from Example* I.35a above. We let $\Gamma_{\mathrm{cocart}}(p) \subseteq \Gamma(p)$ denote the full sub- ∞ -category spanned by sections that take all edges in S to p -cocartesian edges.

I.36. Proposition (Lurie). — *Given a diagram $F: \mathcal{I} \rightarrow \mathrm{Cat}_{\infty}$, we have*

$$\begin{aligned} \mathrm{colim}_{\mathcal{I}} F &\simeq \mathrm{Un}^{\mathrm{cocart}}(F) [\{\mathrm{cocartesian\ edges}\}^{-1}] , \\ \lim_{\mathcal{I}} F &\simeq \Gamma_{\mathrm{cocart}}(\mathrm{Un}^{\mathrm{cocart}}(F)) . \end{aligned}$$

In particular, if $F: \mathcal{I} \rightarrow \mathrm{An}$ takes values in anima, then

$$\mathrm{colim}_{\mathcal{I}} F \simeq |\mathrm{Un}(F)| \quad \text{and} \quad \lim_{\mathcal{I}} F \simeq \Gamma(\mathrm{Un}(F)) .$$

Proof sketch. In [HC_{II}, Theorem XI.23] Fabian gave a proof that's basically the same as what we are going to do now, but more on the model category side (but also less sketchy). We only do the case of limits. Consider the diagram

$$\begin{array}{ccc} & & \mathrm{Fun}(\mathcal{I}, \mathrm{Cat}_{\infty}) \\ & \nearrow \mathrm{const} & \uparrow \\ \mathrm{Cat}_{\infty} & & \mathrm{St} \wr \\ & \searrow & \downarrow \\ & & \mathrm{Cocart}(\mathcal{I}) \end{array}$$

in which the bottom arrow sends $\mathcal{C} \in \mathrm{Cat}_{\infty}$ to the cocartesian fibration $\mathrm{pr}_2: \mathcal{C} \times \mathcal{I} \rightarrow \mathcal{I}$ over \mathcal{I} . Let $p: E \rightarrow \mathcal{I}$ denote the cocartesian unstraightening of F . If $X \simeq \lim_{\mathcal{I}} F$, then we must have equivalences $\mathrm{Nat}(\mathrm{const} \mathcal{C}, F) \simeq \mathrm{Hom}_{\mathrm{Cat}_{\infty}}(\mathcal{C}, X)$ for all ∞ -categories \mathcal{C} . Applying Theorem I.23, this translates into

$$\mathrm{Hom}_{\mathrm{Cocart}(\mathcal{I})}((\mathrm{pr}_2: \mathcal{C} \times \mathcal{I} \rightarrow \mathcal{I}), (p: E \rightarrow \mathcal{I})) \simeq \mathrm{Hom}_{\mathrm{Cat}_{\infty}}(\mathcal{C}, X) ,$$

which we'll prove now is fulfilled for $X \simeq \Gamma_{\mathrm{cocart}}(p)$ (or rather we give an idea why it holds). A map $\varphi: \mathcal{C} \times \mathcal{I} \rightarrow E$ corresponds to a map $\widehat{\varphi}: \mathcal{C} \rightarrow \mathrm{Fun}(\mathcal{I}, E)$ via currying. By inspection, φ commutes with the projection down to \mathcal{I} iff $\widehat{\varphi}$ factors over $\Gamma(p) \rightarrow E$. Since pr_2 -cocartesian edges are precisely those of the form (equivalence in \mathcal{C} , arbitrary morphism in \mathcal{I}),

we see that φ preserves cocartesian edges iff $\widehat{\varphi}$ sends equivalences $f: c \rightarrow c'$ in \mathcal{C} to natural transformations $\eta: \widehat{\varphi}(c) \Rightarrow \widehat{\varphi}(c') \in \text{Fun}(\mathcal{I}, E)_1$ such that for all edges $g: i \rightarrow i'$ in \mathcal{I} the composition (or some choice of it)

$$\widehat{\varphi}(c)(i) \xrightarrow{\eta_i} \widehat{\varphi}(c')(i) \xrightarrow{g_*} \widehat{\varphi}(c')(i')$$

is p -cocartesian in E . Since $f: c \rightarrow c'$ is an equivalence in \mathcal{C} , we get that η_i is an equivalence in the fibre $E \times_{\mathcal{I}} \{i\}$ (even in E), hence automatically p -cocartesian by [HCII, Corollary IX.8]. Thus the composition is p -cocartesian iff g_* is p -cocartesian, in other words, if $\widehat{\varphi}(c')$ sends all morphisms in \mathcal{I} to p -cocartesian edges. Since this holds for all $c' \in \mathcal{C}$ (we can always take $f = \text{id}_{c'}$), we finally obtain that φ preserves pr_2 -cocartesian edges iff $\widehat{\varphi}: \mathcal{C} \rightarrow \Gamma(p)$ factors over $\Gamma_{\text{cocart}}(p)$. “Thus” $X \simeq \Gamma_{\text{cocart}}(p)$.

So far, this is no proof at all, since we only explained why 0-simplices of $\text{Nat}(\text{const } \mathcal{C}, F)$ correspond to 0-simplices of $\text{Hom}_{\text{Cat}_{\infty}}(\mathcal{C}, \Gamma_{\text{cocart}}(p))$ but it's not clear how to proceed for higher simplices, nor how to make this an ∞ -natural transformation. To fix this, first note that we have a map of cocartesian fibrations

$$\begin{array}{ccc} \Gamma_{\text{cocart}}(p) \times \mathcal{I} & \longrightarrow & E \\ \text{pr}_2 \downarrow & & \downarrow p \\ \mathcal{I} & \xlongequal{\quad} & \mathcal{I} \end{array}$$

induced by the evaluation map $\text{Fun}(\mathcal{I}, E) \times \mathcal{I} \rightarrow E$. After straightening, this gives a natural transformation $\eta: \text{const } \Gamma_{\text{cocart}}(p) \Rightarrow F$, hence we get

$$\text{Hom}_{\text{Cat}_{\infty}}(-, \Gamma_{\text{cocart}}(p)) \xrightarrow{\text{const } *} \text{Nat}(\text{const } -, \text{const } \Gamma_{\text{cocart}}(p)) \xrightarrow{\eta_*} \text{Nat}(\text{const } -, F),$$

which is the desired natural transformation. Whether this is an equivalence can be checked pointwise. Since instead of $\text{Nat}(\text{const } -, F)$ we can take the corresponding Hom anima in $\text{Cocart}(\mathcal{I})$, we arrive at the equivalence from the beginning of the proof. Since we know how to compute Hom spaces in Cat_{∞} and in $\text{Cat}_{\infty}/\mathcal{I}$ (the latter by [HCII, Corollary VIII.6]), one easily checks $\text{Hom}_{\text{Cat}_{\infty}}(\mathcal{C}, \Gamma(p)) \simeq \text{Hom}_{\text{Cat}_{\infty}/\mathcal{I}}((\text{pr}_2: \mathcal{C} \times \mathcal{I} \rightarrow \mathcal{I}), (p: E \rightarrow \mathcal{I}))$. Now the Hom anima we are interested in are full subanima of these, so it really suffices to check that their 0-simplices correspond, which we did above. \square

LECTURE 4
10th Nov, 2020

I.37. Example. — Since $\text{core}: \text{Cat}_{\infty} \rightarrow \text{An}$ is right-adjoint to the inclusion $\text{An} \subseteq \text{Cat}_{\infty}$ (Example I.33(a)), we get a natural counit transformation $\text{core} \Rightarrow \text{id}_{\text{Cat}_{\infty}}$, which is given by a map $\Delta^1 \rightarrow \text{Fun}(\text{Cat}_{\infty}, \text{Cat}_{\infty})$. By currying, this yields a functor

$$\begin{aligned} \text{Cat}_{\infty} &\longrightarrow \text{Ar}(\text{Cat}_{\infty}) \\ \mathcal{C} &\longmapsto (\text{core } \mathcal{C} \rightarrow \mathcal{C}). \end{aligned}$$

If $F: W \rightarrow \mathcal{C}$ is an object in $\text{Ar}(\text{Cat}_{\infty})$, a left-adjoint object to F under the above functor is called a *localisation* of \mathcal{C} at F (or rather at the image of $\pi_0 \text{core } \text{Ar}(W) \rightarrow \pi_0 \text{core } \text{Ar}(\mathcal{C})$) of F and denoted $\mathcal{C}[W^{-1}]$. Unwinding, this means that

$$\text{Hom}_{\text{Cat}_{\infty}}(\mathcal{C}[W^{-1}], \mathcal{D}) \subseteq \text{Hom}_{\text{Cat}_{\infty}}(\mathcal{C}, \mathcal{D})$$

is a collection of path components, given by those $G: \mathcal{C} \rightarrow \mathcal{D}$ such that the composite $G \circ F: W \rightarrow \mathcal{D}$ lands in $\text{core } \mathcal{D}$. The localisation $\mathcal{C}[W^{-1}]$ always exists, as one can take

$$\begin{array}{ccc} W & \xrightarrow{F} & \mathcal{C} \\ \downarrow & \lrcorner & \downarrow p \\ |W| & \longrightarrow & \mathcal{C}[W^{-1}] \end{array}$$

(the pushout is taken in Cat_∞ of course). Furthermore, it turns out that

$$p^*: \text{Fun}(\mathcal{C}[W^{-1}], \mathcal{D}) \longrightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$$

is fully faithful, with essential image spanned by the same functors $G: \mathcal{C} \rightarrow \mathcal{D}$ as above (see [HCII, Proposition VIII.7] for a proof). This has the funny consequence that the Yoneda embedding induces fully faithful inclusions $\mathcal{C}[W^{-1}] \subseteq \mathcal{P}(\mathcal{C}[W^{-1}]) \subseteq \mathcal{P}(\mathcal{C})$. In other words, all localisations of \mathcal{C} occur as full subcategories of $\mathcal{P}(\mathcal{C})$, which puts a size bound on how large $\mathcal{C}[W^{-1}]$ can be.


Warning $\triangle!$. — Localisations are difficult! Localisations of locally small ∞ -categories may not be locally small any more. Moreover, the localisation of a 1-category may not be a 1-category any more.

I.38. Example. — Let's compute the pushout

$$\begin{array}{ccc} \Delta^0 & \xrightarrow{1} & \Delta^1 \\ 0 \downarrow & \lrcorner & \downarrow \\ \Delta^1 & \longrightarrow & ? \end{array}$$

in Cat_∞ . One way to do this would be to notice that $0: \Delta^0 \rightarrow \Delta^1$ is a cofibration in the Joyal model structure on sSet (and so is $1: \Delta^0 \rightarrow \Delta^1$, but we don't even need this) and all objects are cofibrant, hence the pushout on the nose, which is the 2-spine I^2 , is also a homotopy pushout. Now $I^2 \subseteq \Delta^2$ is inner anodyne, hence Δ^2 is a fibrant replacement of I^2 in the Joyal model structure, which shows that $? \simeq \Delta^2$ is really the pushout in question by Theorem I.34.

For educational purposes, Fabian presented a less efficient way to show $? \simeq \Delta^2$ using Proposition I.36. Consider the pushout diagram above as a map $F: \Lambda_0^2 \rightarrow \text{Cat}_\infty$. Since all maps in the diagram are nerves of functors of 1-categories, the cocartesian unstraightening $\mathcal{D} = \text{Un}^{\text{cocart}}(F)$ is simply given by the Grothendieck construction. Therefore, the cocartesian fibration $\mathcal{D} \rightarrow \Lambda_0^2$ can be pictured as follows:



The two pink edges are the cocartesian ones and thus sent to equivalences in the localisation $\mathcal{D}[\{\text{cocartesian edges}\}^{-1}] = \mathcal{D}[\{\text{pink edges}\}^{-1}]$. Let \mathcal{D}' be the ∞ -category obtained from \mathcal{D}

by removing the purple and the pink edge on the left, together with the 2-simplex spanned by them. We claim that the restriction

$$\mathrm{Fun}^{\{\text{pink}\}}(\mathcal{D}, \mathcal{E}) \xrightarrow{\sim} \mathrm{Fun}^{\{\text{pink}\}}(\mathcal{D}', \mathcal{E})$$

is an equivalence for any ∞ -category \mathcal{E} , where $\mathrm{Fun}^{\{\text{pink}\}}$ denotes functors that send pink edges to equivalences. Indeed, the purple edge on the left is “superfluous data”, i.e., any map $\mathcal{D} \setminus \{\text{purple edge}\} \rightarrow \mathcal{E}$ has a contractible space of lifts to \mathcal{D} , by Joyal’s lifting theorem. Moreover any map $\mathcal{D}' \rightarrow \mathcal{E}$ has a contractible space of lifts to a map $\mathcal{D} \setminus \{\text{purple edge}\} \rightarrow \mathcal{E}$ that sends the left pink edge to an equivalence. This proves the above equivalence.

A similar argument shows that it doesn’t matter whether we send the pink edge on the right to an equivalence or to an actual identity in \mathcal{E} . Thus

$$\mathrm{Fun}^{\{\text{pink}\}}(\mathcal{D}', \mathcal{E}) \simeq \mathrm{Fun}(\Lambda_1^2, \mathcal{E}) \simeq \mathrm{Fun}(\Delta^2, \mathcal{E}),$$

which proves that $\mathcal{D}[\{\text{cocartesian edges}\}^{-1}] \simeq \Delta^2$, as claimed.

I.39. Lemma. — *If \mathcal{C} is complete or cocomplete, then so is $\mathrm{Fun}(\mathcal{D}, \mathcal{C})$ for any \mathcal{D} . Furthermore, for any $f: \mathcal{E} \rightarrow \mathcal{D}$, the precomposition functor*

$$f^*: \mathrm{Fun}(\mathcal{D}, \mathcal{C}) \longrightarrow \mathrm{Fun}(\mathcal{E}, \mathcal{C})$$

preserves limits and colimits. In particular (taking f to be $\{d\} \hookrightarrow \mathcal{D}$ for any $d \in \mathcal{D}$), limits and colimits in functor categories are computed pointwise.

For the proof we need the following observation.

I.40. Observation. — *If $L: \mathcal{C} \rightleftarrows \mathcal{D} : R$ are adjoint functors, then so are*

$$L_*: \mathrm{Fun}(\mathcal{E}, \mathcal{C}) \rightleftarrows \mathrm{Fun}(\mathcal{E}, \mathcal{D}) : R_* \quad \text{and} \quad R^*: \mathrm{Fun}(\mathcal{D}, \mathcal{E}) \rightleftarrows \mathrm{Fun}(\mathcal{C}, \mathcal{E}) : L^*$$

for any ∞ -category \mathcal{E} .

Proof. Being adjoints can be characterized by the existence of a unit and a counit transformation satisfying the triangle identities (see [HCII, Proposition XI.14], and L_* , R_* inherit these transformations from L , R . Same for R^* , L^* . \square

I.41. Observation. — *Left-adjoint functors preserve colimits, right-adjoint functors preserve limits.*

Proof. Straightforward calculation using Observation I.40. See [HCII, Lemma XI.22] for details. \square

Proof of Lemma I.39. For any diagram shape \mathcal{I} , consider the commutative diagram

$$\begin{array}{ccc} & & \mathrm{Fun}(\mathcal{I}, \mathrm{Fun}(\mathcal{D}, \mathcal{C})) \\ & \nearrow \text{const} & \\ \mathrm{Fun}(\mathcal{D}, \mathcal{C}) & & \\ & \searrow \text{colim}_* & \\ & & \mathrm{Fun}(\mathcal{D}, \mathrm{Fun}(\mathcal{I}, \mathcal{C})) \end{array} \quad \begin{array}{c} \parallel \\ \parallel \end{array}$$

\lim_*

By Observation I.40, colim_* and lim_* are left and right adjoints of const^* respectively, which proves that $\text{const}: \text{Fun}(\mathcal{D}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{I}, \text{Fun}(\mathcal{D}, \mathcal{C}))$ has a left and right adjoint as well. Hence $\text{Fun}(\mathcal{D}, \mathcal{C})$ is complete and cocomplete again. Moreover, the assertion that limits and colimits are computed pointwise follows by unraveling.

It remains to show that $f^*: \text{Fun}(\mathcal{D}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{E}, \mathcal{C})$ preserves limits and colimits. For any diagram $F: \mathcal{I} \rightarrow \text{Fun}(\mathcal{D}, \mathcal{C})$ we get a map

$$\text{colim}_{\mathcal{I}} f^* F \longrightarrow f^* \text{colim}_{\mathcal{I}} F$$

(unique up to contractible choice). Whether this is an equivalence in $\text{Fun}(\mathcal{E}, \mathcal{C})$ can be checked pointwise (by Theorem I.19(b)), i.e., after composition with $x^*: \text{Fun}(\mathcal{E}, \mathcal{C}) \rightarrow \text{Fun}(\{x\}, \mathcal{C}) \simeq \mathcal{C}$ for all $x \in \mathcal{E}$. Since we already know that colimits in $\text{Fun}(\mathcal{E}, \mathcal{C})$ and $\text{Fun}(\mathcal{D}, \mathcal{C})$ are computed pointwise, we see that both $x^* \text{colim}_{\mathcal{I}} f^* F$ and $x^* f^* \text{colim}_{\mathcal{I}} F$ are colimits of the diagram $\mathcal{I} \times \{f(x)\} \rightarrow \mathcal{C}$ induced by currying of F and restriction along $\{f(x)\} \hookrightarrow \mathcal{D}$. \square

I.42. Proposition. — *Let $p: \mathcal{E} \rightarrow \mathcal{C}$ be a cocartesian fibration and $F: \mathcal{E} \rightarrow \mathcal{D}$ a diagram. Suppose that the restrictions $F|_{\mathcal{E}|_c}: \mathcal{E}|_c \rightarrow \mathcal{D}$ of F to the fibres $\mathcal{E}|_c = \mathcal{E} \times_{\mathcal{C}} \{c\}$ of p have colimits for all $c \in \mathcal{C}$. Then these assemble into a functor $G: \mathcal{C} \rightarrow \mathcal{D}$ sending $c \mapsto \text{colim}_{\mathcal{E}|_c} F|_{\mathcal{E}|_c}$, and*

$$\text{colim}_{\mathcal{E}} F \simeq \text{colim}_{\mathcal{C}} G,$$

whenever either exists. In particular (taking p to be the projection $\text{pr}_2: \mathcal{I} \times \mathcal{C} \rightarrow \mathcal{C}$ for some ∞ -category \mathcal{I}), this means that “colimits commute”. Also a dual assertion holds, with p a cartesian fibration and all colimits replaced by limits.

*Proof**. Fabian claims (I think) this follows from [HTT, Proposition 4.3.2.12], but Lurie’s proof is rather laborious, so I’ll give my own proof*. As it turns out, the arguments are pretty much the same as for the proof of Theorems I.51 and I.52 in Fabian’s notes. I’ll freely reference later results (without running into circular arguments, I hope).

We won’t prove the assertion as stated above, but its dualized version. The reason is that the direct proof of the undualized version is quite confusing (for example, neither of the Yoneda embeddings preserves colimits, so we would have to consider $(Y^{\mathcal{D}})^{\text{op}}: \mathcal{D}^{\text{op}} \rightarrow \mathcal{P}(\mathcal{D})^{\text{op}}$) and it seems cleaner to prove the dualized statement and then deduce the undualized one by applying the dual statement to $p^{\text{op}}: \mathcal{E}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$.

So let $p: \mathcal{E} \rightarrow \mathcal{C}$ be cartesian instead. We want to construct G as the right Kan extension $G \simeq p_* F$. Let’s first consider the case where \mathcal{D} is replaced by $\mathcal{P}(\mathcal{D})$. Then the dual of Theorem I.21 implies $\text{Fun}(\mathcal{C}, \mathcal{P}(\mathcal{D})) \simeq \text{Fun}(\mathcal{C} \times \mathcal{D}^{\text{op}}, \text{An}) \simeq \text{Right}(\mathcal{C}^{\text{op}} \times \mathcal{D})$ and same for \mathcal{E} , so we may apply Lemma* I.42a below to the cocartesian fibration $p^{\text{op}} \times \text{id}_{\mathcal{D}}: \mathcal{E}^{\text{op}} \times \mathcal{D} \rightarrow \mathcal{C}^{\text{op}} \times \mathcal{D}$ to see that $p^*: \text{Fun}(\mathcal{C}, \mathcal{P}(\mathcal{D})) \rightarrow \text{Fun}(\mathcal{E}, \mathcal{P}(\mathcal{D}))$ has a right adjoint p_* . Now consider the diagram

$$\begin{array}{ccc} \text{Fun}(\mathcal{C}, \mathcal{P}(\mathcal{D})) & \xrightarrow{p^*} & \text{Fun}(\mathcal{E}, \mathcal{P}(\mathcal{D})) \\ c^* \downarrow & & \downarrow i^* \\ \text{Fun}(\{c\}, \mathcal{P}(\mathcal{D})) & \xrightarrow{p|_c^*} & \text{Fun}(\mathcal{E}|_c, \mathcal{P}(\mathcal{D})) \end{array}$$

where $i: \mathcal{E}|_c \hookrightarrow \mathcal{E}$ denotes the inclusion of the fibre. The second part of Lemma* I.42a ensures that $c^* p_* F \simeq p|_{c,*} i^* F$. By inspection, right Kan extension along the functor $p|_c: \mathcal{E}|_c \rightarrow \{c\}$ is given by taking a functor $G: \mathcal{E}|_c \rightarrow \mathcal{P}(\mathcal{D})$ to $\text{lim}_{\mathcal{E}|_c} G$. This implies $p_* F(c) \simeq \text{lim}_{\mathcal{E}|_c} F|_c$, so

p_*F takes the correct values. Moreover, limits over \mathcal{C} are given by right Kan extension along the unique map $\pi: \mathcal{C} \rightarrow *$, and similar for $\pi \circ p: \mathcal{E} \rightarrow *$, hence

$$\lim_{\mathcal{C}} p_*F \simeq \pi_* p_*F \simeq (\pi \circ p)_*F \simeq \lim_{\mathcal{E}} F$$

since right adjoints compose. This finishes the proof for $\mathcal{P}(\mathcal{D})$.

Now for the case of general \mathcal{D} . The Yoneda embedding $Y^{\mathcal{D}}: \mathcal{D} \rightarrow \mathcal{P}(\mathcal{D})$ is fully faithful, hence an equivalence onto its essential image. To obtain a functor $p_*F: \mathcal{C} \rightarrow \mathcal{D}$, it thus suffices to check that $p_*(Y^{\mathcal{D}} \circ F): \mathcal{C} \rightarrow \mathcal{P}(\mathcal{D})$ has image in the representable functors. By assumption, all limits $\lim_{\mathcal{E}|_{\mathcal{C}}} F|_{\mathcal{C}}$ exist in \mathcal{D} and $Y^{\mathcal{D}}$ preserves limits by Corollary I.50, hence the description of the values $p_*(Y \circ F)(c)$ for $c \in \mathcal{C}$ shows that indeed it takes values in the representables. Thus $p_*F: \mathcal{C} \rightarrow \mathcal{D}$ exists. The additional assertion about limits follows as above, noting that the limits in question exist in \mathcal{D} if and only if the limits taken in $\mathcal{P}(\mathcal{D})$ (where they definitely exist) are representable, since $Y^{\mathcal{D}}$ is fully faithful and preserves limits. \square

I.42a. Lemma*. — *Let $p: \mathcal{C} \rightarrow \mathcal{D}$ be a cocartesian fibration of ∞ -categories. Consider a pullback square (inside Cat_{∞}) as on the left*

$$\begin{array}{ccc} \mathcal{C}' & \xrightarrow{q} & \mathcal{D}' \\ g \downarrow & \lrcorner & \downarrow f \\ \mathcal{C} & \xrightarrow{p} & \mathcal{D} \end{array} \rightsquigarrow \begin{array}{ccc} \text{Right}(\mathcal{C}') & \xleftarrow{q^*} & \text{Right}(\mathcal{D}') \\ g^* \uparrow & \text{---} q_* \text{---} & \uparrow f^* \\ \text{Right}(\mathcal{C}) & \xleftarrow{p^*} & \text{Right}(\mathcal{D}) \\ & \text{---} p_* \text{---} & \end{array}$$

inducing a square as on the right. Then p^ and q^* have right adjoints p_* and q_* as indicated. Moreover, there's a natural equivalence*

$$f^* p_* \xrightarrow{\sim} q_* g^*.$$

Proof.* Without loss of generality all cartesian or right fibrations can be taken in the old sense. Recall (e.g. from the dual of [HTT, §2.1.4]) that there exists the *contravariant model structure* on sSet/\mathcal{C} , in which the cofibrations are monomorphisms of simplicial sets and fibrant objects are given by right fibrations over \mathcal{C} (general fibrations are more complicated). The functor $p^*: \text{sSet}/\mathcal{D} \rightarrow \text{sSet}/\mathcal{C}$ has a right adjoint p_* for abstract reasons: It's straightforward to check that p^* preserves colimits, so Theorem I.2 can be applied to $\text{sSet}/\mathcal{D} \simeq \mathcal{P}((\Delta/\mathcal{D})^{\text{op}})$ (here Δ/S denotes the slice category with respect to the functor $\Delta: \Delta \rightarrow \text{sSet}$ sending $[n] \mapsto \Delta^n$). We claim that

$$p^*: \text{sSet}/\mathcal{D} \rightleftarrows \text{sSet}/\mathcal{C} : p_*$$

is even a Quillen adjunction when p is a cocartesian fibration. Indeed, it's clear that p^* preserves cofibrations and cofibrant objects. For preservation of trivial cofibrations, we refer to [HTT, Proposition 4.1.2.15] or the dual of [HCII, Proposition X.43] for the essential case.

As for any Quillen adjunction of model categories, we get an induced adjunction

$$Lp^*: (\text{sSet}/\mathcal{C})_{\infty} \rightleftarrows (\text{sSet}/\mathcal{D})_{\infty} : Rp_*$$

on underlying ∞ -categories, defined as the localisations at all weak equivalences, or equivalently as $N^c((\text{sSet}/\mathcal{C})^{\text{cf}})$ and $N^c((\text{sSet}/\mathcal{D})^{\text{cf}})$, since sSet/\mathcal{C} and sSet/\mathcal{D} can be made into a Kan simplicial model categories, so [HCII, Digression III Theorem D] applies. The latter

description shows that the underlying ∞ -categories identify with $\text{Right}(\mathcal{C})$ and $\text{Right}(\mathcal{D})$. Hence Lp^* (which we abbreviate to p^* in the following) has indeed an adjoint Rp_* (which we abbreviate to p_*).

So p_* and q_* exist. The counit $p^*p_* \Rightarrow \text{id}$ induces a transformation $q^*f^*p_* \simeq g^*p^*p_* \Rightarrow g^*$, whose adjoint is the transformation $f^*p_* \Rightarrow q_*g^*$. Whether this is an equivalence can be checked on objects. However, explicitly computing p_* and q_* is nasty, so we employ a trick: By Yoneda, it suffices to check that $\text{Hom}_{\text{Right}(\mathcal{D}')}(-, f^*p_*-) \Rightarrow \text{Hom}_{\text{Right}(\mathcal{D}')}(-, q_*g^*-) is a natural equivalence. But f^* and g^* have left adjoints $f_!$ and $g_!$ by Theorem I.47, so we may as well show that $\text{Hom}_{\text{Right}(\mathcal{C})}(p^*f_!-, -) \Rightarrow \text{Hom}_{\text{Right}(\mathcal{C})}(g_!q^*-, -)$ is an equivalence, i.e., that $g_!q^* \Rightarrow p^*f_!$ is an equivalence. This is now easily checked on objects: Unravelling, we need to prove that if $X \twoheadrightarrow \mathcal{D}'$ is a right fibration and $X \hookrightarrow Y \twoheadrightarrow \mathcal{D}$ is a factorisation into a right anodyne and a right fibration, then $X \times_{\mathcal{D}} \mathcal{C} \hookrightarrow Y \times_{\mathcal{D}} \mathcal{C}$ is right anodyne again, which follows from the references above since $p: \mathcal{C} \rightarrow \mathcal{D}$ is cocartesian. $\square$$

I.43. Theorem (Joyal's version of Quillen's Theorem A). — *For a functor $\alpha: \mathcal{I} \rightarrow \mathcal{J}$ the following statements are equivalent.*

- (a) *A functor $F: \mathcal{J} \rightarrow \mathcal{C}$ has a colimit iff the composition $F \circ \alpha: \mathcal{I} \rightarrow \mathcal{C}$ has a colimit, and in this case*

$$\text{colim}_{\mathcal{I}} F \circ \alpha \simeq \text{colim}_{\mathcal{J}} F.$$

- (b) *For all $j \in \mathcal{J}$ we have $|j/\alpha| \simeq *$, i.e., the slice category j/α is weakly contractible. Here we define j/α by the pullback*

$$\begin{array}{ccc} j/\alpha & \longrightarrow & \text{Ar}(\mathcal{J}) \\ \downarrow & \lrcorner & \downarrow (s,t) \\ \{j\} \times \mathcal{I} & \xrightarrow{(\text{incl}, \alpha)} & \mathcal{J} \times \mathcal{J} \end{array}$$

Also observe that $j/\alpha = \mathcal{I} \times_{\mathcal{J}} j/\mathcal{J}$, which is how people usually denote this in the literature.

Proof.* Combine [HTT, Proposition 4.1.1.8] (for more equivalent characterisations) and [HTT, Theorem 4.1.3.1] (for the actual proof). \square

I.44. Definition. — Maps $\alpha: \mathcal{I} \rightarrow \mathcal{J}$ as in Theorem I.43 are called *cofinal*. Dually, if precomposition with α preserves limits and detects their existence, then α is called *final*.

Cofinal maps induce homotopy equivalences $|\alpha|: |\mathcal{I}| \rightarrow |\mathcal{J}|$ of anima, which can be seen via the chain of homotopy equivalences

$$|\mathcal{I}| \simeq \text{colim}_{\mathcal{I}} \text{const } * \simeq \text{colim}_{\mathcal{J}} \text{const } * \simeq |\mathcal{J}|$$

(the colimits in the middle are taken in An). Here we use $|\mathcal{I}| \simeq \text{colim}_{\mathcal{I}} \text{const } *$, which can be seen in several ways: One could use Theorem I.34 and the Bousfield–Kan formula ([HCII, Digression III]). Alternatively, compute $\text{Hom}_{\text{An}}(\text{colim}_{\mathcal{I}} \text{const } *, K) \simeq \lim_{\mathcal{I}} \text{Hom}_{\text{An}}(*, K) \simeq \lim_{\mathcal{I}} K$ for any anima K using Corollary I.50 below, and apply Proposition I.36 to get $\lim_{\mathcal{I}} K \simeq \Gamma(\text{pr}_2: K \times \mathcal{I} \rightarrow \mathcal{I})$. The space of sections of $\text{pr}_2: K \times \mathcal{I} \rightarrow \mathcal{I}$ clearly identifies with $\text{Fun}(\mathcal{I}, K) \simeq \text{core Fun}(\mathcal{I}, K) \simeq \text{Hom}_{\text{Cat}_{\infty}}(\mathcal{I}, K)$, so $\text{colim}_{\mathcal{I}} \text{const } *$ is indeed a left-adjoint object of \mathcal{I} under the inclusion $\text{An} \subseteq \text{Cat}_{\infty}$.

Examples of cofinal maps are given by right-adjoint functors and localisations. Indeed, if $\alpha: \mathcal{I} \rightarrow \mathcal{J}$ is a right adjoint, then consider the diagram

$$\begin{array}{ccc} & \text{colim}_{\mathcal{J}} \dashrightarrow \mathcal{C} \dashleftarrow \text{colim}_{\mathcal{I}} & \\ \text{const} \swarrow & & \searrow \text{const} \\ \text{Fun}(\mathcal{J}, \mathcal{C}) & \xrightarrow{\alpha^*} & \text{Fun}(\mathcal{I}, \mathcal{C}) \end{array}$$

By Observation I.40, $\alpha^*: \text{Fun}(\mathcal{J}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{I}, \mathcal{C})$ is a left-adjoint. Since left-adjoints compose, we obtain $\text{colim}_{\mathcal{J}} \simeq \text{colim}_{\mathcal{I}} \circ \alpha^*$, as required. If $p: \mathcal{C} \rightarrow \mathcal{C}[W^{-1}]$ is a localisation and $F: \mathcal{C}[W^{-1}] \rightarrow \mathcal{D}$ is any diagram, then

$$\text{Nat}(F, \text{const } x) \simeq \text{Nat}(F \circ p, \text{const } x)$$

holds for every $x \in \mathcal{C}$ since $p^*: \text{Fun}(\mathcal{C}[W^{-1}], \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$ is fully faithful, which again proves cofinality by means of Theorem I.43(a).

I.45. Corollary. — *Let $G: \mathcal{I} \rightarrow \text{Cat}_{\infty}$ be a diagram and put $\mathcal{J} \simeq \text{colim}_{\mathcal{I}} G$. Let $F: \mathcal{J} \rightarrow \mathcal{D}$ be another diagram and suppose that the restrictions*

$$G(i) \longrightarrow \mathcal{J} \xrightarrow{F} \mathcal{D}$$

have colimits for all $i \in \mathcal{I}$. Then these colimits assemble into a functor $H: \mathcal{I} \rightarrow \mathcal{D}$ sending $i \mapsto \text{colim}_{G(i)} F$, and we have

$$\text{colim}_{\mathcal{J}} F \simeq \text{colim}_{\mathcal{I}} H$$

if either exists.

Proof. By Proposition I.36, we have $\mathcal{J} \simeq \text{colim}_{\mathcal{I}} G \simeq \text{Un}^{\text{cocart}}(G)[\{\text{cocartesian edges}\}^{-1}]$. Since localisations are cofinal, we obtain

$$\text{colim}_{\mathcal{J}} F \simeq \text{colim}_{\text{Un}^{\text{cocart}}(G)} F.$$

Now recall that $G(i) \simeq \text{Un}^{\text{cocart}}(G) \times_{\mathcal{I}} \{i\}$, so we are done by Proposition I.42. \square

I.46. “Corollary”. — *If \mathcal{C} has coproducts and pushouts, then it is cocomplete. Dually, if \mathcal{C} has products and pullbacks, then it is complete.*

“Proof” sketch. Decompose any test category \mathcal{I} into its skeleta and repeatedly apply Corollary I.45 using $\text{colim}_{\Delta^n} F = F(n)$ since $\{n\} \hookrightarrow \Delta^n$ is cofinal—in fact, all right anodyne maps are cofinal. For a complete proof, check out [HTT, Proposition 4.4.2.6]. \square

Colimits and the Yoneda embedding

We start with an existence result for “Kan extensions” which was already used in the proof* of Proposition I.42.

I.47. Theorem (Joyal). — *Given an arbitrary functor $f: \mathcal{C} \rightarrow \mathcal{D}$ of ∞ -categories, the pullback functor*

$$f^*: \text{Right}(\mathcal{D}) \longrightarrow \text{Right}(\mathcal{C})$$

admits a left-adjoint. Its value on $p: \mathcal{E} \rightarrow \mathcal{C}$ is obtained by factoring $\mathcal{E} \rightarrow \mathcal{C} \rightarrow \mathcal{D}$ as a cofinal map followed by a right fibration $p': \mathcal{E}' \rightarrow \mathcal{D}$. In diagrams

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & \mathcal{E}' \\ p \downarrow & & p' \downarrow \\ \mathcal{C} & \xrightarrow{f} & \mathcal{D} \end{array}$$

where the top arrow is cofinal.

*Proof sketch**. One way to prove this would be to use the contravariant model structures on $\mathbf{sSet}/\mathcal{C}$ and $\mathbf{sSet}/\mathcal{D}$ as in the proof* of Lemma* I.42a: One shows that $f^*: \mathbf{sSet}/\mathcal{D} \rightarrow \mathbf{sSet}/\mathcal{C}$ has a left Quillen adjoint $f_!$, sending $\mathcal{E} \rightarrow \mathcal{C}$ to $\mathcal{E} \rightarrow \mathcal{C} \rightarrow \mathcal{D}$. Since right anodyne maps are cofinal ([HTT, Proposition 4.1.1.3] or [HCII, Theorem XI.28]), a quick unraveling shows that $Lf_!$ can indeed be described as above.

However, Bastiaan has pointed out a simpler proof that doesn't need any model categories. We will proceed in three steps:

- (1) *The pullback functor $f^*: \mathbf{Cat}_\infty/\mathcal{D} \rightarrow \mathbf{Cat}_\infty/\mathcal{C}$ (where we take the pullback along f in \mathbf{Cat}_∞) has a left adjoint. On objects, it is the “forgetful functor” sending $p: \mathcal{E} \rightarrow \mathcal{C}$ to $f \circ p: \mathcal{E} \rightarrow \mathcal{C} \rightarrow \mathcal{D}$.*
- (2) *Under the fully faithful inclusions $\mathbf{Right}(\mathcal{C}) \subseteq \mathbf{Cat}_\infty/\mathcal{C}$ and $\mathbf{Right}(\mathcal{D}) \subseteq \mathbf{Cat}_\infty/\mathcal{D}$, the pullback functor $f^*: \mathbf{Cat}_\infty/\mathcal{D} \rightarrow \mathbf{Cat}_\infty/\mathcal{C}$ restricts to $f^*: \mathbf{Right}(\mathcal{D}) \rightarrow \mathbf{Right}(\mathcal{C})$ from the formulation of the theorem.*
- (3) *The fully faithful inclusion $\mathbf{Right}(\mathcal{D}) \subseteq \mathbf{Cat}_\infty/\mathcal{D}$ has a left adjoint, sending an object $q: \mathcal{E} \rightarrow \mathcal{D}$ to $q': \mathcal{E}' \rightarrow \mathcal{D}$, where $\mathcal{E} \rightarrow \mathcal{E}' \rightarrow \mathcal{D}$ is any factorisation into a cofinal map followed by a right fibration.*

Since left adjoints compose, combining (1), (2), and (3) shows that $f^*: \mathbf{Right}(\mathcal{D}) \rightarrow \mathbf{Right}(\mathcal{C})$ has a left adjoint $f_!$ with the desired description on objects.

To show (1), we use Corollary I.32 to see that it suffices to show that $f \circ p$ is a left adjoint object of p for any $(p: \mathcal{E} \rightarrow \mathcal{C}) \in \mathbf{Cat}_\infty/\mathcal{C}$. If we had the dual of Corollary I.50 below already, this would be almost trivial, but as it is, we need to be a bit careful. Factor $f \circ p$ into a Joyal equivalence $\mathcal{E} \rightarrow \mathcal{E}'$ followed by an isofibration $p': \mathcal{E}' \rightarrow \mathcal{D}$. Then $f^*\mathcal{E}'$ equals the same pullback taken in \mathbf{sSet} by Theorem I.34. Hence there is a natural map $\mathcal{E} \rightarrow f^*\mathcal{E}'$. Together with functoriality of f^* , it induces natural transformations

$$\begin{aligned} \mathrm{Hom}_{\mathbf{Cat}_\infty/\mathcal{D}}(\mathcal{E}, -) &\simeq \mathrm{Hom}_{\mathbf{Cat}_\infty/\mathcal{D}}(\mathcal{E}', -) \implies \mathrm{Hom}_{\mathbf{Cat}_\infty/\mathcal{C}}(f^*\mathcal{E}', f^*(-)) \\ &\implies \mathrm{Hom}_{\mathbf{Cat}_\infty/\mathcal{C}}(\mathcal{E}, f^*(-)). \end{aligned}$$

We are to show that the composite is an equivalence. By Theorem I.19(b), this can be done on objects. Let $T \rightarrow \mathcal{D}$ be an object of $\mathbf{Cat}_\infty/\mathcal{D}$, which we may choose to be an isofibration without restriction, so that $f^*(T)$ agrees with the corresponding pullback in \mathbf{sSet} . Consider $\mathrm{Fun}_{\mathcal{D}}(\mathcal{E}, T) := \mathrm{Fun}(\mathcal{E}, T) \times_{\mathrm{Fun}(\mathcal{E}, \mathcal{D})} \{f \circ p\}$ and define $\mathrm{Fun}_{\mathcal{C}}(\mathcal{E}, f^*(T))$ similarly. Since $T \rightarrow \mathcal{D}$ is an isofibration, it doesn't matter whether this pullback is taken in \mathbf{sSet} or in \mathbf{Cat}_∞ . Moreover, $\mathrm{core}(-)$ transforms pullbacks in \mathbf{Cat}_∞ into pullbacks in \mathbf{An} (it is a right adjoint by Example I.33(a), so Observation I.41 can be applied), hence our computation of Hom anima in slice categories from [HCII, Corollary VIII.6] shows $\mathrm{Hom}_{\mathbf{Cat}_\infty/\mathcal{D}}(\mathcal{E}, T) \simeq \mathrm{core} \mathrm{Fun}_{\mathcal{D}}(\mathcal{E}, T)$. Likewise, $\mathrm{Hom}_{\mathbf{Cat}_\infty/\mathcal{C}}(\mathcal{E}, f^*(T)) \simeq \mathrm{core} \mathrm{Fun}_{\mathcal{C}}(\mathcal{E}, f^*(T))$. Now the universal property of $f^*(T)$ as a pullback in \mathbf{sSet} implies $\mathrm{Fun}_{\mathcal{D}}(\mathcal{E}, T) \simeq \mathrm{Fun}_{\mathcal{C}}(\mathcal{E}, f^*(T))$, which proves step (1).

For (2), observe that f^* of a right fibration in the old sense can also be taken in \mathbf{sSet} by Theorem I.34 again, and right fibrations (in the old sense) are preserved under pullbacks formed in \mathbf{sSet} . Hence right fibrations in the new sense are preserved by pullbacks in \mathbf{Cat}_∞ .

Finally, to show (3), we can argue as (1) to reduce the assertion to $\mathrm{Fun}_\mathcal{D}(\mathcal{E}', T) \simeq \mathrm{Fun}_\mathcal{D}(\mathcal{E}, T)$ for any right fibration $T \rightarrow \mathcal{D}$ and any cofinal map $\mathcal{E}' \rightarrow \mathcal{E}$ over \mathcal{D} . But this is precisely how Lurie defines cofinal maps in [HTT, Definition 4.1.1.1], and his definition is equivalent to our Definition I.44. \square

I.48. Corollary. — *The functor $f^*: \mathcal{P}(\mathcal{D}) \rightarrow \mathcal{P}(\mathcal{C})$ has a left-adjoint $f_!$ (given by left Kan extension) such that the diagram*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{f} & \mathcal{D} \\ Y^{\mathcal{C}} \downarrow & & \downarrow Y^{\mathcal{D}} \\ \mathcal{P}(\mathcal{C}) & \xrightarrow{f_!} & \mathcal{P}(\mathcal{D}) \end{array}$$

commutes in the ∞ -category \mathbf{Cat}_∞ .

Proof. The first statement immediately follows from Theorem I.47 and (the dual of) Theorem I.21, which asserts that the cartesian straightening $\mathrm{St}^{\mathrm{cart}}: \mathrm{Right}(\mathcal{C}) \xrightarrow{\sim} \mathcal{P}(\mathcal{C})$ is an equivalence with inverse the cartesian unstraightening $\mathrm{Un}^{\mathrm{cart}}$.

For the second statement, we have to show that $f_! \mathrm{Hom}_{\mathcal{C}}(-, x) \rightarrow \mathrm{Hom}_{\mathcal{D}}(-, f(x))$ is an equivalence for all $x \in \mathcal{C}$. After unstraightening, this translates to the statement that the diagram

$$\begin{array}{ccc} \mathcal{C}/x & \longrightarrow & \mathcal{D}/f(x) \\ \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{f} & \mathcal{D} \end{array}$$

exhibits $\mathcal{D}/f(x) \rightarrow \mathcal{D}$ as $f_!(\mathcal{C}/x \rightarrow \mathcal{C})$. The right vertical arrow is already a right fibration, so it suffices to check that $\mathcal{C}/x \rightarrow \mathcal{D}/f(x)$ is cofinal. This holds because

$$\begin{array}{ccc} & \Delta^0 & \\ x \swarrow & & \searrow f(x) \\ \mathcal{C}/x & \xrightarrow{f} & \mathcal{D}/f(x) \end{array}$$

has both downward arrows cofinal (as x and $f(x)$ are terminal objects in \mathcal{C}/x and $\mathcal{D}/f(x)$ respectively and thus both sloped arrows are right anodyne by [HCII, Digression I Corollary D.7]). \square

I.49. Corollary. — *Given functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$ of ∞ -categories, we have an equivalence of anima*

$$\mathrm{Nat}(F, G) \simeq \lim_{(x \rightarrow y) \in \mathrm{TwAr}(\mathcal{C})} \mathrm{Hom}_{\mathcal{D}}(F(x), G(y)).$$

Explicitly, the limit is taken over $\mathrm{TwAr}(\mathcal{C}) \rightarrow \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \xrightarrow{F^{\mathrm{op}} \times G} \mathcal{D}^{\mathrm{op}} \times \mathcal{D} \xrightarrow{\mathrm{Hom}_{\mathcal{D}}} \mathbf{An}$.

Note that Corollary I.49 explains why natural transformations are complicated in ∞ -land: π_0 does not commute with arbitrary limits!

Proof of Corollary I.49. By Proposition I.36, the right-hand side can be computed as the anima of sections of $p: P \rightarrow \mathrm{TwAr}(\mathcal{C})$, where P denotes the correct unstraightening. Explicitly, we have a diagram

$$\begin{array}{ccccccc}
 P & \longrightarrow & P' & \longrightarrow & \mathrm{TwAr}(\mathcal{D}) & \longrightarrow & */\mathrm{An} \\
 \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\
 \mathrm{TwAr}(\mathcal{C}) & \xrightarrow{(s,t)} & \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{C} & \xrightarrow{F^{\mathrm{op}} \times G} & \mathcal{D}^{\mathrm{op}} \times \mathcal{D} & \xrightarrow{\mathrm{Hom}_{\mathcal{D}}} & \mathrm{An}
 \end{array}$$

which shows $\Gamma(p) \simeq \mathrm{Hom}_{\mathrm{Cat}_{\infty}/\mathcal{C}^{\mathrm{op}} \times \mathcal{C}}(\mathrm{TwAr}(\mathcal{C}), P')$. But these are both left fibrations over $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$, so the Hom anima on the right-hand side can be equivalently computed as $\mathrm{Nat}(\mathrm{Hom}_{\mathcal{C}}, \mathrm{Hom}_{\mathcal{D}} \circ (F^{\mathrm{op}} \times G))$. Now the currying equality $\mathrm{Fun}(\mathcal{C}^{\mathrm{op}} \times \mathcal{C}, \mathrm{An}) = \mathrm{Fun}(\mathcal{C}, \mathcal{P}(\mathcal{C}))$ sends $\mathrm{Hom}_{\mathcal{C}}$ to $Y^{\mathcal{C}}$ and $\mathrm{Hom}_{\mathcal{D}} \circ (F^{\mathrm{op}} \times G)$ to $F^* \circ Y^{\mathcal{D}} \circ G$, hence the Hom anima under consideration is given by

$$\mathrm{Nat}(Y^{\mathcal{C}}, F^* \circ Y^{\mathcal{D}} \circ G) \simeq \mathrm{Nat}(F! \circ Y^{\mathcal{C}}, Y^{\mathcal{D}} \circ G) \simeq \mathrm{Nat}(Y^{\mathcal{D}} \circ F, Y^{\mathcal{D}} \circ G) \simeq \mathrm{Nat}(F, G),$$

as claimed. For the left equivalence, we use that $F! \circ -$ is an adjoint of $F^* \circ -$ by construction and Observation I.40, the middle equivalence follows from Corollary I.48, and the right one since $Y^{\mathcal{D}}: \mathcal{D} \rightarrow \mathcal{P}(\mathcal{D})$ is fully faithful by Yoneda's lemma (Theorem I.29). \square

I.50. Corollary. — *Let $F: \mathcal{I} \rightarrow \mathcal{C}$ be a functor of ∞ -categories. A natural transformation $\eta: F \Rightarrow \mathrm{const} c$ exhibits $c \in \mathcal{C}$ as the colimit of F if and only if the natural map*

$$\eta^*: \mathrm{Hom}_{\mathcal{C}}(c, x) \xrightarrow{\sim} \lim_{i \in \mathcal{I}^{\mathrm{op}}} \mathrm{Hom}_{\mathcal{C}}(F(i), x)$$

is an equivalence for all $x \in \mathcal{C}$. A similar assertion holds for limits. In particular, the functors $\mathrm{Hom}_{\mathcal{C}}(-, d): \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{An}$ and $Y^{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{P}(\mathcal{C})$ preserve limits.

Proof. We are done if we can show that the right-hand side is just $\mathrm{Nat}(F, \mathrm{const} x)$. From Corollary I.49 we get a map

$$\mathrm{Nat}(F, \mathrm{const} x) \simeq \lim_{(i \rightarrow i') \in \mathrm{TwAr}(\mathcal{I})} \mathrm{Hom}_{\mathcal{C}}(F(i), x) \longrightarrow \lim_{i \in \mathcal{I}^{\mathrm{op}}} \mathrm{Hom}_{\mathcal{C}}(F(i), x).$$

So we are done if we show that $s: \mathrm{TwAr}(\mathcal{I}) \rightarrow \mathcal{I}^{\mathrm{op}}$ is final. By (the dual of) Theorem I.43(b) we must show $|s/i| \simeq *$ for all $i \in \mathcal{I}^{\mathrm{op}}$. Applying Exercise I.50a to the cocartesian fibration $s = \mathrm{pr}_2 \circ (s, t): \mathrm{TwAr}(\mathcal{I}) \rightarrow \mathcal{I}^{\mathrm{op}} \times \mathcal{I} \rightarrow \mathcal{I}^{\mathrm{op}}$, we find that $|s^{-1}\{i\}| \simeq |s/i|$ since adjoints induce inverse homotopy equivalences after applying $|-|$; alternatively one can use that left adjoints are final. Now $s^{-1}\{i\}$ has an initial object which immediately shows $|s^{-1}\{i\}| \simeq *$. In fact, we have $s^{-1}\{i\} \simeq i/\mathcal{I}$ (and id_i is initial in the right-hand side): Depending on your definitions, this is either something one has to check (as in the proof of [HA, Proposition 5.2.1.10]), or follows from the fact that $t: s^{-1}\{i\} \rightarrow \mathcal{I}$ represents the functor $\mathrm{Hom}_{\mathcal{I}}(i, -)$ and is thus given by the left fibration $i/\mathcal{I} \rightarrow \mathcal{I}$. \square

I.50a. Exercise. — Let $p: \mathcal{E} \rightarrow \mathcal{C}$ be a cocartesian fibration (in the old sense; if you want this to work in the new sense as well, all fibre products need to be taken inside Cat_{∞}). Show that the natural map $\mathcal{E} \times_{\mathcal{C}} \{c\} \rightarrow \mathcal{E} \times_{\mathcal{C}} \mathcal{C}/c$ has a left adjoint for all $c \in \mathcal{C}$.

Fabian wrote i/s instead of s/i in the lecture and in [A&HK, Corollar I.50], and the exercise was formulated with c/\mathcal{C} instead of \mathcal{C}/c . I'm pretty sure that we really need the other slices—after all, we need to apply the dual version of Theorem I.43(b), also it will become apparent during the proof* of Exercise I.50a that we really need to use \mathcal{C}/c .

Proof of Exercise I.50a.* Since adjoints can be constructed objectwise by Corollary I.32, we only need to show that every object $(e', \varphi: c' \rightarrow c) \in \mathcal{E} \times_{\mathcal{C}} \mathcal{C}/c$ admits a left-adjoint object. Let $\eta: e' \rightarrow e$ be a p -cocartesian lift of φ . This induces a morphism $\eta: (e', \varphi) \rightarrow (e, \text{id}_c)$ in $\mathcal{E} \times_{\mathcal{C}} \mathcal{C}/c$. We claim that η exhibits e as a left-adjoint object of (e', φ) , so we need to show that the composite

$$\text{Hom}_{\mathcal{E} \times_{\mathcal{C}} \{c\}}(e, e'') \longrightarrow \text{Hom}_{\mathcal{E} \times_{\mathcal{C}} \mathcal{C}/c}((e, \text{id}_c), (e'', \text{id}_c)) \xrightarrow{\eta^*} \text{Hom}_{\mathcal{E} \times_{\mathcal{C}} \mathcal{C}/c}((e', \varphi), (e'', \text{id}_c))$$

is an equivalence for all $e'' \in \mathcal{E} \times_{\mathcal{C}} \{c\}$. Plugging in the homotopy pullback that describes $\text{Hom}_{\mathcal{E}}(e, e'')$ due to $e' \rightarrow e$ being p -cocartesian, as well as the homotopy pullback that computes $\text{Hom}_{\mathcal{C}/c}(c', c)$ due to [HCII, Corollary VIII.6], shows that both the left-hand side and the right-hand side are homotopy equivalent to $\text{Hom}_{\mathcal{E}}(e', e) \times_{\text{Hom}_{\mathcal{C}}(c', c)}^R \{\varphi\}$ and that the above composite induces the identity on that anima. \square

I.50b. Exercise. — Use Corollary I.50 to show that $\mathcal{K}(R)$ and $\mathbf{N}^c(\text{Top})$ are (co)complete.

⚠ Public Service Announcement. — *From now on, all pullbacks of ∞ -categories or anima will be taken in Cat_{∞} or An , unless specified otherwise!* Still, in many cases they will agree with the corresponding pullbacks in \mathbf{sSet} by arguments like Example* I.35a.

Kan Extensions

LECTURE 5
12th Nov, 2020

The first goal for today is to discuss two closely related theorems, which provide an ∞ -analogue of Theorem I.2.

I.51. Theorem. — *For every cocomplete ∞ -category \mathcal{D} and every small ∞ -category \mathcal{C} , the restriction of $Y^*: \text{Fun}(\mathcal{P}(\mathcal{C}), \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$ to colimit-preserving functors in the source is an equivalence. Furthermore, any such functor has a right adjoint. In view of Observation I.41, we thus get an equivalence*

$$Y^{\mathcal{C},*}: \text{Fun}^L(\mathcal{P}(\mathcal{C}), \mathcal{D}) \xrightarrow{\sim} \text{Fun}(\mathcal{C}, \mathcal{D}),$$

where Fun^L denotes the full sub- ∞ -category spanned by left adjoint functors.

I.52. Theorem. — *If \mathcal{C} is a small ∞ -category and \mathcal{D} cocomplete or complete, then for any $f: \mathcal{C} \rightarrow \mathcal{E}$ the functor*

$$f^*: \text{Fun}(\mathcal{E}, \mathcal{D}) \longrightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$$

has a left adjoint Lan_f or a right adjoint Ran_f (sometimes also denoted $f_!$ and f_) which satisfy*

$$\text{Lan}_f F(e) \simeq \text{colim}_{(c, f(c) \rightarrow e) \in f/e} F(c) \quad \text{and} \quad \text{Ran}_f F(e) \simeq \lim_{(c, e \rightarrow f(c)) \in e/f} F(c).$$

In fact, if \mathcal{D} is not necessarily cocomplete or complete, but these colimits or limits happen to exist for all $e \in \mathcal{E}$, then they assemble into a functor $\text{Lan}_f F: \mathcal{E} \rightarrow \mathcal{D}$ or $\text{Ran}_f F: \mathcal{E} \rightarrow \mathcal{D}$ which is a left- or right-adjoint object of F with respect to f^ .*

I.53. Definition. — If $G: \mathcal{E} \rightarrow \mathcal{D}$ is a left or right adjoint object of $F: \mathcal{C} \rightarrow \mathcal{D}$ under $f^*: \text{Fun}(\mathcal{E}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$, then G is called a *left or right Kan extension* of F along f .

A first example is given by $f_! : \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{D})$ from Corollary I.48, which sends a presheaf $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{An}$ to its left Kan extension $f_! F : \mathcal{D}^{\text{op}} \rightarrow \mathbf{An}$ along $\mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$.

In the official lectures notes (starting from [A&HK, Observation I.56]) Fabian outlines two proofs of Theorems I.51 and I.52, one due to Cisinski and one due to Lurie. In these notes we'll present a third proof that Fabian briefly sketched in the lecture. It's perhaps a bit more straightforward than the other two in that it really does the same as for 1-categories, but it's not really shorter.

Proof of Theorem I.52.* Consider the slice category f/\mathcal{E} defined by the pullback

$$\begin{array}{ccc} f/\mathcal{E} & \longrightarrow & \mathbf{Ar}(\mathcal{E}) \\ \downarrow & \lrcorner & \downarrow s \\ \mathcal{C} & \xrightarrow{f} & \mathcal{E} \end{array}$$

(this is not equivalent to the thin or fat slice $\mathcal{E}_{f/}$ or $\mathcal{E}_{f//}$). Informally, f/\mathcal{E} consists of pairs $(c, f(c) \rightarrow e)$ where $c \in \mathcal{C}$, $e \in \mathcal{E}$. It comes with morphisms $s : f/\mathcal{E} \rightarrow \mathcal{C}$ and $t : f/\mathcal{E} \rightarrow \mathcal{E}$ sending such a pair to c or e respectively. Then t is a cocartesian fibration by an easy generalisation of Example I.25(b) and its fibres $t^{-1}\{e\} \simeq f/e$ are the slice categories from the definition of $\text{Lan}_f F(e)$. We may thus apply Proposition I.42 to the cocartesian fibration t and the functor $F \circ s : f/\mathcal{E} \rightarrow \mathcal{D}$ to see that the pointwise formulas indeed assemble into a functor $\text{Lan}_f F$. Going through the proof of Proposition I.42, we see that $(\text{Lan}_f F)^{\text{op}} \simeq t_*^{\text{op}}(F^{\text{op}} \circ s^{\text{op}})$ is a right-adjoint object of $F^{\text{op}} \circ s^{\text{op}}$ with respect to $(t^{\text{op}})^* : \text{Fun}(\mathcal{E}^{\text{op}}, \mathcal{D}^{\text{op}}) \rightarrow \text{Fun}((f/\mathcal{E})^{\text{op}}, \mathcal{D}^{\text{op}})$, hence $\text{Lan}_f F$ is a left-adjoint object of $F \circ s$ under $t^* : \text{Fun}(\mathcal{E}, \mathcal{D}) \rightarrow \text{Fun}(f/\mathcal{E}, \mathcal{D})$. Thus, we only need to show that there is an equivalence

$$\text{Nat}(F, - \circ f) \xrightarrow{\sim} \text{Nat}(F \circ s, - \circ t).$$

Observe that s has a section $r : \mathcal{C} \rightarrow f/\mathcal{E}$, defined by the universal property of pullbacks and the maps $\text{id}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$ and $f^* \circ \text{const} : \mathcal{C} \rightarrow \mathbf{Ar}(\mathcal{E})$. Then $f = t \circ r$ and there are natural transformations $\eta : f \circ s \Rightarrow t$ and $\eta' : r \circ s \Rightarrow \text{id}_{f/\mathcal{E}}$; in diagrams

$$\begin{array}{ccccc} \mathcal{C} & \xrightarrow{r} & f/\mathcal{E} & \xrightarrow{s} & \mathcal{C} \\ & \searrow f & \downarrow t & \swarrow f & \\ & & \mathcal{E} & & \end{array} \quad \begin{array}{c} \eta \\ \eta' \end{array}$$

To get η , we compose $f/\mathcal{E} \times \Delta^1 \rightarrow \mathbf{Ar}(\mathcal{E}) \times \Delta^1$ with the evaluation map $\mathbf{Ar}(\mathcal{E}) \times \Delta^1 \rightarrow \mathcal{E}$. To get $\eta' : f/\mathcal{E} \times \Delta^1 \rightarrow f/\mathcal{E}$, we must choose homotopies $f/\mathcal{E} \times \Delta^1 \rightarrow \mathcal{C}$ and $f/\mathcal{E} \times \Delta^1 \rightarrow \mathbf{Ar}(\mathcal{E})$. The first can be chosen to be constant at s , the second is equivalently given by a map $\Delta^1 \times \Delta^1 \rightarrow \text{Fun}(f/\mathcal{E}, \mathcal{E})$, which we choose to be

$$\begin{array}{ccc} f \circ s & \xlongequal{\quad} & f \circ s \\ \parallel & \searrow \eta & \downarrow \eta \\ f \circ s & \xrightarrow{\quad} & t \end{array}$$

In particular, $t \circ \eta' = \eta$. Now the desired equivalence can be obtained via the composite $\eta_* \circ s^* : \text{Nat}(F, - \circ f) \Rightarrow \text{Nat}(F \circ s, - \circ f \circ s) \Rightarrow \text{Nat}(F \circ s, - \circ t)$. To show that it is an equivalence, we simply check that $r^* : \text{Nat}(F \circ s, - \circ t) \Rightarrow \text{Nat}(F, - \circ f)$ (here we use $s \circ r = \text{id}_{\mathcal{C}}$ and $t \circ r = f$) is an inverse by a straightforward computation. \square

To prove Theorem I.51, we'll check that $\text{Lan}_{Y^{\mathcal{C}}} : \text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}(\mathcal{P}(\mathcal{C}), \mathcal{D})$ constitutes an inverse to $Y^{\mathcal{C},*}$. For this, we need to check that $\text{Lan}_{Y^{\mathcal{C}}}$ really takes values in $\text{Fun}^L(\mathcal{P}(\mathcal{C}), \mathcal{D})$, which needs some preparations. The first is a lemma that I copied straight from Fabians notes [A&HK, Proposition I.51] (but Fabian proves it differently).

I.53a. Lemma*. — *Every element in $\mathcal{P}(\mathcal{C})$ is a colimit of representable presheaves. More precisely, for every $E \in \mathcal{P}(\mathcal{C})$, the canonical map*

$$\text{colim}_{(c, c \rightarrow E) \in Y^{\mathcal{C}}/E} \text{Hom}_{\mathcal{C}}(-, c) \xrightarrow{\sim} E$$

is an equivalence.

Proof.* By Theorem I.19(b), it suffices to check that both sides agree after evaluation at $x \in \mathcal{C}^{\text{op}}$. Since colimits in functor categories are computed pointwise by Lemma I.39, we see that the left-hand side, when evaluated at x , is given by the colimit of

$$Y^{\mathcal{C}}/E \xrightarrow{s} \mathcal{C} \xrightarrow{\text{Hom}_{\mathcal{C}}(x, -)} \text{An}.$$

We know how to compute colimits in An by Proposition I.36. The cocartesian unstraightening of $\text{Hom}_{\mathcal{C}}(x, -)$ is $x/\mathcal{C} \rightarrow \mathcal{C}$, hence the cocartesian unstraightening U of the functor in question is given by the pullback

$$\begin{array}{ccc} U & \longrightarrow & x/\mathcal{C} \\ \downarrow & \lrcorner & \downarrow \\ Y^{\mathcal{C}}/E & \xrightarrow{s} & \mathcal{C} \end{array}$$

We'll see in the proof of Lemma* I.53b that $Y^{\mathcal{C}}/E \rightarrow \mathcal{C}$ is cartesian (even a right fibration), hence the dual of Exercise I.50a provides an adjunction $U \times_{x/\mathcal{C}} \{\text{id}_x\} \rightleftarrows U$. In particular, we get a homotopy equivalence $|U \times_{x/\mathcal{C}} \{\text{id}_x\}| \simeq |U|$ (see [HCII, Corollary XI.17]). Now Proposition I.36 shows that $|U|$ is precisely the colimit we're interested in, and

$$U \times_{x/\mathcal{C}} \{\text{id}_x\} \simeq Y^{\mathcal{C}}/E \times_{\mathcal{C}} \{x\} \simeq \text{Hom}_{\mathcal{P}(\mathcal{C})}(Y^{\mathcal{C}}(x), E) \simeq E(x)$$

holds by a quick unravelling of definitions and Yoneda's lemma. \square

I.53b. Lemma*. — *Let \mathcal{D} be cocomplete. For every functor $F : \mathcal{C} \rightarrow \mathcal{D}$, the left Kan extension $\text{Lan}_{Y^{\mathcal{C}}} F : \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{D}$ commutes with colimits.*

Proof.* One is tempted to say “This follows immediately from the pointwise formula (Theorem I.52) and the fact that colimits commute (Proposition I.42)”, but in reality it is way more subtle than that. If you don't believe me, have a look at Warning* I.53c below. Anyway, let's move onward to the problem at hand.

Let $\Theta : \mathcal{I} \rightarrow \mathcal{P}(\mathcal{C})$ be a diagram with colimit $E \simeq \text{colim}_{\mathcal{I}} \Theta$. Abusively we will also write $\Theta(i) = E_i$ and $E \simeq \text{colim}_{i \in \mathcal{I}} E_i$. Define \mathcal{J} as the pullback

$$\begin{array}{ccccccc} \mathcal{J} & \longrightarrow & Y^{\mathcal{C}}/\mathcal{P}(\mathcal{C}) & \xrightarrow{s} & \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \downarrow & & \lrcorner & & \downarrow t & \nearrow \text{Lan}_{Y^{\mathcal{C}}} F & \\ \mathcal{I} & \xrightarrow{\Theta} & \mathcal{P}(\mathcal{C}) & & & & \end{array}$$

Note that $t: Y^{\mathcal{C}}/\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{C})$ is a cocartesian fibration by an easy generalisation of Example 1.25(b), hence so is $\mathcal{J} \rightarrow \mathcal{I}$. The pointwise formula from Theorem 1.52 yields

$$\mathrm{Lan}_{Y^{\mathcal{C}}} F \left(\mathrm{colim}_{i \in \mathcal{I}} E_i \right) \simeq \mathrm{Lan}_{Y^{\mathcal{C}}} F(E) \simeq \mathrm{colim} \left(Y^{\mathcal{C}}/E \longrightarrow \mathcal{C} \xrightarrow{F} \mathcal{D} \right)$$

Using the pointwise formula for each E_i together with Proposition 1.42, we find that

$$\mathrm{colim}_{i \in \mathcal{I}} \mathrm{Lan}_{Y^{\mathcal{C}}} F(E_i) \simeq \mathrm{colim} \left(\mathcal{J} \longrightarrow \mathcal{C} \xrightarrow{F} \mathcal{D} \right).$$

We obtain a canonical map $\mathcal{J} \rightarrow Y^{\mathcal{C}}/E$ as follows: Extend $\Theta: \mathcal{I} \rightarrow \mathcal{P}(\mathcal{C})$ to its colimit cocone $\Theta^{\flat}: \mathcal{I}^{\flat} \rightarrow \mathcal{P}(\mathcal{C})$. Since $\mathcal{I}^{\flat} = (\mathcal{I} \times \Delta^1)/(\mathcal{I} \times \{1\})$, this can be further extended to a map $\mathcal{I} \times \Delta^1 \rightarrow \mathcal{P}(\mathcal{C})$ which is $\mathrm{const} E$ on $\mathcal{I} \times \{1\}$. Pulling back $t: Y^{\mathcal{C}}/\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{C})$ along $\mathcal{I} \times \Delta^1 \rightarrow \mathcal{P}(\mathcal{C})$ gives a cocartesian fibration over $\mathcal{I} \times \Delta^1$, hence a cocartesian fibration over Δ^1 by composing with $\mathrm{pr}_2: \mathcal{I} \times \Delta^1 \rightarrow \Delta^1$. After applying $\mathrm{St}^{\mathrm{cocart}}$ this gives a map $\mathcal{J} \rightarrow \mathcal{I} \times Y^{\mathcal{C}}/E$, providing the required map $\mathcal{J} \rightarrow Y^{\mathcal{C}}/E$.

Now observe that $\mathcal{J} \rightarrow \mathcal{C}$ is a cartesian fibration and $Y^{\mathcal{C}}/E \rightarrow \mathcal{C}$ is even a right fibration. Indeed, by construction and some abstract nonsense about pullbacks, we get pullback diagrams

$$\begin{array}{ccc} \mathcal{J} & \longrightarrow & \mathcal{C} \\ \downarrow & \lrcorner & \downarrow_{Y^{\mathcal{C}}} \\ \mathcal{P}(\mathcal{C})/\Theta & \xrightarrow{s} & \mathcal{P}(\mathcal{C}) \end{array} \quad \text{and} \quad \begin{array}{ccc} Y^{\mathcal{C}}/E & \longrightarrow & \mathcal{C} \\ \downarrow & \lrcorner & \downarrow_{Y^{\mathcal{C}}} \\ \mathcal{P}(\mathcal{C})/E & \xrightarrow{s} & \mathcal{P}(\mathcal{C}) \end{array}$$

Now $\mathcal{P}(\mathcal{C})/\Theta \rightarrow \mathcal{P}(\mathcal{C})$ is a cartesian fibration by an easy generalisation of the dual of Example 1.25(b) and $\mathcal{P}(\mathcal{C})/E \rightarrow \mathcal{P}(\mathcal{C})$ is a right fibration. The right-hand side shows that $\mathrm{St}^{\mathrm{cart}}(Y^{\mathcal{C}}/E \rightarrow \mathcal{C}) \simeq \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(Y^{\mathcal{C}}(-), E) \simeq E(-)$ as functors $\mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{An}$ by Yoneda's lemma. Hence the map $\mathcal{J} \rightarrow Y^{\mathcal{C}}/E$ constructed above induces a natural transformation

$$\mathrm{St}^{\mathrm{cart}}(\mathcal{J} \rightarrow \mathcal{C}) \Longrightarrow E.$$

Since $\mathbf{An} \subseteq \mathbf{Cat}_{\infty}$ has a left-adjoint $|\cdot|: \mathbf{Cat}_{\infty} \rightarrow \mathbf{An}$, this transformation factors over $|\mathrm{St}^{\mathrm{cart}}(\mathcal{J} \rightarrow \mathcal{C})|$. In fact, we will show that $|\mathrm{St}^{\mathrm{cart}}(\mathcal{J} \rightarrow \mathcal{C})| \simeq E!$. This can be done objectwise, so we need to show $|\mathcal{J} \times_{\mathcal{C}} \{x\}| \simeq E(x)$ for all $x \in \mathcal{C}$. We have $E(x) \simeq \mathrm{colim}_{i \in \mathcal{I}} E_i(x)$ as colimits of presheaves are computed pointwise by Lemma 1.39. The colimit on the right-hand is precisely the colimit over $\mathrm{ev}_x \circ \Theta: \mathcal{I} \rightarrow \mathcal{P}(\mathcal{C}) \rightarrow \mathbf{An}$. This can be computed via Proposition 1.36, so it suffices to show $\mathcal{J} \times_{\mathcal{C}} \{x\} \simeq \mathrm{Un}^{\mathrm{cocart}}(\mathrm{ev}_x \circ \Theta)$. By Yoneda's lemma, the diagram

$$\begin{array}{ccc} \mathcal{I} & \xrightarrow{\mathrm{ev}_x \circ \Theta} & \mathbf{An} \\ \Theta \downarrow & \nearrow & \\ \mathcal{P}(\mathcal{C}) & \xrightarrow{\mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(Y^{\mathcal{C}}(x), -)} & \end{array}$$

commutes, hence $\mathrm{Un}^{\mathrm{cocart}}(\mathrm{ev}_x \circ \Theta)$ is the pullback of the left fibration $Y^{\mathcal{C}}(x)/\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{P}(\mathcal{C})$ along Θ . By abstract pullback nonsense again, this agrees with $\mathcal{J} \times_{\mathcal{C}} \{x\}$, as required.

Now that we know $|\mathrm{St}^{\mathrm{cart}}(\mathcal{J} \rightarrow \mathcal{C})| \simeq E$, we can finally prove that $\mathcal{J} \rightarrow Y^{\mathcal{C}}/E$ is cofinal, which ultimately shows what we want. Since $|\cdot|: \mathbf{Cat}_{\infty} \rightarrow \mathbf{An}$ is a left adjoint of $\mathbf{An} \subseteq \mathbf{Cat}_{\infty}$, we obtain $\mathrm{Hom}_{\mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{Cat}_{\infty})}(\mathrm{St}^{\mathrm{cart}}(\mathcal{J} \rightarrow \mathcal{C}), G) \simeq \mathrm{Hom}_{\mathcal{P}(\mathcal{C})}(E, G)$ for all presheaves $G: \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{An}$, hence

$$\mathrm{Hom}_{\mathrm{Cart}(\mathcal{C})}(\mathcal{J}, X) \simeq \mathrm{Hom}_{\mathrm{Right}(\mathcal{C})}(Y^{\mathcal{C}}/E, X)$$

for right fibrations $X \rightarrow \mathcal{C}$. This easily implies $\mathrm{Fun}_{Y^{\mathcal{C}}/E}(\mathcal{J}, X) \simeq \mathrm{Fun}_{Y^{\mathcal{C}}/E}(Y^{\mathcal{C}}/E, X)$ for all right fibrations $X \rightarrow Y^{\mathcal{C}}/E$, hence $\mathcal{J} \rightarrow Y^{\mathcal{C}}/E$ is indeed cofinal in the sense of [HTT, Definition 4.1.1.1], which is equivalent to our Definition I.44. \square

I.53c. Warning* $\triangle!$. — In the situation of Theorem I.52, there examples where $f: \mathcal{C} \rightarrow \mathcal{E}$ is fully faithful and colimit-dense, but still $\mathrm{Lan}_f F$ doesn't commute with colimits (but of course the functor Lan_f itself does preserve colimits as it is a left adjoint). Here's a counterexample, I think. Let $\mathcal{C} = \mathbb{N} \sqcup \{c\}$ be the poset consisting of \mathbb{N} and a totally unrelated additional point c , such that $\mathrm{Hom}_{\mathcal{C}}(c, n) = \emptyset = \mathrm{Hom}_{\mathcal{C}}(n, c)$ for all $n \in \mathbb{N}$. Let $\mathcal{E} = \mathcal{C}^{\triangleright}$ be the cocone below \mathcal{C} , with tip $e \in \mathcal{E}$, and let $f: \mathcal{C} \rightarrow \mathcal{E}$ be the canonical inclusion. Let $F: \mathcal{C} \rightarrow \mathbf{An}$ be the functor that is constant with value $\emptyset \in \mathbf{An}$ on \mathbb{N} and takes $F(c) = * \in \mathbf{An}$. Then $\mathrm{colim}_{n \in \mathbb{N}} n \simeq e$ and we have

$$\mathrm{colim}_{n \in \mathbb{N}} \mathrm{Lan}_f F(n) \simeq \mathrm{colim}_{n \in \mathbb{N}} \emptyset = \emptyset$$

(here we use Corollary I.54 below). However, $\mathrm{Lan}_f F(e) \neq \emptyset$, because there is a map $F(c) \rightarrow \mathrm{Lan}_f F(e)$ by the pointwise formula.

The key problem in this counterexample is the existence of a map $c \rightarrow e \simeq \mathrm{colim}_{n \in \mathbb{N}} n$ that doesn't factor over any $c \rightarrow n$, and the only reason why this doesn't happen for $Y^{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{P}(\mathcal{C})$ as well is the Yoneda lemma! This shows that Lemma* I.53b critically depends on the Yoneda lemma and perhaps justifies the effort we had to put into its proof.

Proof of Theorem I.51.* From Theorem I.52 we obtain that $Y^{\mathcal{C},*}$ admits a left adjoint $\mathrm{Lan}_{Y^{\mathcal{C}}}: \mathrm{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \mathrm{Fun}(\mathcal{P}(\mathcal{C}), \mathcal{D})$. We have verified in Lemma* I.53b that $\mathrm{Lan}_{Y^{\mathcal{C}}}$ takes values in the full sub- ∞ -category $\mathrm{Fun}^{\mathrm{colim}}(\mathcal{P}(\mathcal{C}), \mathcal{D})$ of colimit-preserving functors. We first show that

$$\mathrm{Lan}_{Y^{\mathcal{C}}}: \mathrm{Fun}(\mathcal{C}, \mathcal{D}) \xrightarrow[\sim]{\sim} \mathrm{Fun}^{\mathrm{colim}}(\mathcal{P}(\mathcal{C}), \mathcal{D}) : Y^{\mathcal{C},*}$$

is an equivalence. It suffices to show that the unit and counit of $(\mathrm{Lan}_{Y^{\mathcal{C}}}, Y^{\mathcal{C},*})$ are equivalences. For the counit, this follows from Corollary I.54 below. For the unit, let $G: \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{D}$ be a colimit-preserving functor. We need to prove that $G \simeq \mathrm{Lan}_{Y^{\mathcal{C}}}(G \circ Y^{\mathcal{C},*})$. But both sides coincide on representable presheaves (i.e. after precomposition with $Y^{\mathcal{C}}$, as we have just verified) and every presheaf can be written as a colimit of representable ones by Lemma* I.53a, hence the claim.

It remains to check that $\mathrm{Fun}^L(\mathcal{P}(\mathcal{C}), \mathcal{D}) \subseteq \mathrm{Fun}^{\mathrm{colim}}(\mathcal{P}(\mathcal{C}), \mathcal{D})$ is an equivalence, i.e. that every colimit-preserving functor $\mathcal{P}(\mathcal{C}) \rightarrow \mathcal{D}$ admits a right adjoint. This can be written down explicitly, which we do in Corollary I.55 below. \square

I.54. Corollary. — *If $f: \mathcal{C} \rightarrow \mathcal{E}$ is fully faithful in the situation of Theorem I.52, then*

$$F \simeq \mathrm{Lan}_f F \circ f \quad \text{and} \quad F \simeq \mathrm{Ran}_f F \circ f$$

holds for all functors $F: \mathcal{C} \rightarrow \mathcal{D}$.

Proof. The natural transformations in question are induced by the unit of the (Lan_f, f^*) -adjunction and the counit of the (f^*, Ran_f) -adjunction, so it suffices to check both of them on objects $c \in \mathcal{C}$. But if f is fully faithful, then the slice categories $f/f(c)$ have $(c, \mathrm{id}_{f(c)})$ as terminal objects, hence colimits over $f/f(c)$ are computed by evaluation at that object, proving $\mathrm{Lan}_f F(f(c)) \simeq F(c)$. Same for $\mathrm{Ran}_f F(f(c))$. \square

I.55. Corollary. — *In the situation of Theorem I.51, let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor corresponding to $\text{Lan}_{Y^{\mathcal{C}}} F: \mathcal{P}(\mathcal{C}) \rightarrow \mathcal{D}$. Then $\text{Lan}_{Y^{\mathcal{C}}} F$ admits a right-adjoint $R: \mathcal{D} \rightarrow \mathcal{P}(\mathcal{C})$, which is given by*

$$R(d) \simeq \text{Hom}_{\mathcal{D}}(F -, d): \mathcal{C}^{\text{op}} \longrightarrow \text{An}$$

on objects $d \in \mathcal{D}$.

Proof. To check that $\text{Lan}_{Y^{\mathcal{C}}} F$ admits a right adjoint, it suffices to check that all $d \in \mathcal{D}$ admit right-adjoint objects under $\text{Lan}_{Y^{\mathcal{C}}} F$ by Corollary I.32. We already have a candidate for $R(d)$, so we need to check that there is an equivalence

$$\text{Hom}_{\mathcal{D}}(\text{Lan}_{Y^{\mathcal{C}}} F -, d) \simeq \text{Nat}(-, R(d))$$

of functors $\mathcal{P}(\mathcal{C})^{\text{op}} \rightarrow \text{An}$. But regarding both sides as functors $\mathcal{P}(\mathcal{C}) \rightarrow \text{An}^{\text{op}}$ instead, note that they preserve colimits and they agree on representables since both become $\text{Hom}_{\mathcal{D}}(F -, d)$ after precomposition with $Y^{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{P}(\mathcal{C})$ (for the left-hand side we need to use Corollary I.54, for the right-hand side use Yoneda's lemma). Hence these two functors must be equivalent by Theorem I.51. To complete the proof that $R(d)$ is a right-adjoint object of d , we need to check that the above equivalence is actually induced by a map $\text{Lan}_{Y^{\mathcal{C}}} F(R(d)) \rightarrow d$, but this we get for free by plugging $R(d)$ into the above equivalence and taking the of $\text{id}_{R(d)} \in \text{Nat}(R(d), R(d))$ in the left-hand side. \square

This finishes the proofs of Theorem I.51 and Theorem I.52. Time for examples!

I.56. Very Long Example. — We will show that algebraic topology is a corollary of Theorem I.51. Let $\mathcal{C} = *$. Then the theorem shows that the evaluation at the point $*$ $\in \text{An}$ is an equivalence

$$\text{ev}_*: \text{Fun}^L(\text{An}, \mathcal{D}) \xrightarrow{\sim} \mathcal{D}.$$

for every cocomplete \mathcal{D} . For example, take $\mathcal{D} = \mathcal{D}_{\geq 0}(\mathbb{Z})$ to be the derived category of abelian groups (which is cocomplete—after “Corollary” I.46 and Corollary I.50 there isn't much to do). Then the equivalence $\mathcal{D}_{\geq 0}(\mathbb{Z}) \simeq \text{Fun}^L(\text{An}, \mathcal{D}_{\geq 0}(\mathbb{Z}))$ from above takes $\mathbb{Z}[0] \in \mathcal{D}_{\geq 0}(\mathbb{Z})$ to the “normalized chain complex” functor

$$C_{\bullet}: \text{An} \longrightarrow \mathcal{D}_{\geq 0}(\mathbb{Z}),$$

that is, the homology of $C_{\bullet}(X)$ gives the unreduced homology of X (as a simplicial set, or equivalently the unreduced cellular/singular homology of the realisation $|X| \in \text{CW}$). The fact that C_{\bullet} preserves coproducts and pushouts precisely gives that simplicial/cellular/singular homology takes disjoint unions to direct sums and satisfies a Mayer–Vietoris sequence.

The right adjoint of C_{\bullet} is

$$K: \mathcal{D}_{\geq 0}(\mathbb{Z}) \longrightarrow \text{An}$$

(this is another K that has nothing to do with K -theory), which is now our official definition of *Eilenberg–MacLane spaces anima*. On objects $C \in \mathcal{D}_{\geq 0}(\mathbb{Z})$ it can be explicitly described as $K(C) \simeq \text{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(\mathbb{Z}[0], C)$ by Corollary I.55. The (C_{\bullet}, K) adjunction can be upgraded to an adjunction

$$\tilde{C}_{\bullet}: */\text{An} \rightleftarrows \mathcal{D}_{\geq 0}(\mathbb{Z}) : K.$$

Here $\tilde{C}_{\bullet}(X, x)$ is the complex computing the *reduced homology* of a pointed anima (X, x) . For example, if \mathbb{S}^i is your favourite model for the i -sphere in anima, then $C_{\bullet}(\mathbb{S}^i, *) \simeq \mathbb{Z}[i]$.

This new adjunction needs a bit of justification, but to not interrupt the flow we'll discuss that later in Remark* [I.56c](#). For now let's compute

$$\begin{aligned}\pi_i K(C) &\simeq \pi_0 \operatorname{Hom}_{*/\mathbf{An}}((\mathbb{S}^i, *), K(C)) \\ &\simeq \pi_0 \operatorname{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(C_\bullet(\mathbb{S}^i, *), C) \\ &\simeq \pi_0 \operatorname{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(\mathbb{Z}[i], C) \\ &\simeq H_i(C).\end{aligned}$$

In fact, we can say more: Let A be an abelian group and $A[n] \in \mathcal{D}_{\geq 0}(\mathbb{Z})$ the complex consisting of A placed in degree n (technically we have to replace $A[n]$ by a projective resolution to be consistent with Example [I.15\(e\)](#)). Then $K(A, n) := K(A[n])$ satisfies

$$\begin{aligned}\pi_i \operatorname{Hom}_{\mathbf{An}}(X, K(A, n)) &\simeq \pi_i \operatorname{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(C_\bullet(X), A[n]) \\ &\simeq H_{i-n} C^\bullet(X, A) \\ &\simeq H^{n-i}(X, A),\end{aligned}$$

where $C^\bullet(X, A) \in \mathcal{D}_{\leq 0}(\mathbb{Z})$ is the complex of simplicial/cellular/singular cochains with coefficients in A . If X is pointed and connected, the same calculation works with reduced cohomology (this only matters for $n = i$). We thus obtain:

I.56a. Theorem (Eilenberg–MacLane). — *The functors $H^n(-, A): \pi \mathbf{An} \rightarrow \operatorname{Set}$ (unreduced cohomology) and $\tilde{H}^n(-, A): \pi(*/\mathbf{An}) \rightarrow \operatorname{Set}$ (reduced cohomology) are represented by $K(A, n)$.*

In particular, if $X \in */\mathbf{An}$ satisfies $\pi_i X = 0$ for $i \neq n$, then $H := \operatorname{Hom}_{*/\mathbf{An}}(X, K(A, n))$ is given by the discrete anima $H^n(X, A)$: Indeed, we have $\pi_i H = \tilde{H}^{n-i}(X, A)$, which vanishes for $1 \leq i \leq n$ by Hurewicz and the universal coefficient theorem, and for $i > n$ by the fact that cohomology is zero in negative degrees. So $H \rightarrow \pi_0 H$ is a homotopy equivalence by Whitehead's theorem. Using universal coefficients again, we can summarize this by

$$\operatorname{Hom}_{*/\mathbf{An}}(X, K(A, n)) \simeq H^n(X, A) \simeq \operatorname{Hom}_{\mathbf{Ab}}(H_n(X), A).$$

For $n \geq 1$, an isomorphism on the right-hand side gives an equivalence on the left-hand side by a combination of Hurewicz and Whitehead (for $n = 1$ we really need that A is abelian for this argument to work). This shows that $K(-, n): \mathbf{Ab} \rightarrow */\mathbf{An}$ is fully faithful for $n \geq 1$.

Now consider the full sub- ∞ -category $*/\mathbf{An}_{\leq n} \subseteq */\mathbf{An}$ spanned by n -truncated anima, i.e. those with vanishing π_i for $i > n$. The inclusion has a left-adjoint $\tau_{\leq n}: */\mathbf{An} \rightarrow */\mathbf{An}_{\leq n}$. One can construct $\tau_{\leq n} X$ by iteratively attaching cells to X to kill higher and higher homotopy groups and then use that adjunctions can be constructed objectwise by Corollary [I.32](#). In fact, one can take $\tau_{\leq n} X = \operatorname{cosk}_{n+1} X$, since the right-hand side is a really brutal way of killing homotopy groups in degrees n and higher. In particular, the n -truncation $\tau_{\leq n} X$ satisfies

$$\pi_i(\tau_{\leq n} X) \simeq \begin{cases} 0 & \text{if } i > n \\ \pi_i X & \text{if } i \leq n \end{cases}.$$

The unit map $X \rightarrow \tau_{\leq n} X$ factors over $\tau_{\leq n+1} X \rightarrow \tau_{\leq n} X$, since $\tau_{\leq n} X$ is already $(n+1)$ -truncated. We claim that the homotopy fibre F (i.e. the fibre taken inside the ∞ -category \mathbf{An}) of this map is equivalent to $K(\pi_{n+1} X, n+1)$. Indeed, the long exact sequence of homotopy

groups shows that $\pi_i F = 0$ for $i \neq n+1$ and that $\pi_{n+1} F \simeq \pi_{n+1} X$. Hence, by the discussion after Theorem I.56a,

$$\mathrm{Hom}_{*/\mathrm{An}}(F, K(\pi_{n+1} X, n+1)) \simeq \mathrm{Hom}_{\mathrm{Ab}}(H_{n+1}(F), \pi_{n+1} X),$$

and an isomorphism on the right-hand side induces an isomorphism on the left-hand side. But $H_{n+1}(F) \simeq \pi_{n+1} X$ by Hurewicz, hence we do obtain an equivalence as desired.

The diagram

$$X \longrightarrow \left(\dots \longrightarrow \tau_{\leq n+1} X \longrightarrow \tau_{\leq n} X \longrightarrow \dots \longrightarrow \tau_{\leq 1} X \right)$$

is called the *Postnikov tower* of X . It's not hard to show that the Postnikov tower induces an equivalence $X \xrightarrow{\sim} \lim_{n \in \mathbb{N}^{\mathrm{op}}} \tau_{\leq n} X$ in $*/\mathrm{An}$ when X is connected (see [Hat02, Corollary 4.68] for example). The Postnikov tower is an important organisational and computational tool, since it “stratifies” $*/\mathrm{An}$ into pieces, whose “difference is controlled by Ab ” via the Eilenberg–MacLane functor $K(-, n): \mathrm{Ab} \rightarrow */\mathrm{An}$.

I.56b. Exercise. — Show that every chain complex over \mathbb{Z} (in fact, over any PID) is quasi-isomorphic via a zigzag to its homology, considered as a chain complex all of whose differentials are 0.

In particular, Exercise I.56b implies that for $C \in \mathcal{D}_{\geq 0}(\mathbb{Z})$ there is a (non-canonical) equivalence

$$C \simeq \bigoplus_{i=0}^{\infty} H_i(C)[i] \simeq \prod_{i=0}^{\infty} H_i(C)[i].$$

Since the right-adjoint functor K preserves limits (Observation I.41), this implies $K(C) \simeq \prod_{i=0}^{\infty} K(H_i(C), i)$. In general, anima X with the property

$$X \simeq \prod_{i=0}^{\infty} K(\pi_i X, i)$$

are called *gems* (a pun on “generalized Eilenberg–MacLane spaces”). It is also easy to check that the following diagram is a pullback in the ∞ -category $\mathcal{D}_{\geq 0}(\mathbb{Z})$:

$$\begin{array}{ccc} \tau_{\geq 0}(C[-1]) & \longrightarrow & 0 \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & C \end{array}$$

Again, since K preserves limits, we get $K(\tau_{\geq 0} C[-1]) \simeq \Omega K(C)$ (indeed if you take the same pullback in An , it will give you the loop space). In particular, $K(A, n-1) \simeq \Omega K(A, n)$ for all $n \geq 1$. And that's our first spectrum!

I.56c. Remark*. — To construct a lift $K: \mathcal{D}_{\geq 0}(\mathbb{Z}) \rightarrow */\mathrm{An}$, Fabian explained that $0 \in \mathcal{D}_{\geq 0}(\mathbb{Z})$ is an initial object and $K(0) \simeq \mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(\mathbb{Z}[0], 0) \simeq *$, so K lifts to a functor $K: \mathcal{D}_{\geq 0}(\mathbb{Z}) \simeq 0/\mathcal{D}_{\geq 0}(\mathbb{Z}) \rightarrow */\mathrm{An}$ on slice categories. I claimed that $K(0) \simeq \emptyset$ in a previous version of these notes, but that's nonsense and Fabian was completely right!

Yet the fact that K has a left adjoint is not entirely automatic; more precisely, it's not an instance where an adjunction automatically descends to slice categories (this will

happen occasionally in the future). The reason is $C_\bullet(*) \simeq \mathbb{Z}[0]$, so C_\bullet doesn't lift to a functor $*/\mathbf{An} \rightarrow 0/\mathcal{D}_{\geq 0}(\mathbb{Z}) \simeq \mathcal{D}_{\geq 0}(\mathbb{Z})$ on slice categories. Instead, we have to define a functor $\tilde{C}_\bullet: */\mathbf{An} \rightarrow \mathcal{D}_{\geq 0}(\mathbb{Z})$ taking a pointed anima (X, x) to the pushout

$$\begin{array}{ccc} \mathbb{Z}[0] \simeq C_\bullet(x) & \longrightarrow & C_\bullet(X) \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & \tilde{C}_\bullet(X, x) \end{array}$$

or in other words, to the *cofibre* $\mathrm{cofib}(C_\bullet(x) \rightarrow C_\bullet(X))$. It's straightforward to check that \tilde{C}_\bullet is indeed left-adjoint to K . Indeed, from Corollary I.50 and the computation of Hom anima in slice categories ([HCII, Proposition VIII.6]) we get

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(C_\bullet(X, x), D) &\simeq \mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(C_\bullet(X), D) \times_{\mathrm{Hom}_{\mathcal{D}_{\geq 0}(\mathbb{Z})}(C_\bullet(x), D)} \{0\} \\ &\simeq \mathrm{Hom}_{\mathbf{An}}(X, K(D)) \times_{\mathrm{Hom}_{\mathbf{An}}(x, K(D))} \{K(0) \simeq * \rightarrow K(D)\} \\ &\simeq \mathrm{Hom}_{*/\mathbf{An}}((X, x), K(D)) \end{aligned}$$

(all pullbacks are taken in \mathbf{An} , as announced on Page 34). The complex $\tilde{C}_\bullet(X, x)$ computes reduced homology. For example, if \mathbb{S}^i denotes the i -sphere, then $C_\bullet(\mathbb{S}^i, *) \simeq \mathrm{cofib}(\mathbb{Z}[0] \rightarrow \mathbb{Z}[0] \oplus \mathbb{Z}[i]) \simeq \mathbb{Z}[i]$, as one would expect.

I.57. Example. — Two more applications of Theorem I.51:

- (a) If \mathcal{D} is an ∞ -category, we will again write $s\mathcal{D}$ and $c\mathcal{D}$ for the functor categories $\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathcal{D})$ and $\mathrm{Fun}(\Delta, \mathcal{D})$ respectively. If \mathcal{D} is cocomplete and $F \in c\mathcal{D}$ a cosimplicial object in \mathcal{D} , we obtain an adjunction

$$| \quad |_F: s\mathbf{An} \rightleftarrows \mathcal{D} : \mathrm{Sing}_F .$$

In the case where \mathcal{D} is a 1-category, we also get an adjunction $| \quad |_F: s\mathbf{Set} \rightleftarrows \mathcal{D} : \mathrm{Sing}_F^1$ from Theorem I.2. To see how these two adjunctions are related, recall that $s\mathbf{Set} \subseteq s\mathbf{An}$ has a left adjoint $\pi_0: s\mathbf{An} \rightarrow s\mathbf{Set}$ by Example I.33(a) and Observation I.40, which yields a factorisation

$$| \quad |_F: s\mathbf{An} \xrightleftharpoons[\mathrm{incl}]{\pi_0} s\mathbf{Set} \xrightleftharpoons[\mathrm{Sing}_F^1]{| \quad |_F} \mathcal{D} : \mathrm{Sing}_F$$

of the adjunction above.

Warning $\triangle!$. — Right now there is another possible definition of π_0 . Recall that $\pi_0: s\mathbf{Set} \rightarrow \mathbf{Set}$ can also be written as $\mathrm{colim}_{\Delta^{\mathrm{op}}} = | \quad |_{\mathrm{const}*}^1$, where the cosimplicial set $\mathrm{const}: \Delta \rightarrow \mathbf{Set}$ sends everything to a point. Analogously, we have a functor $\mathrm{colim}_{\Delta^{\mathrm{op}}} = | \quad |_{\mathrm{const}*}$: $s\mathbf{An} \rightarrow \mathbf{An}$, which arises via Theorem I.51 from the cosimplicial anima $\mathrm{const}: \Delta \rightarrow \mathbf{An}$. We will never call this functor π_0 , but $\mathrm{colim}_{\Delta^{\mathrm{op}}}$ instead, which is what it really is.

- (b) Consider $\Delta \rightarrow \mathrm{Cat}_1 \subseteq \mathrm{Cat}_1^{(2)} \subseteq \mathrm{Cat}_\infty$ sending $[n]$ to itself, considered as a poset. By (a), this cosimplicial ∞ -category induces an adjunction

$$\mathrm{asscat}: s\mathbf{An} \rightleftarrows \mathrm{Cat}_\infty : \mathbf{N}^r$$

(Fabian remarks that “asscat” is short for “associated category” and has nothing to do with “Arschkatzen”). The right adjoint N^r is called the *Rezk nerve* and is a very important construction! If \mathcal{C} is an ∞ -category, Corollary 1.55 provides the explicit description

$$N_n^r(\mathcal{C}) \simeq \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathcal{C}) \simeq \mathrm{core} \mathrm{Fun}([n], \mathcal{C}).$$

Beware that if \mathcal{C} is a 1-category, then $N^r(\mathcal{C})$ is different from the ordinary nerve $N(\mathcal{C}) = ([n] \mapsto \mathrm{Hom}_{\mathrm{Cat}_1}([n], \mathcal{C}))$, since $\mathrm{Cat}_1 \subseteq \mathrm{Cat}_1^{(2)}$ is not fully faithful. In fact, $N(\mathcal{C})$ is a discrete simplicial anima, but $N^r(\mathcal{C})$ is usually not. After a brief detour to Bousfield localisations, we will come back to Rezk nerves in Theorem/Definition 1.64.

Bousfield Localisations

LECTURE 6
17th Nov, 2020

1.58. Proposition/Definition. — *Let $L: \mathcal{C} \rightleftarrows \mathcal{D}: R$ be an adjoint pair.*

- (a) *The functor R is fully faithful iff the counit $c: LR \Rightarrow \mathrm{id}_{\mathcal{D}}$ is an equivalence.*
- (b) *A morphism $f: x \rightarrow y$ in \mathcal{C} is taken to an equivalence by L iff it is a left R -local equivalence, i.e. iff the induced map*

$$f^*: \mathrm{Hom}_{\mathcal{C}}(y, Rd) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(x, Rd)$$

is an equivalence for all $d \in \mathcal{D}$. We will often drop the “left” part and just write “ R -local equivalence”.

Moreover, if the equivalent conditions from (a) hold, then:

- (c) *The unit $c \rightarrow RLc$ is always a left R -local equivalence.*
- (d) *$L: \mathcal{C} \rightarrow \mathcal{D}$ is a localisation at the left R -local equivalences.*

One then says that L is a left Bousfield localisation. In particular, $\mathcal{C}[\{R\text{-local equiv.}\}^{-1}]$ is again locally small. Of course, there's also a dual notion of right Bousfield localisations.

Proof. Fabian decided to prove (b) first. By Yoneda's lemma, for $Lf: Lx \rightarrow Ly$ to be an equivalence, it suffices to show that it induces an equivalence after $\mathrm{Hom}_{\mathcal{D}}(-, d)$ for all $d \in \mathcal{D}$. Using the natural adjunction equivalence $\mathrm{Hom}_{\mathcal{D}}(L-, -) \simeq \mathrm{Hom}_{\mathcal{C}}(-, R-)$, we obtain a homotopy commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(y, Rd) & \xrightarrow{f^*} & \mathrm{Hom}_{\mathcal{C}}(x, Rd) \\ \downarrow \wr & & \downarrow \wr \\ \mathrm{Hom}_{\mathcal{D}}(Ly, d) & \xrightarrow{Lf^*} & \mathrm{Hom}_{\mathcal{D}}(Lx, d) \end{array}$$

in anima, which immediately proves (b).

Next up, we show (a). Bastiaan found a proof which is considerably simpler than the one from the lecture, so we shall include his proof here. Observe that the diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{D}}(d, x) & \xrightarrow{c_d^*} & \mathrm{Hom}_{\mathcal{D}}(LRd, x) \\ R \downarrow & \nearrow \sim & \\ \mathrm{Hom}_{\mathcal{C}}(Rd, Rx) & & \end{array}$$

commutes (in $\pi\mathbf{An}$) for all $d, x \in \mathcal{D}$. Indeed, since Yoneda's lemma tells us that

$$\mathrm{ev}_{\mathrm{id}_d} : \mathrm{Nat}(\mathrm{Hom}_{\mathcal{D}}(d, -), \mathrm{Hom}_{\mathcal{D}}(LRd, -)) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(LRd, d)$$

is an equivalence, it suffices to check commutativity in the special case $x = d$ on the single element $\mathrm{id}_d \in \mathrm{Hom}_{\mathcal{D}}(d, d)$. But id_d is clearly mapped to $c_d \in \mathrm{Hom}_{\mathcal{D}}(LRd, d)$ in both cases, proving that the diagram does indeed commute. By Theorem I.19(b) and Yoneda's lemma again, $c : LR \Rightarrow \mathrm{id}_{\mathcal{D}}$ is an equivalence iff $c_d^* : \mathrm{Hom}_{\mathcal{D}}(d, x) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(LRd, x)$ is an equivalence for all $x \in \mathcal{D}$. By the diagram above, this holds iff $R : \mathrm{Hom}_{\mathcal{D}}(d, x) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{C}}(Rd, Rx)$ is an equivalence for all $d, x \in \mathcal{D}$, i.e. iff R is fully faithful.

Now for (c). By one of the triangle identities, the composition

$$Lx \xrightarrow{Lu_x} LRLx \xrightarrow{c_{Lx}} Lx$$

is equivalent to id_{Lx} for all $x \in \mathcal{C}$. Since we assume that the conditions from (a) hold, c_{Lx} is an equivalence, hence Lu_x is one too. By (b) this means that u_x is an R -local equivalence.

For (d), let \mathcal{E} be any test category and consider the adjunction

$$R^* : \mathrm{Fun}(\mathcal{C}, \mathcal{E}) \rightleftarrows \mathrm{Fun}(\mathcal{D}, \mathcal{E}) : L^*$$

obtained from Observation I.40. Since the counit of (L, R) is an equivalence, the counit of (R^*, L^*) is an equivalence too (it is induced by the former counit). But the direction of the adjunction has switched, so now L^* is fully faithful by (a). We are left to show that the essential image of L^* is $\mathrm{Fun}^{\{R\text{-local equiv.}\}}(\mathcal{C}, \mathcal{E})$, i.e. those functors that invert R -local equivalences. But any $L^*F \simeq F \circ L$ inverts the R -local equivalences by (b), hence the essential image of L^* is contained in $\mathrm{Fun}^{\{R\text{-local equiv.}\}}(\mathcal{C}, \mathcal{E})$. Conversely, for $F : \mathcal{C} \rightarrow \mathcal{E}$ inverting the R -local equivalences, we have $F \simeq FRL \simeq L^*(FR)$ since the unit $u_x : x \rightarrow RLx$ is an R -local equivalence for all $x \in \mathcal{C}$ by (c). Hence F is contained in the essential image of L^* . \square

Now that we know when adjunctions are Bousfield localisations, we would like to study the “converse” question, i.e. when localisations are Bousfield localisations.

I.59. Proposition. — *Let $W \subseteq \pi_0 \mathrm{core} \mathrm{Ar}(\mathcal{C})$ be some collection of morphisms with associated localisation $p : \mathcal{C} \rightarrow \mathcal{C}[W^{-1}]$. Then a map $\tau : px \rightarrow y$ in $\mathcal{C}[W^{-1}]$ exhibits x as a right-adjoint object to y under p iff τ is an equivalence and $\mathrm{Hom}_{\mathcal{C}}(-, x) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{An}$ inverts all morphisms from W .*

In particular, a right-adjoint functor R to p is automatically fully faithful. Note that this also characterizes the essential image of R as the (left) R -local objects, i.e. those $c \in \mathcal{C}$ such that $\mathrm{Hom}_{\mathcal{C}}(-, c) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{An}$ inverts (left) R -local equivalences.

We first prove the following lemma.

I.60. Lemma. — *Let $p : \mathcal{C} \rightarrow \mathcal{C}[W^{-1}]$ be a localisation.*

(a) *For all $c \in \mathcal{C}$, the functor $\mathrm{Hom}_{\mathcal{C}[W^{-1}]}(-, pc)$ is the left Kan extension of $\mathrm{Hom}_{\mathcal{C}}(-, c)$ along p^{op} . In diagrams:*

$$\begin{array}{ccc} \mathcal{C}^{\mathrm{op}} & \xrightarrow{\mathrm{Hom}_{\mathcal{C}}(-, c)} & \mathbf{An} \\ p^{\mathrm{op}} \downarrow & \nearrow \mathrm{Hom}_{\mathcal{C}[W^{-1}]}(-, pc) & \\ \mathcal{C}[W^{-1}]^{\mathrm{op}} & & \end{array}$$

(b) Any functor $F: \mathcal{C}[W^{-1}] \rightarrow \mathcal{D}$ is left Kan extended from $F \circ p$ along p . In diagrams:

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F \circ p} & \mathcal{D} \\ p \downarrow & \nearrow F & \\ \mathcal{C}[W^{-1}] & & \end{array}$$

The same is true for functors $F: \mathcal{C}[W^{-1}]^{\text{op}} \rightarrow \mathcal{D}$ and left Kan extension along p^{op} . In fact, $p^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}[W^{-1}]^{\text{op}}$ is a localisation of \mathcal{C}^{op} at W^{op} .

Warning $\triangle!$. — Don't get too excited and think “Yeah, now Theorem I.52 allows me to compute Hom anima in localisations!”, since the pointwise formula is useless (or rather tautological) here.

Proof of Lemma I.60. For (a), we apply Yoneda's lemma twice to obtain

$$\text{Nat}(\text{Hom}_{\mathcal{C}[W^{-1}]}(-, pc), G) \simeq G(pc) \simeq \text{Nat}(\text{Hom}_{\mathcal{C}}(-, c), G \circ p^{\text{op}})$$

for all $G: \mathcal{C}[W^{-1}]^{\text{op}} \rightarrow \mathcal{D}$. This is precisely what we want from a Kan extension along p^{op} . For (b), recall that $p^*: \text{Fun}(\mathcal{C}[W^{-1}], \mathcal{E}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{E})$ is fully faithful, i.e.

$$p^*: \text{Nat}(F, G) \xrightarrow{\sim} \text{Nat}(F \circ p, G \circ p)$$

is an equivalence for all F and G . This again exhibits F as a left Kan extension, along p this time. For the additional assertion, use the universal property or the explicit construction from [HCII, Theorem VIII.8]. \square

Proof of Proposition I.59. If $\text{Hom}_{\mathcal{C}}(-, x)$ inverts all arrows from W , then it descends to a functor $F: \mathcal{C}[W^{-1}]^{\text{op}} \rightarrow \text{An}$, which is left Kan extended from $\text{Hom}_{\mathcal{C}}(-, x)$ by Lemma I.60(b). But so is $\text{Hom}_{\mathcal{C}[W^{-1}]}(-, px)$ by Lemma I.60(a). Hence

$$\text{Hom}_{\mathcal{C}}(-, x) \simeq \text{Hom}_{\mathcal{C}[W^{-1}]}(p-, px) \xrightarrow{\tau_*} \text{Hom}_{\mathcal{C}[W^{-1}]}(p-, y)$$

is an equivalence since τ is one. This proves that x is indeed a right-adjoint object of y . For the converse, just read the argument backwards. As Sil and Bastiaan pointed out, we have to use that $p: \mathcal{C} \rightarrow \mathcal{C}[W^{-1}]$ is essentially surjective (use its universal property or the construction from [HCII, Theorem VIII.8]), so for τ to be an equivalence it does suffice that $\tau_*: \text{Hom}_{\mathcal{C}[W^{-1}]}(p-, px) \rightarrow \text{Hom}_{\mathcal{C}[W^{-1}]}(p-, y)$ is an equivalence.

For the additional assertions, assume that R is a right adjoint of p . Then the counit $pR \Rightarrow \text{id}$ consists of morphisms τ as above, hence it is an equivalence, hence R is fully faithful by Proposition/Definition I.58(a). A morphism in \mathcal{C} is an R -local equivalence iff it becomes an equivalence in $\mathcal{C}[W^{-1}]$ by Proposition/Definition I.58(b), so it is clear that the essential image of R is contained in the R -local objects by what we showed above. But conversely an R -local object $x \in \mathcal{C}$ is a right-adjoint object of px via the witness $\text{id}_{px}: px \rightarrow px$, hence x is in the essential image of R . \square

I.61. Corollary. — If $L: \mathcal{C} \rightarrow \mathcal{D}$ is a left Bousfield localisation and \mathcal{C} is complete or cocomplete, then so is \mathcal{D} . More precisely, we have

$$\text{colim}_{\mathcal{I}} F \simeq L\left(\text{colim}_{\mathcal{I}} RF\right) \quad \text{and} \quad \lim_{\mathcal{I}} F \simeq L\left(\lim_{\mathcal{I}} RF\right)$$

for any diagram $F: \mathcal{I} \rightarrow \mathcal{D}$.

Proof. We start with the formula for colimits. Compute

$$\mathrm{Hom}_{\mathcal{D}} \left(L \left(\mathrm{colim}_{\mathcal{I}} RF \right), - \right) \simeq \mathrm{Hom}_{\mathcal{C}} \left(\mathrm{colim}_{\mathcal{I}} RF, R- \right) \simeq \mathrm{Nat} \left(RF, \mathrm{const}(R-) \right)$$

Now R is fully faithful by Proposition/Definition I.58(a), hence so is the postcomposition functor $R_* : \mathrm{Fun}(\mathcal{I}, \mathcal{D}) \rightarrow \mathrm{Fun}(\mathcal{I}, \mathcal{C})$, which shows

$$\mathrm{Nat} \left(RF, \mathrm{const}(R-) \right) \simeq \mathrm{Nat}(F, \mathrm{const}-).$$

Hence $L(\mathrm{colim}_{\mathcal{I}} RF)$ satisfies the desired universal property for $\mathrm{colim}_{\mathcal{I}} F$.

For the limit part, first note that R -local objects are closed under limits. Indeed, if $x \simeq \lim_{j \in \mathcal{J}} x_j$ with each x_j an R -local object, then

$$\mathrm{Hom}_{\mathcal{C}}(-, x) \simeq \lim_{j \in \mathcal{J}} \mathrm{Hom}_{\mathcal{C}}(-, x_j)$$

by the dual of Corollary I.50, hence $\mathrm{Hom}_{\mathcal{C}}(-, x)$ inverts R -local equivalences because each $\mathrm{Hom}_{\mathcal{C}}(-, x_j)$ does. Moreover, R -local equivalences between R -local objects are actual equivalences by Yoneda's lemma applied to the full sub- ∞ -category of \mathcal{C} spanned by R -local objects. Now Proposition/Definition I.58 shows that the unit

$$u : \lim_{\mathcal{I}} RF \xrightarrow{\sim} RL \left(\lim_{\mathcal{I}} RF \right)$$

is an R -local equivalence, hence an actual equivalence as RF takes values in the R -local objects by the addendum to Proposition I.59. Using this and the fact that R is fully faithful, we can now compute

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}} \left(-, L \left(\lim_{\mathcal{I}} RF \right) \right) &\simeq \mathrm{Hom}_{\mathcal{C}} \left(R-, RL \left(\lim_{\mathcal{I}} RF \right) \right) \simeq \mathrm{Hom}_{\mathcal{C}} \left(R-, \lim_{\mathcal{I}} RF \right) \\ &\simeq \mathrm{Nat} \left(\mathrm{const}(R-), RF \right) \\ &\simeq \mathrm{Nat}(\mathrm{const}-, F), \end{aligned}$$

so $L(\lim_{\mathcal{I}} RF)$ satisfies the desired universal property for $\lim_{\mathcal{I}} F$. \square

Before we discuss our first examples of Bousfield localisations, we need to discuss a proposition which Fabian originally forgot to put here, but which came up later in the course.

I.61a. Proposition (see [HTT, Proposition 5.2.7.4]). — *Let $L : \mathcal{C} \rightarrow \mathcal{C}$ be a functor together with a natural transformation $\eta : \mathrm{id} \Rightarrow L$ such that both maps*

$$\eta_{Lx} : Lx \xrightarrow{\sim} LLx \quad \text{and} \quad L\eta_x : Lx \xrightarrow{\sim} LLx$$

are equivalences for all $x \in \mathcal{C}$. Then $L : \mathcal{C} \rightarrow \mathrm{im}(L)$ is left-adjoint to the inclusion $\mathrm{im}(L) \subseteq \mathcal{C}$ with unit η . In particular, L is a Bousfield localisation and $L\eta \simeq \eta L$ as natural transformations (by the triangle identities).

Proof. We will prove the assertion by reduction to the case of 1-categories, for which we already know this statement. We have to show that $\eta_c^* : \mathrm{Hom}_{\mathcal{C}}(Lc, Ld) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{C}}(c, Ld)$ is an equivalence for all $c, d \in \mathcal{C}$. For this it suffices to show that for all $K \in \mathrm{An}$, the induced map

$$\pi_0 \mathrm{Hom}_{\mathrm{An}}(K, \mathrm{Hom}_{\mathcal{C}}(Lc, Ld)) \xrightarrow{\sim} \pi_0 \mathrm{Hom}_{\mathrm{An}}(K, \mathrm{Hom}_{\mathcal{C}}(c, Ld))$$

is a bijection. But now consider the functor of 1-categories

$$\pi L_* : \pi \operatorname{Fun}(K, \mathcal{C}) \longrightarrow \pi \operatorname{Fun}(K, \mathcal{C})$$

together with the transformation $\pi \eta_* : \operatorname{id} \Rightarrow \pi L_*$ satisfying that $\pi \eta_* \pi L_* : \pi L_* \xrightarrow{\sim} \pi L_* \pi L_*$ and $\pi L_* \pi \eta_* : \pi L_* \xrightarrow{\sim} \pi L_* \pi L_*$ are natural equivalences. So we get the same situation as before, but now $\pi \operatorname{Fun}(K, \mathcal{C})$ is a 1-category. For 1-categories we already know the statement is true: For example, we've seen this on exercise sheet #2 of Higher Categories I, and also Lurie gives another proof. Plugging in the functors $\operatorname{const} c, \operatorname{const} Lc, \operatorname{const} Ld \in \operatorname{Fun}(K, \mathcal{C})$ then shows that $\pi_0 \operatorname{Hom}_{\operatorname{An}}(K, \operatorname{Hom}_{\mathcal{C}}(Lc, Ld)) \xrightarrow{\sim} \pi_0 \operatorname{Hom}_{\operatorname{An}}(K, \operatorname{Hom}_{\mathcal{C}}(c, Ld))$ is an equivalence, as desired. Thus we have proved that $L : \mathcal{C} \rightarrow \operatorname{im}(L)$ is a Bousfield localisation.

To obtain the additional assertion that $L\eta \simeq \eta L$, write down the triangle identities and note that both sides have the same left inverse, namely the counit evaluated at Lx . \square

I.62. Example. — “My first Bousfield localisations”:

(a) We now recognize all the left pointing arrows in the diagram from Example I.33(a)

$$\begin{array}{ccccc} & & \pi_0 & & | \quad | \\ & \swarrow & \text{---} & \nwarrow & \text{---} \\ \operatorname{Set} & \subseteq & \operatorname{An} & \subseteq & \operatorname{Cat}_{\infty} \\ & & \text{core} & & \end{array}$$

as Bousfield localisations. More precisely, π_0 and $| \quad |$ are left Bousfield localisations, whereas *core* is a right one.

(b) Recall from Example I.15(d) that Top has a Kan enrichment, giving rise to the ∞ -category $N^c(\operatorname{Top})$ of topological spaces. Consider the localisation

$$p : N^c(\operatorname{Top}) \longrightarrow N^c(\operatorname{Top})[\{\text{weak homotopy equiv.}^{-1}\}].$$

The functor $\operatorname{Hom}_{N^c(\operatorname{Top})}(X, -)$ inverts weak homotopy equivalences if X is a CW complex (that's a well-known extended version of Whitehead's theorem) and every topological space admits a weak equivalence from a CW complex (“CW approximation”). Hence the dual of Proposition I.59 implies that p is a right Bousfield localisation.

This also shows that $\operatorname{An} \simeq N^c(\operatorname{CW}) \rightarrow N^c(\operatorname{Top})[\{\text{weak homotopy equiv.}^{-1}\}]$ is an equivalence. In particular, we get for free that CW approximation is ∞ -functorial.

(c) Let R be a ring. Recall from Example I.15(e) that there is an ∞ -category $\mathcal{K}(R) = N^c(\operatorname{Ch}(R))$ of chain complexes over R . We define the *derived ∞ -category of R*

$$\mathcal{D}(R) = \mathcal{K}(R)[\{\text{quasi-isos.}\}^{-1}]$$

as the localisation of $\mathcal{K}(R)$ at the quasi-isomorphisms. Then the fundamental lemma of homological algebra² says that for a bounded below $C \in \mathcal{K}(R)$, any projective resolution

²Here's the complete argument: We need to show that $\operatorname{Hom}_{\mathcal{K}(R)}(P, -)$ inverts quasi-isomorphisms. So let $\varphi : C \rightarrow D$ be one. The fundamental lemma, or at least a strong form of it, says that $\varphi_* : \{\text{homotopy classes of maps } P \rightarrow C\} \xrightarrow{\sim} \{\text{homotopy classes of maps } P \rightarrow D\}$ is bijective. In other words, $\varphi_* : H_0 \underline{\operatorname{Hom}}_R(P, C) \xrightarrow{\sim} H_0 \underline{\operatorname{Hom}}_R(P, D)$ is an isomorphism, where $\underline{\operatorname{Hom}}_R$ is the internal Hom in $\operatorname{Ch}(R)$. Replacing P by its shifts $P[-i]$ shows the same for $\varphi_* : H_i \underline{\operatorname{Hom}}_R(P, C) \xrightarrow{\sim} H_i \underline{\operatorname{Hom}}_R(P, D)$. Now Exercise I.15a shows

$$\pi_i \operatorname{Hom}_{\mathcal{K}(R)}(-, -) \simeq H_i \underline{\operatorname{Hom}}_R(-, -)$$

and we are done.

$P \rightarrow C$ is a left-adjoint object to C . Similarly, for a bounded above C an injective resolution $C \rightarrow I$ is a right-adjoint object. In particular,

$$\mathcal{D}_{\geq 0}^{\text{proj}}(R) \subseteq \mathcal{K}(R) \longrightarrow \mathcal{D}(R)$$

is fully faithful with essential image the non-negative chain complexes (and similarly for $\mathcal{D}_{\leq 0}^{\text{inj}}(R)$). But in fact $\mathcal{K}(R) \rightarrow \mathcal{D}(R)$ admits both adjoints, so its both a left and a right Bousfield localisation! This boils down to the classical result of Spaltenstein that there are *enough* K -projective and K -injective complexes (again a K that has nothing to do with K -theory), see [Spa88] (thanks to Sil for the reference). In particular, combining Exercise I.50b and Corollary I.61 shows that $\mathcal{D}(R)$ is both complete and cocomplete.

Also note that the homotopy category $\pi\mathcal{D}(R)$ is equivalent to the usual derived 1-category $D(R)$ by [HCII, Proposition VIII.12].

- (d) Given a ring morphism $\varphi: R \rightarrow S$, we can look at the functor

$$S \otimes_R -: \text{Ch}(R) \longrightarrow \text{Ch}(S).$$

This is simplicially enriched (basically since the enrichment of $\text{Ch}(R)$ is defined via tensor products, and these are associative), hence it defines a functor $S \otimes_R -: \mathcal{K}(R) \rightarrow \mathcal{K}(S)$. Now define the *derived tensor product* as the composition

$$\varphi_! = S \otimes_R^L -: \mathcal{D}(R) \xrightarrow{K\text{-proj.}} \mathcal{K}(R) \xrightarrow{S \otimes_R -} \mathcal{K}(S) \longrightarrow \mathcal{D}(S).$$

The map $K\text{-proj.}: \mathcal{D}(R) \rightarrow \mathcal{K}(R)$ is a left adjoint of the right Bousfield localisation $p: \mathcal{K}(R) \rightarrow \mathcal{D}(R)$ and given by taking K -projective resolutions. We have seen in (c) that p also admits a right adjoint, but that wouldn't give the correct functor. Also note that $H_i(S \otimes_R^L M) = \text{Tor}_i^R(S, M)$ holds for all R -modules M , by unravelling of constructions.

It follows from general homological nonsense that $\varphi_!$ has a right-adjoint φ^* (the “forgetful functor”), which has another right adjoint $\varphi_* = R\text{Hom}_R(S, -)$. In diagrams:

$$\mathcal{D}(R) \xrightleftharpoons[\varphi_*]{\varphi_!} \mathcal{D}(S) \xrightleftharpoons[\varphi_*]{\varphi^*} \mathcal{D}(R).$$

I.63. Proposition/Definition. — *For a ring morphism $\varphi: R \rightarrow S$, the following conditions are equivalent:*

- (a) $\varphi_! = S \otimes_R^L -: \mathcal{D}(R) \rightarrow \mathcal{D}(S)$ is a left Bousfield localisation.
- (b) $\varphi^* = \text{forget}: \mathcal{D}(S) \rightarrow \mathcal{D}(R)$ is fully faithful.
- (c) $\varphi_* = R\text{Hom}_R(S, -): \mathcal{D}(R) \rightarrow \mathcal{D}(S)$ is a right Bousfield localisation.
- (d) The multiplication map $S \otimes_R^L S \rightarrow S$ is an equivalence in $\mathcal{D}(S)$. As Sil points out, we should actually write $S \otimes_R^L \varphi^* S$, but let's not do that here.
- (e) The cofibre $\text{cofib}(\varphi) = S/\text{}^L R$, i.e. the pushout

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & S \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & S/\text{}^L R \end{array}$$

in $\mathcal{D}(R)$, satisfies $S \otimes_R^L (S/\text{}^L R) \simeq 0$.

If these conditions are satisfied, φ is called a *derived localisation* (or sometimes “Tor-unital”).

Examples. — “My first derived localisations”:

- (a) If $W \subseteq R$ is a multiplicative subset in the commutative ring R , then $R \rightarrow R[W^{-1}]$ is a derived localisation. This easily follows from Proposition/Definition I.63(d) since $R[W^{-1}]$ is flat over R .
- (b) For non-commutative rings, one needs the (left or right, who knows?) *Ore condition* for $R \rightarrow R[W^{-1}]$ to be a derived localisation.
- (c) If $U \subseteq V$ is an open embedding of affine schemes, then $\Gamma(V, \mathcal{O}_V) \rightarrow \Gamma(U, \mathcal{O}_U)$ is a derived localisation (but not necessarily an ordinary localisation). Indeed, the condition from Proposition/Definition I.63(d) easily implies that a ring morphism $\varphi: R \rightarrow S$ is a derived localisation iff $R_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$ is a derived localisation for all prime ideals $\mathfrak{q} \in \text{Spec } S$ and $\mathfrak{p} \in \text{Spec } R$ such that $\mathfrak{p} = \varphi^{-1}(\mathfrak{q})$. In our case, $R_{\mathfrak{p}}$ and $S_{\mathfrak{q}}$ correspond to the local rings $\mathcal{O}_{V,v}$ and $\mathcal{O}_{U,v}$ for $v \in V$, and these are isomorphic since U is an open subscheme of V . Hence the morphisms $\mathcal{O}_{V,v} \xrightarrow{\sim} \mathcal{O}_{U,v}$ for $v \in V$ are derived localisations for obvious reasons.

We will later see that K -theory has exact sequences for derived localisations!

Proof of Proposition/Definition I.63. We already know $(a) \Leftrightarrow (b) \Leftrightarrow (c)$ from Proposition/Definition I.58 and its dual. For $(d) \Leftrightarrow (e)$, first note that $S \otimes_R^L R \xrightarrow{\sim} S$ is an equivalence. Indeed, R can be taken as its own projective resolution, hence $S \otimes_R^L R \simeq S \otimes_R R \simeq S$.

Now $S \otimes_R^L -$, being a left adjoint, commutes with colimits, which implies that

$$\begin{array}{ccc} S & \xrightarrow{S \otimes_R^L \varphi} & S \otimes_R^L S \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & S \otimes_R^L (S / {}^L R) \end{array}$$

is a pushout square. This immediately implies $S \otimes_R^L S \simeq S$ iff $S \otimes_R^L (S / {}^L R) \simeq 0$, as desired.

We finish the proof by showing $(b) \Leftrightarrow (d)$. We must show that the counit $S \otimes_R^L C \rightarrow C$ is an equivalence for all $C \in \mathcal{D}(S)$ iff it is an equivalence for S . The “only if” direction being obvious, let’s assume it is an equivalence for S . Then it is also an equivalence for all shifts $S[i]$. But the $\{S[i]\}_{i \in \mathbb{Z}}$ generate $\mathcal{D}(S)$ under colimits and $S \otimes_R^L -$ commutes with colimits, so we are done. \square

The Rezk Nerve and Complete Segal Spaces

LECTURE 7
19th Nov, 2020

From now on, a good portion of the lecture will take place in the land of simplicial anima. Unfortunately, this leads to “ Δ^n ” having two different meanings: It could mean the ∞ -category $\Delta^n = \mathcal{N}([n])$, or the simplicial set Δ^n , which we consider as a discrete simplicial anima via $\mathbf{sSet} \subseteq \mathbf{sAn}$. To avoid confusion, we adopt the following convention: The ∞ -category will always be denoted $[n]$, whereas the simplicial anima will be denoted Δ^n .

Under the Rezk nerve functor $N^r: \mathbf{Cat}_{\infty} \rightarrow \mathbf{sAn}$ from Example I.57(b), these two are related via $N^r([n]) \simeq \Delta^n$. Indeed, we have

$$N_m^r([n]) \simeq \text{Hom}_{\mathbf{Cat}_{\infty}}([m], [n]) \simeq \text{core Fun}([m], [n]),$$

and the right-hand side is the discrete set $(\Delta^n)_m$ because every isomorphism of functors in $\text{Fun}([m], [n])$ must be an identity (as $[n]$ itself has no non-identity isomorphisms).

I.64. Theorem/Definition (Rezk, Joyal–Tierney, Lurie). — *The Rezk nerve functor $N^r: \text{Cat}_\infty \rightarrow \text{sAn}$ is fully faithful and its essential image consists of the complete Segal spaces/anima. Here a Segal anima is a simplicial anima $X: \Delta^{\text{op}} \rightarrow \text{An}$ such that*

$$X_n \simeq \text{Hom}_{\text{sAn}}(\Delta^n, X) \xrightarrow{\sim} \text{Hom}_{\text{sAn}}(I_n, X) \simeq X \times_{X_0} X_1 \times_{X_0} \cdots \times_{X_0} X_1$$

is an equivalence for all n (here $I_n \subseteq \Delta^n$ denotes the n^{th} spine, i.e. the union of all edges between consecutive 0-simplices of Δ^n). We call X complete if the following equivalent conditions hold:

(a) *Also the restriction $X_0 \simeq \text{Hom}_{\text{sAn}}(\Delta^0, X) \xrightarrow{\sim} \text{Hom}_{\text{sAn}}(J, X)$ along $J \rightarrow \Delta^0$ is an equivalence, where J is the (ordinary nerve of the) free-living isomorphism, considered as a simplicial set/discrete simplicial anima.*

(b) *The diagram*

$$\begin{array}{ccc} X_0 & \xrightarrow{\Delta} & X_0 \times X_0 \\ s \downarrow & \lrcorner & \downarrow (s,s) \\ X_3 & \xrightarrow{(d_{\{0,2\}}, d_{\{1,3\}})} & X_1 \times X_1 \end{array}$$

is cartesian (note that Δ denotes a good old honest diagonal and not some simplicial stuff).

(c) *The degeneracy map $s: X_0 \xrightarrow{\sim} X_1^\times$ is an equivalence, where $X_1^\times \subseteq X_1$ is the collection of path components of those $g \in X_1$ for which the following holds: Let $x = d_1(g)$ and $y = d_0(g)$ and for arbitrary $w, z \in X_0$ put*

$$\begin{array}{ccc} P_{w,z} & \longrightarrow & X_1 \\ \downarrow & \lrcorner & \downarrow (d_1, d_0) \\ * & \xrightarrow{(w,z)} & X_0 \times X_0 \end{array}$$

Moreover, there is a “composition map”

$$\circ: X_1 \times_{d_1, X_0, d_0} X_1 \xleftarrow[\sim]{(d_0, d_2)} X_2 \xrightarrow{d_1} X_1$$

which, after restriction to $P_{w,x} \simeq P_{w,x} \times_{\{x\}} \{g\}$ and $P_{y,z} \simeq \{g\} \times_{\{y\}} P_{y,z}$, induces “post- and precomposition maps”

$$g_*: P_{w,x} \longrightarrow P_{w,y} \quad \text{and} \quad g^*: P_{y,z} \longrightarrow P_{x,z}.$$

Then we put $g \in X_1^\times$ iff g_ and g^* are equivalences for all $w, z \in X_0$.*

(d) *The simplicial subanima X^\times given by*


$$[n] \longmapsto X_n^\times := \left\{ \begin{array}{l} \text{collection of path components in} \\ X_n \text{ such that all edges lie in } X_1^\times \end{array} \right\}$$

is constant. Note that $X_0^\times = X_0$, so it is actually constant on X_0 .

I.65. Some Explanations. — Before we (not so much) talk about the proof, some explanations are in order.

First of, the Segal condition. Just like Segal sets, *Segal anima* should be thought of simplicial anima with “unique spine lifting”, i.e. unique (up to contractible choice) lifting against $I_n \subseteq \Delta^n$. To see why $\mathrm{Hom}_{\mathrm{sAn}}(I^n, X) \simeq X_1 \times_{X_0} \cdots \times_{X_0} X_1$, we must show that I^n is an iterated pushout of copies of Δ^1 and then invoke Corollary I.50. The pushout condition is certainly true in sSet . In particular, the discrete set $(I^n)_m$ is an iterated pushout of copies of $(\Delta^1)_m$, and the maps along the pushouts are taken are injective. Now pushouts in sAn are computed degreewise by Lemma I.39, the simplicial anima I^n , Δ^1 , and Δ^0 are discrete sets in every degree, and $\mathrm{Set} \subseteq \mathrm{An}$ preserves pushouts along injective maps, so in this particular special case, the pushout in sSet agrees with the one in sAn . We’ll also link the Segal condition to monoidal structures soon, see Chapter II.

Now about completeness. As Bastiaan explained to me, the completeness condition is probably best understood in the **words of Emily Riehl**: “[The completeness condition] says that isomorphisms are equivalent to identities.” Although these words were spoken in a slightly different context, they fit here just wonderfully. For conditions (a) and (c) it is intuitive that “isomorphisms are equivalent to identities” is what they’re saying, and condition (d) is basically just the statement that all higher simplices having equivalences as edges are degenerate. To make it more intuitive why also condition (b) is saying the same thing, Fabian drew the following picture of a 3-simplex in X :



What (b) is trying to say is that whenever the $\Delta^{\{0,2\}}$ - and $\Delta^{\{1,3\}}$ -edges are degenerate as indicated, then all edges are degenerate and the 3-simplex itself is equivalent to a degenerate one. Notice that this is just the higher categorical way of phrasing the requirement that identity morphisms are closed under the 2-out-of-6-property. This is also equivalent to all equivalences being identities since the equivalences in an ∞ -category are the smallest collection of arrows containing the identities and closed under the 2-out-of-6-property.

Not a proof of Theorem/Definition I.64. It’s relatively easy to see that $N^r(\mathcal{C})$ for an ∞ -category \mathcal{C} is a complete Segal anima: The Segal condition reduces to the claim that $[n] \simeq [1] \sqcup_{[0]} \cdots \sqcup_{[0]} [1]$ in Cat_∞ , which can be shown as in Example I.38. For completeness, one unwinds that $N_1^r(\mathcal{C})^\times \subseteq \mathrm{core} \, \mathrm{Ar}(\mathcal{C})$ consists of the equivalences, hence $s: N_0^r(\mathcal{C}) \xrightarrow{\sim} N_1^r(\mathcal{C})^\times$ being an equivalence reduces to $[1] \rightarrow [0]$ being a localisation. Fabian also recommends to prove the equivalence of the four conditions as a (not easy, but doable) exercise.

However, the proof that N^r is fully faithful is difficult. The original proof of Joyal–Tierney can be found in [JT07] and Lurie’s poof is in [Lur09]. The key step is to give a description of $\mathrm{asscat}(X)$ for X a (not necessarily complete) Segal anima: First, there is a canonical equivalence

$$\mathrm{core} \, \mathrm{asscat}(X) \simeq |X^\times|, \quad (\text{I.65.1})$$

where $| \cdot | \simeq \mathrm{colim}_{\Delta^{\mathrm{op}}} : \mathrm{sAn} \rightarrow \mathrm{An}$ denotes the colimit functor over Δ^{op} (Fabian points out that this notation is very much consistent with the other kinds of realisation we’ve seen so

far; see Remark I.66 below). Second, one can show that for every (not necessarily complete) Segal anima X there are pullback diagrams

$$\begin{array}{ccc} \mathrm{Hom}_{\mathrm{asscat}(X)}(x, y) & \longrightarrow & X_1 \\ \downarrow & \lrcorner & \downarrow (d_1, d_0) \\ * & \xrightarrow{(x, y)} & X_0 \times X_0 \end{array} \quad (\text{I.65.2})$$

for all $x, y \in X_0$. In particular $\mathrm{Hom}_{\mathrm{asscat}(X)}(x, y) \simeq P_{x, y}$ in the notation of (b).

With (I.65.1) and (I.65.2), the proof of Theorem/Definition I.64 becomes quite easy. To prove that N^r is fully faithful, we must show that the counit $\mathrm{asscat}(N^r(\mathcal{C})) \xrightarrow{\sim} \mathcal{C}$ is an equivalence for all $\mathcal{C} \in \mathrm{Cat}_\infty$. It is an equivalence on cores because

$$\mathrm{core} \mathrm{asscat}(N^r(\mathcal{C})) \simeq |N^r(\mathcal{C})^\times| \quad \text{and} \quad \mathrm{core} \mathcal{C} \simeq N_0^r(\mathcal{C});$$

since $N^r(\mathcal{C})$ is complete, $N^r(\mathcal{C})^\times$ is constant on $N_0^r(\mathcal{C})$, hence the colimit over the weakly contractible category Δ^{op} is just $|N^r(\mathcal{C})^\times| \simeq N_0^r(\mathcal{C})$ by Proposition I.36, as claimed. This shows that the counit is essentially surjective.

To prove that it is fully faithful, i.e. induces equivalences on Hom anima, observe that using (I.65.2) it induces the following equivalence of pullback squares:

$$\begin{array}{ccccc} \mathrm{Hom}_{\mathrm{asscat}(N^r(\mathcal{C}))}(x, y) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(x, y) & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ * & \xrightarrow{(x, y)} & N_1^r(\mathcal{C}) & \xrightarrow{\sim} & \mathrm{core} \mathrm{Ar}(\mathcal{C}) \\ & \lrcorner & \downarrow & & \downarrow \\ & & * & \xrightarrow{(x, y)} & \mathrm{core} \mathcal{C} \\ & & \downarrow & & \\ & & N_0^r(\mathcal{C}) \times N_0^r(\mathcal{C}) & \xrightarrow{\sim} & \mathrm{core} \mathcal{C} \times \mathrm{core} \mathcal{C} \end{array}$$

For the right pullback square, recall the usual pullback diagram for $\mathrm{Hom}_{\mathcal{C}}(x, y)$ can also be taken in Cat_∞ (Example* I.35a(a)) and $\mathrm{core}(-)$ preserves pullbacks because it is a right adjoint. Up to verifying the actual hard stuff ((I.65.1) and (I.65.2)), this finishes the proof that N^r is fully faithful. To show that its image are the complete Segal anima, one can argue similarly for the unit $X \rightarrow N^r(\mathrm{asscat}(X))$. \square

I.66. Remark. — Note that the notation $| \cdot | \simeq \mathrm{colim}_{\Delta^{\mathrm{op}}} \cdot$ is consistent with our notation for the geometric realisation of simplicial sets: If $T: \Delta^{\mathrm{op}} \rightarrow \mathrm{Set} \subseteq \mathrm{An}$ is a simplicial set/discrete simplicial anima, then the homotopy colimit of T (considered as a functor $T: \Delta^{\mathrm{op}} \rightarrow \mathrm{Set} \subseteq \mathrm{sSet}$, where sSet is equipped with the Kan–Quillen model structure) is weakly homotopy equivalent to T , so

$$\mathrm{colim}_{\Delta^{\mathrm{op}}} T \simeq \left(\mathrm{hocolim}_{\mathfrak{C}[\Delta^{\mathrm{op}}]} T \right)^f \simeq T^f \simeq \mathrm{Sing} |T|$$

using Theorem I.34. The notation is also consistent with $| \cdot |: \mathrm{Cat}_\infty \rightarrow \mathrm{An}$ in that generally for a simplicial anima $T: \Delta^{\mathrm{op}} \rightarrow \mathrm{An}$ we have

$$| \mathrm{asscat}(T) | \simeq |T|.$$

Indeed, this follows from Theorem I.51 as both sides are colimit-preserving functors that send $\Delta^n \mapsto * \in \mathrm{An}$ for all $n \in \mathbb{N}$.

I.67. Lemma/Definition. — Let $\mathbf{CSAn} \subseteq \mathbf{sAn}$ denote the full sub- ∞ -category of complete Segal anima. The functor

$$\mathrm{comp}: \mathbf{sAn} \xrightarrow{\mathrm{asscat}} \mathbf{Cat}_\infty \xrightarrow{N^r} \mathbf{CSAn}$$

is called completion and it is a left adjoint to the inclusion $\mathbf{CSAn} \subseteq \mathbf{sAn}$.

*Proof**. Indeed, if $X \in \mathbf{sAn}$ and $Y \in \mathbf{CSAn}$, so $Y \simeq N^r(\mathcal{C})$ for some ∞ -category \mathcal{C} , then $\mathrm{Hom}_{\mathbf{sAn}}(\mathrm{comp} X, Y) \simeq \mathrm{Hom}_{\mathbf{Cat}_\infty}(\mathrm{asscat} X, \mathcal{C}) \simeq \mathrm{Hom}_{\mathbf{sAn}}(X, N^r(\mathcal{C}))$ since N^r is fully faithful and right-adjoint to $\mathrm{asscat}: \mathbf{sAn} \rightarrow \mathbf{Cat}_\infty$. \square

Fabian remarks that $\mathrm{comp}: \mathbf{sAn} \rightarrow \mathbf{CSAn}$ is impossible to control outside of (not necessarily complete) Segal spaces.

I.68. Example. — (a) The ordinary nerve $N(\mathcal{C}) \in \mathbf{sSet} \subseteq \mathbf{sAn}$ of a 1-category is a Segal set and thus also a Segal anima since $\mathbf{Set} \subseteq \mathbf{An}$ preserves limits. But it is usually not complete! In fact, we have

$$\begin{array}{ccc} N_0(\mathcal{C}) & \xlongequal{\quad} & \{\text{set of objects in } \mathcal{C}\} \\ s \downarrow & & \downarrow \\ N_1(\mathcal{C}) & \xlongequal{\quad} & \{\text{set of isomorphisms in } \mathcal{C}\} \end{array}$$

where the right vertical arrow takes $x \in \mathcal{C}$ to id_x . But usually, there are more isomorphisms in a category than just the identities, hence $N_1(\mathcal{C})^\times$ is usually larger than $N(\mathcal{C})_0$. This is also another instance of Emily Riehl’s wise words from I.65: “The completeness condition says that isomorphisms are equivalent to identities.”

Note however that the composition $\mathrm{asscat}(N(\mathcal{C})) \rightarrow \mathrm{asscat}(N^r(\mathcal{C})) \rightarrow \mathcal{C}$ is an equivalence, so $N^r(\mathcal{C})$ is the completion of $N(\mathcal{C})$ in the sense of Lemma/Definition I.67. Indeed, it’s clear from (I.65.2) that $\mathrm{asscat}(N(\mathcal{C})) \rightarrow \mathcal{C}$ induces equivalences on Hom anima, so it remains to see whether it is essentially surjective. By (I.65.1), it suffices to show $|N(\mathcal{C})^\times| \simeq \mathrm{core} \mathcal{C}$. But $N_n(\mathcal{C})^\times$ is given by the set of composable sequences (f_1, \dots, f_n) of isomorphisms in \mathcal{C} , hence it is the ordinary $N(\mathrm{core} \mathcal{C})$. Since this is already a Kan complex, Remark I.66 shows $|N(\mathrm{core} \mathcal{C})| \simeq \mathrm{core}(\mathcal{C})$, as required.

(b) The two functors

$$\begin{array}{ccc} \mathbf{sAn} & \xrightarrow{\mathrm{asscat}} \mathbf{Cat}_\infty & \xrightarrow{\pi} \mathbf{Cat}_1^{(2)} \\ \mathbf{sAn} & \xrightarrow{s\pi_0} \mathbf{sSet} & \xrightarrow{h} \mathbf{Cat}_1 \end{array}$$

are different, even though both are colimit-preserving and take $\Delta^n \mapsto [n]$. The reason is of course that $\mathbf{Cat}_1 \subseteq \mathbf{Cat}_1^{(2)}$ doesn’t preserve colimits. To obtain explicit counterexamples, you can take $\mathrm{const} \mathbb{S}^1$ or $N^r(\{\text{finite-dimensional vector spaces}\})$ for example. Both functors do, however, agree on ordinary nerves $N(\mathcal{C})$ of 1-categories. Nevertheless, Fabian warns you to never look at the lower functor.

(c) Let $\mathbf{sAn}_{\mathrm{const}} \subseteq \mathbf{sAn}$ denote the full subcategory of constant simplicial anima. The adjunction $\mathrm{asscat}: \mathbf{sAn} \rightleftarrows \mathbf{Cat}_\infty : N^r$ restricts to an equivalence

$$\mathrm{ev}_0: \mathbf{sAn}_{\mathrm{const}} \xrightleftharpoons{\sim} \mathbf{An} : \mathrm{const} .$$

Indeed, $N_n^r(K) \simeq \mathrm{core} \mathrm{Fun}([n], K) \simeq \mathrm{Fun}([n], K) \simeq K$ as $[n] \simeq *$.

I.69. Lemma. — *For any ∞ -category \mathcal{C} we can describe $N^r(\mathrm{TwAr}(\mathcal{C}))$ as follows:*

$$N_n^r(\mathrm{TwAr}(\mathcal{C})) \simeq \mathrm{Hom}_{\mathrm{Cat}_\infty}([n]^{\mathrm{op}} \star [n], \mathcal{C}) \simeq \mathrm{Hom}_{\mathrm{sAn}}((\Delta^n)^{\mathrm{op}} \star \Delta^n, N^r(\mathcal{C})).$$

Here $[n]^{\mathrm{op}} \star [n]$ denotes the usual join of categories, and $(\Delta^n)^{\mathrm{op}} \star \Delta^n$ is formed as a join in sSet and then imported into sAn .

Proof. Recall from I.28 that there is a pullback square

$$\begin{array}{ccc} \mathrm{TwAr}(\mathcal{C}) & \longrightarrow & */\mathrm{An} \\ \downarrow & \lrcorner & \downarrow \\ \mathcal{C}^{\mathrm{op}} \times \mathcal{C} & \xrightarrow{\mathrm{Hom}_{\mathcal{C}}} & \mathrm{An} \end{array}$$

(depending on your definitions, this is a pullback both in simplicial sets and in Cat_∞ or only in Cat_∞). Now we do a computation in several steps, each of which will be justified below:

$$\begin{aligned} N_n^r(\mathrm{TwAr}(\mathcal{C})) &\simeq \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathrm{TwAr}(\mathcal{C})) \\ &\stackrel{(1)}{\simeq} \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \times_{\mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathrm{An})} \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], */\mathrm{An}) \\ &\stackrel{(2)}{\simeq} \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \times_{\mathrm{core}(\mathrm{An})} \mathrm{core}(*/\mathrm{An}) \\ &\stackrel{(3)}{\simeq} \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \times_{\mathrm{core}(\mathcal{C}^{\mathrm{op}} \times \mathcal{C})} \mathrm{core}((\mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \times_{\mathrm{An}} */\mathrm{An}) \\ &\stackrel{(4)}{\simeq} (\mathrm{Hom}_{\mathrm{Cat}_\infty}([n]^{\mathrm{op}}, \mathcal{C}) \times \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathcal{C})) \times_{\mathrm{core}(\mathcal{C}^{\mathrm{op}} \times \mathcal{C})} \mathrm{core} \mathrm{TwAr}(\mathcal{C}) \\ &\stackrel{(5)}{\simeq} (\mathrm{Hom}_{\mathrm{Cat}_\infty}([n]^{\mathrm{op}}, \mathcal{C}) \times \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathcal{C})) \times_{\mathrm{core}(\mathcal{C} \times \mathcal{C})} \mathrm{core} \mathrm{Ar}(\mathcal{C}) \\ &\stackrel{(6)}{\simeq} \mathrm{core} \mathrm{Fun}([n]^{\mathrm{op}} \sqcup_{\{0\}} [1] \sqcup_{\{0\}} [n], \mathcal{C}) \\ &\stackrel{(7)}{\simeq} \mathrm{Hom}_{\mathrm{Cat}_\infty}([n]^{\mathrm{op}} \star [n], \mathcal{C}) \\ &\stackrel{(8)}{\simeq} \mathrm{Hom}_{\mathrm{sAn}}((\Delta^n)^{\mathrm{op}} \star \Delta^n, N^r(\mathcal{C})). \end{aligned}$$

For step (1), we plug in $\mathrm{TwAr}(\mathcal{C}) \simeq (\mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \times_{\mathrm{An}} */\mathrm{An}$ and use that $\mathrm{Hom}_{\mathrm{Cat}_\infty}([n], -)$ preserves pullbacks by the dual of Corollary I.50.

For step (2), we observe that restriction along $\{0\} \hookrightarrow [n]$, i.e. “evaluation at 0”, induces morphisms $\mathrm{ev}_0: \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], */\mathrm{An}) \rightarrow \mathrm{Hom}_{\mathrm{Cat}_\infty}(\{0\}, */\mathrm{An}) \simeq \mathrm{core}(*/\mathrm{An})$ and $\mathrm{ev}_0: \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathrm{An}) \rightarrow \mathrm{Hom}_{\mathrm{Cat}_\infty}(\{0\}, \mathrm{An}) \simeq \mathrm{core}(\mathrm{An})$. We claim that these fit into a pullback diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], */\mathrm{An}) & \xrightarrow{\mathrm{ev}_0} & \mathrm{core}(*/\mathrm{An}) \\ \downarrow & \lrcorner & \downarrow \\ \mathrm{Hom}_{\mathrm{Cat}_\infty}([n], \mathrm{An}) & \xrightarrow{\mathrm{ev}_0} & \mathrm{core}(\mathrm{An}) \end{array}$$

in An . To see this, note that $\mathrm{Fun}([n], */\mathrm{An}) \simeq \mathrm{Fun}_*([n+1], \mathrm{An})$ holds by the join-slice adjunction (or by inspection), where Fun_* denotes the full sub- ∞ -categories of functors taking $0 \mapsto * \in \mathrm{An}$. Similarly $*/\mathrm{An} \simeq \mathrm{Fun}_*([1], \mathrm{An})$. Now $[n+1]$ can be written as a pushout $[1] \sqcup_{\{1\}} [1, n+1]$ in Cat_∞ (where $[1, n+1]$ denotes the poset $\{1, \dots, n+1\}$). Indeed, by

Theorem 1.34 we may compute this pushout as a fibrant replacement of the corresponding homotopy pushout in \mathbf{sSet} . Since $\{1\} \hookrightarrow \Delta^1$ is a cofibration, the homotopy pushout is just the ordinary pushout, and $\Delta^1 \sqcup_{\{1\}} \Delta^{\{1, \dots, n+1\}} \hookrightarrow \Delta^{n+1}$ is inner anodyne, so $[n+1]$ is indeed the correct pushout in \mathbf{Cat}_∞ . Since $\mathrm{Hom}_{\mathbf{Cat}_\infty}(-, \mathbf{An}) \simeq \mathrm{core}\mathrm{Fun}(-, \mathbf{An})$ takes pushouts to pullbacks by Corollary 1.50, we get

$$\mathrm{core}\mathrm{Fun}([n+1], \mathbf{An}) \simeq \mathrm{core}\mathrm{Fun}([1], \mathbf{An}) \times_{\mathrm{core}\mathrm{Fun}(\{1\}, \mathbf{An})} \mathrm{core}\mathrm{Fun}([1, n+1], \mathbf{An}),$$

and this restricts to an equivalence

$$\mathrm{core}\mathrm{Fun}_*([n+1], \mathbf{An}) \simeq \mathrm{core}\mathrm{Fun}_*([1], \mathbf{An}) \times_{\mathrm{core}\mathrm{Fun}(\{1\}, \mathbf{An})} \mathrm{core}\mathrm{Fun}([1, n+1], \mathbf{An}).$$

This shows that the pullback diagram above is correct. Plugging it into (1) gives (2).

Step (3) is a formal pullback manipulation together with the fact that $\mathrm{core}(-)$ commutes with pullbacks since it is a right adjoint.

For (4), note that $\mathrm{Hom}_{\mathbf{Cat}_\infty}([n], \mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \simeq \mathrm{Hom}_{\mathbf{Cat}_\infty}([n], \mathcal{C}^{\mathrm{op}}) \times \mathrm{Hom}_{\mathbf{Cat}_\infty}([n], \mathcal{C})$ and we have $\mathrm{Fun}([n], \mathcal{C}^{\mathrm{op}}) \simeq \mathrm{core}\mathrm{Fun}([n]^{\mathrm{op}}, \mathcal{C})^{\mathrm{op}}$. But the $(-)^{\mathrm{op}}$ vanishes upon taking cores, hence

$$\mathrm{Hom}_{\mathbf{Cat}_\infty}([n], \mathcal{C}^{\mathrm{op}}) \simeq \mathrm{Hom}_{\mathbf{Cat}_\infty}([n]^{\mathrm{op}}, \mathcal{C}).$$

This explains the left factor in the pullback. For the right factor, we simply plug in $\mathrm{TwAr}(\mathcal{C}) \simeq (\mathcal{C}^{\mathrm{op}} \times \mathcal{C}) \times_{\mathbf{An}} \mathbf{An}$.


Step (5) follows straight from Exercise 1.28a. Step (6) is immediate from Corollary 1.50, and for (7) we need to check that $[n]^{\mathrm{op}} \sqcup_{\{0\}} [1] \sqcup_{\{0\}} [n]$, when taken as a pushout in \mathbf{Cat}_∞ , is given by $[n]^{\mathrm{op}} \star [n]$. This can be done as before: Take the pushout in \mathbf{sSet} and verify that the canonical map to $[n]^{\mathrm{op}} \star [n]$ is inner anodyne.

Finally, (8) is immediate once we convinced ourselves that

$$N^r([n]^{\mathrm{op}} \star [n]) \simeq (\Delta^n)^{\mathrm{op}} \star \Delta^n,$$

since N^r is fully faithful by Theorem/Definition 1.64. To obtain the equivalence above, note that $\mathrm{Hom}_{\mathbf{Cat}_\infty}([m], [n]^{\mathrm{op}} \star [n]) \simeq \mathrm{Hom}_{\mathbf{Cat}_1}([m], [n]^{\mathrm{op}} \star [n])$, even though $\mathbf{Cat}_1 \subseteq \mathbf{Cat}_\infty$ is not fully faithful in general. Here it works, since every isomorphism of functors in $\mathrm{Fun}([m], [n]^{\mathrm{op}} \star [n])$ must be an identity (as $[n]^{\mathrm{op}} \star [n]$ has no non-identity isomorphisms). Thus $N_m^r([n]^{\mathrm{op}} \star [n])$ is given by the discrete set $\mathrm{Hom}_{\mathbf{sSet}}(\Delta^m, (\Delta^n)^{\mathrm{op}} \star \Delta^n)$ since $N: \mathbf{Cat}_1 \rightarrow \mathbf{sSet}$ is fully faithful and compatible with $- \star -$ and $(-)^{\mathrm{op}}$. This shows that $N^r([n]^{\mathrm{op}} \star [n])$ indeed agrees with $(\Delta^n)^{\mathrm{op}} \star \Delta^n$. \square

1.70. The Quillen Q -Construction. — Let \mathcal{C} be an ∞ -category and let $Q_n(\mathcal{C}) \subseteq \mathrm{Fun}(\mathrm{TwAr}([n])^{\mathrm{op}}, \mathcal{C})$ be the full sub- ∞ -category consisting of those functors that take all “squares” in $\mathrm{TwAr}([n])^{\mathrm{op}}$ to pullbacks. To make sense of this, recall our description of $\mathrm{TwAr}([n])$ from 1.28. In that way, a functor $\mathrm{TwAr}([n])^{\mathrm{op}} \rightarrow \mathcal{C}$ can be pictured as a diagram



in \mathcal{C} (shown here for $n = 3$), and to be in $Q_n(\mathcal{C})$ we require that all squares are pullbacks as indicated. Note that then in fact all rectangles you can find in this picture must be pullbacks.

The $Q_n(\mathcal{C})$ assemble into a simplicial ∞ -category $Q(\mathcal{C}): \Delta^{\text{op}} \rightarrow \text{Cat}_{\infty}$. To see this, one first shows that $\text{Fun}(\text{TwAr}([-])^{\text{op}}, \mathcal{C}): \Delta^{\text{op}} \rightarrow \text{Cat}_{\infty}$ gives a functor. It's clear that all its ingredients are functorial, except perhaps for $\text{TwAr}(-)$. But $\text{TwAr}(-): \text{sSet} \rightarrow \text{sSet}$ is right-adjoint to $(-)^{\text{op}} \star (-): \text{sSet} \rightarrow \text{sSet}$ (indeed, this guy has a right adjoint by Theorem I.2, now unravel that $\text{TwAr}(-)$ as defined in I.26(b) fits the general description of such right adjoints), which one can show is a left Quillen functor for the Joyal model structure (by [HCII, Proposition D.5] for example).

Now that we know $\text{Fun}(\text{TwAr}([-])^{\text{op}}, \mathcal{C}): \Delta^{\text{op}} \rightarrow \text{Cat}_{\infty}$ is a functor, we only need to check that all the boundary and degeneracy maps preserve the full subcategories $Q_n(\mathcal{C})$ described above. We leave this as an exercise. As you might have guessed, assigning to \mathcal{C} the simplicial ∞ -category $Q(\mathcal{C})$ is called the *Quillen Q -construction*.

LECTURE 8
24th Nov, 2020


I.71. Proposition/Definition. — *Let \mathcal{C} be an ∞ -category with pullbacks. The simplicial anima*

$$\text{core } Q(\mathcal{C}): \Delta^{\text{op}} \longrightarrow \text{An}$$

is a complete Segal space. Its associated category $\text{Span}(\mathcal{C}) \simeq \text{asscat}(\text{core } Q(\mathcal{C}))$ is called the ∞ -category of spans in \mathcal{C} .

Proof. We will actually prove that $Q(\mathcal{C})$ itself satisfies the Segal and completeness conditions for simplicial ∞ -categories rather than anima (really what this says is that there's an ∞ -double category $\text{Span}^{(2)}(\mathcal{C})$, but never mind that). Then all assertions for $\text{core } Q(\mathcal{C})$ will follow from the fact that $\text{core}: \text{Cat}_{\infty} \rightarrow \text{An}$ preserves limits as it is a right-adjoint.

Recall that the 0-simplices in $\text{TwAr}([n])^{\text{op}}$ are given by morphisms in the poset $[n]$, i.e. by relations $(i \leq j)$. Now let $J_n \subseteq \text{TwAr}([n])^{\text{op}}$ be the subposet spanned by all 0-simplices $(i \leq j)$ with $j \leq i + 1$. In pictures, J_n (for $n = 3$) looks as follows:



Then one checks from the pointwise formula for Kan extensions (Theorem I.52) that a functor $F: \text{TwAr}([n])^{\text{op}} \rightarrow \mathcal{C}$ is in $Q_n(\mathcal{C})$ iff it is right-Kan extended from $F|_{J_n}$. To see this in the example above, note that Kan extensions can be computed in steps (because right adjoints compose), so we may first Kan extend to the 0-simplex $(0 \leq 2)$, then to $(1 \leq 3)$ and then finally to $(0 \leq 3)$. In each step, the Kan extension is given by pullback, as one can see by restricting the limit from Theorem I.52 to a suitable cofinal sub- ∞ -category of the respective slice categories. The case for general n works precisely the same.

In particular, since $J_n \subseteq \text{TwAr}([n])^{\text{op}}$ is fully faithful, we see that restriction to J_n induces an equivalence

$$Q_n(\mathcal{C}) \xrightarrow{\sim} \text{Fun}(J_n, \mathcal{C})$$

by Corollary I.54. The reason we don't work with the J_n right away is that they do not form a cosimplicial ∞ -category! So we have to include some redundant junk. However, the J_n are closed under the maps induced by the Segal maps $e_i: [1] \rightarrow [n]$ (sending $[1]$ to $\{i, i+1\} \subseteq [n]$). These maps induce an equivalence $J_n \simeq J_1 \sqcup_{J_0} \cdots \sqcup_{J_0} J_1$ in Cat_∞ , hence

$$\begin{aligned} Q_n(\mathcal{C}) &\simeq \text{Fun}(J_1 \sqcup_{J_0} \cdots \sqcup_{J_0} J_1, \mathcal{C}) \simeq \text{Fun}(J_1, \mathcal{C}) \times_{\text{Fun}(J_0, \mathcal{C})} \cdots \times_{\text{Fun}(J_0, \mathcal{C})} \text{Fun}(J_1, \mathcal{C}) \\ &\simeq Q_1(\mathcal{C}) \times_{Q_0(\mathcal{C})} \cdots \times_{Q_0(\mathcal{C})} Q_1(\mathcal{C}) \end{aligned}$$

by Corollary I.50, which proves that $Q(\mathcal{C})$ is indeed a Segal object in Cat_∞ .

For completeness, we have to check the condition from Theorem/Definition I.64(b), i.e. we need to show that

$$\begin{array}{ccc} Q_0(\mathcal{C}) & \xrightarrow{\Delta} & Q_0(\mathcal{C}) \times Q_0(\mathcal{C}) \\ s \downarrow & \lrcorner & \downarrow (s, s) \\ Q_3(\mathcal{C}) & \xrightarrow{(d_{\{0,2\}}, d_{\{1,3\}})} & Q_1(\mathcal{C}) \times Q_1(\mathcal{C}) \end{array}$$

is cartesian. So let P be the pullback. We first check that the induced map $Q_0(\mathcal{C}) \rightarrow P$ is fully faithful. To see this, first note that the canonical map $\text{TwAr}([n])^{\text{op}} \rightarrow \text{TwAr}([0])^{\text{op}}$ induces a fully faithful map $s: \text{Fun}(\text{TwAr}([0])^{\text{op}}, \mathcal{C}) \rightarrow \text{Fun}(\text{TwAr}([n])^{\text{op}}, \mathcal{C})$ for all n . Indeed, $\text{TwAr}([0])^{\text{op}} \simeq *$ is an anima, hence s factors over the full subcategory $\text{Fun}(|\text{TwAr}([n])^{\text{op}}|, \mathcal{C})$. But $|\text{TwAr}([n])^{\text{op}}| \simeq *$ because $\text{TwAr}([n])^{\text{op}}$ contains $(0 \leq n)$ as an initial element, and now it's clear that s is indeed fully faithful. Hence so is $s: Q_0(\mathcal{C}) \rightarrow Q_n(\mathcal{C})$ for all n .

In particular, this is true for $n = 3$. Moreover, $P \rightarrow Q_3(\mathcal{C})$ is fully faithful too because it is a pullback of the fully faithful map $(s, s): Q_0(\mathcal{C}) \times Q_0(\mathcal{C}) \rightarrow Q_1(\mathcal{C}) \times Q_1(\mathcal{C})$. Now the diagram

$$\begin{array}{ccc} Q_0(\mathcal{C}) & \xrightarrow{\quad} & P \\ & \searrow s & \downarrow \\ & & Q_3(\mathcal{C}) \end{array}$$

and the corresponding two-out-of-three property for fully faithful maps show that $Q_0(\mathcal{C}) \rightarrow P$ is fully faithful too.

Now for essential surjectivity. The actual image of $Q_0(\mathcal{C}) \rightarrow P$ consists of the constant diagrams $F: \text{TwAr}([3])^{\text{op}} \rightarrow \mathcal{C}$, hence (by an easy argument) the essential image contains all diagrams that map all edges to equivalences. To see that every 0-simplex in P is a diagram $F: \text{TwAr}([3])^{\text{op}} \rightarrow \mathcal{C}$ of that form, we have to show the following:

$$\begin{array}{c} F(0 \leq 3) \\ \swarrow (1) \quad \searrow (1) \\ F(0 \leq 2) \quad \smile \quad F(1 \leq 3) \\ \swarrow (1) \quad \searrow (3) \quad \swarrow (3) \quad \searrow (1) \\ F(0 \leq 1) \quad \smile \quad F(1 \leq 2) \quad \smile \quad F(2 \leq 3) \\ \swarrow (1) \quad \searrow (2) \quad \swarrow (4) \quad \searrow (4) \quad \swarrow (2) \quad \searrow (1) \\ F(0 \leq 0) \quad F(1 \leq 1) \quad F(2 \leq 2) \quad F(3 \leq 3) \end{array}$$

Suppose we are given a diagram $F: \text{TwAr}([3])^{\text{op}} \rightarrow \mathcal{C}$ in which all squares are pullbacks as indicated and such that the four dashed purple arrows are equivalences. Then all arrows are equivalences.

Indeed, then the two dashed pink arrows $F(0 \leq 3) \rightarrow F(0 \leq 1)$ and $F(0 \leq 3) \rightarrow F(2 \leq 3)$ are equivalences as well, since they are pullbacks the purple arrows. By two-out-of-six, we see that all arrows labelled “(1)” must be equivalences.

Now the commutative square

$$\begin{array}{ccc} F(0 \leq 3) & \xrightarrow{\sim} & F(1 \leq 3) \\ \downarrow \wr & \text{///} & \downarrow \wr \\ F(0 \leq 1) & \longrightarrow & F(1 \leq 1) \end{array}$$

shows that $F(0 \leq 1) \rightarrow F(1 \leq 1)$ is an equivalence. An analogous argument applies to $F(2 \leq 3) \rightarrow F(2 \leq 2)$. In other words, the two arrows labelled “(2)” are equivalences. This implies that the two arrows labelled “(3)” are equivalences, since they are pullbacks of the former. Finally, the two remaining arrows, labelled “(4)”, must be equivalences by two-out-of-three. We are done. \square

CHAPTER II.

II

Symmetric Monoidal and Stable ∞ -Categories

\mathbb{E}_1 -Monoids and \mathbb{E}_1 -Groups

We start things off slowly by only considering (not necessarily symmetric) monoidal ∞ -categories and \mathbb{E}_1 -spaces.

II.0. Stasheff’s Definition of Coherently Associative Monoids. — The naive way to define a monoid in \mathbf{An} would be to have an object $M \in \mathbf{An}$ together with a multiplication map

$$\mu: M \times M \longrightarrow M.$$

Now μ induces two different maps $M^3 \rightarrow M$ (“two ways of bracketing”), so there ought to be a homotopy $H: \mu \circ (\mu \times \text{id}_M) \simeq \mu \circ (\text{id}_M \times \mu)$ between them. But then there are five maps $M^4 \rightarrow M$ (“five ways of bracketing”), and we see that H induces a loop in $\text{Hom}_{\mathbf{An}}(M^4, M)$. Since M should be associative up to *coherent homotopy*, this loop has to be filled. The story goes on: There are a number of maps $M^5 \rightarrow M$ (too lazy to count them) and now there are some 3-simplices to be filled and so on.

Although this looks horrible, it is possible to turn these considerations into a precise definition, and one obtains Stasheff’s \mathbb{A}_∞ -spaces. But we can take a simpler route! Indeed, thanks to our efforts so far, we now have the luxury of saying: “Well, a coherently associative monoid is just an ∞ -category with only one object.” This leads to the following definition.

II.1. Definition. — Let \mathcal{C} be an ∞ -category with finite products (in particular, \mathcal{C} has a final object $*$ in \mathcal{C}). A *cartesian monoid in \mathcal{C}* is a functor $X: \Delta^{\text{op}} \rightarrow \mathcal{C}$ such that:

- (a) $X_0 \simeq *$.
- (b) It satisfies the Segal condition, i.e., the Segal maps $e_i: [1] \rightarrow [n]$ define an equivalence

$$X_n \xrightarrow{\sim} \prod_{i=1}^n X_1$$

Let $\text{Mon}(\mathcal{C}) \subseteq \mathbf{sC}$ be the full sub- ∞ -category spanned by cartesian monoids. If $\mathcal{C} = \mathbf{An}$, these are also called \mathbb{E}_1 -monoids, \mathbb{A}_∞ -spaces, *coherent monoids*, *special Δ -spaces*, For $\mathcal{C} = \mathbf{Cat}_\infty$ we simply call them *monoidal ∞ -categories* (note that these can also be encoded as cocartesian fibrations over Δ^{op}).

To make sense of the condition from Definition II.1(b), note that since $X_0 \simeq *$ is terminal, we have $X_1 \times_{X_0} \cdots \times_{X_0} X_1 \simeq \prod_{i=1}^n X_1$, so this condition really *is* the Segal condition. Somewhat weird though is that we don’t impose any completeness conditions, and in fact,

cartesian monoids in \mathbf{An} are usually *not* complete Segal spaces! This is in some sense fixed by the following proposition.

II.2. Proposition. — *The completion functor $\text{comp}: \mathbf{sAn} \rightarrow \mathbf{CSAn}$ from Lemma/Definition 1.67 restricts to a fully faithful functor*

$$\text{comp}: \mathbf{Mon}(\mathbf{An}) \longrightarrow \mathbf{*/CSAn}$$

*with essential image those pointed complete segal spaces (X, x) with $\pi_0 X_0 = *$.*

In other words, monoids really are categories with one object (up to equivalence, but not up to contractible choice, as Bastiaan pointed out in the lecture), since the target is also pointed categories with connected base.

Proof of Proposition II.2. The proof is not hard, but quite lengthy, so we break it down into four major steps. Be aware that Step 3 wasn't done in the lecture, so any errors in it are my fault.

Step 1. *There exists a reasonable candidate $\text{decomp}: \mathbf{*/CSAn} \rightarrow \mathbf{Mon}(\mathbf{An})$ of an inverse functor (“decompletion”).*

If (X, x) is a pointed complete Segal space, and if $Y \in \mathbf{Mon}(\mathbf{An})$ is the “universal way” to make (X, x) connected and still satisfy the Segal condition, then it's reasonable to expect that Y_n sits inside a pullback

$$\begin{array}{ccc} Y_n & \longrightarrow & X_n \\ \downarrow & \lrcorner & \downarrow \\ * & \xrightarrow{(x, \dots, x)} & X_0^{n+1} \end{array}$$

for all $[n] \in \Delta^{\text{op}}$. We will see that this gives indeed the correct functor.

Let's first address the elephant in the room, i.e. how to make the pointwise-defined Y_n into a functor $Y: \Delta^{\text{op}} \rightarrow \mathbf{An}$, which in turn should be functorial in X . Since taking pullbacks is functorial, we'll have solved both problems at once if we show that the cospan diagram $* \rightarrow X_0^{n+1} \leftarrow X_n$ is functorial in (X, x) and in $[n]$. To this end consider X as a functor $\Delta \rightarrow \mathbf{An}^{\text{op}}$ and colimit-extend it to a functor

$$| \quad |_X: \mathbf{sAn} \longrightarrow \mathbf{An}^{\text{op}}.$$

This can be done functorially in X , as Theorem 1.51 shows. In more abstract terms, what we do here is to identify $\mathbf{sAn} \simeq \mathbf{Fun}(\Delta, \mathbf{An}^{\text{op}})^{\text{op}} \simeq \mathbf{Fun}^L(\mathbf{sAn}, \mathbf{An}^{\text{op}})^{\text{op}}$. We define two functors $\Delta, \Delta_0: \Delta \rightarrow \mathbf{sSet} \subseteq \mathbf{sAn}$ via

$$\Delta([n]) = \Delta^n \quad \text{and} \quad \Delta_0([n]) = \coprod_{i=0}^n \Delta^0.$$

Since Δ and Δ_0 are functors between 1-categories (before we compose them with the inclusion $\mathbf{sSet} \subseteq \mathbf{sAn}$), we don't need to check any higher coherences and immediately obtain that Δ and Δ_0 are indeed functors. Moreover, the canonical map $\coprod_{i=0}^n \Delta^0 \rightarrow \Delta^n$ sending the i^{th} component to the i^{th} vertex of Δ^n is clearly functorial in $[n] \in \Delta$, hence it induces a natural transformation $\Delta_0 \Rightarrow \Delta$ (which exhibits Δ_0 as the “0-skeleton” of Δ , hence the notation). Now consider the two composites

$$\mathbf{sAn} \simeq \mathbf{Fun}(\Delta, \mathbf{An}^{\text{op}})^{\text{op}} \simeq \mathbf{Fun}(\mathbf{sAn}, \mathbf{An}^{\text{op}})^{\text{op}} \xrightleftharpoons[\Delta_0^*]{\Delta^*} \mathbf{Fun}(\Delta, \mathbf{An}^{\text{op}})^{\text{op}} \simeq \mathbf{sAn},$$

which we denote $|\Delta|_{(-)}$ and $|\Delta_0|_{(-)}$. The transformation $\Delta_0 \Rightarrow \Delta$ induces a transformation $|\Delta|_{(-)} \Rightarrow |\Delta_0|_{(-)}$ going in the other direction, since there has been an $(-)^{\text{op}}$ in between. Upon closer inspection, we find that $\Delta: \Delta \rightarrow \text{sAn}$ is actually nothing else but the Yoneda embedding, hence Theorem I.51 shows that $|\Delta|_{(-)} \simeq \text{id}$. Moreover, since $|-|_X$ preserves colimits, we easily obtain

$$X_0^{n+1} \simeq \left| \prod_{i=0}^n \Delta^0 \right|_X,$$

hence the X_0^{n+1} can be organized into a simplicial anima $|\Delta_0|_X$. The transformation $\text{id} \simeq |\Delta|_{(-)} \Rightarrow |\Delta_0|_{(-)}$ gives a natural map $X \rightarrow |\Delta_0|_X$.

This shows that the maps $X_0^{n+1} \leftarrow X_n$ can indeed be made functorial in X and $[n]$. A similar argument applies shows that the maps $* \rightarrow X_0^{n+1}$, which assemble into a functorial map $\Delta^0 \rightarrow |\Delta_0|_X$. We conclude that $Y \simeq X \times_{|\Delta_0|_X} \Delta^0$ as above is indeed a simplicial anima and functorial in X . We have $Y_0 \simeq *$ by construction and also $Y_n \simeq Y_1 \times_{Y_0} \cdots \times_{Y_0} Y_1$ follows from the fact that X itself satisfies the Segal conditions together with some abstract pullback nonsense (basically that “limits commute”, see the dual of Proposition I.42). So $Y \in \text{Mon}(\text{An})$ and we can finally write down the functor

$$\text{decomp}: */\text{CSAn} \longrightarrow \text{Mon}(\text{An}).$$

This finishes Step 1.

Step 2. The completion functor $\text{comp}: \text{Mon}(\text{An}) \rightarrow \text{CSAn}$ from Lemma/Definition I.67 factors over $/\text{CSAn} \rightarrow \text{CSAn}$ and its essential image is contained in those pointed complete Segal spaces (X, x) with $\pi_0 X_0 = *$.*

To get the factorisation, observe that $\text{Mon}(\text{An})$ has an initial object, namely $\Delta^0 \in \text{sSet} \subseteq \text{sAn}$ (in the lecture we called it “const*”, but it’s really just Δ^0). Indeed, if $T \in \text{Mon}(\text{An})$, then

$$\text{Hom}_{\text{Mon}(\text{An})}(\Delta^0, T) \simeq \text{Hom}_{\text{sAn}}(\Delta^0, T) \simeq T_0 \simeq *$$

by Definition II.1(a). Also $\text{comp}(\Delta^0) \simeq *$, so the completion functor really lifts to a functor $\text{Mon}(\text{An}) \simeq \Delta^0/\text{Mon}(\text{An}) \rightarrow */\text{CSAn}$, as required.

To check that $\pi_0(\text{comp } T)_0 = *$, recall that $(\text{comp } T)_0 \simeq \text{core asscat } T \simeq |T^\times|$ by (I.65.1). Now we have to use the following general fact:

(*) *For all $X \in \text{sAn}$ the map $\pi_0 X_0 \rightarrow \pi_0 |X|$ is surjective.*

This shows what we want since $(T^\times)_0 \simeq T_0 \simeq *$ holds by assumption. To see where (*) comes from, observe that π_0 , being a left adjoint (Example I.33(a)), commutes with colimits. This shows $\pi_0 |X| \simeq \pi_0 \text{colim}_{\Delta^{\text{op}}} X \simeq \text{colim}_{\Delta^{\text{op}}} \pi_0 X$, where the colimit on the right-hand side is a good old colimit taken in sSet . Now $\{d_0, d_1: [1] \rightrightarrows [0]\} \subseteq \Delta^{\text{op}}$ is 1-cofinal (but not ∞ -cofinal!), hence

$$\text{colim}_{\Delta^{\text{op}}} \pi_0 X = \text{Coeq} \left(\pi_0 X_1 \xrightleftharpoons[d_0]{d_1} \pi_0 X_0 \right).$$

This immediately implies (*) and thus Step 2 is done.

Step 3. The functors from Step 1 and Step 2 fit into an adjunction

$$\text{comp}: \text{Mon}(\text{An}) \rightleftarrows */\text{CSAn} : \text{decomp}$$

Consider $T \in \text{Mon}(\text{An})$, which is naturally pointed via the unique (up to contractible choice) map $1_T: \Delta^0 \rightarrow T$, and let (X, x) be a pointed complete Segal space. Then

$$\begin{aligned} \text{Hom}_{*/\text{CSAn}}(\text{comp } T, (X, x)) &\simeq \text{Hom}_{*/\text{sAn}}((T, 1_T), (X, x)) \\ &\simeq \text{Hom}_{\text{sAn}}(T, X) \times_{\text{Hom}_{\text{sAn}}(\text{const } 1_T, X)} \{\text{const } x\} \\ &\simeq \text{Hom}_{\text{sAn}}(T, X) \times_{\text{Hom}_{\text{An}}(1_T, X_0)} \{x\}. \end{aligned}$$

The first equivalence follows from Lemma/Definition I.67 (and the fact that adjunctions extend to slice categories), the second follows from [HCII, Corollary VIII.6]. The last one follows by inspection. All of them are functorial in (X, x) . Moreover, we have

$$\begin{aligned} \text{Hom}_{\text{Mon}(\text{An})}(T, \text{decomp}(X, x)) &\simeq \text{Hom}_{\text{sAn}}(T, X \times_{|\Delta_0|_X} \Delta^0) \\ &\simeq \text{Hom}_{\text{sAn}}(T, X) \times_{\text{Hom}_{\text{sAn}}(T, |\Delta_0|_X)} \text{Hom}_{\text{sAn}}(T, \Delta^0). \end{aligned}$$

Since $\text{id} \Rightarrow |\Delta_0|_{(-)}$ is a natural transformation, one quickly checks that the morphism $\text{Hom}_{\text{sAn}}(T, X) \rightarrow \text{Hom}_{\text{sAn}}(T, |\Delta_0|_X)$ factors over $\text{Hom}_{\text{sAn}}(|\Delta_0|_T, |\Delta_0|_X)$. But $T_0 \simeq *$, hence $|\Delta_0|_T \simeq \Delta^0$. Also $\text{Hom}_{\text{sAn}}(T, \Delta^0) \simeq *$. Putting everything together, we can rewrite

$$\text{Hom}_{\text{Mon}(\text{An})}(T, \text{decomp}(X, x)) \simeq \text{Hom}_{\text{sAn}}(T, X) \times_{\text{Hom}_{\text{An}}(T_0, X_0)} \{x\},$$

and after another inspection, we see that our calculations of $\text{Hom}_{*/\text{CSAn}}(\text{comp } T, (X, x))$ and $\text{Hom}_{\text{Mon}(\text{An})}(T, \text{decomp}(X, x))$ agree, as they are supposed to. This finishes Step 3.

Step 4. The restriction $\text{decomp}: (/\text{CSAn})_{\pi_0=*} \rightarrow \text{Mon}(\text{An})$ is indeed an inverse to comp , where $(*/\text{CSAn})_{\pi_0=*} \subseteq */\text{CSAn}$ denotes the full subcategory spanned by those (X, x) with $\pi_0 X_0 = *$.*

Thanks to Step 3, we already have unit and counit transformations $\text{id} \Rightarrow \text{decomp} \circ \text{comp}$ and $\text{comp} \circ \text{decomp} \Rightarrow \text{id}$, so we only need to show that they are equivalences. Let's start with the counit. We must show

$$\text{comp}(\text{decomp}(X, x)) \simeq (X, x)$$

for all $(X, x) \in (*/\text{CSAn})_{\pi_0=*}$. Observe that $\text{comp } X \simeq X$ because X is already complete. Hence it suffices to show $\text{asscat}(\text{decomp}(X, x)) \simeq \text{asscat } X$ (and check that the chosen base points on both sides correspond, but that's easy) because $\text{comp} \simeq N^r \circ \text{asscat}$ by Lemma/Definition I.67 and N^r is fully faithful by Theorem/Definition I.64. Using (I.65.1) and the fact that X^\times is a constant simplicial anima with value X_0 as X is complete, we see that

$$\pi_0 \text{core}(\text{asscat } X) \simeq \pi_0 |X^\times| \simeq \pi_0 X_0 \simeq *,$$

so $\text{asscat}(\text{decomp}(X, x)) \rightarrow \text{asscat } X$ is essentially surjective for trivial reasons. Moreover, we have $\text{decomp}(X, x)_0 \simeq \{x\}$ by construction, and (I.65.2) (which is applicable here because $\text{decomp}(X, x)$ is Segal by construction) easily shows

$$\text{Hom}_{\text{asscat}(\text{decomp}(X, x))}(x, x) \simeq P_{x, x} \simeq \text{Hom}_{\text{asscat } X}(x, x),$$

so $\text{asscat}(\text{decomp}(X, x)) \rightarrow \text{asscat } X$ is fully faithful too. This shows that the counit is an equivalence, as claimed. The argument for the unit is similar: We have to show

$$T \simeq \text{decomp}(\text{comp } T)$$

for all $T \in \text{Mon}(\text{An})$. Since both sides are functors $\Delta^{\text{op}} \rightarrow \text{An}$ and a natural transformation between them is already given, we can do this degree-wise by Theorem I.19(b). Since both sides are cartesian monoids, it suffices to check that the map in degree 1 is an equivalence, i.e. that $T_1 \simeq \text{decomp}(\text{comp } T)_1$. Write $(X, x) \simeq \text{comp } T$. By construction we have

$$\text{decomp}(X, x)_1 \simeq P_{x,x} \quad \text{and} \quad T_1 \simeq P_{1_T, 1_T},$$

where as usual $1_T: \Delta^0 \rightarrow T$ is the natural pointing of T . As observed at the very beginning of this proof, the spaces $P_{y,z}$ aren't affected by completion, hence $P_{x,x} \simeq P_{1_T, 1_T}$ and we're done. \square

LECTURE 9
26th Nov, 2020

Under the equivalence $\text{Cat}_\infty \simeq \text{CSAn}$ from Theorem/Definition I.64, the construction of decomp from the proof of Proposition II.2 corresponds to a functor $*/\text{Cat}_\infty \rightarrow \text{Mon}(\text{An})$ which becomes an equivalence when restricted to the full subcategory $(*/\text{Cat}_\infty)_{\geq 1} \subseteq */\text{Cat}_\infty$ spanned by those \mathcal{C} with $\pi_0 \text{core } \mathcal{C} \simeq *$. In diagrams,

$$\begin{array}{ccc} */\text{Cat}_\infty & \longrightarrow & \text{Mon}(\text{An}) \\ \cup \! \! \! \downarrow & \nearrow \sim & \\ (* / \text{Cat}_\infty)_{\geq 1} & & \end{array}$$

Explicitly, this functor sends a pointed ∞ -category $(\{x\} \rightarrow \mathcal{C})$ to $\text{Hom}_{\mathcal{C}}(x, x)$ in degree 1 (and then all other degrees are determined by the conditions from Definition II.1). We thus obtain:

II.3. Corollary. — *If \mathcal{C} is an ∞ -category and $x \in \mathcal{C}$, then $\text{Hom}_{\mathcal{C}}(x, x)$ carries a canonical structure of an \mathbb{E}_1 -monoid.*

II.4. Definition. — Let \mathcal{C} be an ∞ -category with finite products. A cartesian monoid X in \mathcal{C} is called a *cartesian group* (or *\mathbb{E}_1 -group* in the case $\mathcal{C} = \text{An}$) if the map

$$\begin{aligned} (\text{pr}_1, \circ): X_1 \times X_1 &\xrightarrow{\sim} X_1 \times X_1 \\ (f, g) &\longmapsto (f, f \circ g) \end{aligned}$$

is an equivalence. Here “ \circ ” is the composition as defined in Theorem/Definition I.64(c). We denote by $\text{Grp}(\mathcal{C}) \subseteq \text{Mon}(\mathcal{C})$ the full sub- ∞ -category of cartesian groups.

II.5. Proposition. — *The equivalences $\text{Mon}(\text{An}) \simeq (* / \text{Cat}_\infty) \simeq (* / \text{CSAn})_{\pi_0=*}$ restrict to equivalences*

$$\begin{array}{ccc} \text{Mon}(\text{An}) & \supseteq & \text{Grp}(\text{An}) \\ \downarrow \wr & & \downarrow \wr \\ (* / \text{Cat}_\infty)_{\geq 1} & \supseteq & (* / \text{An})_{\geq 1} \\ \downarrow \wr & & \downarrow \wr \\ (* / \text{CSAn})_{\pi_0=*} & \supseteq & (* / \text{sAn}_{\text{const}})_{\pi_0=*} \end{array}$$

In the lower right corner we use the notation from Example I.68(c).

Proof sketch. In the lecture we noted only that $X \in \text{Mon}(\text{An})$ is an \mathbb{E}_1 -group iff the set $\pi_0 X_1$ with its induced ordinary monoid structure is an ordinary group.

To elaborate on this a bit more, first note that we may assume $X \simeq \text{Hom}_{\mathcal{C}}(x, x)$ for some ∞ -category \mathcal{C} with $\pi_0 \text{core } \mathcal{C} \simeq *$, as was argued above. If X is an \mathbb{E}_1 -group, then $\pi_0 \text{Hom}_{\mathcal{C}}(x, x)$ must be an ordinary group, which implies that all endomorphisms of x are equivalences, hence all morphisms in \mathcal{C} are equivalences as $\pi_0 \text{core } \mathcal{C} \simeq *$. This proves that \mathcal{C} is an anima by Theorem I.13. Conversely, if \mathcal{C} is an anima, either inclusion $[1] \hookrightarrow J$ into the free-living isomorphism induces a trivial fibration $\text{Fun}(J, \mathcal{C}) \xrightarrow{\sim} \text{Ar}(\mathcal{C})$. Choosing a section and composing it with the other map $\text{Fun}(J, \mathcal{C}) \rightarrow \text{Ar}(\mathcal{C})$ (induced by the other inclusion $[1] \hookrightarrow J$) yields an “inversion” map $(-)^{-1}: \text{Hom}_{\mathcal{C}}(x, x) \rightarrow \text{Hom}_{\mathcal{C}}(x, x)$. From this we can construct an inverse of the map in Definition II.4 \square

II.6. Some Explicit Calculations. — The equivalence $\text{Grp}(\text{An}) \simeq (*/\text{An})_{\geq 1}$ is explicitly given as follows:

$$B = | \cdot | : \text{Grp}(\text{An}) \xrightarrow{\sim} (*/\text{An})_{\geq 1} : \Omega.$$

The use B to denote the functor $| \cdot | = \text{colim}_{\Delta^{\text{op}}}$ is common in the literature, as it is short for “bar construction”. Fabian doesn’t like it, since he finds it stupid to name something after the way it was typeset in the pre-L^AT_EX age (where people had to resort to bars to denote their tensor products). Amusingly though, his preferred notation consists of actual bars ... In these notes, however, I’ll show my bad taste and consistently use B .

Anyway, back to the matters at hand. To make sense of why the above equivalence is indeed given by $B = | \cdot |$ and Ω , recall that the composition

$$\text{Mon}(\text{An}) \xrightarrow{\text{comp}} \text{CSAn} \xrightarrow{\text{ev}_0} \text{An}$$

sends $X \mapsto |X^\times|$, as follows from (I.65.1) and some unravelling. If X is an \mathbb{E}_1 -group, one easily verifies $X^\times = X$ and thus $X \in \text{Grp}(\text{An})$ is sent to $BX = |X|$, as claimed. Conversely, the composition

$$(*/\text{An})_{\geq 1} \xrightarrow{\text{decomp}} \text{Grp}(\text{An}) \xrightarrow{\text{ev}_1} \text{An}$$

sends a pointed connected anima (K, k) to $\text{Hom}_K(k, k)$, as argued previously. So to justify why $(*/\text{An})_{\geq 1} \xrightarrow{\sim} \text{Grp}(\text{An})$ should be thought of as a “loop space functor”, we must explain why $\Omega_k K \simeq \text{Hom}_K(k, k)$. This can be done by several arguments and I will present a different one than in the lecture. The loop space is usually defined by the left of the following pullback diagrams in An :

$$\begin{array}{ccc} \Omega_k K & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow k \\ * & \xrightarrow{k} & K \end{array} \xrightarrow{\sim} \begin{array}{ccc} \text{Hom}_K^L(k, k) & \longrightarrow & K_{k/} \\ \downarrow & \lrcorner & \downarrow \\ * & \xrightarrow{k} & K \end{array}$$

The map $* \rightarrow K$ can be factored into the left anodyne map $* \hookrightarrow K_{k/}$ followed by the left fibration $K_{k/} \rightarrow K$, which is even a Kan fibration since K is an anima. Replacing one of the $* \rightarrow K$ accordingly, we obtain the pullback diagram on the right, which is already a pullback in sSet as one leg is a Kan fibration (see Theorem I.34). Its pullback is thus $\text{Hom}_K^L(k, k) \simeq \text{Hom}_K(k, k)$ as left-Hom anima coincide with the regular Hom anima by [HCII, Digression I Corollary D.2].

In particular, we have proved another result from algebraic topology (or rather deduced it from the much more general statement of Theorem/Definition I.64, which we didn't prove, but never mind that).

II.7. Corollary (Recognition Principle for Loop Spaces, Stasheff). — *The loop functor $\Omega: */\text{An} \rightarrow \text{An}$ lifts to an equivalence*

$$\Omega: (*/\text{An})_{\geq 1} \xrightarrow{\sim} \text{Grp}(\text{An}).$$

Note that $\text{ev}_1: \text{Mon}(\text{An}) \rightarrow \text{An}$ preserves limits since $\text{Mon}(\text{An})$ is closed under limits in sAn (the Segal condition is given by a limit and limits commute by the dual of Proposition I.42). Colimits however are bloody complicated, and even though one can show that $\text{Mon}(\text{An})$ is cocomplete, colimits in it are *by far not* just colimits of underlying simplicial anima, same as colimits of ordinary monoids are much more complicated than colimits in sets.

II.8. Proposition. — *The inclusion $\text{Grp}(\text{An}) \subseteq \text{Mon}(\text{An})$ has a left adjoint called group completion and denoted*

$$(-)^{\infty\text{-grp}}: \text{Mon}(\text{An}) \longrightarrow \text{Grp}(\text{An}).$$

It is given by $X^{\infty\text{-grp}} \simeq \Omega BX$, where $B: \text{Mon}(\text{An}) \rightarrow (/\text{An})_{\geq 1}$ denotes a functor extending the B from II.6.*

Proof. We know that $\text{An} \subseteq \text{Cat}_{\infty}$ has a left-adjoint $| |: \text{Cat}_{\infty} \rightarrow \text{An}$. The unit and counit are equivalences on $*$ $\in \text{An}$, hence the adjunction passes to slices under it. Moreover the $(-)^{\geq 1}$ -parts on both sides are clearly preserved, thus $| |$ descends to a left adjoint

$$| |: (*/\text{Cat}_{\infty})_{\geq 1} \longrightarrow (*/\text{An})_{\geq 1}$$

of the obvious inclusion in the other direction. Now use Proposition II.5 to get the desired left-adjoint $(-)^{\infty\text{-grp}}: \text{Mon}(\text{An}) \rightarrow \text{Grp}(\text{An})$. The explicit description follows from our calculations in II.6, and Remark I.66, which ensures that the different meanings of $| |$ ($\text{colim}_{\Delta^{\text{op}}}$ or localisation of an ∞ -category at all its morphisms) are actually compatible. \square

A consequence that wasn't explicitly mentioned, but used later in the course, is the following:

II.8a. Corollary*. — *Without restricting them as in II.6, the functors B and Ω form an adjunction*

$$B: \text{Mon}(\text{An}) \rightleftarrows */\text{An}: \Omega.$$

Proof.* For a pointed anima (X, x) , let X_x denote the connected component containing x . Then $\Omega_x X \simeq \Omega_x X_x$, so the equivalence $\Omega: (*/\text{An})_{\geq 1} \xrightarrow{\sim} \text{Grp}(\text{An})$ from II.6 extends to a functor $\Omega: */\text{An} \rightarrow \text{Grp}(\text{An})$. Now if $M \in \text{Mon}(\text{An})$ is an \mathbb{E}_1 -monoid, we can compute

$$\begin{aligned} \text{Hom}_{\text{Mon}(\text{An})}(M, \Omega_x X) &\simeq \text{Hom}_{\text{Grp}(\text{An})}(M^{\infty\text{-grp}}, \Omega_x X_x) \\ &\simeq \text{Hom}_{\text{Grp}(\text{An})}(\Omega BM, \Omega_x X_x) \\ &\simeq \text{Hom}_{(*/\text{An})_{\geq 1}}(BM, X_x) \\ &\simeq \text{Hom}_{*/\text{An}}(BM, X), \end{aligned}$$

which establishes the desired adjunction. The second equivalence follows from Proposition II.8, the third equivalence uses that $\Omega: (*/\text{An})_{\geq 1} \xrightarrow{\sim} \text{Grp}(\text{An})$ is an equivalence, and the fourth equivalence follows from the fact that BM is connected (by construction), hence any pointed map $BM \rightarrow X$ only hits the connected component X_x . \square

Next we will discuss some examples. We begin with free monoids and groups.

II.9. Proposition. — *The evaluation functor $\text{ev}_1: \text{Mon}(\text{An}) \rightarrow \text{An}$ has a left adjoint $\text{Free}^{\text{Mon}}: \text{An} \rightarrow \text{Mon}(\text{An})$. It is given explicitly by the “anima of words of arbitrary length”*

$$\text{Free}^{\text{Mon}}(K)_1 \simeq \coprod_{n \geq 0} K^n.$$

We postpone the proof of Proposition II.9 indefinitely (but Fabian’s notes give a sketch in the much more general setting of algebras over ∞ -operads, see [A&HK, Chapter II pp. 128–132]). By the explicit construction of Free^{Mon} , the diagram

$$\begin{array}{ccc} \text{Set} & \xrightarrow{\text{Free}} & \text{Mon}(\text{Set}) \\ \downarrow & & \downarrow \\ \text{An} & \xrightarrow{\text{Free}^{\text{Mon}}} & \text{Mon}(\text{An}) \end{array}$$

commutes. But be aware that this is not automatic, and in fact we’ll see it very much breaks once we impose commutativity (see pages 86 and 99). As some first examples, one has $\text{Free}^{\text{Mon}}(*) \simeq \mathbb{N}$ and $\text{Free}^{\text{Mon}}(\emptyset) \simeq *$, which is initial in $\text{Mon}(\text{An})$.

II.10. Proposition. — *The evaluation functor $\text{ev}_1: \text{Grp}(\text{An}) \rightarrow \text{An}$ has a left-adjoint $\text{Free}^{\text{Grp}}: \text{An} \rightarrow \text{Grp}(\text{An})$ too. Explicitly, it is given by the composite*

$$\begin{aligned} \text{An} &\longrightarrow */\text{An} \xrightarrow{\Sigma} */\text{An} \xrightarrow{\Omega} \text{Grp}(\text{An}) \\ X &\longmapsto X \sqcup *. \end{aligned}$$

In other words, the free \mathbb{E}_1 -group on X is $\text{Free}^{\text{Grp}}(X) \simeq \Omega \Sigma X_+$, where $X_+ = X \sqcup *$. As usual, the suspension functor $\Sigma: \text{An} \rightarrow \text{An}$ is given by the pushout diagram

$$\begin{array}{ccc} X & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \Sigma X \end{array}$$

II.10a. Remark*. — For the proof, we’ll need a functor $\Sigma: */\text{An} \rightarrow */\text{An}$ instead. It is induced by its unpointed variant, of course, but we can also define it by taking the above pushout in $*/\text{An}$. In fact, it doesn’t matter whether the pushout is taken in $*/\text{An}$ or An since $*/\text{An} \rightarrow \text{An}$ commutes with weakly contractible colimits (but not with arbitrary ones, which I erroneously claimed in an earlier version of these notes; for example, it doesn’t preserve the initial object). Indeed, if \mathcal{I} is any diagram shape, then a quick calculation shows $\text{Fun}(\mathcal{I}, */\text{An}) \simeq (\text{const } *) / \text{Fun}(\mathcal{I}, \text{An})$. For the adjunction

$$\text{colim}_{\mathcal{I}}: \text{Fun}(\mathcal{I}, \text{An}) \rightleftarrows \text{An} : \text{const}$$

to descend to slice categories below $\text{const } *$ and $*$, we need that these objects are mapped to each other and that unit/counit are equivalences on them. If \mathcal{I} is weakly contractible, then $\text{colim}_{\mathcal{I}}(\text{const } *) \simeq |\mathcal{I}| \simeq *$ by Proposition I.36 and the required conditions are easily checked. Therefore, the adjunction above descends to an adjunction

$$\text{colim}_{\mathcal{I}}: \text{Fun}(\mathcal{I}, */\text{An}) \simeq (\text{const } *) / \text{Fun}(\mathcal{I}, \text{An}) \rightleftarrows */\text{An} : \text{const}$$

in this case. If you think about this for a moment, this is exactly what we need to show that $*/\text{An} \rightarrow \text{An}$ commutes with colimits over \mathcal{I} .

Similarly, once we have chosen a basepoint $x \in X$, it doesn't matter whether we take the defining pullback of $\Omega_x X$ in $*/\text{An}$ or in An (and then equip it with its natural basepoint). This can be seen as above, or simply by the fact that $*/\text{An} \rightarrow \text{An}$ has a left adjoint, so we can apply Observation I.41. However, in contrast to Σ , we can't define Ω as a functor on An , since there is no canonical choice of basepoint.

Proof sketch of Proposition II.10. The existence of a left adjoint would be immediate from Propositions II.8 and II.9, but we can give a direct argument that also yields the explicit description. We have seen in II.6 that the diagram

$$\begin{array}{ccc} \text{Grp}(\text{An}) & \xrightarrow{\text{ev}_1} & \text{An} \\ B \downarrow & \nearrow \Omega & \\ (*/\text{An})_{\geq 1} & & \end{array}$$

commutes. Observe that $(-)_+ = - \sqcup *: \text{An} \rightarrow */\text{An}$ is a left adjoint of the forgetful functor $*/\text{An} \rightarrow \text{An}$ (this is basically trivial) and that $\Sigma: */\text{An} \rightleftarrows */\text{An} : \Omega$ are adjoints (this we leave as an exercise—just play around with the defining pushouts/pullbacks, which can be taken in $*/\text{An}$ by Remark* II.10a). Finally, $\Omega: (*/\text{An})_{\geq 1} \rightleftarrows \text{Grp}(\text{An}) : B$ is an adjunction (even an equivalence) by II.6. The assertion now follows from the fact that left adjoints compose. \square

II.11. Example. — (a) The evaluation functor $\text{ev}_1: \text{Mon}(\text{An}) \rightarrow \text{An}$ factors canonically over $*/\text{An} \rightarrow \text{An}$, and as it turns out, the resulting functors

$$\text{ev}_1: \text{Mon}(\text{An}) \longrightarrow */\text{An} \quad \text{and} \quad \text{ev}_1: \text{Grp}(\text{An}) \longrightarrow */\text{An}$$

both have left adjoints. For a pointed anima (X, x) , we denote the corresponding left adjoint objects by $\text{Free}^{\text{Mon}}(X, x)$ and $\text{Free}^{\text{Grp}}(X, x)$ and think of them as the “free \mathbb{E}_1 -monoid/group with unit x ”. We have

$$\text{Free}^{\text{Mon}}(X, x)^{\infty\text{-grp}} \simeq \text{Free}^{\text{Grp}}(X, x) \simeq \Omega \Sigma X$$

The first equivalence follows from the fact that left adjoints compose. The second equivalence can be seen along the lines of the proof of Proposition II.10, without us needing to know that $\text{Free}^{\text{Mon}}(X, x)$ exists.

If (X, x) is connected, one actually has $\text{Free}^{\text{Mon}}(X, x) \simeq \text{Free}^{\text{Grp}}(X, x)$. Indeed, as we've seen in the proof of Proposition II.5, we only need to check that the ordinary monoid $\pi_0 \text{Free}^{\text{Mon}}(X, x)_1$ is already an ordinary group. Playing around with universal properties, we see that $\pi_0 \text{Free}^{\text{Mon}}(X, x)_1$ is the free ordinary monoid on the set $\pi_0 X$ with unit $[x]$, hence just $\{[x]\}$ itself since X is connected, hence it already is a group.

- (b) By Proposition II.9, $\text{Free}^{\text{Mon}}(\mathbb{S}^1) \simeq \coprod_{n \geq 0} \mathbb{T}^n$ is an infinite union of tori $\mathbb{T}^n = \prod_{i=1}^n \mathbb{S}^1$ and thus a reasonable geometric object. However, Proposition II.10 implies

$$\text{Free}^{\text{Grp}}(\mathbb{S}^1) \simeq \Omega \Sigma(\mathbb{S}_+^1) \simeq \Omega(\mathbb{S}^2 \vee \mathbb{S}^1),$$

and the homotopy groups of the right-hand side are not fully known to this day. Similarly, using the pointed variant from (a) above, we get

$$\text{Free}^{\text{Mon}}(\mathbb{S}^1, *) \simeq \text{Free}^{\text{Grp}}(\mathbb{S}^1, *) \simeq \Omega \Sigma(\mathbb{S}^1, *) \simeq \Omega \mathbb{S}^2$$

LECTURE 10
1st Dec, 2020

- (c) It is completely formal to see that $\text{ev}_1: \text{Mon}(\text{An}) \rightarrow \text{An}$ and $\text{ev}_1: \text{Grp}(\text{An}) \rightarrow \text{An}$ are represented by $\text{Free}^{\text{Mon}}(*)$ and $\text{Free}^{\text{Grp}}(*) \simeq \text{Free}^{\text{Mon}}(*)^{\infty\text{-grp}}$ respectively. What's not formal, however, is that $\text{Free}^{\text{Mon}}(*) \simeq \mathbb{N}$ and $\text{Free}^{\text{Grp}}(*) \simeq \Omega\mathbb{S}^1 \simeq \mathbb{Z}$, regarded as discrete simplicial anima. This really needs Propositions II.9 and II.10.

Moreover, we'll see (pages 86 and 99) that the analogous functors are no longer represented by \mathbb{N} and \mathbb{Z} once we impose commutativity, even though they are the free \mathbb{E}_1 -monoid/group on a point and happen to be commutative. In fact they are “too commutative to be free commutative monoids” (**sphere spectrum intensifies**).

- (d) Let's for some reason compute the pushout

$$\begin{array}{ccc} \mathbb{N} & \xrightarrow{\cdot 2} & \mathbb{N} \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \mathbb{N} // 2 \end{array}$$

in $\text{Mon}(\text{An})$. The answer is $\mathbb{N} // 2 \simeq \Omega\mathbb{RP}^2$. To see this, first observe that the functor $\pi_0: \text{Mon}(\text{An}) \rightarrow \text{Mon}(\text{Set})$ commutes with colimits, so $\pi_0(\mathbb{N} // 2) = \mathbb{Z} / 2$ since we know how to compute pushouts in $\text{Mon}(\text{Set})$. But $\mathbb{Z} / 2$ is a group, so $\mathbb{N} // 2 \in \text{Grp}(\text{An})$ by Proposition II.5. In particular, since the group completion functor $(-)^{\infty\text{-grp}}$ commutes with colimits, we get $\mathbb{N} // 2 \simeq \mathbb{Z} // 2$, i.e., the diagram

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{\cdot 2} & \mathbb{Z} \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \mathbb{N} // 2 \end{array}$$

is a pushout in $\text{Grp}(\text{An})$. Translating to $*/\text{An}$ via Proposition II.5 gives a diagram

$$\begin{array}{ccc} \mathbb{S}^1 & \xrightarrow{\cdot 2} & \mathbb{S}^1 \\ \downarrow & \lrcorner & \downarrow \\ \mathbb{D}^2 & \longrightarrow & \mathbb{RP}^2 \end{array}$$

(we have replaced $*$ by the disk \mathbb{D}^2 to make one leg of the pushout into a cofibration so that it can be computed by an ordinary pushout of CW complexes by Theorem I.34) and translating back using II.6 shows $\mathbb{N} // 2 \simeq \Omega\mathbb{RP}^2$, as claimed.

- (e) Every monoidal 1-category (\mathcal{C}, \otimes) gives rise to a monoidal ∞ -category, i.e. an object in $\text{Mon}(\text{Cat}_1^{(2)}) \subseteq \text{Mon}(\text{Cat}_\infty)$. We will produce this by straightening a suitable cocartesian fibration $p^\otimes: \mathcal{C}^\otimes \rightarrow \Delta^{\text{op}}$ of 1-categories. Define $\mathcal{C}^\otimes \in \text{Cat}_1^{(2)}$ as follows: Its objects are

$$\text{obj}(\mathcal{C}^\otimes) = \coprod_{n \geq 0} \text{obj}(\mathcal{C})^n,$$

i.e. given by tuples (n, x_1, \dots, x_n) with $[n] \in \Delta$ and $x_i \in \mathcal{C}$. We'll usually abbreviate $x = (x_1, \dots, x_n)$ and simply write (n, x) . Morphisms are defined as

$$\text{Hom}_{\mathcal{C}^\otimes}((n, x), (m, y)) = \left\{ (\alpha, f) \left| \begin{array}{l} \alpha: [m] \rightarrow [n] \text{ in } \Delta \text{ and } f = (f_1, \dots, f_m) \\ \text{where } f_j: x_{\alpha(j-1)+1} \otimes \dots \otimes x_{\alpha(j)} \rightarrow y_j \\ \text{are morphisms in } \mathcal{C} \text{ for all } j = 1, \dots, m \end{array} \right. \right\}.$$

If $\alpha(j-1) + 1 > \alpha(j)$ for some j , then $x_{\alpha(j-1)+1} \otimes \cdots \otimes x_{\alpha(j)}$ is the empty tensor product, which is the tensor unit $1_{\mathcal{C}} \in \mathcal{C}$ by definition.

The composition of $(\alpha, f): (n, x) \rightarrow (m, y)$ and $(\beta, g): (m, y) \rightarrow (k, z)$ is given as follows: Its first component is $\alpha \circ \beta: [k] \rightarrow [n]$. For the second component, observe that we can write

$$x_{\alpha\beta(j-1)+1} \otimes \cdots \otimes x_{\alpha\beta(j)} = \bigotimes_{i=1}^{\beta(j)-\beta(j-1)} \left(x_{\alpha(\beta(j-1)+i-1)+1} \otimes \cdots \otimes x_{\alpha(\beta(j-1)+i)} \right).$$

The i^{th} tensor factor on the right maps to $y_{\beta(j-1)+i}$ via $f_{\beta(j-1)+i}$. Tensoring them together and postcomposing with $g_j: y_{\beta(j-1)+1} \otimes \cdots \otimes z_j$ provides the desired composition of f and g . This finishes the construction of \mathcal{C}^{\otimes} .

The functor $p^{\otimes}: \mathcal{C}^{\otimes} \rightarrow \Delta^{\text{op}}$ just extracts $[n]$ from (n, x) and α from (α, f) . The fibres \mathcal{C}_n^{\otimes} of p^{\otimes} are then obviously given by \mathcal{C}^n . To show that p is a cocartesian fibration of 1-categories, we must provide cocartesian lifts of morphisms in Δ^{op} . Given $\alpha: [m] \rightarrow [n]$ in Δ and $(n, x) \in \mathcal{C}^{\otimes}$, we claim that the identities can be regarded as a morphism

$$\alpha_*: (n, x) \longrightarrow (m, x_{\alpha(0)+1} \otimes \cdots \otimes x_{\alpha(1)}, \dots, x_{\alpha(m-1)+1} \otimes \cdots \otimes x_{\alpha(m)}).$$

It's straightforward to check that α_* is a cocartesian lift of α (note that the daunting homotopy pullbacks in Definition I.24 become just good old pullbacks of sets/discrete anima, since we are dealing with 1-categories).

It remains to check that $\text{St}(p^{\otimes})$ satisfies the conditions from Definition II.1. As seen above, $\text{St}(p^{\otimes})_0 \simeq \mathcal{C}_0^{\otimes} \simeq \mathcal{C}^0 \simeq *$ is a point, as desired. To verify the Segal condition, let's organize the world a little bit. Call a map in Δ *inert* if it is the inclusion of an interval and *active* if it preserves the largest and the smallest element. The Segal maps $e_i: [1] \rightarrow [n]$ are precisely the inert maps with source $[1]$. We claim that for a general inert map $\alpha: [m] \rightarrow [n]$, the induced functor (given by taking cocartesian lifts)

$$\alpha_*: \mathcal{C}^n \simeq \mathcal{C}_n^{\otimes} \longrightarrow \mathcal{C}_m^{\otimes} \simeq \mathcal{C}^m$$

simply sends $(x_1, \dots, x_n) \in \mathcal{C}^n$ to $(x_{\alpha(1)}, \dots, x_{\alpha(m)}) \in \mathcal{C}^m$ with no tensor products occurring. Thus the Segal condition follows immediately.

As a slogan, “inerts induce forgetful functors on fibres”, whereas “actives don't forget, they just tensor”. Generally, any map in Δ can be uniquely factored into an inert followed by an active.

- (f) The construction from (e) also works if \mathcal{C} is Kan enriched. In this case one can again construct a cocartesian fibration $p^{\otimes}: \mathcal{C}^{\otimes} \rightarrow \Delta^{\text{op}}$, but of Kan-enriched categories this time (where Δ^{op} receives its discrete enrichment). Then $N^c(\mathcal{C}^{\otimes}) \rightarrow N^c(\Delta^{\text{op}}) \simeq \Delta^{\text{op}}$ straightens to a monoidal ∞ -category as well. Fabian wouldn't tell us what a cocartesian fibration of Kan-enriched categories is though, since this would take—not a lot, but too much—time.

Example II.11(e) allows us to construct many interesting examples. All examples will turn out to be commutative (see Example II.21a), but that doesn't matter for now.

II.12. Algebraic K -Theory. — Let R be a ring and consider $\text{Mod}(R)$ as a monoidal category under \oplus . Moreover, let $\text{Proj}(R)$ denote the sub-groupoid of spanned by finite

projective R -modules (“vector bundles on $\mathrm{Spec} R$ ”) and their isomorphisms. This inherits a (symmetric) monoidal structure by Example II.11(e), so we obtain

$$\mathrm{Proj}(R) \in \mathrm{Mon}(\mathrm{Grpd}_1^{(2)}) \subseteq \mathrm{Mon}(\mathrm{An}).$$

II.12a. Definition (Quillen). — The *projective class anima* (or *algebraic K -theory space*) of a ring R is defined as

$$k(R) := \mathrm{Proj}(R)^{\infty\text{-grp}} \in \mathrm{Grp}(\mathrm{An}),$$

and the *higher projective class groups* (or *higher K -groups*) of R are given by

$$K_i(R) = \pi_i(k(R)_1, *).$$

The reason we write $k(R)$ is that the notation $K(R)$ is reserved for the K -theory *spectrum* which we will eventually define. Note immediately that

$$K_0(R) = \pi_0(\mathrm{Proj}(R)^{\infty\text{-grp}}) = (\pi_0 \mathrm{Proj}(R))^{\mathrm{grp}} = \left\{ \begin{array}{l} \text{iso. classes of finite} \\ \text{projective } R\text{-modules} \end{array} \right\}^{\mathrm{grp}},$$

where $(-)^{\mathrm{grp}}: \mathrm{Mon}(\mathrm{Set}) \rightarrow \mathrm{Grp}(\mathrm{Set})$ denotes the ordinary group completion. This shows that $K_0(R)$ is precisely what we wanted it to be way back in the introduction, Definition 0.6. We will soon(ish, Corollary III.7) see that also $K_1(R)$ can be described as in Definition 0.8.

II.13. Hermitian K -Theory/Grothendieck–Witt Theory. — Let R be a ring, M an $R \otimes R$ -module and $\sigma: M \rightarrow M$ an involution that is flip-linear, i.e. $\sigma((a \otimes b)m) = (b \otimes a)\sigma(m)$. Moreover, assume that M is finite projective when considered as an R -module via the map $R \simeq \mathbb{Z} \otimes R \rightarrow R \otimes R$ (one can check that it doesn’t matter if we choose this map or $R \simeq R \otimes \mathbb{Z} \rightarrow R \otimes R$) and that the map

$$\begin{aligned} R &\xrightarrow{\sim} \mathrm{Hom}_R(M, M) \\ r &\longmapsto (m \mapsto (r \otimes 1)m) \end{aligned}$$

is an isomorphism. A pair (M, σ) subject to these conditions is called an *invertible module with involution* over R (beware that this is non-standard terminology).

For a finite projective R -module $P \in \mathrm{Proj}(R)$ consider the group $\mathrm{Hom}_{R \otimes R}(P \otimes P, M)$ of M -valued R -bilinear forms on P . This is acted upon by $\mathbb{Z}/2$ via conjugation with the flip of $P \otimes P$ and σ on M . Define the groups of *symmetric* and *quadratic forms* on P as

$$\begin{aligned} \mathrm{Sym}_R(P, M) &:= \mathrm{Hom}_{R \otimes R}(P \otimes P, M)^{\mathbb{Z}/2} \\ \mathrm{Quad}_R(P, M) &:= \mathrm{Hom}_{R \otimes R}(P \otimes P, M)_{\mathbb{Z}/2}, \end{aligned}$$

where $(-)^{\mathbb{Z}/2}$ denotes the invariants of the $\mathbb{Z}/2$ -action and $(-)_{\mathbb{Z}/2}$ denotes the coinvariants (i.e. take the coequalizer rather than the equalizer). Finally, let the group of *even forms* on P be

$$\mathrm{Even}_R(P, M) := \mathrm{im}(\mathrm{Nm}: \mathrm{Quad}_R(P, M) \rightarrow \mathrm{Sym}_R(P, M)),$$

where Nm denotes the *norm map* which can be defined on general abelian groups with a $\mathbb{Z}/2$ -action: If $X \in \mathbb{Z}/2\text{-Ab}$ is an abelian group with $\mathbb{Z}/2$ acting via an involution $\tau: X \rightarrow X$,

$$\begin{aligned} \mathrm{Nm}: X_{\mathbb{Z}/2} &\longrightarrow X^{\mathbb{Z}/2} \\ [x] &\longmapsto x + \tau(x). \end{aligned}$$

A symmetric form $q \in \text{Sym}_R(P, M)$ is called *unimodular* if the induced map

$$q_*: P \xrightarrow{\sim} \text{Hom}_R(P, M) =: D_M(P)$$

is an isomorphism. The right-hand side should be thought of the “ M -valued dual” of P . This is reasonable because the assumptions on M are chosen precisely to ensure that $D_M: \text{Proj}(R)^{\text{op}} \xrightarrow{\sim} \text{Proj}(R)$ is an equivalence with inverse D_M^{op} .

Similarly, an even or quadratic form is called *unimodular* if their image in $\text{Sym}_R(P, M)$ (via the inclusion or via Nm) is unimodular. For $r \in \{\text{sym}, \text{quad}, \text{even}\}$ we denote

$$\text{Unimod}^r(P, M) = \left\{ \begin{array}{l} \text{groupoid of unimodular } r\text{-forms, i.e. pairs} \\ (P, q), \text{ where } q \text{ is a unimodular } r\text{-form on } P, \\ \text{and isomorphisms between them} \end{array} \right\}$$

This is a symmetric monoidal 1-groupoid via \oplus , hence induces a symmetric monoidal anima by Example II.11(e).

II.13a. Definition (Karoubi). — For $r \in \{\text{sym}, \text{quad}, \text{even}\}$, the *Grothendieck–Witt anima* of (R, M) is given by

$$\text{gw}^r(R, M) := \text{Unimod}^r(R, M)^{\infty\text{-grp}}.$$

LECTURE 11
3rd Dec, 2020

To make the rather technical definitions of symmetric, quadratic, and even forms a bit clearer, we discussed some examples. Choose R a commutative ring, let $M = R$ and make it an $R \otimes R$ -module via the multiplication map $\mu: R \otimes R \rightarrow R$. Choosing either $\sigma = \text{id}_R$ or $\sigma = -\text{id}_R$, we see that $\text{Sym}_R(P, M)$ returns the usual notion of symmetric or skew-symmetric R -bilinear forms on P .

If $R = \mathbb{C}$, we can also equip $M = \mathbb{C}$ with a $\mathbb{C} \otimes \mathbb{C}$ -module structure via

$$\mathbb{C} \otimes \mathbb{C} \xrightarrow{\text{id} \otimes \overline{(-)}} \mathbb{C} \otimes \mathbb{C} \xrightarrow{\mu} \mathbb{C},$$

where $\overline{(-)}: \mathbb{C} \rightarrow \mathbb{C}$ denotes complex conjugation, as usual. In this case, $\text{Sym}_{\mathbb{C}}(P, \mathbb{C})$ amounts to hermitian or skew-hermitian forms on P , depending on whether $\sigma = \text{id}_{\mathbb{C}}$ or $\sigma = -\text{id}_{\mathbb{C}}$.

II.13b. Exercise. — Show that $\text{Quad}_R(P, M)$ is in bijection with the set of all pairs $(b, q) \in \text{Sym}_R(P, M) \times \text{Hom}_{\text{Set}}(P, M_{\mathbb{Z}/2})$ satisfying

$$b(p, p) = \text{Nm}(q(p)) \quad \text{and} \quad q(rp) = (r \otimes r)q(p)$$

for all $p \in P$ and $r \in R$. Under this bijection, the map $\text{Nm}: \text{Quad}_R(P, M) \rightarrow \text{Sym}_R(P, M)$ simply forgets q .

What Exercise II.13b means depends a little on R . We always assume that R is commutative.

- (a) If $\sigma = \text{id}_M$, then $M_{\mathbb{Z}/2} = M = M^{\mathbb{Z}/2}$ and Nm is simply multiplication by 2. So if M has no 2-torsion, then q is already defined by b and the equation $b(p, p) = \text{Nm}(q(p))$. Hence the norm map $\text{Nm}: \text{Quad}_R(P, M) \rightarrow \text{Sym}_R(P, M)$ is an isomorphism onto its image, which means that quadratic forms and even forms are in bijection in this case.

- (b) If $\sigma = -\text{id}_M$, then $M_{\mathbb{Z}/2} = M/2$ and $M^{\mathbb{Z}/2} = M[2] = 2\text{-torsion of } M$. Hence $\text{Nm}: M_{\mathbb{Z}/2} \rightarrow M^{\mathbb{Z}/2}$ is the zero map. If $M[2] = 0$, then $b(p, p) = 0$ for all $b \in \text{Sym}_R(P, M)$ and all $p \in P$, hence $\text{Nm}: \text{Quad}_R(P, M) \rightarrow \text{Sym}_R(P, M)$ is surjective, because under the identification from Exercise II.13b, $(b, 0)$ is always a preimage of b . Thus all symmetric forms are even, but in general an even form may have lots of quadratic forms that map to it.
- (c) If $2 \in R^\times$ is a unit and $\sigma = \text{id}_M$, then $\text{Nm}: \text{Quad}_R(P, M) \rightarrow \text{Sym}_R(P, M)$ is an isomorphism and the notions of symmetric, quadratic, and even forms all agree.

II.14. Topological K -Theory. — Let's define four ordinary categories as follows:

$$\begin{aligned} \text{Vect}_{\mathbb{R}} &= \left\{ \begin{array}{l} \text{finite-dim. } \mathbb{R}\text{-vector spaces,} \\ \mathbb{R}\text{-linear homeomorphisms} \end{array} \right\}, & \text{Eucl}_{\mathbb{R}} &= \left\{ \begin{array}{l} \text{finite-dim. } \mathbb{R}\text{-vector spaces,} \\ \text{arbitrary homeomorphisms} \end{array} \right\}, \\ \text{Vect}_{\mathbb{C}} &= \left\{ \begin{array}{l} \text{finite-dim. } \mathbb{C}\text{-vector spaces,} \\ \mathbb{C}\text{-linear homeomorphisms} \end{array} \right\}, & \text{Sph}_{\mathbb{R}} &= \left\{ \begin{array}{l} \text{finite-dim. } \mathbb{R}\text{-vector spaces,} \\ \text{proper homotopy equiv's} \end{array} \right\}. \end{aligned}$$

We didn't include $\text{Vect}_{\mathbb{C}}$ in the lecture, but in hindsight we really should have. These four categories can be Kan-enriched: For each category \mathcal{C} among them, we let $\text{F}_{\mathcal{C}}(U, V)_n$ be the set of continuous maps $|\Delta^n| \times U \rightarrow V$ such that $\{d\} \times U \rightarrow V$ is a morphism in \mathcal{C} for all points $d \in |\Delta^n|$ of the topological n -simplex. Moreover, all of them carry a Kan-enriched (symmetric) monoidal structure via the usual product \times . Hence we may apply the construction from Example II.11(f), to obtain monoidal anima

$$\mathcal{V}\text{ect}_{\mathbb{R}}, \quad \mathcal{V}\text{ect}_{\mathbb{C}}, \quad \mathcal{E}\text{ucl}_{\mathbb{R}}, \quad \text{and} \quad \mathcal{S}\text{ph}_{\mathbb{R}}.$$

Observe that while $\text{Vect}_{\mathbb{R}}$, $\text{Vect}_{\mathbb{C}}$, and $\text{Eucl}_{\mathbb{R}}$ are already groupoids, $\text{Sph}_{\mathbb{R}}$ is not. But it becomes an anima after taking the coherent nerve, since homotopy equivalences have homotopy inverses. Also observe that the one-point compactification $(-)^*$ of topological spaces induces equivalences

$$\text{Hom}_{\mathcal{S}\text{ph}_{\mathbb{R}}}(U, V) \simeq \text{Hom}_{\text{core}(*/\text{An})}(U^*, V^*)$$

for all finite-dimensional \mathbb{R} -vector spaces $U, V \in \mathcal{S}\text{ph}_{\mathbb{R}}$. Writing $U = \mathbb{R}^n$, $V = \mathbb{R}^m$, we get that $U^* = \mathbb{S}^n$ and $V^* = \mathbb{S}^m$ are actually spheres. Hence the name $\mathcal{S}\text{ph}_{\mathbb{R}}$.

II.14a. Definition. — Define the following \mathbb{E}_1 -groups:

$$ko = \mathcal{V}\text{ect}_{\mathbb{R}}^{\infty\text{-grp}}, \quad ku = \mathcal{V}\text{ect}_{\mathbb{C}}^{\infty\text{-grp}}, \quad k\text{top} = \mathcal{E}\text{ucl}_{\mathbb{R}}^{\infty\text{-grp}}, \quad \text{and} \quad ksph = \mathcal{S}\text{ph}_{\mathbb{R}}^{\infty\text{-grp}}.$$

Before we indulge ourselves in their properties, I'd like to do two reality checks. The first one is an alternative construction of $\mathcal{V}\text{ect}_{\mathbb{R}}$ and $\mathcal{V}\text{ect}_{\mathbb{C}}$.

II.14b. Lemma*. — Let $O(n) \subseteq \text{GL}_n(\mathbb{R})$ and $U(n) \subseteq \text{GL}_n(\mathbb{C})$ be the n^{th} orthogonal and the n^{th} unitary group. Then

$$\mathcal{V}\text{ect}_{\mathbb{R}} \simeq \coprod_{n \geq 0} BO(n) \quad \text{and} \quad \mathcal{V}\text{ect}_{\mathbb{C}} \simeq \coprod_{n \geq 0} BU(n).$$

The groups $O(n)$, $\text{GL}_n(\mathbb{R})$, $U(n)$ and $\text{GL}_n(\mathbb{C})$ all admit CW structures, hence they define \mathbb{E}_1 -groups via

$$\text{Grp}(\text{CW}) \longrightarrow \text{Grp}(\text{N}^c(\text{CW})) \simeq \text{Grp}(\text{An}).$$

Thus it makes sense to talk of $BO(n)$, $B\text{GL}_n(\mathbb{R})$, $BU(n)$, and $B\text{GL}_n(\mathbb{C})$. We can also find similar descriptions of $\mathcal{E}\text{ucl}_{\mathbb{R}}$ and $\mathcal{S}\text{ph}_{\mathbb{R}}$, but let me not get into that.

Proof of Lemma II.14b*.* It's well-known that $O(n)$ and $U(n)$ are deformation retracts of $GL_n(\mathbb{R})$ and $GL_n(\mathbb{C})$, respectively (see [Hat02, Section 3.D] for example). Hence it suffices to prove stuff for the latter.

Let $\mathcal{V}_{\mathbb{R}}^n \subseteq \mathcal{V}_{\mathbb{R}}$ denote the connected component corresponding to n -dimensional vector spaces. By II.6 and Theorem I.14, constructing an equivalence $BGL_n(\mathbb{R}) \xrightarrow{\sim} \mathcal{V}_{\mathbb{R}}^n$ is the same as constructing an equivalence

$$GL_n(\mathbb{R}) \xrightarrow{\sim} \Omega_{\mathbb{R}^n} \mathcal{V}_{\mathbb{R}}^n \simeq \text{Hom}_{\mathcal{V}_{\mathbb{R}}}(\mathbb{R}^n, \mathbb{R}^n) \simeq F_{\mathcal{V}_{\mathbb{R}}}(\mathbb{R}^n, \mathbb{R}^n).$$

But we easily get $F_{\mathcal{V}_{\mathbb{R}}}(\mathbb{R}^n, \mathbb{R}^n)_i = \text{Hom}_{\text{Top}}(|\Delta^i|, |GL_n(\mathbb{R})|)$ from the definition above (where we write $|GL_n(\mathbb{R})|$ to indicate that we really mean the topological space, not the associated anima), hence $F_{\mathcal{V}_{\mathbb{R}}}(\mathbb{R}^n, \mathbb{R}^n) = \text{Sing } |GL_n(\mathbb{R})|$, hence the unit map

$$GL_n(\mathbb{R}^n) \xrightarrow{\sim} \text{Sing } |GL_n(\mathbb{R})|.$$

provides the desired equivalence. This is also an equivalence of \mathbb{E}_1 -groups since both Sing and $| \cdot |$ preserve finite products, hence the Segal condition. The complex case works completely analogous. \square

II.14c. Lemma*. — *If G is a topological group with the homotopy type of a CW complex, then BG (as constructed in II.6) and BG (the classifying space for principal G -bundles) coincide.*

Proof.* There are probably lots of ways to see this, but what convinced me was that $\Omega BG \simeq G$ also holds on the topological side of things (see [Die08, Example (14.4.7)]), hence both BG s are mapped to G under the equivalence $\Omega: (*/\text{An})_{\geq 1} \xrightarrow{\sim} \text{Grp}(\text{An})$. \square

From Lemmas II.14b, II.14c, and the classification of principal G -bundles we find that

$$\pi_0 \text{Hom}_{\text{An}}(X, \mathcal{V}_{\mathbb{R}}) = \pi_0 \mathcal{V}_{\mathbb{R}}(X) \quad \text{and} \quad \pi_0 \text{Hom}_{\text{An}}(X, \mathcal{V}_{\mathbb{C}}) = \pi_0 \mathcal{V}_{\mathbb{C}}(X)$$

are the sets of isomorphism classes of real and complex vector bundles on X whenever X is a CW complex (it doesn't really make sense to talk about vector bundles over anima). Taking this a step further, we can generalise our constructions from above and produce monoidal anima

$$\mathcal{V}_{\mathbb{R}}(X), \quad \mathcal{V}_{\mathbb{C}}(X), \quad \mathcal{E}\text{ucl}_{\mathbb{R}}(X), \quad \text{and} \quad \mathcal{S}\text{ph}_{\mathbb{R}}(X),$$

of real vector bundles, complex vector bundles, \mathbb{R}^n -fibre bundles, and spherical fibrations over X , respectively. These satisfy (either by construction and the argument above, or simply by definition; it's up to you to decide)

$$\begin{aligned} \mathcal{V}_{\mathbb{R}}(X) &\simeq \text{Hom}_{\text{An}}(X, \mathcal{V}_{\mathbb{R}}), & \mathcal{E}\text{ucl}_{\mathbb{R}}(X) &\simeq \text{Hom}_{\text{An}}(X, \mathcal{E}\text{ucl}_{\mathbb{R}}), \\ \mathcal{V}_{\mathbb{C}}(X) &\simeq \text{Hom}_{\text{An}}(X, \mathcal{V}_{\mathbb{C}}), & \mathcal{S}\text{ph}_{\mathbb{R}}(X) &\simeq \text{Hom}_{\text{An}}(X, \mathcal{S}\text{ph}_{\mathbb{R}}). \end{aligned}$$

Now observe that $\text{Hom}_{\text{An}}(X, -): \text{An} \rightarrow \text{An}$ preserves products, hence Segal conditions. Thus it preserves \mathbb{E}_1 -monoids, and by an easy check also \mathbb{E}_1 -groups (since we only need that $\pi_0 \text{Hom}_{\text{An}}(X, -)$ sends \mathbb{E}_1 -groups to ordinary groups). Hence Definition II.14a and the universal property of group completion induces comparison maps

$$\begin{aligned} \mathcal{V}_{\mathbb{R}}(X)^{\infty\text{-grp}} &\longrightarrow \text{Hom}_{\text{An}}(X, ko), & \mathcal{E}\text{ucl}_{\mathbb{R}}(X)^{\infty\text{-grp}} &\longrightarrow \text{Hom}_{\text{An}}(X, k\text{top}), \\ \mathcal{V}_{\mathbb{C}}(X)^{\infty\text{-grp}} &\longrightarrow \text{Hom}_{\text{An}}(X, ku), & \mathcal{S}\text{ph}_{\mathbb{R}}(X)^{\infty\text{-grp}} &\longrightarrow \text{Hom}_{\text{An}}(X, ks\text{ph}). \end{aligned}$$

Be aware that in general these aren't equivalences in general! This is only true when X is a finite CW complex (or a retract of such), as we will see in Corollary III.15 once we have the group completion theorem available.

Now put $ko^*(X) := \pi_{-*} \text{Hom}_{\text{An}}(X, ko)$ and define $ku^*(X)$, $k\text{top}^*(X)$, and $ksph^*(X)$ similarly. These groups are pretty important invariants in geometric topology! For example, $ko^*(X)$ and $ku^*(X)$ are what's usually called *real* and *complex K-theory*. In the special case $X = *$ they are completely known, thanks to the famous *Bott periodicity theorem* (which we'll see again in a refined version in Theorem III.16):

$$\pi_n(ku) \simeq \begin{cases} \mathbb{Z} & \text{if } n \equiv 0 \pmod{2} \\ 0 & \text{if } n \equiv 1 \pmod{2} \end{cases} \quad \text{and} \quad \pi_n(ko) \simeq \begin{cases} \mathbb{Z} & \text{if } n \equiv 0 \pmod{8} \\ \mathbb{Z}/2 & \text{if } n \equiv 1 \pmod{8} \\ \mathbb{Z}/2 & \text{if } n \equiv 2 \pmod{8} \\ 0 & \text{if } n \equiv 3 \pmod{8} \\ \mathbb{Z} & \text{if } n \equiv 4 \pmod{8} \\ 0 & \text{if } n \equiv 5 \pmod{8} \\ 0 & \text{if } n \equiv 6 \pmod{8} \\ 0 & \text{if } n \equiv 7 \pmod{8} \end{cases}.$$

The homotopy groups of $ksph$ are also “known” in that $\pi_n(ksph) \simeq \pi_n^s$ is the n^{th} stable homotopy group of spheres. Finally, we won't give a description of the homotopy groups of $k\text{top}$, but we can at least their “differences” to those of ko and $ksph$: For $n \neq 4$ we have that

$$\pi_n(\text{fib}(ko \rightarrow k\text{top})) \simeq \Theta^n$$

is the group of exotic spheres in dimension n , i.e. exotic smooth structures on \mathbb{S}^n . Moreover,

$$\pi_n(\text{fib}(k\text{top} \rightarrow ksph)) \simeq L_n^q(\mathbb{Z}) \simeq \begin{cases} \mathbb{Z} & \text{if } n \equiv 0 \pmod{4} \\ 0 & \text{if } n \equiv 1 \pmod{4} \\ \mathbb{Z}/2 & \text{if } n \equiv 2 \pmod{4} \\ 0 & \text{if } n \equiv 3 \pmod{4} \end{cases}.$$

yields Wall's quadratic L -groups of the integers.

\mathbb{E}_∞ -Monoids and \mathbb{E}_∞ -Groups

The “idea” how to impose commutativity is to replace Δ^{op} by some other category \mathbb{I}^{op} which has “flips” in it. Lurie uses Fin_*^{op} to denote this category, but Fabian decided to honour Segal's original notation.

II.15. Definition. — We denote by \mathbb{I}^{op} the 1-category whose objects are finite sets and whose morphisms are partially defined maps. Write $\langle n \rangle = \{1, \dots, n\}$. There is a functor

$$\text{Cut}: \Delta^{\text{op}} \longrightarrow \mathbb{I}^{\text{op}}$$

defined on objects via $\text{Cut}([n]) = \langle n \rangle$. On a morphism $\alpha: [m] \rightarrow [n]$ in Δ , which then corresponds to a morphism α^{op} in Δ^{op} pointing in the other direction, $\text{Cut}(\alpha^{\text{op}}): \langle n \rangle \rightarrow \langle m \rangle$ is defined via

$$\text{Cut}(\alpha^{\text{op}})(i) = \begin{cases} \text{undefined} & \text{if } i \leq \alpha(0) \\ j & \text{if } \alpha(j-1) < i \leq \alpha(j) \\ \text{undefined} & \text{if } \alpha(n) < i \end{cases}.$$

In more invariant words, Cut sends a finite non-empty totally ordered set $I \in \Delta^{\text{op}}$ to its set of *Dedekind cuts* (i.e. partitions into two non-empty intervals), and a map $\alpha: I \rightarrow J$ in Δ is sent to $\text{Cut}(\alpha^{\text{op}}) = \alpha^*: \text{Cut}(J) \rightarrow \text{Cut}(I)$ which is given by taking preimages (whenever these are non-empty, otherwise it's undefined).

If now \mathcal{C} is an ∞ -category with finite products, then a *cartesian commutative monoid* in \mathcal{C} is a functor $X: \mathbb{I}^{\text{op}} \rightarrow \mathcal{C}$ such that $X \circ \text{Cut}: \Delta^{\text{op}} \rightarrow \mathcal{C}$ is a cartesian monoid in the sense of Definition II.1. If it is even a cartesian group, then X is called a *cartesian commutative group* in \mathcal{C} . We let $\text{CMon}(\mathcal{C}), \text{CGrp}(\mathcal{C}) \subseteq \text{Fun}(\mathbb{I}^{\text{op}}, \mathcal{C})$ denote the full sub- ∞ -categories spanned by cartesian commutative monoids/groups.

II.15a. — Time to unwind! The functor $\text{Cut}: \Delta^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$ takes the Segal maps $e_i: [1] \rightarrow [n]$ in Δ to the maps $\rho_i: \langle n \rangle \rightarrow \langle 1 \rangle$ defined by

$$\rho_i(j) = \begin{cases} 1 & \text{if } i = j \\ \text{undefined} & \text{else} \end{cases}.$$

Moreover, the face maps $d_0, d_1, d_2: [1] \rightarrow [2]$ in Δ are sent to $l, m, r: \langle 2 \rangle \rightarrow \langle 1 \rangle$, the unique maps defined on $\{1\}, \{1, 2\}$, and $\{2\}$ respectively. So composition in a cartesian commutative monoid becomes

$$\mu: X_1 \times X_1 \xleftarrow[\sim]{(l,r)} X_2 \xrightarrow{m} X_1.$$

But consider the flip map $\text{flip}: \langle 2 \rangle \rightarrow \langle 2 \rangle$ that swaps 1 and 2. Then $m \circ \text{flip} = m$, $r \circ \text{flip} = l$, and $l \circ \text{flip} = r$ hold in \mathbb{I}^{op} , so the diagram

$$\begin{array}{ccc} X_1 \times X_1 & \xrightarrow{\mu} & X_1 \\ \text{flip factors} \downarrow & \text{///} & \uparrow \\ X_1 \times X_1 & \xrightarrow{\mu} & X_1 \end{array}$$

commutes. So it really makes sense to think of X as commutative!

II.15b. Exercise. — If \mathcal{C} is a 1-category, show that $\text{Cut}^*: \text{Fun}(\mathbb{I}^{\text{op}}, \mathcal{C}) \rightarrow \text{Fun}(\Delta^{\text{op}}, \mathcal{C})$ restricts to a fully faithful functor $\text{CMon}(\mathcal{C}) \rightarrow \text{Mon}(\mathcal{C})$. Note that this is false for general ∞ -categories, even for $\mathcal{C} = \text{Cat}_1^{(2)}$!

II.15c. — Similar to Example II.11(e), any symmetric monoidal 1-category (\mathcal{C}, \otimes) gives rise to an object in $\text{CMon}(\text{Cat}_1^{(2)}) \subseteq \text{CMon}(\text{Cat}_\infty)$. Similar to the monoidal case, this is constructed from a cocartesian fibration $p^\otimes: \mathcal{C}^\otimes \rightarrow \mathbb{I}^{\text{op}}$.

Let's briefly sketch the construction. Objects of \mathcal{C}^\otimes are pairs (n, x_1, \dots, x_n) , where $\langle n \rangle \in \mathbb{I}^{\text{op}}$ and $x_1, \dots, x_n \in \mathcal{C}$. Morphisms $(n, x) \rightarrow (m, y)$ comprise the data of a map $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ in \mathbb{I}^{op} together with maps $f_j: \bigotimes_{i \in \alpha^{-1}(j)} x_i \rightarrow y_j$ for all $j = 1, \dots, m$. Composition is defined in the “obvious” way, using the braiding σ , and to see that composition is associative one has to use $\sigma^2 = \text{id}$. If you are confused by this braiding stuff: In a nutshell, a *braiding* of a monoidal category is a choice of isomorphisms $\sigma_{x,y}: x \otimes y \xrightarrow{\sim} y \otimes x$ subject to some naturality conditions. In this terminology, a symmetric monoidal category is a braided monoidal category for which $\sigma_{x,y} \circ \sigma_{y,x} = \text{id}$.

One can check that the cocartesian fibration $p_{\text{mon}}^\otimes: \mathcal{C}_{\text{mon}}^\otimes \rightarrow \Delta^{\text{op}}$ from Example II.11(e) fits into a pullback diagram

$$\begin{array}{ccc} \mathcal{C}_{\text{mon}}^\otimes & \longrightarrow & \mathcal{C}^\otimes \\ p_{\text{mon}}^\otimes \downarrow & \lrcorner & \downarrow p^\otimes \\ \Delta^{\text{op}} & \xrightarrow{\text{Cut}} & \mathbb{I}^{\text{op}} \end{array}$$

It turns out that the diagram

$$\begin{array}{ccc} \text{CMon}(\text{Cat}_1^{(2)}) & \xrightarrow{\text{Cut}^*} & \text{Mon}(\text{Cat}_1^{(2)}) \\ \downarrow \wr & & \downarrow \wr \\ \text{SymMonCat}_1^{(2)} & \xrightarrow{\text{forget}} & \text{MonCat}_1^{(2)} \end{array}$$

commutes (we never defined the bottom categories, but if you sit down for a while you will surely come up with suitable a definition of the 2-category of symmetric/arbitrary monoidal 1-categories). However, a monoidal category can have many symmetries, so the horizontal functors are not at all fully faithful! For example, let (\mathcal{C}, \otimes) be the category of \mathbb{Z} -graded R -modules equipped with the graded tensor product. Any unit $u \in R^\times$ defines a braiding τ_u on \mathcal{C} via $\tau_u: X \otimes Y \xrightarrow{\sim} Y \otimes X$ given by $\tau_u(x \otimes y) = u^{|x||y|} y \otimes x$ on elementary tensors. If $u^2 = 1$, then this braiding defines a symmetric monoidal structure. In particular, there can be many of these! So commutativity is a *structure*, not a *property*.

LECTURE 12
8th Dec, 2020

II.15d. Semi-Additive and Additive ∞ -Categories. — An ∞ -category is called *semi-additive* if it admits finite products and finite coproducts, its initial and terminal object agree, and for any $x, y \in \mathcal{C}$ the natural map

$$\begin{pmatrix} \text{id}_x & 0 \\ 0 & \text{id}_y \end{pmatrix}: x \sqcup y \xrightarrow{\sim} x \times y$$

is an equivalence. Here $0: x \rightarrow 0 \rightarrow y$ denotes the unique (up to contractible choice) map in $\text{Hom}_{\mathcal{C}}(x, y)$ factoring through an initial/terminal object which we also denote $0 \in \mathcal{C}$. If \mathcal{C} is semi-additive, then we'll usually denote its unified product and coproduct by \oplus .

We call \mathcal{C} *additive* if also the shear map

$$\begin{pmatrix} \text{id}_x & \text{id}_x \\ 0 & \text{id}_x \end{pmatrix}: x \oplus x \xrightarrow{\sim} x \oplus x$$

is an equivalence for all $x \in \mathcal{C}$. The typical source of examples is everything that has to do with abelian groups, so for example $\mathcal{K}(\mathbb{Z})$ and the derived category $\mathcal{D}(\mathbb{Z})$ are additive.

II.16. Proposition. — *If \mathcal{C} is a semi-additive ∞ -category, then the forgetful maps*

$$\text{CMon}(\mathcal{C}) \xrightarrow{\sim} \text{Mon}(\mathcal{C}) \xrightarrow{\sim} \mathcal{C}$$

are equivalences. If \mathcal{C} is additive, then the same holds true for

$$\text{CGrp}(\mathcal{C}) \xrightarrow{\sim} \text{Grp}(\mathcal{C}) \xrightarrow{\sim} \mathcal{C}.$$

Before we get into the proof, let's show a consequence.

II.17. Corollary. — *If \mathcal{C} is an additive ∞ -category, then there is a canonical lift*

$$\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \longrightarrow \mathrm{CGrp}(\mathrm{An}).$$

If \mathcal{C} is only semi-additive, one still gets a lift to $\mathrm{CMon}(\mathrm{An})$.

*Proof**. In general, if $F: \mathcal{C} \rightarrow \mathcal{D}$ is a finite product-preserving functor, then the induced functor $F_*: \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathcal{C}) \rightarrow \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathcal{D})$ preserves the Segal condition and also the condition from Definition II.4, hence it restricts to functors $F_*: \mathrm{CMon}(\mathcal{C}) \rightarrow \mathrm{CMon}(\mathcal{D})$ and $F_*: \mathrm{CGrp}(\mathcal{C}) \rightarrow \mathrm{CGrp}(\mathcal{D})$.

We can't apply this directly to our situation, since $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathrm{An}$ doesn't preserve finite products—or rather it is too good at respecting products, as

$$\mathrm{Hom}_{\mathcal{C}} \left(\bigoplus_{i=1}^n x_i, \bigoplus_{i=1}^n y_i \right) \simeq \prod_{i,j=1}^n \mathrm{Hom}_{\mathcal{C}}(x_i, y_j),$$

of which the diagonal $\prod_{i=1}^n \mathrm{Hom}_{\mathcal{C}}(x_i, y_i)$ is only a factor.

At least $\mathrm{Hom}_{\mathcal{C}}(x, -): \mathcal{C} \rightarrow \mathrm{An}$ preserves products for all $x \in \mathcal{C}$, whence it induces a functor $\mathrm{Hom}_{\mathcal{C}}(x, -)_*: \mathrm{CGrp}(\mathcal{C}) \rightarrow \mathrm{CGrp}(\mathrm{An})$. But if \mathcal{C} is additive, then $\mathrm{CGrp}(\mathcal{C}) \simeq \mathcal{C}$ by Proposition II.16, which provides the desired lift

$$\mathrm{Hom}_{\mathcal{C}}(x, -): \mathcal{C} \longrightarrow \mathrm{CGrp}(\mathrm{An}).$$

To paste these lifts together into a functor $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathrm{CGrp}(\mathrm{An})$, consider the composite

$$\mathrm{Fun}(\mathcal{C}, \mathrm{An}) \longrightarrow \mathrm{Fun}(\mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathcal{C}), \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathrm{An})) \longrightarrow \mathrm{Fun}(\mathrm{CGrp}(\mathcal{C}), \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathrm{An})).$$

If we restrict to the full subcategory $\mathrm{Fun}^{\times}(\mathcal{C}, \mathrm{An}) \subseteq \mathrm{Fun}(\mathcal{C}, \mathrm{An})$ of finite product-preserving functors, it lands in $\mathrm{Fun}(\mathrm{CGrp}(\mathcal{C}), \mathrm{CGrp}(\mathrm{An}))$, as argued above. Moreover, if \mathcal{C} is additive, then $\mathrm{CGrp}(\mathcal{C}) \simeq \mathcal{C}$ by Proposition II.16. Now $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Fun}(\mathcal{C}, \mathrm{An})$ has image in $\mathrm{Fun}^{\times}(\mathcal{C}, \mathrm{An})$, hence we may compose it with the above to obtain the desired functor

$$\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \longrightarrow \mathrm{Fun}(\mathcal{C}, \mathrm{CGrp}(\mathrm{An})).$$

If \mathcal{C} is only semi-additive, then the argument still works with $\mathrm{CMon}(-)$ instead of $\mathrm{CGrp}(-)$ everywhere. Moreover, one can check (but we won't do that here) that if we use the (semi-)additive structure on $\mathcal{C}^{\mathrm{op}}$ instead to do the constructions above, we still end up with the same functor $\mathrm{Hom}_{\mathcal{C}}: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathrm{CGrp}(\mathrm{An})$. \square

Proof sketch of Proposition II.16. The key to the proof is the following claim:

(\boxtimes) *Let $\mathcal{I} \in \{\Delta^{\mathrm{op}}, \mathbb{I}^{\mathrm{op}}\}$. Given a functor $X: \mathcal{I} \rightarrow \mathcal{C}$ satisfying $X_0 \simeq 0$, then X satisfies the Segal condition iff it is left-Kan extended from $X|_{\mathcal{I}_{\leq 1}}$ along $p: \mathcal{I}_{\leq 1} \rightarrow \mathcal{I}$.*

To conclude the statement of the proposition from (\boxtimes), write $\mathrm{Fun}(\mathcal{I}_{\leq 1}, \mathcal{C})_0 \subseteq \mathrm{Fun}(\mathcal{I}_{\leq 1}, \mathcal{C})$ for the full subcategory of functors F satisfying $F(0) \simeq 0$. Since $0 \in \mathcal{C}$ is initial and terminal, one easily checks that $\mathcal{C} \xrightarrow{\sim} \mathrm{Fun}(\mathcal{I}_{\leq 1}, \mathcal{C})_0$, sending $x \in \mathcal{C}$ to the unique (up to contractible choice) functor F_x satisfying $F_x(0) \simeq 0$ and $F_x(1) \simeq x$, is an equivalence. Moreover,

$$\mathrm{Fun}(\mathcal{I}_{\leq 1}, \mathcal{C})_0 \subseteq \mathrm{Fun}(\mathcal{I}_{\leq 1}, \mathcal{C}) \xrightarrow{\mathrm{Lan}_p} \mathrm{Fun}(\mathcal{I}, \mathcal{C})$$

is fully faithful by Corollary I.54, and (\boxtimes) precisely says that its essential image is $\text{Mon}(\mathcal{C})$ for $\mathcal{I} = \Delta^{\text{op}}$ or $\text{CMon}(\mathcal{C})$ for $\mathcal{I} = \mathbb{T}^{\text{op}}$. This shows $\text{Mon}(\mathcal{C}) \simeq \mathcal{C} \simeq \text{CMon}(\mathcal{C})$, as required. If \mathcal{C} is additive, then the image of $x \in \mathcal{C}$ under the above composite is automatically a cartesian group: Indeed, after an easy unravelling, the condition from Definition II.4 translates into the condition that the shear map on x be an equivalence, which is guaranteed by \mathcal{C} being additive.


To prove (\boxtimes) , first note that we get a natural counit transformation $\text{Lan}_p X|_{\mathcal{I}_{\leq 1}} \Rightarrow X$ for free, so whether this is an equivalence can be checked on objects. Recall from Theorem I.52 that

$$\text{Lan}_p X|_{\mathcal{I}_{\leq 1}}(n) \simeq \text{colim}_{i \in \mathcal{I}_{\leq 1}/n} X_i$$

(it will follow from the discussion below that these colimits exist in \mathcal{C} , so the theorem is indeed applicable). Therefore we have to understand the slice categories


$$\mathbb{T}_{\leq 1}^{\text{op}}/\langle n \rangle \quad \text{and} \quad \Delta_{\leq 1}^{\text{op}}/[n] \simeq ([n]/\Delta_{\leq 1})^{\text{op}}.$$

The slice category $[n]/\Delta_{\leq 1}$. Let's describe $[n]/\Delta_{\leq 1}$ first in words and then in a picture (whichever you start with will probably confuse you, and then the other thing will hopefully unconfuse you). Its vertices are given by maps $[n] \rightarrow [1]$ and $[n] \rightarrow [0]$ in Δ . There's only one of the latter kind and it is denoted 0 (in the picture it's depicted as a black dot on the right). There are $n+2$ maps $[n] \rightarrow [1]$: One that is constantly 0, one that is constantly 1 (depicted on the top and on the bottom), and n surjective maps (depicted on the left). Moreover, there's a bunch of non-degenerate edges. The pink ones are those induced by $0: [0] \rightarrow [1]$ and $0: [1] \rightarrow [1]$, the purple ones are those induced by $1: [0] \rightarrow [1]$ and $1: [1] \rightarrow [1]$, and finally the black ones are induced by the unique morphism $[1] \rightarrow [0]$.



However, the punchline is that all of this doesn't really matter, since the dotted part, consisting of the vertices on the left and 0 on the right, turns out to be final (in the lecture we claimed that already the vertices on the left are final, but I think that's not true). To prove this, verify the criterion from Theorem I.43(b) and use that the slice categories in question (in fact, slice categories between slice categories) contain the respective arrow starting at 0 as a terminal object. After passing to $\Delta_{\leq 1}^{\text{op}}/[n]$ instead of $[n]/\Delta_{\leq 1}$, the dotted part becomes cofinal, hence the colimit in question can be taken over the dotted part only. But the 0-vertex on the right is mapped to $X_0 \simeq 0$ and there are no edges between the vertices on the left, hence it's easy to see that the colimit over the dotted part is just $X_1^{\oplus n}$. This is precisely what we wanted to show!

The slice category $\mathbb{I}_{\leq 1}^{\text{op}}/\langle n \rangle$. The vertices of $\mathbb{I}_{\leq 1}^{\text{op}}/\langle n \rangle$ consist of partially defined maps $\langle 0 \rangle \rightarrow \langle n \rangle$ and $\langle 1 \rangle \rightarrow \langle n \rangle$. As $\langle 0 \rangle = \emptyset$, there is only one such map of the first kind, depicted by the vertex on the right in the picture below. The vertex in the middle represents the unique nowhere-defined map $\langle 1 \rangle \rightarrow \langle n \rangle$, and the vertices on the left correspond to the maps $\langle 1 \rangle \rightarrow \langle n \rangle$ which are actually defined on 1.



Again, the dotted part (i.e. everything except the vertex in the middle) is cofinal and the colimit over it evaluates to $X_1^{\oplus n}$ since the vertex on the right is sent to $X_0 \simeq 0$. \square

Since $\mathcal{D}_{\geq 0}(\mathbb{Z})$ is additive, a simple consequence of Proposition II.16 is that the Eilenberg–MacLane functor from Very Long Example I.56 lifts to a functor

$$K: \mathcal{D}_{\geq 0}(\mathbb{Z}) \longrightarrow \text{CGrp}(\text{An}).$$

Indeed, K is a right-adjoint by construction, hence it preserves products, so we can use the argument from the proof of Corollary II.17 to produce the desired lift. We will soon (in Example II.25(c)) extend this even further!

So Eilenberg–MacLane anima give rise to \mathbb{E}_∞ -groups. Another source of examples for \mathbb{E}_∞ -monoids/groups is

$$\text{CMon}(\text{Kan}) \longrightarrow \text{CMon}(\text{An}) \quad \text{and} \quad \text{CGrp}(\text{Kan}) \longrightarrow \text{CGrp}(\text{An}),$$

(again note that $\text{Kan} \rightarrow \text{An}$ preserves products), but these are essentially the same examples: It is a theorem of Dold–Thom, that a connected commutative monoid in Kan complexes has underlying anima a *gem* (“generalized Eilenberg–MacLane space”; we defined these things in Very Long Example I.56). The same then holds for not necessarily connected commutative groups in Kan complexes, since all connected components are equivalent.

II.18. Proposition. — *The realisation functor $|\cdot|: \text{sAn} \rightarrow \text{An}$ preserves finite products.*

Proof. We compute

$$\begin{aligned} |M \times N| &\simeq \text{colim}_{[n] \in \Delta^{\text{op}}} (M_n \times N_n) \simeq \text{colim}_{([n], [m]) \in \Delta^{\text{op}} \times \Delta^{\text{op}}} (M_n \times N_m) \\ &\simeq \text{colim}_{[n] \in \Delta^{\text{op}}} \left(M_n \times \text{colim}_{[m] \in \Delta^{\text{op}}} N_m \right) \\ &\simeq \left(\text{colim}_{[n] \in \Delta^{\text{op}}} M_n \right) \times \left(\text{colim}_{[m] \in \Delta^{\text{op}}} N_m \right) \\ &\simeq |M| \times |N|. \end{aligned}$$

For the second equivalence see Exercise II.18a below. The third equivalence follows from Proposition I.42 together with the fact that $M_n \times -$ commutes with arbitrary colimits. Indeed, for any $X \in \text{An}$ the functor $X \times -: \text{An} \rightarrow \text{An}$ is a left adjoint of $\text{Hom}_{\text{An}}(X, -)$. Applying the same argument to $- \times \text{colim}_{\Delta^{\text{op}}} N_m$ gives the fourth equivalence. \square

II.18a. Exercise. — Show that the diagonal $\Delta: \Delta^{\text{op}} \rightarrow \Delta^{\text{op}} \times \Delta^{\text{op}}$ is cofinal as a map of ∞ -categories. In other words, Δ^{op} is *sifted*.

*Proof**. By Theorem I.43(b) we need to check that $\Delta^{\text{op}} \times_{\Delta^{\text{op}} \times \Delta^{\text{op}}} ([m], [n]) / (\Delta^{\text{op}} \times \Delta^{\text{op}})$ is weakly contractible for all $[m], [n] \in \Delta^{\text{op}}$. Note that we may form the slice category as a 1-category, since the nerve functor commutes with limits. Also note that we may replace the slice category by its opposite $\mathcal{S}_{m,n} := \Delta \times_{\Delta \times \Delta} (\Delta \times \Delta) / ([m], [n])$ for convenience.

Unravelling definitions, we find that objects in $\mathcal{S}_{m,n}$ can be thought of as maps of posets $\alpha: [i] \rightarrow [m] \times [n]$. Any such map factors uniquely into a composition $[i] \rightarrow [j] \hookrightarrow [m] \times [n]$ of a surjective and an injective map. This defines a functor $r: \mathcal{S}_{m,n} \rightarrow \mathcal{C}_{m,n}$ into the category $\mathcal{C}_{m,n}$ of non-empty linearly ordered subposets of $[m] \times [n]$ ($\mathcal{C}_{m,n}$ is partially ordered by inclusion). But conversely, any such subposet is given by an injective map $[j] \hookrightarrow [m] \times [n]$ for some $j \leq m + n + 1$, hence we get an inclusion $s: \mathcal{C}_{m,n} \rightarrow \mathcal{S}_{m,n}$ in the reverse direction. Clearly $r \circ s = \text{id}_{\mathcal{C}_{m,n}}$, whereas the surjections $[i] \twoheadrightarrow [j]$ define a natural transformation $\text{id}_{\mathcal{S}_{m,n}} \Rightarrow s \circ r$.

This shows that $\mathcal{S}_{m,n}$ and $\mathcal{C}_{m,n}$ are weakly (and even strongly) homotopy equivalent after taking nerves. Now if we recall [HC_I, Definition V.4.7], we recognize $\mathcal{C}_{m,n}$ as the barycentric subdivision $\text{sd}(\Delta^m \times \Delta^n)$, which is weakly contractible because $\Delta^m \times \Delta^n$ is. \square

II.18b. — Recall from Corollary* II.8a that we have an adjunction

$$B: \text{Mon}(\text{An}) \rightleftarrows * / \text{An} : \Omega.$$

Now $B = | \cdot |$ and Ω both preserve finite products—the former by Proposition II.18, the latter since it is a right-adjoint. So both functors preserve the Segal condition. Therefore, if we apply $\text{Fun}(\mathbb{I}^{\text{op}}, -)$ to both sides of the adjunction above, then the resulting adjunction restricts to another adjunction

$$B: \text{CMon}(\text{Mon}(\text{An})) \rightleftarrows \text{CMon}(* / \text{An}) : \Omega \quad (\text{II.18b.1})$$

Now it is easy to see that the forgetful map $\text{CMon}(* / \mathcal{C}) \rightarrow \text{CMon}(\mathcal{C})$ is an equivalence for any ∞ -category \mathcal{C} with finite products (and thus a terminal object $* \in \mathcal{C}$). We claim that also

$$\begin{array}{ccc} \text{CMon}(\text{Mon}(\mathcal{C})) & \xrightarrow{\text{forget}_*} & \text{CMon}(\mathcal{C}) \\ \downarrow \wr & \text{///} & \uparrow \sim \\ \text{Mon}(\text{CMon}(\mathcal{C})) & \xrightarrow{\text{forget}} & \end{array} \quad (\text{II.18b.2})$$

commutes (the forgetful functors are given by evaluation in degree 1 of course). Once we have this, we will obtain the following statement (which comes only later in Fabian’s notes, hence the gap in numbering).

II.21. Corollary. — *There is a commutative diagram of horizontal adjunctions*

$$\begin{array}{ccc} \text{CMon}(\text{An}) & \xrightleftharpoons[\Omega]{B} & \text{CMon}(\text{An}) \\ \text{Cut}^* \downarrow & & \downarrow \text{ev}_1 \\ \text{Mon}(\text{An}) & \xrightleftharpoons[\Omega]{B} & * / \text{An} \end{array}$$

Moreover, we have the following analogues of II.6 and Proposition II.8:

- (a) Restricting B to the full subcategory $\mathrm{CGrp}(\mathrm{An}) \subseteq \mathrm{CMon}(\mathrm{An})$ gives a fully faithful functor $B: \mathrm{CGrp}(\mathrm{An}) \rightarrow \mathrm{CMon}(\mathrm{An})$.
- (b) Both functors $B, \Omega: \mathrm{CMon}(\mathrm{An}) \rightarrow \mathrm{CMon}(\mathrm{An})$ actually take values in $\mathrm{CGrp}(\mathrm{An})$.
- (c) $\Omega B: \mathrm{CMon}(\mathrm{An}) \rightarrow \mathrm{CGrp}(\mathrm{An})$ is left-adjoint to $\mathrm{CGrp}(\mathrm{An}) \subseteq \mathrm{CMon}(\mathrm{An})$.

*Proof**. We construct the diagram and prove commutativity first. The bottom row is already known. After identifying $\mathrm{CMon}(*/\mathrm{An}) \simeq \mathrm{CMon}(\mathrm{An})$ and $\mathrm{CMon}(\mathrm{An}) \simeq \mathrm{CMon}(\mathrm{Mon}(\mathrm{An}))$, the upper row becomes (II.18b.1). To see that the diagram commutes, we first check that these identifications transform the forgetful functor $\mathrm{CMon}(\mathrm{Mon}(\mathrm{An})) \rightarrow \mathrm{Mon}(\mathrm{An})$ into $\mathrm{Cut}^*: \mathrm{CMon}(\mathrm{An}) \rightarrow \mathrm{Mon}(\mathrm{An})$. Indeed, this follows easily from (II.18b.2). Now both vertical arrows are of the form $\mathrm{forget}: \mathrm{CMon}(-) \rightarrow (-)$, from which commutativity of the diagram is clear.

Next we check (b). Let $M \in \mathrm{CMon}(\mathrm{An})$. To check $\Omega M \in \mathrm{CGrp}(\mathrm{An})$, it suffices to check $\mathrm{Cut}^* \Omega M \in \mathrm{Grp}(\mathrm{An})$ (that's just how \mathbb{E}_∞ -groups are defined). But $\mathrm{Cut}^* \circ \Omega \simeq \Omega \circ \mathrm{ev}_1$ and the right-hand side takes values in $\mathrm{Grp}(\mathrm{An})$ by II.6. To show $BM \in \mathrm{CGrp}(\mathrm{An})$, it similarly suffices to check that the ordinary monoid $\pi_0(BM)_1$ is an ordinary group, as argued in the proof of Proposition II.5. But in fact $\pi_0(BM)_1 \simeq *$, since $B: \mathrm{Mon}(\mathrm{An}) \rightarrow */\mathrm{An}$ takes values in $(*/\mathrm{An})_{\geq 1}$ and the diagram above commutes. This shows (b).

In particular, the top row adjunction restricts to an adjunction

$$B: \mathrm{CGrp}(\mathrm{An}) \rightleftarrows \mathrm{CGrp}(\mathrm{An}) : \Omega.$$

To show (a), it suffices therefore to check that the unit $\mathrm{id} \Rightarrow \Omega B$ is an equivalence. Equivalences of \mathbb{E}_∞ -groups can be detected on underlying \mathbb{E}_1 -groups (in fact, even on underlying anima, i.e. after applying ev_1 , since the Segal condition will do the rest). So it suffices that $\mathrm{Cut}^* \Rightarrow \mathrm{Cut}^* \Omega B$ is an equivalence. But $\mathrm{Cut}^* \Omega B \simeq \Omega \mathrm{ev}_1 B \simeq \Omega B \mathrm{Cut}^*$, and if G is an \mathbb{E}_∞ -group, then

$$\Omega B \mathrm{Cut}^* G \simeq (\mathrm{Cut}^* G)^{\infty\text{-grp}} \simeq \mathrm{Cut}^* G$$

by Proposition II.8. This finishes the proof of (a).

Let $\mathcal{C} \subseteq \mathrm{CGrp}(\mathrm{An})$ denote the essential image of $B|_{\mathrm{CGrp}(\mathrm{An})}$. Then $B: \mathrm{CGrp}(\mathrm{An}) \xrightarrow{\sim} \mathcal{C}$ is an equivalence, hence its right adjoint $\Omega: \mathcal{C} \rightarrow \mathrm{CGrp}(\mathrm{An})$ is an equivalence too. For $M \in \mathrm{CMon}(\mathrm{An})$ and $G \in \mathrm{CGrp}(\mathrm{An})$ we may thus compute

$$\begin{aligned} \mathrm{Hom}_{\mathrm{CGrp}(\mathrm{An})}(\Omega BM, G) &\simeq \mathrm{Hom}_{\mathrm{CGrp}(\mathrm{An})}(\Omega BM, \Omega BG) \\ &\simeq \mathrm{Hom}_{\mathcal{C}}(BM, BG) \\ &\simeq \mathrm{Hom}_{\mathrm{CMon}(\mathrm{An})}(M, \Omega BG) \\ &\simeq \mathrm{Hom}_{\mathrm{CMon}(\mathrm{An})}(M, G), \end{aligned}$$

where we also used $G \simeq \Omega BG$ twice. This shows (c). \square

II.21a. Example. — Some consequences:

- (a) Corollary II.21(c) shows that group completion in $\mathrm{Mon}(\mathrm{An})$ and $\mathrm{CMon}(\mathrm{An})$ is compatible. Hence all examples from II.12 to II.14, i.e.

$$k(R), \quad \mathrm{gw}^r(R), \quad ko, \quad ku, \quad k\mathrm{top}, \quad \text{and} \quad k\mathrm{sph},$$

are canonically \mathbb{E}_∞ -groups. Indeed, they arise as group completions of certain \mathbb{E}_1 -monoids, all of which have canonical \mathbb{E}_∞ -monoid refinements by II.15c.

- (b) It follows from Corollary II.21(a) that $\Omega BX \simeq X$ for all $X \in \mathbf{CGrp}(\mathbf{An})$. Since it's well-known that Ω shifts homotopy groups down, we see that B shifts homotopy groups up. That is, if $e \in X_1$ denotes the identity element, then

$$\pi_i((BX)_1) \simeq \begin{cases} \pi_{i-1}(X_1, e) & \text{if } i > 0 \\ * & \text{if } i = 0 \end{cases}$$

(in particular, we don't need to specify a basepoint for $(BX)_1$ since it is connected).

- (c) We get a commutative solid diagram

$$\begin{array}{ccc} \mathcal{D}_{\geq 0}(\mathbb{Z}) & \xrightarrow{K} & \mathrm{CGrp}(\mathrm{An}) \\ [1]=\Sigma \downarrow \hat{\Omega} & \text{///} & B \downarrow \hat{\Omega} \\ \mathcal{D}_{\geq 0}(\mathbb{Z}) & \xrightarrow{K} & \mathrm{CGrp}(\mathrm{An}) \end{array}$$

in Cat_∞ . So B preserves Eilenberg–MacLane anima and only shifts the homotopy groups up by 1.

To see commutativity, first note that the diagram with the dotted vertical arrows commutes (where $\Omega = (\tau_{\geq 1})[-1] : \mathcal{D}_{\geq 0}(\mathbb{Z}) \rightarrow \mathcal{D}_{\geq 0}(\mathbb{Z})$ is defined as in Philosophical Nonsense I.18). Indeed, Ω is given by a pullback which is preserved by the right-adjoint functor K (being a right adjoint is still true for the lift $K : \mathcal{D}_{\geq 0}(\mathbb{Z}) \rightarrow \mathbf{CGrp}(\mathbf{An})$, simply by construction and our Observation I.40 that adjunctions extend to functor categories). By abstract nonsense, commutativity of the dotted diagram induces a natural transformation $BK \Rightarrow K(-[1])$. Whether this is an equivalence can be checked on objects $C \in \mathcal{D}_{\geq 0}(\mathbb{Z})$. But whether $B(KC) \rightarrow K(C[1])$ is an equivalence can in turn be checked on homotopy groups of underlying anima. But from (b) we see that homotopy groups of both $B(KC)$ and $K(C[1])$ are those of KC , shifted up by one.

Now towards the proof of (II.18b.2). Let's denote by $\text{Cat}_\infty^\times \subseteq \text{Cat}_\infty$ the (non-full) sub- ∞ -category spanned by ∞ -categories with finite products and functors preserving these. Let furthermore $\text{Cat}_\infty^{\text{add}} \subseteq \text{Cat}_\infty^{\text{semi-add}} \subseteq \text{Cat}_\infty^\times$ denote the full sub- ∞ -categories spanned by the additive and semi-additive ∞ -categories respectively.

II.19. Theorem. — *If an ∞ -category \mathcal{C} has products, then $\mathbf{CMon}(\mathcal{C})$ and $\mathbf{CGrp}(\mathcal{C})$ are semi-additive and additive, respectively. Furthermore, the functors*

$$\begin{aligned} \mathbf{CMon} &: \mathbf{Cat}_\infty^\times \longrightarrow \mathbf{Cat}_\infty^{\text{semi-add}} \\ \mathbf{CGrp} &: \mathbf{Cat}_\infty^\times \longrightarrow \mathbf{Cat}_\infty^{\text{add}} \end{aligned}$$

are right-adjoint to the inclusions $\text{Cat}_\infty^{\text{semi-add}} \subseteq \text{Cat}_\infty^\times$ and $\text{Cat}_\infty^{\text{add}} \subseteq \text{Cat}_\infty^\times$ respectively.

Using Theorem II.19 we get an easy proof of (II.18b.2). First note that the two maps

$$\begin{array}{ccc} & \xrightarrow[\sim]{\text{forget}_*} & \\ \text{CMon}(\text{CMon}(\mathcal{C})) & & \text{CMon}(\mathcal{C}) \\ & \xrightarrow[\text{forget}]{\sim} & \end{array}$$

agree and are equivalences. Indeed, $\text{CMon}(-) \rightarrow (-)$ is the counit of the adjunction from Theorem II.19, hence an equivalence since the left adjoint $\text{Cat}_{\infty}^{\text{semi-add}} \subseteq \text{Cat}_{\infty}^{\times}$ is fully faithful.

This shows that both functors are equivalences. Moreover, by writing down the triangle identities we find that both maps are inverses (once from the right, once from the left) to the unit $\mathrm{CMon}(\mathcal{C}) \rightarrow \mathrm{CMon}(\mathrm{CMon}(\mathcal{C}))$, hence they must agree. This almost shows (II.18b.2), except that we want to replace one CMon by a Mon . To do this, note that Proposition II.16 implies $\mathrm{CMon}(\mathrm{CMon}(\mathcal{C})) \simeq \mathrm{Mon}(\mathrm{CMon}(\mathcal{C}))$. Moreover $\mathrm{CMon}(\mathrm{Mon}(\mathcal{C})) \simeq \mathrm{Mon}(\mathrm{CMon}(\mathcal{C}))$ follows by straightforward inspection, as both sides can be interpreted as full subcategories of $\mathrm{Fun}(\Delta^{\mathrm{op}} \times \mathbb{I}^{\mathrm{op}}, \mathcal{C})$. Hence we are done.

Note that our argument can not be extended to show that the two forgetful functors $\mathrm{Mon}(\mathrm{Mon}(\mathcal{C})) \rightrightarrows \mathrm{Mon}(\mathcal{C})$ agree. This is still true, but much harder to show. Moreover, if \mathcal{C} is a 1-category, then $\mathrm{Mon}(\mathrm{Mon}(\mathcal{C})) \simeq \mathrm{CMon}(\mathcal{C})$, a fact that is usually called the *Eckmann–Hilton trick*. This is false for ∞ -categories! What is true instead is

$$\mathrm{CMon}(\mathcal{C}) \simeq \lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Mon}^{(n)}(\mathcal{C}),$$

where $\mathrm{Mon}^{(n)}(-) = \mathrm{Mon}(\mathrm{Mon}(\dots(\mathrm{Mon}(-))\dots))$ is the n -fold iteration of $\mathrm{Mon}(-)$. The transition morphisms $\mathrm{Mon}^{(n)}(\mathcal{C}) \rightarrow \mathrm{Mon}^{(n-1)}(\mathcal{C})$ are given by forgetting one Mon . There are n choices to do so, but it turns out that they all agree, generalizing our claim about $\mathrm{Mon}(\mathrm{Mon}(\mathcal{C})) \rightrightarrows \mathrm{Mon}(\mathcal{C})$.

For the proof of Theorem II.19, we need a criterion for semi-additivity.

II.20. Lemma (compare [HA, Proposition 2.4.3.19], which is stronger). — *Let \mathcal{C} be a category with finite products. Then \mathcal{C} is semi-additive if the following conditions hold:*

- (a) *The terminal object $*$ $\in \mathcal{C}$ also is initial.*
- (b) *We have a natural transformation $\mu: (\Delta: x \mapsto x \times x) \Rightarrow \mathrm{id}_{\mathcal{C}}$ such that both compositions*

$$\begin{aligned} x &\simeq x \times * \xrightarrow{\mathrm{id}_x \times 0} x \times x \xrightarrow{\mu_x} x \\ x &\simeq * \times x \xrightarrow{0 \times \mathrm{id}_x} x \times x \xrightarrow{\mu_x} x \end{aligned}$$

are homotopic to id_x for all $x \in \mathcal{C}$, and the diagram

$$\begin{array}{ccc} (x \times x) \times (y \times y) & \xrightarrow{\mathrm{id}_x \times \mathrm{flip} \times \mathrm{id}_y} & (x \times y) \times (x \times y) \\ & \searrow \mu_x \times \mu_y & \swarrow \mu_x \times \mu_y \\ & x \times y & \end{array}$$

commutes up to homotopy for all $x, y \in \mathcal{C}$.

II.20a. Remark*. — Lurie only requires that the first of the two compositions in Lemma II.20(b) is homotopic to id_x , and Fabian may or may not have done the same in the lecture. However, the second one doesn't follow from the first, and we'll really need both to be homotopic to id_x in the proof. In fact, if only the first composition were to be homotopic to id_x , we could take μ_x to be $\mathrm{pr}_1: x \times x \rightarrow x$ for all $x \in \mathcal{C}$ (it's clear that this also makes the diagram above commute), and then we could show that every ∞ -categories with finite products and a zero object is semi-additive. This can't be true!

Proof of Lemma II.20. Let $c, d \in \mathcal{C}$. We want to show that the maps

$$c \simeq c \times * \xrightarrow{\mathrm{id}_c \times 0} c \times d \quad \text{and} \quad d \simeq * \times d \xrightarrow{0 \times \mathrm{id}_d} c \times d$$

exhibit $c \times d$ as a coproduct of c and d . That is, we need to show that the natural transformation

$$\varphi: \text{Hom}_{\mathcal{C}}(c \times d, -) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(c, -) \times \text{Hom}_{\mathcal{C}}(d, -)$$

induced by these maps is an equivalence. This can be checked on objects. Given $a \in \mathcal{C}$, we construct an inverse ψ_a to φ_a as follows:

$$\psi_a: \text{Hom}_{\mathcal{C}}(c, a) \times \text{Hom}_{\mathcal{C}}(d, a) \xrightarrow{\times} \text{Hom}_{\mathcal{C}}(c \times d, a \times a) \xrightarrow{\mu_a} \text{Hom}_{\mathcal{C}}(c \times d, a)$$

Note that we do not claim that the ψ_a combine into a natural transformation ψ which is inverse to φ , only that ψ_a is inverse to φ_a , since it suffices to have φ a pointwise equivalence. Let's prove $\varphi_a \circ \psi_a \simeq \text{id}$ first. This follows essentially from the first condition in (b), but since this sparked some confusion in the lecture, I'll try to give a much more detailed proof than in the lecture and be as precise as possible (hopefully this doesn't make things worse). We begin by drawing a tiny diagram:

$$\begin{array}{ccccc}
 & & \text{Hom}_{\mathcal{C}}(c, a) \times \text{Hom}_{\mathcal{C}}(d, a) & & \\
 & \swarrow \times & \downarrow \wr & \searrow \wr \text{id} & \\
 & & \text{Hom}_{\mathcal{C}}(c \times *, a \times *) \times \text{Hom}_{\mathcal{C}}(* \times d, * \times a) & & \\
 & & \text{(I)} & & \text{(III)} \\
 & & \downarrow & & \\
 \text{Hom}_{\mathcal{C}}(c \times d, a \times a) & \rightarrow & \text{Hom}_{\mathcal{C}}(c \times *, a \times a) \times \text{Hom}_{\mathcal{C}}(* \times d, a \times a) & & \\
 \downarrow \mu_a & & \downarrow \mu_a \times \mu_a & & \\
 \text{Hom}_{\mathcal{C}}(c \times d, a) & \longrightarrow & \text{Hom}_{\mathcal{C}}(c \times *, a) \times \text{Hom}_{\mathcal{C}}(* \times d, a) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{C}}(c, a) \times \text{Hom}_{\mathcal{C}}(d, a)
 \end{array}$$

If we can show that this diagram commutes up to homotopy, then walking around its perimeter will show $\varphi_a \circ \psi_a \simeq \text{id}$. The cell labelled (III) commutes up to homotopy because the composition of the three downward pointing arrows and the rightward pointing arrow is induced by $\mu_a \circ (\text{id}_a \times 0): a \simeq a \times * \rightarrow a \times a \rightarrow a$ in the first factor and by $\mu_a \circ (0 \times \text{id}_a): a \simeq * \times a \rightarrow a \times a \rightarrow a$ in the second factor, both of which are homotopic to id_a by assumption.

To see that the cell labelled (II) commutes up to homotopy, it suffices to check this after composition with $\text{pr}_1: \text{Hom}_{\mathcal{C}}(c \times *, a) \times \text{Hom}_{\mathcal{C}}(* \times d, a) \rightarrow \text{Hom}_{\mathcal{C}}(c \times *, a)$ and after composition with $\text{pr}_2: \text{Hom}_{\mathcal{C}}(c \times *, a) \times \text{Hom}_{\mathcal{C}}(* \times d, a) \rightarrow \text{Hom}_{\mathcal{C}}(* \times d, a)$. We only give the argument for pr_1 , since the argument for pr_2 will be analogous. After composition with pr_1 , we end up with two maps $\text{Hom}_{\mathcal{C}}(c \times d, a \times a) \rightarrow \text{Hom}_{\mathcal{C}}(c \times *, a)$. Both are given by precomposition with $c \times * \rightarrow c \times d$ and postcomposition with $\mu_a: a \times a \rightarrow a$, but in different orders. However, it doesn't matter (up to homotopy) in which order we postcompose and precompose, which proves commutativity of (II).

Finally, to see that (I) commutes, we look at the diagram (and its obvious analogue where $c \times *$ is replaced by $* \times d$)

$$\begin{array}{ccc}
 \text{Hom}_{\mathcal{C}}(c, a) \times \text{Hom}_{\mathcal{C}}(d, a) & \xrightarrow{\times} & \text{Hom}_{\mathcal{C}}(c \times d, a \times a) \\
 \downarrow & & \downarrow \\
 \text{Hom}_{\mathcal{C}}(c, a) \times \text{Hom}_{\mathcal{C}}(*, a) & \xrightarrow{\times} & \text{Hom}_{\mathcal{C}}(c \times *, a \times a) \\
 \wr \uparrow & & \uparrow \\
 \text{Hom}_{\mathcal{C}}(c, a) \times \text{Hom}_{\mathcal{C}}(*, *) & \xrightarrow{\times} & \text{Hom}_{\mathcal{C}}(c \times *, a \times *)
 \end{array}$$

This commutes “by naturality of the \times -construction on morphism anima”. The bottom left vertical arrow is an equivalence since $*$ is initial, so $\mathrm{Hom}_{\mathcal{C}}(*, -) \simeq *$. In particular, this shows $\mathrm{Hom}_{\mathcal{C}}(c, a) \times \mathrm{Hom}_{\mathcal{C}}(*, *) \simeq \mathrm{Hom}_{\mathcal{C}}(c, a)$. Under this identification, the bottom row becomes the morphism $\mathrm{Hom}_{\mathcal{C}}(c, a) \rightarrow \mathrm{Hom}_{\mathcal{C}}(c \times *, a \times *)$ from (I) and the two left vertical arrows compose to the projection $\mathrm{pr}_1: \mathrm{Hom}_{\mathcal{C}}(c, a) \times \mathrm{Hom}_{\mathcal{C}}(d, a) \rightarrow \mathrm{Hom}_{\mathcal{C}}(c, a)$. This shows that (I) commutes indeed and therefore that $\varphi_a \circ \psi_a \simeq \mathrm{id}$.

Now we prove $\psi_a \circ \varphi_a \simeq \mathrm{id}$. First we check that $\psi_a \circ \varphi_a$ is homotopic to the map induced by precomposition with

$$c \times d \simeq (c \times *) \times (* \times d) \longrightarrow (c \times d) \times (c \times d) \xrightarrow{\mu_{c \times d}} c \times d.$$

To show this, we draw another moderately large diagram:

$$\begin{array}{ccccc} \mathrm{Hom}_{\mathcal{C}}(c, a) \times \mathrm{Hom}_{\mathcal{C}}(d, a) & \longleftarrow & \mathrm{Hom}_{\mathcal{C}}(c \times d, a) & & \\ \downarrow & & \downarrow \Delta & & \searrow \mu_{c \times d} \\ \mathrm{Hom}_{\mathcal{C}}(c \times d, a \times a) & \cdots & \mathrm{Hom}_{\mathcal{C}}((c \times d) \times (c \times d), a \times a) & & \\ \downarrow \mu_a & & \downarrow & & \searrow \mu_a \\ \mathrm{Hom}_{\mathcal{C}}(c \times d, a) & \longleftarrow & \mathrm{Hom}_{\mathcal{C}}((c \times *) \times (* \times d), a) & \longleftarrow & \mathrm{Hom}_{\mathcal{C}}((c \times d) \times (c \times d), a) \end{array}$$

(IV) (V) (VI)

The cell labelled (V) commutes up to homotopy because $\mu: \Delta \Rightarrow \mathrm{id}$ is a natural transformation (and I believe Fabian is right that we really need this here—if we were to use Lurie’s weaker condition, it would only commute on π_0). If we choose the dotted arrow to be induced by $c \times d \simeq (c \times *) \times (* \times d) \rightarrow (c \times d) \times (c \times d)$, then also (VI) commutes up to homotopy because it doesn’t matter in which order precomposition with said morphism and postcomposition with μ_a are applied. Finally, (IV) commutes by inspection. Walking around the perimeter of this diagram now proves that $\psi_a \circ \varphi_a$ is really induced by precomposition with the map from above.

Now we finally use the second condition from (b) to obtain a homotopy-commutative diagram

$$\begin{array}{ccccc} c \times d & \xrightarrow{\sim} & (c \times *) \times (* \times c) & \longrightarrow & (c \times d) \times (c \times d) \xrightarrow{\mu_{c \times d}} c \times d \\ & \searrow \sim & \parallel & \uparrow \mathrm{id}_c \times \mathrm{flip} \times \mathrm{id}_c \wr & \nearrow \mu_c \times \mu_d \\ & & (c \times *) \times (* \times d) & \longrightarrow & (c \times c) \times (d \times d) \end{array}$$

in \mathcal{C} . Using the first condition from (b) we see that the composition of the lower three arrows is homotopic to $\mathrm{id}_c \times \mathrm{id}_d$. Hence the morphism we precompose with is homotopic to the identity on $c \times d$, which shows $\psi_a \times \varphi_a \simeq \mathrm{id}$, as required. \square

Proof of Theorem II.19. We first prove that $\mathrm{CMon}(\mathcal{C})$ is semi-additive. Of course we would like to apply Lemma II.20, so we must produce the corresponding data on $\mathrm{CMon}(\mathcal{C})$. First of all, note that $\mathrm{const} *: \mathbb{T}^{\mathrm{op}} \rightarrow \mathcal{C}$ is both terminal and initial. It is even terminal in $\mathrm{Fun}(\mathbb{T}^{\mathrm{op}}, \mathcal{C})$ because $*$ $\in \mathcal{C}$ is terminal and limits in functor categories are computed pointwise (Lemma I.39). To see that it is initial, let $X \in \mathrm{CMon}(\mathcal{C})$ and compute

$$\mathrm{Nat}(\mathrm{const} *, X) \simeq \mathrm{Hom}_{\mathcal{C}}\left(*, \lim_{\langle n \rangle \in \mathbb{T}^{\mathrm{op}}} X_n\right) \simeq \mathrm{Hom}_{\mathcal{C}}(*, X_0) \simeq \mathrm{Hom}_{\mathcal{C}}(*, *) \simeq *,$$

where we used that $\langle 0 \rangle \in \mathbb{T}^{\text{op}}$ is a terminal object and thus spans a final subset.

Next we need to construct functorial maps $\mu_M: M \times M \rightarrow M$ for all $M \in \text{CMon}(\mathcal{C})$. Consider the map $\times: \mathbb{T}^{\text{op}} \times \mathbb{T}^{\text{op}} \rightarrow \mathbb{T}^{\text{op}}$ given by crossing the sets and crossing the morphisms (note that this wouldn't work with Δ^{op}). It induces a functor $\text{Fun}(\mathbb{T}^{\text{op}}, \mathcal{C}) \rightarrow \text{Fun}(\mathbb{T}^{\text{op}}, \text{Fun}(\mathbb{T}^{\text{op}}, \mathcal{C}))$, and one easily checks that the Segal condition is preserved, i.e. we get a functor

$$\text{Double}: \text{CMon}(\mathcal{C}) \longrightarrow \text{CMon}(\text{CMon}(\mathcal{C})).$$

Unravelling definitions, we find that $\text{Double}(-)_1: \text{CMon}(\mathcal{C}) \rightarrow \text{CMon}(\mathcal{C})$ is naturally equivalent to the identity and $\text{Double}(-)_2: \text{CMon}(\mathcal{C}) \rightarrow \text{CMon}(\mathcal{C})$ is naturally equivalent to the diagonal $\Delta: M \mapsto M \times M$. Hence we can take $\mu_M: M \times M \rightarrow M$ to be the multiplication $m: \text{Double}(M)_2 \rightarrow \text{Double}(M)_1$, i.e. the map induced by $m: \langle 2 \rangle \rightarrow \langle 1 \rangle$ in \mathbb{T}^{op} (see Definition II.15). Since the μ_M are induced by a map in \mathbb{T}^{op} , it's clear that they assemble into a natural transformation μ . Also the required conditions for μ are straightforward to check.

This shows that $\text{CMon}(\mathcal{C})$ is semi-additive by Lemma II.20. Hence the same holds for its full subcategory $\text{CGrp}(\mathcal{C}) \subseteq \text{CMon}(\mathcal{C})$ since this is closed under finite products. To show that $\text{CGrp}(\mathcal{C})$ is additive, we need to check that the shear map is an equivalence for all objects $G \in \text{CGrp}(\mathcal{C})$. But in view of the fact that the codiagonal $G \times G \rightarrow G$ is induced by μ_G (this follows by unravelling the explicit inverse in the proof of Lemma II.20), hence by multiplication on $\text{Double}(G)$, this gets translated into the fact that $\text{Double}(G)$ is a cartesian commutative group itself, which is again easy to check.

Finally, we need to show that $\text{CMon}: \text{Cat}_\infty^\times \rightarrow \text{Cat}_\infty^{\text{semi-add}}$ and $\text{CGrp}: \text{Cat}_\infty^\times \rightarrow \text{Cat}_\infty^{\text{add}}$ are right adjoints. If we denote $R = \text{CMon}$ and $\eta: R \Rightarrow \text{id}$ the forgetful natural transformation (given by evaluation in degree 1), then Proposition II.16 and the fact that $\text{CMon}(\mathcal{C})$ is semi-additive show that $\eta R: R \circ R \xrightarrow{\sim} R$ and $R\eta: R \circ R \xrightarrow{\sim} R$ are natural equivalences. Thus we may apply the dual of Proposition I.61a to see that R is a right adjoint to the inclusion $\text{im}(R) \subseteq \text{Cat}_\infty^\times$ of its essential image. But $\text{im}(R) = \text{Cat}_\infty^{\text{semi-add}}$: Indeed, “ \subseteq ” was checked above, whereas “ \supseteq ” follows from Proposition II.16 again. The argument for CGrp is completely analogous. \square

LECTURE 13
10th Dec, 2020

A Few Comments/Examples. — The forgetful functor $\text{CMon}(\text{An}) \rightarrow \text{An}$ as a left adjoint $\text{Free}^{\text{CMon}}: \text{An} \rightarrow \text{CMon}(\text{An})$. Like Proposition II.9, we won't prove this, but Fabian gives a sketch in [A&HK, Chapter II pp.128–132], after our venture into the land of ∞ -operads. On objects $X \in \text{An}$, the left adjoint is given by

$$\text{Free}^{\text{CMon}}(X)_1 \simeq \coprod_{n \geq 0} X_{h\mathfrak{S}_n}^n,$$

i.e. we take free words in X , but we need to homotopy-quotient out the action of the symmetric group \mathfrak{S}_n , which the notation $(-)_{h\mathfrak{S}_n}$ is supposed to do. In general, given a functor $F: BG \rightarrow \mathcal{C}$, where \mathcal{C} is an ∞ -category with colimits and G is a discrete group, we put

$$F_{hG} \simeq \text{colim}_{BG} F.$$

On X^n , the \mathfrak{S}_n -action is given by “permuting factors”. More precisely, we may apply II.6 to see that giving a functor $B\mathfrak{S}_n \rightarrow \text{core}(\text{An})$ sending the unique 0-simplex of $B\mathfrak{S}_n$ to X^n is the same as giving a functor $\mathfrak{S}_n \rightarrow \Omega_{X^n} \text{core}(\text{An}) \simeq \text{Hom}_{\text{core}(\text{An})}(X^n, X^n)$. But each factor permutation is a map in $\text{Hom}_{\text{core}(\text{An})}(X^n, X^n)$, hence there's indeed a canonical map from the discrete anima \mathfrak{S}_n into it.

As an example,

$$\mathrm{Free}^{\mathrm{CMon}}(*)_1 \simeq \coprod_{n \geq 0} B\mathfrak{S}_n \simeq (\{\text{finite sets, bijections}\}, \sqcup) =: \mathfrak{S},$$

where the right-hand side is the symmetric monoidal 1-groupoid of finite sets and bijections between them, whose tensor product is given by disjoint union. In particular, even though $\mathrm{Free}^{\mathrm{Mon}}(*) \simeq \mathbb{N}$ is the free monoid on a point and commutative, it's not the free commutative monoid on a point.

Spectra and Stable ∞ -Categories

Let $\mathrm{CGrp}(\mathrm{An})_{\geq i} \subseteq \mathrm{CGrp}(\mathrm{An})$ be the full sub- ∞ -category spanned by those X with $\pi_j X_1 = 0$ for $j < i$. We have seen in Corollary II.21 that $B: \mathrm{CGrp}(\mathrm{An}) \rightarrow \mathrm{CGrp}(\mathrm{An})$ is fully faithful with right adjoint Ω . Moreover, we've seen during the proof that B actually has essential image in $\mathrm{CGrp}(\mathrm{An})_{\geq 1}$. In fact, its essential image is $\mathrm{CGrp}(\mathrm{An})_{\geq 1}$! To see this, it suffices to check that the counit $c: B\Omega X \xrightarrow{\sim} X$ is an equivalence for all $X \in \mathrm{CGrp}(\mathrm{An})_{\geq 1}$. This can be checked on underlying anima, i.e. after ev_1 , and then again on homotopy groups. Since Ω shifts homotopy groups down and B shifts them up, we get that $c_*: \pi_i(B\Omega X)_1 \xrightarrow{\sim} \pi_i X_1$ is an isomorphism for $i > 0$. For $i = 0$ it is one as well as both sides are connected.

Iterating this argument, we find that the i -fold application $B^{(i)} = B \circ \cdots \circ B$ is an equivalence

$$B^{(i)}: \mathrm{CGrp}(\mathrm{An}) \xrightarrow{\sim} \mathrm{CGrp}(\mathrm{An})_{\geq i}.$$

Moreover, since $\Omega B \simeq \mathrm{id}$ by Corollary II.21(a), we get $\Omega B^{(i+1)} \simeq B^{(i)}$. This yields a fully faithful functor

$$B^\infty: \mathrm{CGrp}(\mathrm{An}) \longrightarrow \lim_{\mathbb{N}^{\mathrm{op}}} \left(\cdots \xrightarrow{\Omega} \mathrm{CGrp}(\mathrm{An}) \xrightarrow{\Omega} \mathrm{CGrp}(\mathrm{An}) \xrightarrow{\Omega} \mathrm{CGrp}(\mathrm{An}) \right)$$

(the limit being taken in Cat_∞ of course) with essential image

$$\lim_{\mathbb{N}^{\mathrm{op}}} \left(\cdots \xrightarrow{\Omega} \mathrm{CGrp}(\mathrm{An})_{\geq 2} \xrightarrow{\Omega} \mathrm{CGrp}(\mathrm{An})_{\geq 1} \xrightarrow{\Omega} \mathrm{CGrp}(\mathrm{An}) \right).$$

II.22. Definition. — We say that an ∞ -category \mathcal{C} has *finite limits* if it has finite products and pullbacks (one can check that this is equivalent to having limits over all finite simplicial sets). If this is the case, then a suspension functor $\Omega: */\mathcal{C} \rightarrow */\mathcal{C}$ can be defined and we put

$$\mathrm{Sp}(\mathcal{C}) := \lim_{\mathbb{N}^{\mathrm{op}}} \left(\cdots \xrightarrow{\Omega} */\mathcal{C} \xrightarrow{\Omega} */\mathcal{C} \right)$$

called the ∞ -category of *spectrum objects* in \mathcal{C} . If $\mathcal{C} = \mathrm{An}$, we obtain $\mathrm{Sp} = \mathrm{Sp}(\mathrm{An})$, the ∞ -category of *spectra*.

II.23. Proposition. — Suppose \mathcal{C} has finite limits. Then the forgetful functor

$$\mathrm{Sp}(\mathrm{CGrp}(\mathcal{C})) \xrightarrow{\sim} \mathrm{Sp}(\mathcal{C})$$

is an equivalence.

Proof sketch. Topologists hate this proof. First observe that $\mathrm{Sp}(\mathrm{CGrp}(\mathcal{C})) \simeq \mathrm{CGrp}(\mathrm{Sp}(\mathcal{C}))$. This is because

$$\lim_{\mathbb{N}^{\mathrm{op}}} \left(\dots \xrightarrow{\Omega} \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, */\mathcal{C}) \xrightarrow{\Omega} \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, */\mathcal{C}) \right) \simeq \mathrm{Fun} \left(\mathbb{I}^{\mathrm{op}}, \lim_{\mathbb{N}^{\mathrm{op}}} \left(\dots \xrightarrow{\Omega} */\mathcal{C} \xrightarrow{\Omega} */\mathcal{C} \right) \right)$$

(here we use that $\mathrm{Fun}(\mathcal{D}, -): \mathrm{Cat}_{\infty} \rightarrow \mathrm{Cat}_{\infty}$ commutes with limits since it is right-adjoint to $\mathcal{D} \times -$ and also that Ω in functor categories can be computed pointwise by Lemma I.39) and one checks that the Segal condition and the group condition are preserved.

Therefore, we will be done by Proposition II.16 once we show that $\mathrm{Sp}(\mathcal{C})$ is an additive category. Of course we will employ Lemma II.20 again. Clearly $(*, *, \dots) \in \mathrm{Sp}(\mathcal{C})$ is both initial and terminal. Also $\mathrm{Sp}(\mathcal{C})$ again has finite limits, inherited from $*/\mathcal{C}$ because limits in limits can be computed degreewise (combine the fact that Hom anima in limits are limits of Hom anima with Corollary I.50 and Proposition I.42). Moreover, we know that $\Omega: \mathrm{Sp}(\mathcal{C}) \xrightarrow{\sim} \mathrm{Sp}(\mathcal{C})$ is an equivalence by construction and factors as

$$\Omega: \mathrm{Sp}(\mathcal{C}) \longrightarrow \mathrm{Grp}(\mathrm{Sp}(\mathcal{C})) \xrightarrow{\mathrm{ev}_1} \mathrm{Sp}(\mathcal{C})$$

(I didn't find this quite obvious, so I decided to sketch a proof in Remark* II.23a below). Finally, to apply Lemma II.20, we must produce functorial morphisms $\mu_X: X \times X \rightarrow X$ with the properties stated there. For this, we use $X \simeq \Omega(\Omega^{-1}X)_1$ and take μ_X to be induced by the multiplication on $\Omega(\Omega^{-1}X) \in \mathrm{Grp}(\mathrm{Sp}(\mathcal{C}))$.

Up to checking the various conditions, this shows that $\mathrm{Sp}(\mathcal{C})$ is semi-additive. For additivity we must check that the shear map is an equivalence on every $X \in \mathrm{Sp}(\mathcal{C})$. Similar to the proof of Theorem II.19 this follows from the fact that $\Omega(\Omega^{-1}X)$ is a cartesian group rather than just a monoid. \square

II.23a. Remark*. — For any ∞ -category \mathcal{D} with finite limits and a zero object $*$ $\in \mathcal{D}$ (i.e. an object that is both initial and terminal) the functor $\Omega: \mathcal{D} \rightarrow \mathcal{D}$ factors as

$$\Omega: \mathcal{D} \longrightarrow \mathrm{Grp}(\mathcal{D}) \xrightarrow{\mathrm{ev}_1} \mathcal{D}$$

This can be bootstrapped from the case $\mathcal{D} = */\mathrm{An}$ which we already know from II.6. The first step is to enhance the Yoneda embedding to a fully faithful functor $\mathcal{D} \rightarrow \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An})$. To this end we consider $\mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, \mathrm{An}) \rightarrow \mathrm{Fun}(\mathrm{Ar}(\mathcal{D}^{\mathrm{op}}), \mathrm{Ar}(\mathrm{An})) \rightarrow \mathrm{Fun}(*/\mathcal{D}^{\mathrm{op}}, \mathrm{Ar}(\mathrm{An}))$ and check that its image lands in the “correct fibre”, i.e. in $\mathrm{Fun}(*/\mathcal{D}^{\mathrm{op}}, */\mathrm{An})$ if we restrict the source to finite limits-preserving functors (where the Yoneda embedding lands).

On $\mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An})$ we have loop functor $\Omega: \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An}) \rightarrow \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An})$ as well, and it is simply given by postcomposition with $\Omega: */\mathrm{An} \rightarrow */\mathrm{An}$. Hence it factors as

$$\begin{array}{ccc} & \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, \mathrm{Grp}(*/\mathrm{An})) & \xrightarrow{\sim} \mathrm{Grp}(\mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An})) \\ & \uparrow \Omega & \downarrow \mathrm{ev}_1 \\ \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An}) & \xrightarrow{\Omega} & \mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An}) \end{array}$$

Since the Yoneda embedding preserves limits, we get a lift $\Omega: \mathcal{D} \rightarrow \mathrm{Grp}(\mathrm{Fun}(\mathcal{D}^{\mathrm{op}}, */\mathrm{An}))$ and one checks that it lands in the full sub- ∞ -category $\mathrm{Grp}(\mathcal{D})$, as required.

II.23b. Homotopy Groups of Spectra. — Because $\mathbb{N}^{\text{op}} \subseteq \mathbb{Z}^{\text{op}}$ is final in the sense of Definition I.44, we can also write $\text{Sp}(\mathcal{C}) \simeq \lim_{\mathbb{Z}^{\text{op}}} (* / \mathcal{C} \supset \Omega)$. This gives evaluation functors

$$\Omega^{\infty-i}: \text{Sp}(\mathcal{C}) \longrightarrow \mathcal{C}.$$

for all $i \in \mathbb{Z}$. Note $\Omega\Omega^{\infty-i}X \simeq \Omega^{\infty-(i-1)}X \simeq \Omega^{\infty-i}\Omega X$, so this somewhat weird notation actually makes sense. In the case $\mathcal{C} = \text{An}$, we can now define homotopy groups for $X \in \text{Sp}$ as

$$\pi_i(X) = \pi_0(\Omega^{\infty+i}X).$$

If we think of a spectrum X as a sequence $X = (X_0, X_1, X_2, \dots)$, where $X_i = \Omega^{\infty-i}X$, together with equivalences $X_i \simeq \Omega X_{i+1}$, then we see that $\pi_i(X) \simeq \pi_0(X_{-i}) \simeq \pi_j(X_{j-i})$ for all $j \geq 0$ (we don't need to specify any base points, since the X_i are pointed anima) because $\Omega: */\text{An} \rightarrow */\text{An}$ shifts homotopy groups down.

It's probably clear to all of you, but let me also mention that equivalences of spectra can be detected on homotopy groups:

II.23c. Lemma*. — *A map $f: X \rightarrow Y$ of spectra is an equivalence in Sp iff it induces isomorphisms $f_*: \pi_*(X) \xrightarrow{\sim} \pi_*(Y)$ on homotopy groups.*

Proof.* Write $X_i = \Omega^{\infty-i}$ and $Y_i = \Omega^{\infty-i}$ as above. Then f is an equivalence in Sp iff each $\Omega^{\infty-i}f: X_i \rightarrow Y_i$ is an equivalence in An . Indeed, the inverses of $\Omega^{\infty-i}f$ will again assemble into a compatible system of morphisms and hence into an inverse of f . Whether $\Omega^{\infty-i}f$ is an equivalence in An can be detected on homotopy groups. In fact, both X_i is an element of $\text{CGrp}(\text{An})$, so all its connected components are equivalent, and the same is true for Y_i . Hence, instead of considering homotopy groups with arbitrary base points, it suffices to check that $\pi_0(X) \rightarrow \pi_0(Y)$ is a bijection and $\pi_j(X_i) \rightarrow \pi_j(Y_i)$ (with the canonical choice of base points) are group isomorphisms for all $j > 0$.

Now $\pi_j(X_i) \simeq \pi_{j-i}(X)$ for all $j \geq 0$, and the same holds for Y , hence it suffices to have isomorphisms $f_*: \pi_*(X) \xrightarrow{\sim} \pi_*(Y)$, as claimed. \square

II.23d. Summary of What We Know So Far. — Let's explicitly record the following facts we already learned in the proof of Proposition II.23 and afterwards:

- (a) $\text{Sp}(\mathcal{C})$ has finite limits (because $*/\mathcal{C}$ has) and limits in $\text{Sp}(\mathcal{C})$ are computed degreewise, i.e. $\Omega^{\infty-i}: \text{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$ preserves limits.
- (b*) The only reason this doesn't work for colimits as well is that the degreewise colimits might not be compatible with the Ω -maps any more, so they don't necessarily define an element of $\text{Sp}(\mathcal{C})$. If that happens to be the case, then the degreewise colimit is in fact the colimit in Sp , by the same argument as for limits (combine the fact that Hom anima in limits are limits of Hom anima with Corollary I.50 and Proposition I.42).
- (c) $\Omega: \text{Sp}(\mathcal{C}) \xrightarrow{\sim} \text{Sp}(\mathcal{C})$ is an equivalence.
- (d) $\text{Sp}(\mathcal{C})$ is additive.

II.24. Corollary (Recognition principle for infinite loop spaces, Boardman–Vogt, May, Segal). — *The functor*

$$B^\infty: \text{CGrp}(\text{An}) \longrightarrow \text{Sp}$$

*is fully faithful with essential image those $X \in \text{Sp}$ with $\pi_i X = 0$ for $i < 0$ (as a slogan, \mathbb{E}_∞ -groups are spectra with no negative homotopy groups). Also $\pi_i(B^\infty X) = \pi_i(X_1, *)$ for all \mathbb{E}_∞ -groups $X \in \text{CGrp}(\text{An})$ and all $i \geq 0$.*

*Proof**. The target of B^∞ is $\mathrm{Sp}(\mathrm{CGrp}(\mathrm{An})) \simeq \mathrm{Sp}$ by Proposition II.23. We have seen that B^∞ is fully faithful and identified its essential image before Definition II.22. Clearly, any element X of the limit characterising the essential image satisfies $\pi_i X = 0$ for $i < 0$. Conversely, if X satisfies this condition, then $\pi_j \Omega^{\infty-i} X \simeq \pi_{j-i} X$, which vanishes for $j < i$. Remembering $\mathrm{Sp} \simeq \mathrm{Sp}(\mathrm{CGrp}(\mathrm{An}))$, this shows $\Omega^{\infty-i} X \in \mathrm{CGrp}(\mathrm{An})_{\geq i}$, so X lies in the essential image of B^∞ . The final assertion is clear since $\pi_i(B^\infty X) \simeq \pi_i(\Omega^\infty B^\infty X) \simeq \pi_i(X_1, *)$ holds by definition. \square

II.25. Example. — Let \mathcal{C} always have finite limits. Here are “my first spectra”:

- (a) If $\Omega: */\mathcal{C} \xrightarrow{\sim} */\mathcal{C}$ is an equivalence, then $\mathrm{Sp}(\mathcal{C}) \simeq \mathcal{C}$ (because in that case we’re taking a constant limit over \mathbb{N}^{op} , which is weakly contractible, so that limit is \mathcal{C} by Proposition I.36). For example, $\mathrm{Sp}(\mathcal{D}(\mathbb{Z})) \simeq \mathcal{D}(\mathbb{Z})$ or, even more silly, $\mathrm{Sp}(\mathrm{Sp}) \simeq \mathrm{Sp}$.
- (b) Let’s compute $\mathrm{Sp}(\mathcal{D}_{\geq 0}(R))$ for a ring R . Recall that the loop functor on $\mathcal{D}_{\geq 0}(R)$ is defined as $\Omega C \simeq \tau_{\geq 0}(C[-1]) \in \mathcal{D}_{\geq 0}(R)$. Then the functor

$$\begin{aligned} \mathcal{D}(R) &\xrightarrow{\sim} \mathrm{Sp}(\mathcal{D}_{\geq 0}(R)) \\ C &\mapsto (\tau_{\geq 0}C, (\tau_{\geq -1}C)[1], (\tau_{\geq -2}C)[2], \dots) \end{aligned}$$

is an equivalence. The inverse functor sends a sequence (C_0, C_1, \dots) with equivalences $\sigma_i: C_i \simeq (\tau_{\geq 1}C_{i+1})[-1]$ to $\mathrm{colim}_{\mathbb{N}} C_i[-i]$. The structure maps $C_i[-i] \rightarrow C_{i+1}[-(i+1)]$ in this colimit are constructed from the σ_i in the obvious way.

- (c) Consider the Eilenberg–MacLane functor $K: \mathcal{D}_{\geq 0}(\mathbb{Z}) \rightarrow \mathrm{An}$. Being a right adjoint, it preserves finite limits, and in particular loop spaces. Hence it upgrades canonically to a functor

$$\begin{array}{ccc} \mathrm{Sp}(\mathcal{D}_{\geq 0}(\mathbb{Z})) & \xrightarrow{\mathrm{Sp}(K)} & \mathrm{Sp}(\mathrm{An}) \\ \parallel & & \parallel \\ \mathcal{D}(\mathbb{Z}) & \xrightarrow{H} & \mathrm{Sp} \end{array}$$

the *Eilenberg–MacLane spectrum functor*. By construction, the diagram

$$\begin{array}{ccc} \mathcal{D}(\mathbb{Z}) & \xrightarrow{H} & \mathrm{Sp} \\ & \searrow & \swarrow \\ & K(\tau_{\geq -i-})[i] & \Omega^{\infty-i} \\ & & \mathrm{CGrp}(\mathrm{An}) \end{array}$$

commutes. If $C \in \mathcal{D}_{\geq 0}(\mathbb{Z})$, then $HC = (K(C), K(C[1]), K(C[2]), \dots)$ and we have $\pi_i(HC) = H_i(C)$. If C is an abelian group A concentrated in degree 0, we obtain the *Eilenberg–MacLane spectrum* $HA = (K(A, 0), K(A, 1), K(A, 2), \dots)$.

- (d) We have $\mathrm{Sp}(\mathrm{An}^{\mathrm{op}}) \simeq *$, because it is a sequential limit over $(\emptyset/\mathrm{An}^{\mathrm{op}}) \simeq (\mathrm{An}/\emptyset)^{\mathrm{op}} \simeq *$ (since there is no map $X \rightarrow \emptyset$ in An except $\mathrm{id}_\emptyset: \emptyset \rightarrow \emptyset$). Also

$$\begin{aligned} \mathrm{Sp}((*/\mathrm{An})^{\mathrm{op}}) &\simeq \lim_{\mathbb{N}^{\mathrm{op}}} \left(\dots \xrightarrow{\Omega} (* / \mathrm{An})^{\mathrm{op}} \xrightarrow{\Omega} (* / \mathrm{An})^{\mathrm{op}} \right) \\ &\simeq \lim_{\mathbb{N}^{\mathrm{op}}} \left(\dots \xrightarrow{\Sigma} */\mathrm{An} \xrightarrow{\Sigma} */\mathrm{An} \right)^{\mathrm{op}} \simeq *, \end{aligned}$$

since only contractible spaces can be an infinite suspension (as all homotopy groups must be vanish).

II.26. Homology and Cohomology of Spectra. — Via the Eilenberg–MacLane spectrum functor H , the adjunction $\tilde{C}_\bullet : * / \text{An} \rightleftarrows \mathcal{D}_{\geq 0}(\mathbb{Z}) : K$ from Very Long Example 1.56 can be upgraded to another adjunction

$$C_\bullet : \text{Sp} \rightleftarrows \mathcal{D}(\mathbb{Z}) : H.$$

The new C_\bullet is defined by the formula $C_\bullet(X) \simeq \text{colim}_{i \in \mathbb{N}} \tilde{C}_\bullet(\Omega^{\infty-i} X)[-i]$ on objects. To see where the transition maps in this colimit come from, recall that $\Omega = [-1]$ is the loop functor in $\mathcal{D}(\mathbb{Z})$. Together with $\tilde{C}(\ast) \simeq 0$, this provides a natural transformation $\tilde{C}_\bullet(\Omega -) \Rightarrow \tilde{C}_\bullet(-)[-1]$, which provide the required transition maps.

To check that C_\bullet is a left adjoint, it suffices to do so on objects by Corollary 1.32 (in particular, this will show that C_\bullet is a functor at all), where we can compute

$$\begin{aligned} \text{Hom}_{\mathcal{D}(\mathbb{Z})}(C_\bullet(X), D) &\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \text{Hom}_{\mathcal{D}(\mathbb{Z})}(\tilde{C}_\bullet(\Omega^{\infty-i} X), D[i]) \\ &\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \text{Hom}_{*/\text{An}}(\Omega^{\infty-i} X, K(D[i])) \\ &\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \text{Hom}_{*/\text{An}}(\Omega^{\infty-i} X, \Omega^{\infty-i} HD), \end{aligned}$$

and the last term is $\text{Hom}_{\text{Sp}}(X, HD)$ since Hom anima in Sp (in fact, in arbitrary limits of ∞ -categories) can be computed as a limit over the Hom anima in each degree. Also all equivalences are clearly functorial in $D \in \mathcal{D}(\mathbb{Z})$.

Now define the *homology* and the *cohomology* of $X \in \text{Sp}$ with coefficients in $A \in \text{Ab}$ to be

$$\begin{aligned} H_*(X, A) &:= H_*(C_\bullet(X) \otimes_{\mathbb{Z}}^L A) = \text{colim}_{i \in \mathbb{N}} H_*(\Omega^{\infty-i} X, A), \\ H^*(X, A) &:= H_{-*}(R\text{Hom}_{\mathbb{Z}}(C_\bullet(X), A)). \end{aligned}$$

In the special case $X \simeq \mathbb{S}[Y]$ for some anima Y , it turns out that $H_*(\mathbb{S}[Y], A) = H_*(Y, A)$ and $H^*(\mathbb{S}[Y], A) = H^*(Y, A)$ coincide with singular cohomology. See Lemma II.31c for the case of homology, and Corollary* II.55b for the case of cohomology. In the latter corollary* we'll also find out that $H_*(X, A) = \text{Tor}_*^{\mathbb{S}}(X, HA)$ and $H^*(X, A) \simeq \text{Ext}_{\mathbb{S}}^*(X, HA)$ can be interpreted as “Tor and Ext over the sphere spectrum”.

⚠ Warning. — Don't confuse the homology or cohomology of a spectrum with the homology or cohomology theory *associated* to a spectrum, which we'll introduce in Definition II.35 below. Also, neither $H_*(HC, A)$ for a complex $C \in \mathcal{D}(\mathbb{Z})$ nor $H_*(B^\infty M, A)$ for some $M \in \text{CGrp}(\text{An})$ are easy to compute in general!

II.27. Proposition/Definition. — Suppose \mathcal{C} is an ∞ -category with a zero object (i.e. a simultaneously initial and terminal object $0 \in \mathcal{C}$). Then the following conditions are equivalent:

- (a) \mathcal{C} has finite limits and $\Omega: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$ is an equivalence.
- (b) \mathcal{C} has finite colimits and $\Sigma: \mathcal{C} \xrightarrow{\sim} \mathcal{C}$ is an equivalence.
- (c) \mathcal{C} has finite limits and finite colimits and a commutative square in \mathcal{C} is a pushout square iff it is a pullback square.

Such ∞ -categories are called *stable ∞ -categories*.

Proof. The implications (c) \Rightarrow (a), (b) are clear: Applying the pushout-pullback condition to the pushout square defining Σ and the pullback square defining Ω shows that the unit

$\text{id} \xRightarrow{\sim} \Omega\Sigma$ and counit $\Sigma\Omega \xRightarrow{\sim} \text{id}$ are natural equivalences, so Σ and Ω must be equivalences of ∞ -categories.

For $(a) \Rightarrow (c)$, the same argument as in the proof of Proposition II.23 shows that \mathcal{C} is additive (write X as ΩY in $\text{Grp}(\mathcal{C})$ and apply Lemma II.20). So we only need to check that pushouts exist and coincide with pullbacks. Let $P \subseteq \text{Fun}(\Delta^1 \times \Delta^1, \mathcal{C})$ be the full subcategory spanned by pullback squares. We claim:

(\boxtimes) *The restriction $P \xrightarrow{\sim} \text{Fun}(\Lambda_0^2, \mathcal{C})$ is an equivalence.*

To prove (\boxtimes), we construct an inverse functor. Given a diagram $F: \Lambda_0 \rightarrow \mathcal{C}$, which we can view as a span $c \leftarrow a \rightarrow b$ in \mathcal{C} , we construct the following moderately large diagram:

$$\begin{array}{ccccccc}
 \Omega a & \longrightarrow & \Omega c & \longrightarrow & 0 & & \\
 \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & & \\
 \Omega b & \longrightarrow & x & \longrightarrow & g & \longrightarrow & 0 \\
 \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\
 0 & \longrightarrow & f & \longrightarrow & a & \longrightarrow & b \\
 & & \downarrow & \lrcorner & \downarrow & & \\
 & & 0 & \longrightarrow & c & &
 \end{array}$$

All squares are pullbacks as indicated. The fact that Ωa , Ωb , and Ωc appear in the top left corner follows by combining suitable pullback squares into larger pullback rectangles.

The inverse functor now sends

$$F = \left(\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & & \\ c & & \end{array} \right) \mapsto \Omega^{-1} \left(\begin{array}{ccc} \Omega a & \longrightarrow & \Omega b \\ \downarrow & \lrcorner & \downarrow \\ \Omega c & \longrightarrow & x \end{array} \right)$$

(technically we have only defined the inverse on objects, but its clear how to make it functorial since limits are functorial).

To construct pushouts using (\boxtimes), let $F: \Lambda_0^2 \rightarrow \mathcal{C}$ be a span $c \leftarrow a \rightarrow b$ as above. We know that it can be uniquely (up to contractible choice) extended to a pullback square, where the bottom right corner is some object $d \in \mathcal{C}$. The same goes for the trivial span consisting of identities $x = x = x$ for some $x \in \mathcal{C}$, and in this case the object we have to add is x again of course. Hence

$$\text{Nat} \left(\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & & \\ c & & \end{array}, \begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & & \\ x & & \end{array} \right) \simeq \text{Nat} \left(\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & \lrcorner & \downarrow \\ c & \longrightarrow & d \end{array}, \begin{array}{ccc} x & \xlongequal{\quad} & x \\ \parallel & \lrcorner & \parallel \\ x & \xlongequal{\quad} & x \end{array} \right) \simeq \text{Hom}_{\mathcal{C}}(d, x)$$

because $\Delta^1 \times \Delta^1$ has terminal vertex $(1, 1)$. This computation shows that d is a pushout of the given span. Moreover, we get the property that pushouts agree with pullbacks for free. This finishes the proof of $(a) \Rightarrow (c)$. For $(b) \Rightarrow (c)$ just dualize everything. \square

An immediate consequence of Proposition/Definition II.27 is that $\text{Sp}(\mathcal{C})$ is stable for all ∞ -categories \mathcal{C} with finite limits. In particular, it holds for Sp and $\mathcal{D}(R)$ for any ring R by Example II.25(b) (but it's also true for $\mathcal{K}(R)$). Moreover, $\text{Fun}(\mathcal{I}, \mathcal{C})$ is stable whenever \mathcal{C} is.

LECTURE 14
15th Dec, 2020**II.28. Corollary.** — *For a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ between stable ∞ -categories the following are equivalent:*

- (a) *F preserves finite limits.*
- (b) *F preserves finite colimits.*

Proof. Since \mathcal{C} and \mathcal{D} are additive, F preserves finite coproducts iff it preserves finite products. Since also pushout squares and pullback squares coincide, F preserves the former iff it preserves the latter. This suffices to get all finite colimits and limits. \square

II.29. Definition. — (a) Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor between ∞ -categories with finite limits. We call F *left exact* if it preserves finite limits.

- (b) We denote by $\text{Cat}_\infty^{\text{lex}} \subseteq \text{Cat}_\infty$ the (non-full) sub- ∞ -category spanned by these. Dually, $\text{Cat}_\infty^{\text{rex}} \subseteq \text{Cat}_\infty$ is the full sub- ∞ -category spanned by *right-exact* functors, i.e. finite colimit-preserving functors between ∞ -categories with finite colimits.
- (c) Finally, we denote by $\text{Cat}_\infty^{\text{st}}$ the full subcategory of both $\text{Cat}_\infty^{\text{lex}}$ and $\text{Cat}_\infty^{\text{rex}}$ spanned by the stable ∞ -categories.

For example, the functor $\Omega^\infty: \text{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$ is left exact for every \mathcal{C} with finite limits. Also note that Ω^∞ factors canonically over $\text{CGrp}(\mathcal{C})$. This needs doesn't need an argument as in Remark* II.23a, simply recall that $\text{Sp}(\mathcal{C}) \simeq \text{Sp}(\text{CGrp}(\mathcal{C}))$ by Proposition II.23.

II.29a. Remark*. — Before we move on, I would like to point out a subtlety that's perhaps easy to overlook: In Theorem II.30 below, we'll investigate the functor

$$\text{Sp}(-): \text{Cat}_\infty^{\text{lex}} \longrightarrow \text{Cat}_\infty^{\text{st}}.$$

But why is it a functor though? My own knee-jerk reaction was “well, limits are functorial”, but that's not the problem. The problem is that $\Omega: */\mathcal{C} \rightarrow */\mathcal{C}$ is also supposed to be functorial in \mathcal{C} ! Essentially this leads to the following question:

- (\boxtimes) *Let $\text{Cat}_\infty^{\mathcal{I}\text{-ex}} \subseteq \text{Cat}_\infty$ be the (non-full) subcategory spanned by ∞ -categories with \mathcal{I} -shaped limits and functors preserving these. Why do the maps $\lim_{\mathcal{I}}: \text{Fun}(\mathcal{I}, \mathcal{C}) \rightarrow \mathcal{C}$ for $\mathcal{C} \in \text{Cat}_\infty^{\mathcal{I}\text{-ex}}$ assemble into a natural transformation*

$$\lim_{\mathcal{I}}: \text{Fun}(\mathcal{I}, -) \Longrightarrow (-)$$

of functors $\text{Cat}_\infty^{\mathcal{I}\text{-ex}} \rightarrow \text{Cat}_\infty$?

There are a few more steps involved to show that Ω is natural too. Namely one has to produce natural transformations $(-) \Rightarrow */(-) \Rightarrow \text{Fun}(\Lambda_2^2, */(-))$, but I'll leave that to you.

To show (\boxtimes), let's first produce the “adjoint” transformation $\text{const}: (-) \Rightarrow \text{Fun}(\mathcal{I}, -)$. Since the adjunction

$$- \times \mathcal{I}: \text{Cat}_\infty \rightleftarrows \text{Cat}_\infty : \text{Fun}(\mathcal{I}, -)$$

extends to $\text{Fun}(\text{Cat}_\infty^{\mathcal{I}\text{-ex}}, \text{Cat}_\infty)$ by Observation I.40, we may equivalently give a transformation $- \times \mathcal{I} \Rightarrow (-)$. Here the projection to the first factor is an obvious candidate and it is indeed the correct one.

We may view the natural transformation const as a functor $\text{Cat}_\infty^{\mathcal{I}\text{-ex}} \rightarrow \text{Ar}(\text{Cat}_\infty)$. Its image lies in the full subcategory spanned by those $[1] \rightarrow \text{Cat}_\infty$ that represent left-adjoint functors (because $\lim_{\mathcal{I}}$ is a pointwise right adjoint). But in fact, more is true (and that's

the crucial point). Consider objects $(L: \mathcal{C} \rightarrow \mathcal{D})$ and $(L': \mathcal{C}' \rightarrow \mathcal{D}')$ in $\text{Ar}(\text{Cat}_\infty)$ as well as a morphism between them, which we view as a solid diagram

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \mathcal{C}' \\ L \downarrow \wr & R & L' \downarrow \wr \\ \mathcal{D} & \longrightarrow & \mathcal{D}' \end{array}$$

If L and L' are left adjoints, then the dotted arrows exist, but the resulting new square only commutes up to a natural transformation, not up to natural equivalence in general. We let $\text{Ar}^L(\text{Cat}_\infty)$ be the non-full sub- ∞ -category spanned by those squares for which the adjoint square does in fact commute. Then $\text{Cat}_\infty^{\mathcal{I}\text{-ex}} \rightarrow \text{Ar}(\text{Cat}_\infty)$ already factors over $\text{Ar}^L(\text{Cat}_\infty)$ since morphisms in $\text{Cat}_\infty^{\mathcal{I}\text{-ex}}$ preserve \mathcal{I} -shaped limits.

Now note that under the equivalence $\text{Un}^{\text{cocart}}: \text{Ar}(\text{Cat}_\infty) \xrightarrow{\sim} \text{Cocart}([1])$, the non-full sub- ∞ -category $\text{Ar}^L(\text{Cat}_\infty)$ corresponds to $\text{Bicart}([1])$, the ∞ -category of bicartesian fibrations over Δ^1 with morphisms that preserve both cocartesian and cartesian edges (we know from [HCII, Proposition XI.10] that bicartesian fibrations correspond to adjunctions, and the additional condition unravels to the condition that cartesian edges are preserved as well). Now we can apply the cartesian straightening $\text{St}^{\text{cart}}: \text{Bicart}([1]) \rightarrow \text{Fun}([1]^{\text{op}}, \text{Cat}_\infty)$ to get a functor $\text{Cat}_\infty^{\mathcal{I}\text{-ex}} \rightarrow \text{Fun}([1]^{\text{op}}, \text{Cat}_\infty)$. After currying around, this gives a map $[1]^{\text{op}} \rightarrow \text{Fun}(\text{Cat}_\infty^{\mathcal{I}\text{-ex}}, \text{Cat}_\infty)$ which finally provides the desired transformation.

II.30. Theorem. — *The functor $\text{Sp}(-): \text{Cat}_\infty^{\text{lex}} \rightarrow \text{Cat}_\infty^{\text{st}}$ is right-adjoint to the inclusion $\text{Cat}_\infty^{\text{st}} \subseteq \text{Cat}_\infty^{\text{lex}}$, the counit being given by $\Omega^\infty: \text{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$. In particular, if \mathcal{C} is a stable ∞ -category, we get a canonical lift*

$$\begin{array}{ccc} & & \text{Sp} \\ & \nearrow \text{hom}_{\mathcal{C}} & \downarrow \Omega^\infty \\ \mathcal{C}^{\text{op}} \times \mathcal{C} & \xrightarrow{\text{Hom}_{\mathcal{C}}} & \text{An} \end{array}$$

of the $\text{Hom}_{\mathcal{C}}$ functor.

Fabian remarks that $\text{Cat}_\infty^{\text{st}} \subseteq \text{Cat}_\infty^{\text{lex}}$ also has a left adjoint given by $\text{colim}(* / \mathcal{C} \supset \Omega)$ but we probably won't ever use this. More interestingly, we can apply Theorem II.30 to the stable ∞ -categories $\mathcal{D}(R)$ and $\mathcal{K}(R)$ for any ring R :

II.30a. Corollary. — *The Hom functors on $\mathcal{D}(R)$ and $\mathcal{K}(R)$ have canonical refinements over Sp , given by*

$$\text{hom}_{\mathcal{D}(R)}(C, D) \simeq H(R\text{Hom}_R(C, D)) \quad \text{and} \quad \text{hom}_{\mathcal{K}(R)}(C, D) \simeq H(\underline{\text{Hom}}_R(C, D)).$$

Proof sketch.* To prove the left equivalence, it suffices to check that we get $\text{Hom}_{\mathcal{D}(R)}(C, D)$ after applying Ω^∞ (and that everything is functorial in C and D , but this will be clear). So we compute

$$\begin{aligned} \Omega^\infty H(R\text{Hom}_R(C, D)) &\simeq K(R\text{Hom}_R(C, D)) \simeq \text{Hom}_{\text{An}}(*, K(R\text{Hom}_R(C, D))) \\ &\simeq \text{Hom}_{\mathcal{D}(\mathbb{Z})}(\mathbb{Z}[0], R\text{Hom}_R(C, D)) \\ &\simeq \text{Hom}_{\mathcal{D}(R)}(\mathbb{Z}[0] \otimes_{\mathbb{Z}}^L C, D) \\ &\simeq \text{Hom}_{\mathcal{D}(R)}(C, D), \end{aligned}$$

using $C_\bullet(*) \simeq \mathbb{Z}$ and the derived tensor-Hom adjunction. The right equivalence can be proved analogously, but one should first check that $C_\bullet: \mathbf{An} \rightarrow \mathcal{D}_{\geq 0}(\mathbb{Z})$ already factors over $\mathcal{K}_{\geq 0}(\mathbb{Z})$ and gives rise to the same Eilenberg–MacLane functor $K: \mathcal{K}_{\geq 0}(\mathbb{Z}) \rightarrow \mathbf{An}$. We haven’t done this, but one can simply modify the construction in Very Long Example I.56 (details left to the reader). \square

Proof of Theorem II.30. By the dual of Proposition I.61a, all we need to show is that the two functors $\Omega^\infty: \mathbf{Sp}(\mathbf{Sp}(\mathcal{C})) \xrightarrow{\sim} \mathbf{Sp}(\mathcal{C})$ and $\Omega_*^\infty: \mathbf{Sp}(\mathbf{Sp}(\mathcal{C})) \rightarrow \mathbf{Sp}(\mathcal{C})$ are equivalences. But we know that Ω^∞ is one by Example II.25(a). Also Ω_*^∞ is an equivalence too because it is a “coordinate flip” of Ω^∞ . By that, we mean the following: Plugging in the definitions, we can write $\mathbf{Sp}(\mathbf{Sp}(\mathcal{C}))$ as a limit indexed by $\mathbb{N}^{\text{op}} \times \mathbb{N}^{\text{op}}$. Using the dual of Proposition I.42, we can flip that diagram around, swapping Ω^∞ and Ω_*^∞ , which proves the latter is an equivalence as well.

For the “in particular”, we need another small argument similar to the proof* of Corollary II.17, since $\text{Hom}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{An}$ isn’t left exact. Every $\text{Hom}_{\mathcal{C}}(x, -): \mathcal{C} \rightarrow \mathbf{An}$ is, however, hence the adjunction provides us with canonical lifts $\text{hom}_{\mathcal{C}}(x, -) \rightarrow \mathbf{Sp}$ for all $x \in \mathcal{C}$ when \mathcal{C} is a stable ∞ -category. To show that these combine into a functor $\text{hom}_{\mathcal{C}}: \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{An}$, consider the full subcategory $\text{Fun}^{\text{lex}}(\mathcal{C}, \mathbf{An}) \subseteq \text{Fun}(\mathcal{C}, \mathbf{An})$ spanned by left exact functors. We claim that

$$\Omega_*^\infty: \text{Fun}^{\text{ex}}(\mathcal{C}, \mathbf{Sp}) \xrightarrow{\sim} \text{Fun}^{\text{lex}}(\mathcal{C}, \mathbf{An})$$

is an equivalence. Indeed, it becomes one after applying $\text{core}(-)$ by the adjunction we’ve just proved. To show that it is an equivalence after $\text{core Ar}(-)$, we can use the same argument once we convince ourselves that $\text{Ar Fun}^{\text{ex}}(\mathcal{C}, \mathbf{Sp}) \simeq \text{Fun}^{\text{ex}}(\mathcal{C}, \text{Ar}(\mathbf{Sp}))$ (use that limits in arrow categories are pointwise by Lemma I.39) and that $\text{Ar}(\mathbf{Sp}) \simeq \mathbf{Sp}(\text{Ar}(\mathbf{An}))$ (use that $\text{Ar}(-)$ commutes with limits).

Now the Yoneda embedding $\mathcal{C}^{\text{op}} \rightarrow \text{Fun}(\mathcal{C}, \mathbf{An})$ has image in $\text{Fun}^{\text{lex}}(\mathcal{C}, \mathbf{An}) \simeq \text{Fun}^{\text{ex}}(\mathcal{C}, \mathbf{Sp})$ and we get our desired $\text{hom}_{\mathcal{C}}$ after currying. \square

Spectra and Prespectra. — We already know that \mathbf{Sp} is complete and $\Omega^{\infty-i}: \mathbf{Sp} \rightarrow \mathbf{An}$ preserves limits for all i , see the summary before Corollary II.24. We also know from Proposition/Definition II.27 that \mathbf{Sp} has finite colimits because it is stable. But in fact, \mathbf{Sp} is even cocomplete! To see this, we will write \mathbf{Sp} as a Bousfield localisation of a suitable cocomplete ∞ -category of *prespectra* and apply Corollary I.61.

Prespectra can be introduced in general. So let \mathcal{C} be an ∞ -category with finite limits and (thus) a terminal object $*$ $\in \mathcal{C}$. We let

$$\mathbf{PSp}(\mathcal{C}) \subseteq \text{Fun}(\mathbb{N}^2, */\mathcal{C})$$

be the full subcategory of functors that “vanish away from the diagonal”. So the objects of $\mathbf{PSp}(\mathcal{C})$ can be pictured as lattice diagrams

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & \\ & \uparrow & & \uparrow & & \uparrow & \\ * & \longrightarrow & * & \longrightarrow & X_2 & \longrightarrow & \cdots \\ & \uparrow & & \uparrow & & \uparrow & \\ * & \longrightarrow & X_1 & \longrightarrow & * & \longrightarrow & \cdots \\ & \uparrow & & \uparrow & & \uparrow & \\ X_0 & \longrightarrow & * & \longrightarrow & * & \longrightarrow & \cdots \end{array}$$

The diagonal squares induce morphisms $\Sigma X_i \rightarrow X_{i+1}$ (provided \mathcal{C} has pushouts) or equivalently $X_i \rightarrow \Omega X_{i+1}$. If the latter were to be isomorphisms, we would recover $\mathrm{Sp}(\mathcal{C})$, as we'll see in a moment.

To construct a fully faithful embedding $\mathrm{Sp}(\mathcal{C}) \hookrightarrow \mathrm{P}\mathrm{Sp}(\mathcal{C})$, write $\mathbb{N}^2 = \mathrm{colim}_{n \in \mathbb{N}} \mathbb{N}_{\leq(n,n)}^2$ as an ascending union of finite pieces $\mathbb{N}_{\leq(n,n)}^2$ and consider the diagram

$$\begin{array}{ccc} \mathrm{P}\mathrm{Sp}(\mathcal{C}) & \subseteq & \lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Fun}(\mathbb{N}_{\leq(n,n)}^2, */\mathcal{C}) \\ \uparrow \text{dashed} & & \uparrow \eta = \lim \eta_n \\ \mathrm{Sp}(\mathcal{C}) & \xrightarrow{\sim} & \lim_{n \in \mathbb{N}^{\mathrm{op}}} */\mathcal{C} \end{array}$$

We need to explain where the upward-pointing arrow $\eta = \lim \eta_n$ comes from: The morphisms

$$\eta_n: */\mathcal{C} \longrightarrow \mathrm{Fun}(\mathbb{N}_{\leq(n,n)}^2, */\mathcal{C})$$

send an object $X \in */\mathcal{C}$ to the right Kan extension of the following functor:

$$\left. \begin{array}{ccccccc} * & \longrightarrow & \cdots & \longrightarrow & * & \longrightarrow & X \\ \uparrow & & & & \uparrow & & \uparrow \\ \vdots & \longrightarrow & * & \longrightarrow & \text{dashed} & \longrightarrow & * \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ * & \longrightarrow & \text{dashed} & \longrightarrow & * & \longrightarrow & \vdots \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \text{dashed} & \longrightarrow & * & \longrightarrow & \cdots & \longrightarrow & * \end{array} \right\} \begin{array}{l} \\ \\ \\ \\ \\ \\ \end{array} \left. \begin{array}{l} \\ \\ \\ \\ \\ \\ \end{array} \right\} n+1$$

$\underbrace{\hspace{10em}}_{n+1}$

Note the blank spots along the diagonal, except for the $(n, n)^{\mathrm{th}}$ entry! So we right-Kan extend along the inclusion $\mathbb{N}_{\leq(n,n)}^2 \setminus \{(0, 0), \dots, (n-1, n-1)\} \subseteq \mathbb{N}_{\leq(n,n)}^2$. Since this inclusion is fully faithful, Corollary I.54 (combined with Proposition/Definition I.58(a)) shows that η_n is fully faithful, and thus also the limit η over all η_n gives a fully faithful map between the limits. One easily checks that the image of η lands in the full subcategory $\mathrm{P}\mathrm{Sp}(\mathcal{C})$ and thus we get the desired fully faithful functor $\mathrm{Sp}(\mathcal{C}) \hookrightarrow \mathrm{P}\mathrm{Sp}(\mathcal{C})$.

As another consequence of the maps η_n being defined by right Kan extension, we obtain canonical equivalences

$$\mathrm{Hom}_{\mathrm{P}\mathrm{Sp}(\mathcal{C})}(X, Y) \simeq \lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Hom}_{*/\mathcal{C}}(X_n, Y_n)$$

for all $X \in \mathrm{P}\mathrm{Sp}(\mathcal{C})$ and $Y \in \mathrm{Sp}(\mathcal{C})$ (just compute the left-hand side as a Hom anima in $\lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Fun}(\mathbb{N}_{\leq(n,n)}^2, */\mathcal{C})$ and use the universal property of right Kan extension).

Note that $\mathrm{P}\mathrm{Sp}(\mathcal{C})$ inherits all limits and colimits from $*/\mathcal{C}$. Also $\mathrm{Sp}(\mathcal{C}) \subseteq \mathrm{P}\mathrm{Sp}(\mathcal{C})$ is closed under limits, but not colimits. To show that $\mathrm{Sp}(\mathcal{C})$ is cocomplete (under appropriate conditions), we need to show that it is in fact a Bousfield localisation of $\mathrm{P}\mathrm{Sp}(\mathcal{C})$. This is done by the following proposition.

II.31. Proposition. — *If $*/\mathcal{C}$ has sequential colimits and $\Omega: */\mathcal{C} \rightarrow */\mathcal{C}$ commutes with them, then the inclusion $\mathrm{Sp}(\mathcal{C}) \subseteq \mathrm{PSp}(\mathcal{C})$ has a left adjoint $(-)^{\mathrm{sp}}: \mathrm{PSp}(\mathcal{C}) \rightarrow \mathrm{Sp}(\mathcal{C})$. On objects it is given by*

$$\Omega^{\infty-i}(X^{\mathrm{sp}}) \simeq \operatorname{colim}_{n \in \mathbb{N}} \Omega^n X_{n+i},$$

where the transition maps are induced by the morphisms $X_i \rightarrow \Omega X_{i+1}$ that were constructed on the previous page.

Before we prove Proposition II.31, we'll discuss some consequences. First, to be able to apply it in the case $\mathcal{C} = \mathrm{An}$, we show:

II.31a. Lemma. — *The loop space functor $\Omega: */\mathrm{An} \rightarrow */\mathrm{An}$ for pointed anima commutes with sequential colimits. In particular, Sp is a Bousfield localisation of PSp and therefore cocomplete by Corollary I.61.*

Since we are already at it, here are two more useful assertions that weren't mentioned in the lecture.

- (a*) *The functors $\Omega^{\infty-i}: \mathrm{Sp} \rightarrow */\mathrm{An}$ commute with sequential colimits for all $i \in \mathbb{Z}$.*
- (b*) *The functors $\pi_i: \mathrm{Sp} \rightarrow \mathrm{Ab}$ for $i \in \mathbb{Z}$ and $\pi_i: */\mathrm{An} \rightarrow \mathrm{Set}$ for $i \geq 0$ commute with sequential colimits. In the latter case, also recall that sequential colimits in Grp (for $i = 1$) or Ab (for $i \geq 2$) are computed on underlying sets.*

Proof.* First of, \mathbb{N} is weakly contractible, hence the colimits can be formed in An instead (Remark* II.10a). We can replace any sequential diagram $(X_1 \rightarrow X_2 \rightarrow \dots)$ in An by a diagram $(X'_1 \hookrightarrow X'_2 \hookrightarrow \dots)$ Kan such that all transition maps are cofibrations; then Theorem I.34 allows us to compute the colimit in Kan instead. Now write $\Omega \simeq F_*((\mathbb{S}^1, *), -)$ as the pointed mapping space out of \mathbb{S}^1 (or any simplicial model for it, it doesn't even need to be Kan) and observe that the right-hand side commutes with filtered colimits in Kan because \mathbb{S}^1 is compact. This shows the first assertion.

Now that we know $\Omega: */\mathrm{An} \rightarrow */\mathrm{An}$ commutes with filtered colimits, we may apply the summary before Corollary II.24 to see that sequential colimits in Sp can be computed degreewise, proving (a*). Hence in (b*) it suffices to show the assertion for $*/\mathrm{An}$. We can write $\pi_i \simeq \pi_0: F_*((\mathbb{S}^i, *), -)$. As in the first part, $F_*((\mathbb{S}^i, *), -)$ commutes with filtered colimits in Kan , and $\pi_0: \mathrm{Kan} \rightarrow \mathrm{Set}$ preserves arbitrary colimits since it is a left adjoint. \square

II.31b. Remark*. — Both assertions of Lemma II.31a remain true if we replace “sequential colimits” by “filtered colimits”, where an indexing ∞ -category \mathcal{I} is *filtered* if every map $K \rightarrow \mathcal{I}$ from a finite simplicial set K extends to a map $K^\triangleright \rightarrow \mathcal{I}$ from its cone.

In fact, the proof can more or less be carried over: Filtered ∞ -categories are weakly contractible (one can use the filteredness condition to show $\pi_i(\mathrm{Ex}^\infty \mathcal{I}, *) = 0$ for all i and all basepoints), we can again replace the given diagram by some cofibrant diagram in the correct model structure on $\mathrm{Fun}(\mathcal{C}[\mathcal{I}], \mathrm{sSet})$, and then use that $F_*((\mathbb{S}^i, *), -)$ preserves filtered colimits by compactness.

Similarly to our construction of the embedding $\mathrm{Sp}(\mathcal{C}) \subseteq \mathrm{PSp}(\mathcal{C})$, one can show that we have an adjunction

$$\tilde{\Sigma}^\infty: */\mathcal{C} \rightleftarrows \mathrm{PSp}(\mathcal{C}) : \mathrm{ev}_{(0,0)}.$$

if \mathcal{C} has finite colimits. On objects $c \in */\mathcal{C}$ (which is all we need by Corollary I.32), we define $\tilde{\Sigma}^\infty c$ as follows: Recall

$$\mathrm{PSp}(\mathcal{C}) \subseteq \lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Fun}(\mathbb{N}_{\leq (n,n)}^2, */\mathcal{C}),$$

so we may imagine $\tilde{\Sigma}^\infty c$ as a sequence $((\tilde{\Sigma}^\infty c)_0, (\tilde{\Sigma}^\infty c)_1, (\tilde{\Sigma}^\infty c)_2, \dots)$. The n^{th} piece $(\tilde{\Sigma}^\infty c)_n$ is then defined to be the left Kan extension of the of the following picture:

$$\mathbb{N}_{\leq (n,n)}^2 \left\{ \begin{array}{ccccccc} * & \longrightarrow & \cdots & \longrightarrow & * & \longrightarrow & \text{---} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \vdots & \longrightarrow & * & \longrightarrow & \text{---} & \longrightarrow & * \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ * & \longrightarrow & \text{---} & \longrightarrow & * & \longrightarrow & \vdots \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ c & \longrightarrow & * & \longrightarrow & \cdots & \longrightarrow & * \end{array} \right\} \begin{array}{l} n+1 \\ n+1 \end{array}$$

Again, note the blank spots along the diagonal, so the Kan extension is along the inclusion $\mathbb{N}_{\leq (n,n)}^2 \setminus \{(1,1), \dots, (n,n)\} \subseteq \mathbb{N}_{\leq (n,n)}^2$ this time.

Combining this with Proposition II.31, we see that $\Omega^\infty: \text{Sp} \rightarrow */\text{An}$ has a left adjoint $\Sigma^\infty = (\tilde{\Sigma}^\infty)^{\text{sp}}: */\text{An} \rightarrow \text{Sp}$. Plugging in the explicit formulas for Kan extension (Theorem I.52) to compute $\tilde{\Sigma}^\infty$, we get

$$\Omega^\infty \Sigma^\infty(X, x) \simeq \text{colim}_{n \in \mathbb{N}} \Omega^n \Sigma^n(X, x)$$

for all $(X, x) \in */\text{An}$. The left-hand side is sometimes denoted $Q(X, x)$, but that has nothing to do with the Quillen Q -construction from page 55).

We will denote the composite

$$\mathbb{S}[-]: \text{An} \xrightarrow{(-)_+} */\text{An} \xrightarrow{\Sigma^\infty} \text{Sp}.$$

In particular, we obtain $\mathbb{S} := \mathbb{S}[*] \simeq \Sigma^\infty(\mathbb{S}^0, *)$, the legendary *sphere spectrum*. Plugging in the adjunctions we know gives $\text{Hom}_{\text{Sp}}(\mathbb{S}, -) \simeq \text{Hom}_{\text{An}}(*, \Omega^\infty -) \simeq \Omega^\infty$, so \mathbb{S} represents the functor $\Omega^\infty: \text{Sp} \rightarrow \text{An}$. Using Lemma II.31a(b), the homotopy groups of \mathbb{S} are given by

$$\pi_i(\mathbb{S}) = \text{colim}_{n \in \mathbb{N}} \pi_i \Omega^n \Sigma^n(\mathbb{S}^0, *) = \text{colim}_{n \in \mathbb{N}} \pi_{i+n}(\mathbb{S}^n, *),$$

i.e. the *stable homotopy groups of spheres*. In contrast to the vastly complicated problem of computing these, the homology of \mathbb{S} (defined as in II.26) is rather boring:

II.31c. Lemma. — *For any abelian group A , the homology of \mathbb{S} with coefficients in A is given by*

$$H_i(\mathbb{S}, A) = \begin{cases} A & \text{if } i = 0 \\ 0 & \text{else} \end{cases}.$$

More generally, we have $C_\bullet(\mathbb{S}[X]) \simeq C_\bullet(X)$ for all $X \in \text{An}$ (where the $C_\bullet(-)$ on the left-hand side is the one from II.26, and the $C_\bullet(-)$ on the right-hand side was defined in Very Long Example I.56), so in particular,

$$H_*(\mathbb{S}[X], A) \simeq H_*(X, A) \quad \text{and} \quad H^*(\mathbb{S}[X], A) \simeq H^*(X, A).$$

*Proof**. Fabian claimed the proof needs a bit of homotopy theory to compute $H_*(\Omega^n \mathbb{S}^n, A)$ in low degrees, but I believe there is also a straightforward way that doesn't need anything. By construction (see II.26) we have

$$C_\bullet \mathbb{S} \simeq \operatorname{colim}_{i \in \mathbb{N}} \tilde{C}_\bullet(\Omega^{\infty-i} \mathbb{S})[-i] \simeq \operatorname{colim}_{i \in \mathbb{N}} \operatorname{colim}_{n \in \mathbb{N}_{\geq i}} \tilde{C}_\bullet(\Omega^{n-i} \mathbb{S}^n)[-i],$$

where we used that $\tilde{C}_\bullet: * / \mathbf{An} \rightarrow \mathcal{D}(\mathbb{Z})$ commutes with colimits. Moreover, the homology functors $H_*: \mathcal{D}(\mathbb{Z}) \rightarrow \mathbf{Ab}$ commute with filtered colimits, hence

$$\begin{aligned} H_*(\mathbb{S}, A) &\simeq H_*(C_\bullet(\mathbb{S}) \otimes_{\mathbb{Z}}^L A) \simeq \operatorname{colim}_{i \in \mathbb{N}} \operatorname{colim}_{n \in \mathbb{N}_{\geq i}} \tilde{H}_{*+i}(\Omega^{n-i} \mathbb{S}^n, A) \\ &\simeq \operatorname{colim}_{n \in \mathbb{N}} \operatorname{colim}_{i \in \mathbb{N}_{\leq n}} \tilde{H}_{*+i}(\Omega^{n-i} \mathbb{S}^n, A) \\ &\simeq \operatorname{colim}_{n \in \mathbb{N}} \tilde{H}_{*+n}(\mathbb{S}^n, A), \end{aligned}$$

where we used that $i = n$ is terminal in $\mathbb{N}_{\leq n}$ to get the third isomorphism. The bottom term clearly vanishes for $* \neq 0$ and equals A for $* = 0$, as we wished to show. The additional assertion follows from the fact that $C_\bullet(-): \mathbf{An} \rightarrow \mathcal{D}(\mathbb{Z})$ and $C_\bullet(\mathbb{S}[-]): \mathbf{An} \rightarrow \mathcal{D}(\mathbb{Z})$ both preserve colimits and agree on $* \in \mathbf{An}$ (as we just checked), hence they must be equal by Theorem I.51. \square

II.32. Corollary (Baratt–Priddy–Quillen). — *The functor $\operatorname{ev}_1: \mathbf{CGrp}(\mathbf{An}) \rightarrow * / \mathbf{An}$ has a left adjoint $\operatorname{Free}^{\mathbf{CGrp}}: * / \mathbf{An} \rightarrow \mathbf{CGrp}(\mathbf{An})$ given by*

$$\operatorname{Free}^{\mathbf{CMon}}(X, x)^{\infty\text{-grp}} \simeq \operatorname{Free}^{\mathbf{CGrp}}(X, x) \simeq \Omega^\infty \Sigma^\infty(X, x).$$

Here $\Omega^\infty: \mathbf{Sp}_{\geq 0} \xrightarrow{\sim} \mathbf{CGrp}(\mathbf{An})$ denotes the inverse of $B^\infty: \mathbf{CGrp}(\mathbf{An}) \xrightarrow{\sim} \mathbf{Sp}_{\geq 0}$, which we recall is an equivalence by Corollary II.24.

*Proof**. We didn't spell this out in the lecture since there isn't much to do: It's evident that $\Sigma^\infty: * / \mathbf{An} \rightarrow \mathbf{Sp}$ lands in the full subcategory $\mathbf{Sp}_{\geq 0}$, hence it's also a left-adjoint to $\Omega^\infty: \mathbf{Sp}_{\geq 0} \rightarrow * / \mathbf{An}$. After applying the other Ω^∞ (i.e. the inverse of B^∞) we obtain the second equivalence. For the first equivalence, recall that Corollary II.21 actually shows that the \mathbb{E}_1 -group completion of an \mathbb{E}_∞ -monoid has a canonical \mathbb{E}_∞ -group structure. \square

Applying Corollary II.32 to the unnumbered example on page 86, we see that

$$\mathfrak{S}^{\infty\text{-grp}} = (\{\text{finite sets, bijections}\}, \sqcup)^{\infty\text{-grp}} \simeq \operatorname{Free}^{\mathbf{CMon}}(*)^{\infty\text{-grp}} \simeq \Omega^\infty \mathbb{S},$$

an example that Fabian already terrified us with in the introduction. In particular, we see that even though $\operatorname{Free}^{\mathbf{Grp}}(*) \simeq \mathbb{Z}$ is a free group on a point and commutative, it fails rather drastically to be the free commutative group on a point.

Proof of Proposition II.31. By Corollary I.32, it suffices to check that $X^{\mathbf{sp}}$ as defined there is really a spectrum and a left-adjoint object to X for all $X \in \mathbf{PSp}(\mathcal{C})$. To check that $X^{\mathbf{sp}}$ is a spectrum, we compute

$$\Omega X_{i+1}^{\mathbf{sp}} \simeq \Omega \operatorname{colim}_{n \in \mathbb{N}} \Omega^n X_{n+i+1} \simeq \operatorname{colim}_{n \in \mathbb{N}} \Omega^{n+1} X_{n+i+1} \simeq \operatorname{colim}_{n \in \mathbb{N}} \Omega^n X_{n+i} \simeq X_i^{\mathbf{sp}}$$

(and all of this is functorial) since Ω commutes with sequential colimits. To check that X^{sp} is really a left-adjoint object of X , we compute

$$\begin{aligned}
\text{Hom}_{\text{Sp}(\mathcal{C})}(X^{\text{sp}}, Y) &\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \text{Hom}_{*/\mathcal{C}}(X_i^{\text{sp}}, Y_i) \\
&\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \left(\text{colim}_{n \in \mathbb{N}} \Omega^n X_{n+i}, Y_i \right) \\
&\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \text{Hom}_{*/\mathcal{C}} \left(\text{colim}_{n \in \mathbb{N}_{\geq i}} \Omega^{n-i} X_n, \Omega^{n-i} Y_n \right) \\
&\simeq \lim_{i \in \mathbb{N}^{\text{op}}} \lim_{n \in \mathbb{N}_{\geq i}^{\text{op}}} \text{Hom}_{*/\mathcal{C}}(\Omega^{n-i} X_n, \Omega^{n-i} Y_n) \\
&\simeq \lim_{n \in \mathbb{N}^{\text{op}}} \lim_{j \in \mathbb{N}_{\leq n}} \text{Hom}_{*/\mathcal{C}}(\Omega^j X_n, \Omega^j Y_n) \\
&\simeq \lim_{n \in \mathbb{N}^{\text{op}}} \text{Hom}_{*/\mathcal{C}}(X_n, Y_n) \\
&\simeq \text{Hom}_{\text{PSp}(\mathcal{C})}(X, Y).
\end{aligned}$$

The first two equivalences are just plugging in definitions. For the third equivalence, we do an index shift by $-i$ and use $\Omega^{n-i} Y_n \simeq Y_i$ since Y is a spectrum. The fourth equivalence is Corollary I.50. For the fifth equivalence, we substitute $j = n - i$ and use that limits commute (by the dual of Proposition I.42). Since j has “negative slope in i ”, we really take the limit over $j \in \mathbb{N}_{\leq n}$; the missing $(-)^{\text{op}}$ is not a typo (and crucial for the next argument). The sixth equivalence now follows from the fact that $j = 0$ is initial in $0 \in \mathbb{N}_{\leq n}$. Finally, the seventh equivalence is the formula for $\text{Hom}_{\text{PSp}(\mathcal{C})}(X, Y)$ that we deduced just before the formulation of Proposition II.31. \square

Our next goal is to define a more “Segal style” model for $\text{Sp}(\mathcal{C})$. To do this, we construct another ∞ -category \mathbb{F}^{op} . Fabian remarks that this symbol is not the `\mathbf{F}`-version of the Roman letter F, but of the archaic Greek letter Digamma, which was the letter after Gamma and Delta in the ancient Greek alphabet, so \mathbb{F}^{op} , Δ^{op} , and \mathbb{F}^{op} fit together nicely. Wikipedia claims, however, that Digamma was only the sixth letter and that Epsilon came between it and Delta. Nevertheless, I’ll obediently stick to Fabian’s notation.

II.33. Definition. — Let $\mathbb{F}^{\text{op}} \subseteq */\text{An}$ be the smallest sub- ∞ -category containing \mathbb{S}^0 and closed under finite colimits. There’s a functor $\mathbb{F}^{\text{op}} \rightarrow \mathbb{F}^{\text{op}}$ sending $\langle n \rangle$ to $\langle n \rangle_+$ (i.e. we add a basepoint to the discrete anima $\langle n \rangle$) and any partially defined map $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ to the map $\alpha_+: \langle m \rangle_+ \rightarrow \langle n \rangle_+$ which is α wherever it is defined and sends everything else to the basepoint.

Finally, if \mathcal{C} is an ∞ -category with finite limits and (thus) a terminal object $*$ in \mathcal{C} , we define $\widetilde{\text{Sp}}(\mathcal{C}) \subseteq \text{Fun}(\mathbb{F}^{\text{op}}, */\mathcal{C})$ to be the full subcategory of *excisive* and *reduced* functors, i.e. those taking pushouts to pullbacks and $*$ to $\langle 0 \rangle_+$ to $*$ in $*/\mathcal{C}$. We might as well define $\widetilde{\text{Sp}}(\mathcal{C})$ as the full sub- ∞ -category of $\text{Fun}(\mathbb{F}^{\text{op}}, \mathcal{C})$ (rather than $\text{Fun}(\mathbb{F}^{\text{op}}, */\mathcal{C})$) spanned by reduced and excisive functors, since every reduced functor $F: \mathbb{F}^{\text{op}} \rightarrow \mathcal{C}$ lifts canonically to $*/\mathcal{C}$.

Note that $\widetilde{\text{Sp}}(\mathcal{C})$ is stable: It has finite limits inherited from $*/\mathcal{C}$ and Ω is an equivalence with inverse induced by precomposition with $\Sigma: \mathbb{F}^{\text{op}} \rightarrow \mathbb{F}^{\text{op}}$. Hence the criterion from Proposition/Definition II.27(a) is fulfilled.

II.34. Proposition. — *With assumptions as in Definition II.33, there is a natural equivalence of ∞ -categories*

$$\begin{aligned} \widetilde{\mathrm{Sp}}(\mathcal{C}) &\xrightarrow{\sim} \mathrm{Sp}(\mathcal{C}) \\ F &\longmapsto (F(\mathbb{S}^0), F(\mathbb{S}^1), \dots). \end{aligned}$$

Proof sketch. There are multiple ways to prove this. One can directly show that $\widetilde{\mathrm{Sp}}(\mathcal{C})$ satisfies the universal property from Theorem II.30. Or one can simply check that both

$$\mathrm{ev}_{\mathbb{S}^0}: \widetilde{\mathrm{Sp}}(\widetilde{\mathrm{Sp}}(\mathcal{C})) \xrightarrow{\sim} \widetilde{\mathrm{Sp}}(\mathcal{C}) \quad \text{and} \quad \mathrm{ev}_{\mathbb{S}^0,*}: \widetilde{\mathrm{Sp}}(\widetilde{\mathrm{Sp}}(\mathcal{C})) \xrightarrow{\sim} \widetilde{\mathrm{Sp}}(\mathcal{C})$$

are equivalences and then apply the same reasoning as in the proof of that theorem.

We will sketch a third way by constructing an inverse functor. For this, we use the following general construction: Let $\mathrm{An}^{\mathrm{fin}} \subseteq \mathrm{An}$ denote the smallest sub- ∞ -category containing $*$ and closed under finite colimits. Let \mathcal{D} be any ∞ -category with finite limits and finite colimits. For $d \in \mathcal{D}$ we put

$$X \otimes d := \mathrm{colim}_X \mathrm{const} d \quad \text{and} \quad d^X := \lim_X \mathrm{const} d.$$

These exist by assumption and are functorial in X and d . If $(X, x) \in \mathbb{F}^{\mathrm{op}}$ and \mathcal{D} is *pointed*, i.e. has a zero object, then we put

$$(X, x) \otimes d := \mathrm{cofib} \left(\mathrm{colim}_{\{x\}} \mathrm{const} d \rightarrow \mathrm{colim}_X d \right).$$

The inverse functor $\mathrm{Sp}(\mathcal{C}) \rightarrow \widetilde{\mathrm{Sp}}(\mathcal{C})$ now sends a spectrum $E \in \mathrm{Sp}(\mathcal{C})$ to the functor $\Omega^\infty(- \otimes E): \mathbb{F}^{\mathrm{op}} \rightarrow */\mathcal{C}$. Note that $\mathrm{Sp}(\mathcal{C})$ has finite colimits since it is stable, so the functor $- \otimes E: \mathbb{F}^{\mathrm{op}} \rightarrow \mathrm{Sp}(\mathcal{C})$ is well-defined. \square

Here's a proposition that was added in the 16th lecture.

II.34a. Proposition. — *In addition to the assumptions from Definition II.33, suppose that $*/\mathcal{C}$ has sequential colimits and that $\Omega: */\mathcal{C} \rightarrow */\mathcal{C}$ commutes with them. Then the inclusion $\widetilde{\mathrm{Sp}}(\mathcal{C}) \subseteq \mathrm{Fun}_*(\mathbb{F}^{\mathrm{op}}, */\mathcal{C})$ into the reduced functors has a left adjoint, sending a reduced functor $F: \mathbb{F}^{\mathrm{op}} \rightarrow */\mathcal{C}$ to*

$$F^{\mathrm{sp}} \simeq \mathrm{colim}_{n \in \mathbb{N}} \Omega^n F(\Sigma^n -).$$

The transition maps in this colimit are induced by the natural transformations $F \Rightarrow \Omega F(\Sigma -)$, which exist since F is reduced.

Proof sketch.* This essential argument is copied from Fabian's notes, [A&HK, Chapter II p. 71]. As usual, by Corollary I.32 all we need to do is to verify that the formula above indeed defines a left-adjoint object of F . If $E: \mathbb{F}^{\mathrm{op}} \rightarrow */\mathcal{C}$ is already reduced and excisive, then $E \simeq \Omega E(\Sigma -)$. Using this and a calculation as in the proof of Proposition II.31, one easily shows $\mathrm{Nat}(F, E) \simeq \mathrm{Nat}(F^{\mathrm{sp}}, E)$.

It remains to see, however, that F^{sp} is indeed excisive (and that's the hard part of the proof). Since Ω commutes with filtered colimits, we get $F^{\text{sp}} \simeq \Omega F^{\text{sp}}(\Sigma -)$. Now consider an arbitrary pushout square in \mathbb{F}^{op} and extend it to a diagram

$$\begin{array}{ccccccc}
 A & \longrightarrow & C & \longrightarrow & * & & \\
 \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & & \\
 B & \longrightarrow & D & \longrightarrow & Q & \longrightarrow & * \\
 \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\
 * & \longrightarrow & P & \longrightarrow & \Sigma A & \longrightarrow & \Sigma B \\
 & & \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\
 & & * & \longrightarrow & \Sigma C & \longrightarrow & \Sigma D
 \end{array}$$

The upper left 2×2 -square induces a map $F^{\text{sp}}(B) \times_{F^{\text{sp}}(D)} F^{\text{sp}}(C) \rightarrow \Omega F^{\text{sp}}(\Sigma A) \simeq F^{\text{sp}}(A)$ and one can check that this is an inverse to the canonical map in the other direction. This shows that F^{sp} turns pushouts into pullbacks, as required. \square

The functors $- \otimes E$ and $E^{(-)}$ for spectra that were constructed in the proof of Proposition II.34 are interesting on their own. Observe for example that $X \otimes HA \simeq H(C_{\bullet}(X) \otimes_{\mathbb{Z}}^L A)$, since both sides are colimit-preserving functors $\text{An} \rightarrow \text{Sp}$ with value HA on $* \in \text{An}$, hence they are equal by Theorem I.51 and the fact that Sp is cocomplete. This motivates the following definition.

II.35. Definition. — Let $E \in \text{Sp}$ and $X \in \text{An}$. The *homology* and *cohomology theory* associated to E are defined by

$$E_*(X) = \pi_*(X \otimes E) = \pi_* \left(\text{colim}_X E \right) \quad \text{and} \quad E^*(X) = \pi_{-*}(E^X) = \pi_{-*} \left(\lim_X E \right).$$

A little bit of foreshadowing: We can also extend the tensor product (or *smash product*) $- \otimes -$ to all of Sp by defining

$$E \otimes E' := \text{colim}_{i \in \mathbb{N}} ((\Omega^{\infty-i} E') \otimes E)[-i]$$

(where $[-i] = \Omega^i$ is the intrinsic loop functor on Sp). It turns out that this defines a symmetric monoidal structure on Sp behaving like $- \otimes_{\mathbb{Z}}^L -$ on $\mathcal{D}(\mathbb{Z})$, although not even symmetry is in any way obvious from the definition. Proving this will be the content of the next few lectures.

In the end (see II.53) we will see that $X \otimes E \simeq \mathbb{S}[X] \otimes E$ and $E^X \simeq \text{hom}_{\text{Sp}}(\mathbb{S}[X], E)$ for all $X \in \text{An}$. If X is an \mathbb{E}_{∞} -monoid, it's usually very hard to understand its group completion $X^{\infty\text{-grp}}$. But we can sometimes understand $\mathbb{S}[X^{\infty\text{-grp}}] \in \text{Sp}$. This is still good enough to compute $H_*(X^{\infty\text{-grp}}, A)$ or even $E_*(X^{\infty\text{-grp}})$, which will be the content of the “group completion theorem”. Once we have this, we're in business to compute some K -groups!

An Interlude on ∞ -Operads

LECTURE 15
17th Dec, 2020

Recall that a morphism in \mathbb{I}^{op} is *inert* if it is a bijection (where it is defined) and *active* if it is defined everywhere. This yields two subcategories $\mathbb{I}_{\text{act}}^{\text{op}}$ and $\mathbb{I}_{\text{int}}^{\text{op}}$. Recall also that symmetric monoidal ∞ -categories are precisely the objects of $\text{CMon}(\text{Cat}_{\infty})$, which is a full subcategory of $\text{Cocart}(\mathbb{I}^{\text{op}})$ via cocartesian unstraightening.

II.36. Definition. — A (*multicoloured and symmetric*) ∞ -operad (*in anima*) is a functor $p: \mathcal{O} \rightarrow \mathbb{I}^{\text{op}}$ of ∞ -categories such that the following conditions hold:

- (a) Every inert in \mathbb{I}^{op} has cocartesian lifts with arbitrary sources (in the invariant sense of Definition I.24(b)).
- (b) The functor $\mathcal{O}: \mathbb{I}_{\text{int}}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ arising by cocartesian straightening satisfies the Segal condition. That is, $\mathcal{O}_0 \simeq *$ and $\mathcal{O}_n \simeq \prod_{i=1}^n \mathcal{O}_1$ via the Segal maps $\rho_1, \dots, \rho_n: \langle n \rangle \rightarrow \langle 1 \rangle$.
- (c) Let $x, y \in \mathcal{O}$ with $p(x) = \langle m \rangle$ and $p(y) = \langle n \rangle$. By (b), we can write $y \simeq (y_1, \dots, y_n)$ for some $y_i \in \mathcal{O}_1$. Then we require that the diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{O}}(x, y) & \xrightarrow{(\rho_1, \dots, \rho_n)} & \prod_{i=1}^n \text{Hom}_{\mathcal{O}}(x, y_i) \\ \downarrow p & \lrcorner & \downarrow p \\ \text{Hom}_{\mathbb{I}^{\text{op}}}(\langle m \rangle, \langle n \rangle) & \xrightarrow{(\rho_1, \dots, \rho_n)} & \prod_{i=1}^n \text{Hom}_{\mathbb{I}^{\text{op}}}(\langle m \rangle, \langle 1 \rangle) \end{array}$$

is a pullback. Note that since the bottom arrow is an injective map of discrete sets, the top arrow must be an inclusion of path components!

We call a morphism $f: x \rightarrow y$ in \mathcal{O} *inert* if it is a p -cocartesian lift of an inert morphism in \mathbb{I} . We call f *active* if it is a (not necessarily p -cocartesian) lift of an active morphism. We also call \mathcal{O}_1 the *underlying ∞ -category* of \mathcal{O} .

Finally, the ∞ -category Op_{∞} of ∞ -operads is the (non-full) sub- ∞ -category of $\text{Cat}_{\infty}/\mathbb{I}^{\text{op}}$ spanned by the ∞ -operads and by functors which preserve inert morphisms.

II.37. Example. — (a) The cocartesian unstraightening induces a functor

$$(-)^{\otimes} = \text{Un}^{\text{cocart}}: \text{CMon}(\text{Cat}_{\infty}) \longrightarrow \text{Op}_{\infty}.$$

To see this, note that conditions Definition II.36(a) and (b) are trivially satisfied, so we only have to check (c). We've already remarked that the bottom row of the diagram in question is an injection of discrete anima, hence we must show that the top row is an inclusion of the corresponding path components. So let $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ be some morphism and denote by $\text{Hom}_{\mathcal{O}}^{\alpha}(x, y)$ the path component over α . Let $x \rightarrow \alpha_*(x)$ be a p -cocartesian lift of α and consider the diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{O}_n}(\alpha_*x, y) & \xrightarrow{\sim} & \prod_{i=1}^n \text{Hom}_{\mathcal{O}_1}((\alpha_*x)_i, y_i) \\ \downarrow \wr & & \downarrow \wr \\ \text{Hom}_{\mathcal{O}}^{\alpha}(x, y) & \longrightarrow & \prod_{i=1}^n \text{Hom}_{\mathcal{O}}^{\rho_i \circ \alpha}(x, y_i) \end{array}$$

That the vertical arrows are equivalences follows easily from the fact that $x \rightarrow \alpha_*x$ is p -cocartesian and the pullback diagram in Definition I.24(a). The top arrow is an equivalence by the Segal condition. Hence the bottom arrow is an equivalence as

well, which is what we wanted to show. Thus $(-)^{\otimes}$ has its image in contained in $\text{Op}_{\infty} \subseteq \text{Cat}_{\infty}/\mathbb{I}^{\text{op}}$, as claimed.

However, $(-)^{\otimes}$ is not fully faithful. In $\text{CMon}(\text{Cat}_{\infty})$ a morphism is a *strongly* symmetric monoidal functor, i.e. it has to preserve the tensor product. We define a *lax symmetric monoidal functor* $\mathcal{C} \rightarrow \mathcal{D}$ to be a map of ∞ -operads $\mathcal{C}^{\otimes} \rightarrow \mathcal{D}^{\otimes}$. We can now define ∞ -categories $\text{SymMonCat}_{\infty} \simeq \text{CMon}(\text{Cat}_{\infty})$ and $\text{SymMonCat}_{\infty}^{\text{lax}} \subseteq \text{Op}_{\infty}$ to be the essential image of $(-)^{\otimes}$.

How do we recognize the essential image of $(-)^{\otimes}$? By the Segal condition and the fact that $p: \mathcal{O} \rightarrow \mathbb{I}^{\text{op}}$ has p -cocartesian lifts for all inert morphisms, it suffices to check that it has p -cocartesian lifts for the unique active maps $f_n: \langle n \rangle \rightarrow \langle 1 \rangle$ (we leave it as an exercise to work out a proper argument). This can in turn be reformulated as follows: For elements $(a_1, \dots, a_n) \in \mathcal{O}_n$ and $b \in \mathcal{O}_1$ define the *multi-morphism space*

$$\begin{array}{ccc} \text{Hom}_{\mathcal{O}}^{\text{act}}((a_1, \dots, a_n), b) & \longrightarrow & \text{Hom}_{\mathcal{O}}((a_1, \dots, a_n), b) \\ \downarrow & \lrcorner & \downarrow \\ * & \xrightarrow{f_n} & \text{Hom}_{\mathbb{I}^{\text{op}}}(\langle n \rangle, \langle 1 \rangle) \end{array}$$

If a morphism $g: (a_1, \dots, a_n) \rightarrow a$ over f_n is p -cocartesian, then the precomposition $g^*: \text{Hom}_{\mathcal{O}_1}(a, -) \xrightarrow{\sim} \text{Hom}_{\mathcal{O}}^{\text{act}}((a_1, \dots, a_n), -)$ is an equivalence of functors $\mathcal{O}_1 \rightarrow \text{An}$. The converse, however, is *not quite* true in the sense that if g^* is an equivalence, then g is only *locally* p -cocartesian by the criterion from [HCII, Chapter IX p. 23]. The way to think about this is that “ $a \simeq a_1 \otimes \dots \otimes a_n$ ”. So if p is a locally cocartesian fibration, then these tensor products exist. But for them to be associative, we need (after some unravelling) that locally p -cocartesian lifts compose. By [HCII, Proposition IX.13], this is precisely the condition we need for the locally cocartesian fibration p to be a cocartesian fibration!

The moral of this story is that, intuitively, an ∞ -operad \mathcal{O} is the data you need to make a “ ∞ -category in which you specify what a map out of a tensor product should be”. Whether such a category exists should be equivalent to whether $\mathcal{O} \simeq \mathcal{C}^{\otimes}$ for some $\mathcal{C} \in \text{CMon}(\text{Cat}_{\infty})$.

- (b) There is an entirely dual theory of *dual ∞ -operads* (be aware that’s not the same thing as ∞ -cooperads). These are the functors to \mathbb{I} satisfying Definition II.36(a), (b), and (c) for cartesian lifts. The resulting ∞ -category dOp_{∞} is equivalent to Op_{∞} via $(-)^{\text{op}}: \text{Op}_{\infty} \xrightarrow{\sim} \text{dOp}_{\infty}$. Note, however, that this doesn’t preserve underlying ∞ -categories. Instead it fits into a diagram

$$\begin{array}{ccc} \text{Op}_{\infty} & \xrightarrow{\sim} & \text{dOp}_{\infty} \\ (-)_1 \downarrow & & \downarrow (-)_1 \\ \text{Cat}_{\infty} & \xrightarrow{(-)^{\text{op}}} & \text{Cat}_{\infty} \end{array}$$

The image of the cartesian unstraightening $\text{Un}^{\text{cart}}: \text{CMon}(\text{Cat}_{\infty}) \rightarrow \text{dOp}_{\infty}$ is denoted $\text{SymMonCat}_{\infty}^{\text{oplax}}$, thought of as symmetric monoidal ∞ -categories with oplax monoidal functors.

- (c) Some trivial examples for ∞ -operads are given by the functors $\mathbb{I}_{\text{int}}^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}, * \xrightarrow{\langle 0 \rangle} \mathbb{I}^{\text{op}}$, and $\text{id}: \mathbb{I}^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$.

- (d) Let $\mathcal{O} \rightarrow \mathbb{I}^{\text{op}}$ be an ∞ -operad and $\mathcal{D} \subseteq \mathcal{O}_1$ a full sub- ∞ -category. Define $\mathcal{D}^{\otimes} \subseteq \mathcal{O}$ to be the full sub- ∞ -category spanned by the $d \simeq (d_1, \dots, d_n)$ with $d_i \in \mathcal{D}$. Then $\mathcal{D}^{\otimes} \rightarrow \mathbb{I}^{\text{op}}$ is an ∞ -operad with underlying ∞ -category \mathcal{D} .

Since I didn't find this completely obvious that \mathcal{D}^{\otimes} is an ∞ -operad, here's a full argument. The conditions from Definition II.36(b) and (c) are immediately inherited from \mathcal{O} . To check the remaining condition (a), we must check that whenever $d \rightarrow d'$ is a cocartesian lift of an inert morphism in \mathbb{I}^{op} with $d \in \mathcal{D}^{\otimes}$, then also $d' \in \mathcal{D}^{\otimes}$. Say $d \in \mathcal{D}_n^{\otimes}$ and let $d \simeq (d_1, \dots, d_n)$. Then the components $d_i \in \mathcal{D}_1^{\otimes}$ are precisely the endpoints of cocartesian lifts $d \rightarrow d_i$ of $\rho_i: \langle n \rangle \rightarrow \langle 1 \rangle$. Since cocartesian lifts compose (see [HCII, Proposition IX.5]), we see that the components of d' are a subset of the components of d , hence $d' \in \mathcal{D}^{\otimes}$ as well. This shows that \mathcal{D}^{\otimes} is indeed an ∞ -operad.

In the case where $\mathcal{O} \simeq \mathcal{C}^{\otimes}$ is a symmetric monoidal ∞ -category, it's naturally to ask whether \mathcal{D}^{\otimes} is one as well. Fabian gave two criteria for this in the lecture. Since these deserve to be stated and proved properly, I outsourced them to Lemma II.37a below.

- (e) Let's already apply Lemma II.37a(a) below to derived tensor products. Let R be a commutative ring. One can check that $\text{Ch}(R)$ is symmetric monoidal under $- \otimes_R -$ as a Kan-enriched category, hence its coherent nerve $\mathcal{K}(R)$ is symmetric monoidal. Now recall from Example I.62(c) that $\mathcal{D}(R)$ is a Bousfield localisation of $\mathcal{K}(R)$. More precisely, $\mathcal{D}(R)$ can be embedded fully faithfully into $\mathcal{K}(R)$ as the full sub- ∞ -category $\mathcal{D}(R)^{K\text{-proj}} \subseteq \mathcal{K}(R)$ of K -projective complexes, i.e. those C such that $\text{Hom}_{\mathcal{K}(R)}(C, -)$ inverts quasi-isomorphisms. Clearly the tensor unit $R[0]$ is K -projective (as is any bounded below degree-wise projective complex) and if C and D are K -projective, then so is $C \otimes_R D$ as $\text{Hom}_{\mathcal{K}(R)}(C \otimes_R D, -) \simeq \text{Hom}_{\mathcal{K}(R)}(C, \underline{\text{Hom}}_R(D, -))$ inverts quasi-isomorphisms too.

We may thus apply Lemma II.37a(a) to obtain a symmetric monoidal structure $- \otimes_R^L -$ on $\mathcal{D}(R)$. The localisation functor $\mathcal{K}(R) \rightarrow \mathcal{D}(R)$ ("taking K -projective resolutions") is lax symmetric monoidal in the sense explained in (a). In particular, there are canonical maps $C \otimes_R^L D \rightarrow C \otimes_R D$ for all $C, D \in \mathcal{K}(R)$.

- (f) In general, there is an equivalence of ∞ -categories

$$\{\text{oplax monoidal left adjoints } \mathcal{C} \rightarrow \mathcal{D}\} \simeq \{\text{lax monoidal right adjoints } \mathcal{D} \rightarrow \mathcal{C}\}.$$

This is not trivial since lax and oplax structures are defined in different fibration pictures. The first written proofs appeared in November 2020 by Rune Haugseng [Hau20] and independently by Fabian, Sil, and Joost Nuiten [HLN20].

Here's the lemma that we outsourced above:

II.37a. Lemma. — *Let $\mathcal{D}^{\otimes} \subseteq \mathcal{C}^{\otimes}$ be defined as in Example II.37(d).*

- (a) *Suppose that $d \otimes d' \in \mathcal{D}$ for all $d, d' \in \mathcal{D}$ and $1_{\mathcal{C}} \in \mathcal{D}$. Then \mathcal{D}^{\otimes} is a symmetric monoidal ∞ -category and the inclusion $\mathcal{D} \subseteq \mathcal{C}$ is strongly monoidal. If it has, moreover, a right adjoint $R: \mathcal{C} \rightarrow \mathcal{D}$, then R can be extended to a map of ∞ -operads $R^{\otimes}: \mathcal{C}^{\otimes} \rightarrow \mathcal{D}^{\otimes}$ such that R^{\otimes} is right-adjoint to the inclusion $\mathcal{D}^{\otimes} \subseteq \mathcal{C}^{\otimes}$. In particular, R is lax symmetric monoidal.*
- (b) *Suppose that $\mathcal{D} \subseteq \mathcal{C}$ has a left adjoint $L: \mathcal{C} \rightarrow \mathcal{D}$ with the property that whenever $L(f): L(x) \xrightarrow{\sim} L(y)$ is an equivalence, then $L(f \otimes \text{id}_z): L(x \otimes z) \xrightarrow{\sim} L(y \otimes z)$ is an equivalence too for all $z \in \mathcal{C}$. Then \mathcal{D}^{\otimes} is symmetric monoidal and L extends to a map $L^{\otimes}: \mathcal{C}^{\otimes} \rightarrow \mathcal{D}^{\otimes}$ of ∞ -operads such that L^{\otimes} is left-adjoint to $\mathcal{D}^{\otimes} \subseteq \mathcal{C}^{\otimes}$. Moreover, L is strongly symmetric monoidal and $\mathcal{D} \subseteq \mathcal{C}$ is lax symmetric monoidal.*

*Proof**. For (a), the fact that \mathcal{D}^\otimes is a symmetric monoidal ∞ -category and that $\mathcal{D} \subseteq \mathcal{C}$ is a strongly monoidal functor (i.e. $\mathcal{D}^\otimes \subseteq \mathcal{C}^\otimes$ preserves all cocartesian edges) follows from the recognition criterion sketched in Example II.37(a). To construct R^\otimes , we employ Corollary I.32 as usual to see that it can be constructed pointwise. So let $c \in \mathcal{C}^\otimes$ and write $c = (c_1, \dots, c_n)$. We put $R^\otimes(c) = (R(c_1), \dots, R(c_n))$ and claim that the natural map $\eta_c: c \rightarrow R^\otimes(c)$ (induced by the unit maps $\eta_{c_i}: c_i \rightarrow R(c_i)$ for R) witnesses it as a right-adjoint object of c . That is, we must show that

$$\eta_{c,*}: \mathrm{Hom}_{\mathcal{C}^\otimes}(d, c) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}^\otimes}(d, R^\otimes(c))$$

is an equivalence for all $d \in \mathcal{D}^\otimes$. By Definition II.36(c) it suffices to show the same for $\eta_{c_i,*}: \mathrm{Hom}_{\mathcal{C}^\otimes}(d, c_i) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}^\otimes}(d, R(c_i))$. But whether this is an equivalence of anima can be checked on (derived) fibres over $\mathrm{Hom}_{\mathbb{T}^\mathrm{op}}(\langle n \rangle, \langle 1 \rangle)$. Given $\alpha: \langle n \rangle \rightarrow \langle 1 \rangle$ in \mathbb{T}^op , choose a cocartesian lift $d \rightarrow \alpha_* d$ to \mathcal{D}^\otimes , which is also cocartesian in \mathcal{C}^\otimes by construction. Then the fibres over α are given by $\mathrm{Hom}_{\mathcal{C}^\otimes}^\alpha(d, c_i) \simeq \mathrm{Hom}_{\mathcal{C}}(\alpha_* d, c_i)$ and $\mathrm{Hom}_{\mathcal{D}^\otimes}^\alpha(d, R(c_i)) \simeq \mathrm{Hom}_{\mathcal{D}}(\alpha_* d, c_i)$, hence $\eta_{c_i,*}$ is indeed an equivalence since $R: \mathcal{C} \rightarrow \mathcal{D}$ is a right adjoint of the inclusion $\mathcal{D} \subseteq \mathcal{C}$. This shows that R^\otimes exist as a map in Cat_∞ . It's clear from the construction that it's also a map in $\mathrm{Cat}_\infty/\mathbb{T}^\mathrm{op}$. Finally, since cocartesian lifts of inert morphisms just “forget” some of the factors, they are preserved by R^\otimes , whence it is a map of ∞ -operads. This shows (a).

Now for (b). First observe that the condition on L can be strengthened as follows: If we are given equivalences $L(f_i): L(x_i) \xrightarrow{\sim} L(y_i)$ in \mathcal{D} for all $i = 1, \dots, n$, then also $L(f_1 \otimes \dots \otimes f_n): L(x_1 \otimes \dots \otimes x_n) \xrightarrow{\sim} L(y_1 \otimes \dots \otimes y_n)$ is an equivalence. To show that \mathcal{D}^\otimes is a symmetric monoidal ∞ -category, we use the recognition criterion from Example II.37(a): Given $(d_1, \dots, d_n) \in \mathcal{D}_n^\otimes$ and any $d \in \mathcal{D}$, we have

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}^\otimes}^{\mathrm{act}}((d_1, \dots, d_n), d) &\simeq \mathrm{Hom}_{\mathcal{C}^\otimes}^{\mathrm{act}}((d_1, \dots, d_n), d) \\ &\simeq \mathrm{Hom}_{\mathcal{C}}(d_1 \otimes \dots \otimes d_n, d) \\ &\simeq \mathrm{Hom}_{\mathcal{D}}(L(d_1 \otimes \dots \otimes d_n), d), \end{aligned}$$

hence composing $(d_1, \dots, d_n) \rightarrow d_1 \otimes \dots \otimes d_n$ with the unit map $d_1 \otimes \dots \otimes d_n \rightarrow L(d_1 \otimes \dots \otimes d_n)$ in \mathcal{C}^\otimes gives a locally cocartesian lift of $f_n: \langle n \rangle \rightarrow \langle 1 \rangle$ to \mathcal{D}^\otimes . Hence $\mathcal{D}^\otimes \rightarrow \mathbb{T}^\mathrm{op}$ is a locally cocartesian fibration. To check that the locally cocartesian edges compose, we must verify that $L(L(x \otimes y) \otimes z) \simeq L(x \otimes y \otimes z)$. But $x \otimes y \rightarrow L(x \otimes y)$ becomes an equivalence after applying L , as $L^2 \simeq L$, hence this follows from the assumption on L . Hence $\mathcal{D}^\otimes \rightarrow \mathbb{T}^\mathrm{op}$ is even a cocartesian fibration by [HCII, Proposition IX.13], which precisely says that \mathcal{D}^\otimes is a symmetric monoidal ∞ -category.

To construct L^\otimes , we proceed as in (a) and define it pointwise via $L^\otimes(c_1, \dots, c_n) = (L(c_1), \dots, L(c_n))$. By the same tricks as above, showing that this is indeed a right-adjoint object of (c_1, \dots, c_n) reduces to showing

$$\mathrm{Hom}_{\mathcal{C}^\otimes}^\alpha((c_1, \dots, c_n), d) \simeq \mathrm{Hom}_{\mathcal{D}^\otimes}^\alpha((L(c_1), \dots, L(c_n)), d)$$

for all $d \in \mathcal{D}$ and all $\alpha: \langle n \rangle \rightarrow \langle 1 \rangle$. Let $(c_1, \dots, c_n) \rightarrow c$ be a cocartesian lift of α to \mathcal{C}^\otimes . Then $\mathrm{Hom}_{\mathcal{C}^\otimes}^\alpha((c_1, \dots, c_n), d) \simeq \mathrm{Hom}_{\mathcal{C}}(c, d)$ and also $c \simeq c_{i_1} \otimes \dots \otimes c_{i_m}$, where $i_1, \dots, i_m \in \langle n \rangle$ are the elements where α is defined. The assumption on L then implies

$$L(c) \simeq L(L(c_{i_1}) \otimes \dots \otimes L(c_{i_m})),$$

because each $c_{i_j} \rightarrow L(c_{i_j})$ becomes an equivalence after applying L . Hence a cocartesian lift of α to \mathcal{D}^\otimes is given by $(L(c_1), \dots, L(c_n)) \rightarrow L(c)$ and $\mathrm{Hom}_{\mathcal{D}^\otimes}^\alpha((L(c_1), \dots, L(c_n)), d) \simeq$

$\mathrm{Hom}_{\mathcal{D}}(L(c), d)$. Now $\mathrm{Hom}_{\mathcal{C}}(c, d) \simeq \mathrm{Hom}_{\mathcal{D}}(L(c), d)$ because $L: \mathcal{C} \rightarrow \mathcal{D}$ is left-adjoint to the inclusion $\mathcal{D} \subseteq \mathcal{C}$, which finally shows the desired equivalence. It's clear from the construction that L^{\otimes} preserves cocartesian edges, whence L is indeed strongly symmetric monoidal. \square

The following lemma is closely related, but wasn't mentioned in the lecture, although we'll need it later. It roughly says that whether a left adjoint is strongly monoidal can be checked on objects.

II.37b. Lemma*. — *Let $R^{\otimes}: \mathcal{D}^{\otimes} \rightarrow \mathcal{C}^{\otimes}$ be a map of ∞ -operads (hence a lax symmetric monoidal functor between \mathcal{D} and \mathcal{C}). If the underlying functor $R: \mathcal{D} \rightarrow \mathcal{C}$ has a left adjoint $L: \mathcal{C} \rightarrow \mathcal{D}$ satisfying $L(1_{\mathcal{C}}) \simeq 1_{\mathcal{D}}$ and $L(c_1 \otimes_{\mathcal{C}} c_2) \simeq L(c_1) \otimes_{\mathcal{D}} L(c_2)$ for all $c, c' \in \mathcal{C}$, then L extends to a map $L^{\otimes}: \mathcal{C}^{\otimes} \rightarrow \mathcal{D}^{\otimes}$ of ∞ -operads, which is left-adjoint to R^{\otimes} and witnesses L as a strongly symmetric monoidal functor.*

Proof sketch.* The condition on L implies $L(c_1 \otimes_{\mathcal{C}} \cdots \otimes_{\mathcal{C}} c_n) \simeq L(c_1) \otimes_{\mathcal{D}} \cdots \otimes_{\mathcal{D}} L(c_n)$ for all $c_1, \dots, c_n \in \mathcal{C}$. We define L^{\otimes} pointwise as in Lemma II.37a(b). By the same arguments as given there, all we need to check is that

$$\mathrm{Hom}_{\mathcal{C}^{\otimes}}((c_1, \dots, c_n), R(d)) \simeq \mathrm{Hom}_{\mathcal{D}^{\otimes}}((L(c_1), \dots, L(c_n)), d)$$

for all $c_1, \dots, c_n \in \mathcal{C}$, $d \in \mathcal{D}$ and all $\alpha: \langle n \rangle \rightarrow \langle 1 \rangle$. If α is defined at $i_1, \dots, i_m \in \langle n \rangle$, then $(c_1, \dots, c_n) \rightarrow c_{i_1} \otimes_{\mathcal{C}} \cdots \otimes_{\mathcal{C}} c_{i_m}$ is a cocartesian lift of α to \mathcal{C}^{\otimes} , hence

$$\mathrm{Hom}_{\mathcal{C}^{\otimes}}((c_1, \dots, c_n), R(d)) \simeq \mathrm{Hom}_{\mathcal{C}}(c_{i_1} \otimes_{\mathcal{C}} \cdots \otimes_{\mathcal{C}} c_{i_m}, R(d)).$$

The same argument for \mathcal{D}^{\otimes} together with the condition on L implies

$$\mathrm{Hom}_{\mathcal{D}^{\otimes}}((L(c_1), \dots, L(c_n)), d) \simeq \mathrm{Hom}_{\mathcal{D}}(L(c_1 \otimes_{\mathcal{C}} \cdots \otimes_{\mathcal{C}} c_n), d),$$

whence we are done by the fact that L and R are adjoints. This shows that L^{\otimes} is an adjoint of R^{\otimes} , as desired. It's straightforward to check that L^{\otimes} is a map over \mathbb{T}^{op} and preserves cocartesian edges, hence L is indeed strongly symmetric monoidal. \square

Towards the Tensor Product of Spectra

Our next goal is to make cartesian and cocartesian monoidal structures on ∞ -categories with products, and to stabilize them to eventually define the symmetric monoidal structure on Sp . While Lurie writes down explicit simplicial sets to construct the desired ∞ -operads, we will cheat a bit and use complete Segal spaces instead. This is based on the following lemma.

II.38. Lemma. — *Let \mathcal{E} be an ∞ -category. For a presheaf $X \in \mathcal{P}(\mathcal{E})$, consider the slice category \mathcal{E}/X with respect to the Yoneda embedding $Y^{\mathcal{E}}: \mathcal{E} \rightarrow \mathcal{P}(\mathcal{E})$ and $\{X\} \rightarrow \mathcal{P}(\mathcal{E})$ (in the notation of Lemma* I.53a this would have been $Y^{\mathcal{E}}/X$ instead). The canonical functor*

$$\mathcal{P}(\mathcal{E}/X) \xrightarrow{\sim} \mathcal{P}(\mathcal{E})/X,$$

constructed as the unique colimit-preserving extension of $Y^{\mathcal{E}}: \mathcal{E}/X \rightarrow \mathcal{P}(\mathcal{E})/X$ via Theorem I.51, is an equivalence of ∞ -categories.

Proof. Before we start, observe that $\mathcal{P}(\mathcal{E})/X$ is indeed cocomplete since one immediately checks that it inherits colimits from $\mathcal{P}(\mathcal{E})$. We will use the following criterion:

(\boxtimes) Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor from a small ∞ -category to a cocomplete ∞ -category. Let $| _F: \mathcal{P}(\mathcal{C}) \rightleftarrows \mathcal{D} : \text{Sing}_F$ denote the adjunction resulting from Theorem I.51. Then $| _F$ is an equivalence if the following three conditions hold:

- (a) F is fully faithful.
- (b) $\text{Hom}_{\mathcal{D}}(F(c), -): \mathcal{D} \rightarrow \mathbf{An}$ commutes with colimits for all $c \in \mathcal{C}$.
- (c) $\text{Sing}_F: \mathcal{D} \rightarrow \mathcal{P}(\mathcal{C})$ is conservative.

We prove (\boxtimes) first. Condition (b) implies that Sing_F commutes with colimits. Indeed, this can be checked pointwise by Theorem I.19(b) since there always is a natural transformation $\text{colim}_{\mathcal{I}} \text{Sing}_F \Rightarrow \text{Sing}_F \text{colim}_{\mathcal{I}}$. Moreover, colimits in $\mathcal{P}(\mathcal{C})$ are computed pointwise. Plugging in the explicit description of Sing_F that Corollary I.55 offers us, we must thus check that $\text{colim}_{\mathcal{I}} \text{Hom}_{\mathcal{D}}(F(c), d_i) \simeq \text{Hom}_{\mathcal{D}}(F(c), \text{colim}_{\mathcal{I}} d_i)$ for all $c \in \mathcal{C}$ and $d_i \in \mathcal{D}$. This is (b).

Now since F is fully faithful, the unit $\text{id}_{\mathcal{P}(\mathcal{C})} \Rightarrow \text{Sing}_F | _F$ is an equivalence on representables. Indeed, the right-hand side sends $c \in \mathcal{C}$ to $\text{Hom}_{\mathcal{D}}(F(c), F(c)) \simeq \text{Hom}_{\mathcal{C}}(c, c)$ by Corollary I.55 again. Now both sides preserve colimits and every presheaf on \mathcal{C} can be written as a colimit of representables by Lemma* I.53a, hence the unit is an equivalence everywhere. By the triangle identities and the fact that Sing_F is conservative by (c), this implies that the counit $| _F \text{Sing}_F \Rightarrow \text{id}_{\mathcal{D}}$ is an equivalence as well. This finishes the proof of (\boxtimes).

Now to check that (\boxtimes) is applicable for $\mathcal{C} = \mathcal{E}/X$ and $\mathcal{D} = \mathcal{P}(\mathcal{E})/X$. We already argued that $\mathcal{P}(\mathcal{E})/X$ is cocomplete. It's easily checked that $\mathcal{E}/X \rightarrow \mathcal{P}(\mathcal{E})/X$ is fully faithful since it's induced by the fully faithful Yoneda embedding, so (a) holds. To check (c), let $\alpha: F \rightarrow F'$ be a morphism in $\mathcal{P}(\mathcal{E})/X$ such that $\alpha_*: \text{Hom}_{\mathcal{P}(\mathcal{E}/X)}(-, F) \rightarrow \text{Hom}_{\mathcal{P}(\mathcal{E}/X)}(-, F')$ is an equivalence on representable presheaves. We must check that α is an equivalence itself. But if α_* is an equivalence on representables, then the Yoneda lemma implies that $\alpha: F(e) \rightarrow F'(e)$ is an equivalence for all $e \in \mathcal{E}$ (except those $e \in \mathcal{E}$ for which $X(e) = \emptyset$ since these aren't contained in the image of $\mathcal{E}/X \rightarrow \mathcal{E}$, but in that case we have $F(e) = \emptyset = F'(e)$ too since F and F' come with a map to X), hence α is an equivalence of presheaves.

Finally, we have to check (b). Given a representable object $\text{Hom}_{\mathcal{E}}(e, -) \rightarrow X$ and an arbitrary object $F \rightarrow X$ of $\mathcal{P}(\mathcal{E})/X$, the Yoneda lemma and [HCII, Corollary VIII.6] provide a pullback square

$$\begin{array}{ccccc} \text{Hom}_{\mathcal{P}(\mathcal{E})/X}(\text{Hom}_{\mathcal{E}}(e, -), F) & \longrightarrow & \text{Hom}_{\mathcal{P}(\mathcal{E})}(\text{Hom}_{\mathcal{E}}(e, -), F) & \xrightarrow{\sim} & F(e) \\ \downarrow & \lrcorner & \downarrow & & \downarrow \\ * & \longrightarrow & \text{Hom}_{\mathcal{P}(\mathcal{E})}(\text{Hom}_{\mathcal{E}}(e, -), X) & \xrightarrow{\sim} & X(e) \end{array}$$

Clearly $F(e)$ commutes with colimits in F because colimits in functor categories are computed pointwise (Lemma I.39). Hence it suffices to check that colimits in \mathbf{An} commute with pullbacks. In other words, if $f: K \rightarrow L$ is any morphism of anima, then $f^*: \mathbf{An}/L \rightarrow \mathbf{An}/K$ preserves colimits. But $\text{Right}(K) \simeq \mathbf{An}/K$ since every morphism to K can be factored into a right anodyne and a right fibration, and likewise for L , so we are done since we verified in Lemma* I.42a that $f^*: \text{Right}(L) \rightarrow \text{Right}(K)$ has a right adjoint f_* . \square

II.38a. — Applying Lemma II.38 to $\mathcal{E} = \Delta$ and $X = N^r(\mathcal{C}) \in \mathbf{sAn}$ for some ∞ -category \mathcal{C} , we obtain $\mathcal{P}(\Delta/N^r(\mathcal{C})) \simeq \mathbf{sAn}/N^r(\mathcal{C})$. We will usually abbreviate the left-hand side as $\mathcal{P}(\Delta/\mathcal{C})$. If you think about it, this makes a lot of sense: Objects of $\Delta/N^r(\mathcal{C})$ are maps $\Delta^n \rightarrow N^r(\mathcal{C})$ of simplicial anima. But $N^r: \text{Cat}_{\infty} \rightarrow \mathbf{sAn}$ is fully faithful by Theorem/Definition I.64, so we may equivalently think of maps $[n] \rightarrow \mathcal{C}$.

We can now specify objects of (or functors to) $\text{Cat}_\infty/\mathcal{C}$ as objects of (or functors to) $\mathcal{P}(\Delta/\mathcal{C})$ and then compose with

$$\mathcal{P}(\Delta/\mathcal{C}) \simeq \text{sAn}/N^r(\mathcal{C}) \xrightarrow{\text{asscat}} \text{Cat}_\infty/\mathcal{C}.$$

This will be done quite a number of times on the next few pages.

LECTURE 16
22nd Dec, 2020

II.38b. — To check that an object on the left is complete Segal, we will check the *relative Segal condition*. A map $Y \rightarrow X$ in sAn is called *relative Segal* if the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{sAn}}(\Delta^n, Y) & \longrightarrow & \text{Hom}_{\text{sAn}}(I^n, Y) \\ \downarrow & \lrcorner & \downarrow \\ \text{Hom}_{\text{sAn}}(\Delta^n, X) & \longrightarrow & \text{Hom}_{\text{sAn}}(I^n, X) \end{array}$$

is a pullback in An . If X is a Segal anima, then the lower row is an equivalence. Thus Y is a Segal anima too if $Y \rightarrow X$ is relative Segal.

Likewise, a morphism $Y \rightarrow X$ is called *relatively complete Segal* if it is relative Segal and the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{sAn}}(\Delta^0, Y) & \longrightarrow & \text{Hom}_{\text{sAn}}(\Delta^3/(\Delta^{\{0,2\}} \cup \Delta^{\{1,3\}}), Y) \\ \downarrow & \lrcorner & \downarrow \\ \text{Hom}_{\text{sAn}}(\Delta^0, X) & \longrightarrow & \text{Hom}_{\text{sAn}}(\Delta^3/(\Delta^{\{0,2\}} \cup \Delta^{\{1,3\}}), X) \end{array}$$

Again, if this holds and X is a complete Segal anima, then so is Y (here we use a slight reformulation of Theorem/Definition I.64(b)).

With that let's dive into cocartesian monoidal structure:

II.39. Construction. — Let $\mathbb{I}^{\text{op}}_\sqcup \subseteq \langle 1 \rangle / \mathbb{I}^{\text{op}}$ be the full subcategory spanned by the “defined maps” (for every n there's a unique nowhere-defined map $\langle 1 \rangle \rightarrow \langle n \rangle$; this is the only one we exclude). For any ∞ -category \mathcal{C} define $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ via II.38a applied to the presheaf $F_{\mathcal{C}^\sqcup}: \Delta/\mathbb{I}^{\text{op}} \rightarrow \text{An}$, which is given by

$$F_{\mathcal{C}^\sqcup}([n] \rightarrow \mathbb{I}^{\text{op}}) \simeq \text{Hom}_{\text{Cat}_\infty}([n] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}_\sqcup, \mathcal{C})$$

on objects (and turned into a functor of ∞ -categories in the obvious way).

II.40. Proposition. — *With notation as in Construction II.39:*

- (a) *The presheaf $F_{\mathcal{C}^\sqcup}$ defines a complete Segal anima over $N^r(\mathbb{I}^{\text{op}})$ via Lemma II.38.*
- (b) *$p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ is always an ∞ -operad with underlying category \mathcal{C} .*
- (c) *An active morphism $(a_1, \dots, a_n) \rightarrow b$ in \mathcal{C}^\sqcup is locally p^\sqcup -cocartesian iff the maps $a_i \rightarrow b$ exhibit b as the coproduct of the a_i .*
- (d) *In particular, $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ is a symmetric monoidal category iff \mathcal{C} has finite coproducts.*

Proof sketch. The proof of (a) was left as an exercise, so here's what I figured out. To show that $F_{\mathcal{C}^\sqcup}$ is Segal, consider the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{sAn}}(\Delta^n, F_{\mathcal{C}^\sqcup}) & \longrightarrow & \text{Hom}_{\text{sAn}}(I^n, F_{\mathcal{C}^\sqcup}) \\ \downarrow & & \downarrow \\ \text{Hom}_{\text{sAn}}(\Delta^n, N^r(\mathbb{I}^{\text{op}})) & \xrightarrow{\sim} & \text{Hom}_{\text{sAn}}(I^n, N^r(\mathbb{I}^{\text{op}})) \end{array}$$

The bottom row is an equivalence since $N^r(\mathbb{I}^{\text{op}})$ is Segal. To show that the top row is an equivalence as well, it suffices to check that it induces equivalences on all (derived) fibres. Unravelling what that means (using the computation of Hom anima in slice categories from [HCII, Corollary VIII.6]), we need to check that

$$\text{Hom}_{\text{sAn}/N^r(\mathbb{I}^{\text{op}})}(\Delta^n, F_{\mathcal{C}^\sqcup}) \xrightarrow{\sim} \text{Hom}_{\text{sAn}/N^r(\mathbb{I}^{\text{op}})}(I^n, F_{\mathcal{C}^\sqcup}) \simeq \prod_{i=1}^n \text{Hom}_{\text{sAn}/N^r(\mathbb{I}^{\text{op}})}(\Delta^1, F_{\mathcal{C}^\sqcup})$$

is an equivalence for all choices of $\Delta^n \rightarrow N^r(\mathbb{I}^{\text{op}})$. Recall that $\text{sAn}/N^r(\mathbb{I}^{\text{op}}) \simeq \mathcal{P}(\Delta/\mathbb{I}^{\text{op}})$ by Lemma II.38. Using this together with the Yoneda lemma and the definition of $F_{\mathcal{C}^\sqcup}$, our assertion translates into

$$\text{Hom}_{\text{Cat}_\infty}([n] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) \xrightarrow{\sim} \prod_{i=1}^n \text{Hom}_{\text{Cat}_\infty}([1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}).$$

In other words, we must prove that $[n] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}$ is an iterated pushout of $[1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}$ in Cat_∞ . One way to do this is to simply check by hand that

$$I^n \times_{N(\mathbb{I}^{\text{op}})} N(\mathbb{I}^{\text{op}}) \hookrightarrow \Delta^n \times_{N(\mathbb{I}^{\text{op}})} N(\mathbb{I}^{\text{op}})$$

is an inner anodyne map of simplicial sets. This shows that $F_{\mathcal{C}^\sqcup}$ is Segal. Completeness can be shown analogously, whence we have proved (a).

To prove (b), we must investigate the fibres of $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$. So fix some $\langle n \rangle \in N^r(\mathbb{I}^{\text{op}})$. Since $F_{\mathcal{C}^\sqcup}$ is complete Segal by (a), we get $F_{\mathcal{C}^\sqcup} \simeq N^r(\text{asscat } F_{\mathcal{C}^\sqcup}) \simeq N^r(\mathcal{C}^\sqcup)$. Because N^r preserves limits, we may as well compute the corresponding fibre of $F_{\mathcal{C}^\sqcup} \rightarrow N^r(\mathbb{I}^{\text{op}})$ and then take $\text{asscat}(-)$ again. To do this, we claim that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{P}(\Delta/\mathbb{I}^{\text{op}}) & \xrightarrow{\sim} & \text{sAn}/N^r(\mathbb{I}^{\text{op}}) \\ & \searrow \text{evaluation at} & \swarrow - \times_{N^r(\mathbb{I}^{\text{op}})} \{\langle n \rangle\} \\ & \text{constant maps} & \text{sAn} \end{array}$$

The right diagonal arrow sends a simplicial anima X over $N^r(\mathbb{I}^{\text{op}})$ to $X \times_{N^r(\mathbb{I}^{\text{op}})} \{\langle n \rangle\}$ (i.e. to the fibre we're interested in). The left diagonal arrow sends a presheaf $F: \Delta/\mathbb{I}^{\text{op}} \rightarrow \text{An}$ to a simplicial anima Y defined by $Y_k \simeq F(\text{const } \langle n \rangle: [k] \rightarrow \mathbb{I}^{\text{op}})$. To show commutativity, we may use Theorem I.51, whence it suffices to check that both ways around the diagram agree on representable presheaves and preserve colimits. The former is straightforward to check (although it took me some time to wrap my head around this). For the latter, it's clear that the left diagonal arrow preserves colimits, and for the right diagonal arrow we can use that colimits in anima commute with pullbacks (as seen in the proof of Lemma II.38) and that colimits and pullbacks in sAn are computed pointwise.

Let's apply our newfound knowledge to $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$. It's straightforward to check that the diagram

$$\begin{array}{ccc} [k] \times \langle n \rangle & \longrightarrow & \mathbb{I}^{\text{op}} \\ \text{pr}_1 \downarrow & \lrcorner & \downarrow \\ [k] & \xrightarrow{\text{const } \langle n \rangle} & \mathbb{I}^{\text{op}} \end{array}$$

is a pullback diagram. In particular, the fibre of $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ over $\langle n \rangle$ is the associated category of a simplicial anima Y given by

$$\begin{aligned} Y_k &\simeq F_{\mathcal{C}^\sqcup}(\text{const } \langle n \rangle: [k] \rightarrow \mathbb{I}^{\text{op}}) \simeq \text{Hom}_{\text{Cat}_\infty}([k] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) \\ &\simeq \text{Hom}_{\text{Cat}_\infty}([k] \times \langle n \rangle, \mathcal{C}) \\ &\simeq \text{Hom}_{\text{Cat}_\infty}([k], \mathcal{C}^n) \end{aligned}$$

Up to checking some functoriality stuff, this shows $Y \simeq N^r(\mathcal{C}^n)$, so the fibre of $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ over $\langle n \rangle$ is \mathcal{C}^n . This shows that p^\sqcup satisfies Definition II.36(b).

Two more conditions are to check. Let $x, y \in \mathcal{C}^\sqcup$ and let $\alpha: p^\sqcup(x) \rightarrow p^\sqcup(y)$ be some morphism in \mathbb{I}^{op} . Then the fibre $\text{Hom}_{\mathcal{C}^\sqcup}^\alpha(x, y) := \text{Hom}_{\mathcal{C}^\sqcup}(x, y) \times_{\text{Hom}_{\mathbb{I}^{\text{op}}}(p^\sqcup(x), p^\sqcup(y))} \{\alpha\}$ of $\text{Hom}_{\mathcal{C}^\sqcup}(x, y)$ over α has a particularly simple description: If we think of α as a functor $\alpha: [1] \rightarrow \mathbb{I}^{\text{op}}$, then $\text{Hom}_{\mathcal{C}^\sqcup}^\alpha(x, y)$ fits into a pullback

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}^\sqcup}^\alpha(x, y) & \longrightarrow & \text{Hom}_{\text{Cat}_\infty}([1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) \\ \downarrow & \lrcorner & \downarrow (d_1, d_0) \\ * & \xrightarrow{(x, y)} & \text{Hom}_{\text{Cat}_\infty}([0] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) \times \text{Hom}_{\text{Cat}_\infty}([0] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) \end{array}$$

To prove this, use Yoneda's lemma, the computation of Hom anima in slice categories from [HCII, Corollary VIII.6], and the fact that $N^r: \text{Cat}_\infty \rightarrow \text{sAn}$ is fully faithful to compute

$$\begin{aligned} \text{Hom}_{\text{Cat}_\infty}([1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) &\simeq F_{\mathcal{C}^\sqcup}(\alpha: [1] \rightarrow \mathbb{I}^{\text{op}}) \\ &\simeq \text{Hom}_{\mathcal{P}(\Delta/\mathbb{I}^{\text{op}})}(\alpha: [1] \longrightarrow \mathbb{I}^{\text{op}}, F_{\mathcal{C}^\sqcup}) \\ &\simeq \text{Hom}_{\text{sAn}/N^r(\mathbb{I}^{\text{op}})}(N^r(\alpha): \Delta^1 \rightarrow N^r(\mathbb{I}^{\text{op}}), F_{\mathcal{C}^\sqcup}) \\ &\simeq \text{Hom}_{\text{sAn}}(\Delta^1, F_{\mathcal{C}^\sqcup}) \times_{\text{Hom}_{\text{sAn}}(\Delta^1, N^r(\mathbb{I}^{\text{op}}))} \{N^r(\alpha)\} \\ &\simeq \text{Hom}_{\text{Cat}_\infty}([1], \mathcal{C}^\sqcup) \times_{\text{Hom}_{\text{Cat}_\infty}([1], \mathbb{I}^{\text{op}})} \{\alpha\}. \end{aligned}$$

With this and an analogous computation for $\text{Hom}_{\text{Cat}_\infty}([0] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C})$, the above pullback square is straightforward to verify. Let's use this to check that $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ has cocartesian lifts for inert morphisms. So let $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ be inert, and $x \in \mathcal{C}^\sqcup$ a lift of $\langle m \rangle$. Then x corresponds to a map $[0] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}} \rightarrow \mathcal{C}$ and a lift of α corresponds to a map $[1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}} \rightarrow \mathcal{C}$, where the pullback is formed using $\alpha: [1] \rightarrow \mathbb{I}^{\text{op}}$. So we need to solve a lifting problem

$$\begin{array}{ccc} [0] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}} & \longrightarrow & \mathcal{C} \\ d_1 \downarrow & \nearrow & \\ [1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}} & & \end{array}$$

in such a way that the solution is a p^\sqcup -cocartesian lift of α . Since α is inert, i.e. an isomorphism where it is defined, $[1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}$ consists of a disjoint union of copies of $[1]$ (their number corresponds to the number of elements of $\langle m \rangle$ where α is defined) and $[0]$ (corresponding to the elements of $\langle m \rangle$ where α isn't). We define the desired lift $[1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}} \rightarrow \mathcal{C}$ to be degenerate on every copy of $[1]$, noting that it is already defined on their starting points and on the additional copies of $[0]$.

To show that this gives a cocartesian lift, we must check the condition from Definition I.24(a), i.e. that

$$\text{Hom}_{\mathcal{C}^\sqcup}(y, z) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}^\sqcup}(x, z) \times_{\text{Hom}_{\mathbb{I}^{\text{op}}}(\langle n \rangle, \langle k \rangle)} \text{Hom}_{\mathbb{I}^{\text{op}}}(\langle m \rangle, \langle k \rangle)$$

is an isomorphism for all $z \in \mathcal{C}^\sqcup$ and $p^\sqcup(z) = \langle k \rangle$. This can be checked on (derived) fibres over any $\beta \in \text{Hom}_{\mathbb{I}^{\text{op}}}(\langle m \rangle, \langle k \rangle)$. The fibre of the left-hand side is $\text{Hom}_{\mathcal{C}^\sqcup}^\beta(y, z)$. The fibre of the right-hand side is the same as that of $\text{Hom}_{\mathcal{C}^\sqcup}(x, z) \rightarrow \text{Hom}_{\mathbb{I}^{\text{op}}}(\langle n \rangle, \langle k \rangle)$ over $\beta \circ \alpha$, which is $\text{Hom}_{\mathcal{C}^\sqcup}^{\beta \circ \alpha}(x, z)$. That these two are equivalent is straightforward to check from the constructions, using the pullback diagram above. This proves that $p^\sqcup: \mathcal{C}^\sqcup \rightarrow \mathbb{I}^{\text{op}}$ satisfies Definition II.36(a). The condition from Definition II.36(c) can be checked in the same way. This proves (b).

For (c), it suffices to check the condition from Example II.37(a). By definition, we have $\text{Hom}_{\mathcal{C}^\sqcup}^{\text{act}}((a_1, \dots, a_n), -) \simeq \text{Hom}_{\mathcal{C}^\sqcup}^{f_n}((a_1, \dots, a_n), -)$. The pullback $[1] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^\sqcup$ formed using $f_n: [1] \rightarrow \mathbb{I}^{\text{op}}$ is precisely the cone over a discrete set with n elements. Using this and the pullback diagram above, one easily verifies

$$\text{Hom}_{\mathcal{C}^\sqcup}^{f_n}((a_1, \dots, a_n), -) \simeq \prod_{i=1}^n \text{Hom}_{\mathcal{C}}(a_i, -).$$

Hence any representing object of $\text{Hom}_{\mathcal{C}^\sqcup}^{\text{act}}((a_1, \dots, a_n), -)$ is a coproduct of the a_i , whence (c) follows. Part (d) is an immediate consequence of (c) and the discussion in Example II.37, as coproducts are clearly associative. \square

Proposition II.40 allows us to define the cocartesian monoidal structure on an ∞ -category with finite coproducts. To construct the cartesian monoidal structure, one can consider $((\mathcal{C}^{\text{op}})^\sqcup)^{\text{op}} \in \text{dOp}_\infty$. If \mathcal{C} has finite products, so \mathcal{C}^{op} has finite coproducts, then the dual ∞ -operad $((\mathcal{C}^{\text{op}})^\sqcup)^{\text{op}}$ defines a symmetric monoidal ∞ -category and we can define $p^\times: \mathcal{C}^\times \rightarrow \mathbb{I}^{\text{op}}$ via

$$\mathcal{C}^\times = \text{Un}^{\text{cocart}}(\text{St}^{\text{cart}}((\mathcal{C}^{\text{op}})^\sqcup)^{\text{op}}).$$

II.41. Construction. — There is also a direct construction of $p^\times: \mathcal{C}^\times \rightarrow \mathbb{I}^{\text{op}}$, but this is a bit tricky, which perhaps isn't very surprising since ∞ -operads would like to specify maps out of a tensor product, whereas products would like to have maps into them specified. We define a 1-category $\mathbb{I}_\times^{\text{op}}$ as follows: Its objects are pairs (n, S) , where $S \subseteq \langle n \rangle$ is a subset, and morphisms $(n, S) \rightarrow (m, T)$ are given by morphisms $\alpha: \langle n \rangle \rightarrow \langle m \rangle$ in \mathbb{I}^{op} such that $\alpha^{-1}(T) = S$. Let $\tilde{p}^\times: \tilde{\mathcal{C}}^\times \rightarrow \mathbb{I}^{\text{op}}$ be given by II.38a applied to the presheaf $F_{\mathcal{C}^\times} \Delta / \mathbb{I}^{\text{op}} \rightarrow \text{An}$, which is in turn given by

$$F_{\mathcal{C}^\times}([n] \rightarrow \mathbb{I}^{\text{op}}) \simeq \text{Hom}_{\text{Cat}_\infty}([n] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}_\times^{\text{op}}, \mathcal{C}).$$

The same analysis as in the proof of Proposition II.40(a) shows that the fibre of $\tilde{p}^\times: \tilde{\mathcal{C}}^\times \rightarrow \mathbb{I}^{\text{op}}$ over $\langle n \rangle$ is given by

$$\tilde{\mathcal{C}}_n^\times \simeq \text{Fun}((\text{power set of } \langle n \rangle)^{\text{op}}, \mathcal{C}).$$

Now let \mathcal{C}^\times be the full subcategory of $\tilde{\mathcal{C}}^\times$ which is fibrewise spanned by those functors $F: (\text{power set of } \langle n \rangle)^{\text{op}} \rightarrow \mathcal{C}$ satisfying $F(S) \simeq \prod_{i=1}^n F(\{i\})$. Then $p^\times: \mathcal{C}^\times \rightarrow \mathbb{I}^{\text{op}}$ is an ∞ -operad satisfying $\mathcal{C}_n^\times \simeq \mathcal{C}^n$ and it represents the cartesian monoidal structure if \mathcal{C} has enough products. An active morphism $(a_1, \dots, a_n) \rightarrow b$ is p^\times -cocartesian iff it exhibits b as a product of the a_i .

II.42. Proposition. — *With notation as in Construction II.39:*

- (a) *The presheaf $F_{\mathcal{C}^\times}$ defines a complete Segal anima via Lemma II.38.*
- (b) *$p^\times: \mathcal{C}^\times \rightarrow \mathbb{I}^{\text{op}}$ is always an ∞ -operad with underlying category \mathcal{C} (and thus $\mathcal{C}_n^\times \simeq \mathcal{C}^n$).*

- (c) If \mathcal{C} has finite products, then $p^\times: \mathcal{C}^\times \rightarrow \mathbb{I}^{\text{op}}$ is a symmetric monoidal category and $\text{Hom}_{\mathcal{C}^\times}^{\text{act}}((a_1, \dots, a_n), -) \simeq \text{Hom}_{\mathcal{C}^\times}(a_1 \times \dots \times a_n, -)$.

Proof. Similar to Proposition II.40 and omitted. Fabian's script [A&HK, Proposition II.42] has some details. \square

Monoids over an ∞ -Operad. — If $p: \mathcal{O} \rightarrow \mathbb{I}^{\text{op}}$ and $p': \mathcal{O}' \rightarrow \mathbb{I}^{\text{op}}$ are ∞ -operads, let $\text{Fun}^{\text{Op}\infty}(\mathcal{O}, \mathcal{O}')$ be the full sub- ∞ -category of $\text{Fun}_{\mathbb{I}^{\text{op}}}(\mathcal{O}, \mathcal{O}')$ spanned by the ∞ -operad maps. Here $\text{Fun}_{\mathbb{I}^{\text{op}}}(\mathcal{O}, \mathcal{O}')$ is given by the pullback

$$\begin{array}{ccc} \text{Fun}_{\mathbb{I}^{\text{op}}}(\mathcal{O}, \mathcal{O}') & \longrightarrow & \text{Fun}(\mathcal{O}, \mathcal{O}') \\ \downarrow & \lrcorner & \downarrow p'_* \\ * & \xrightarrow{p} & \text{Fun}(\mathcal{O}, \mathbb{I}^{\text{op}}) \end{array}$$

as usual. If \mathcal{C} has products, then $\text{Fun}^{\text{Op}\infty}(\mathbb{I}^{\text{op}}, \mathcal{C}^\times) \simeq \text{CMon}(\mathcal{C})$. In fact, this can be stated in much greater generality, which we'll do in Theorem II.43 below: Define the ∞ -category of \mathcal{O} -monoids to be the full subcategory $\mathcal{O}\text{Mon}(\mathcal{C}) \subseteq \text{Fun}(\mathcal{O}, \mathcal{C})$ spanned by those functors $F: \mathcal{O} \rightarrow \mathcal{C}$ satisfying the Segal condition. That is, the maps $(a_1, \dots, a_n) \rightarrow a_i$ in \mathcal{O} are supposed to induce equivalences $F(a_1, \dots, a_n) \xrightarrow{\sim} \prod_{i=1}^n F(a_i)$.

Also note that there is a section $\mathbb{I}^{\text{op}} \rightarrow \mathbb{I}_\times^{\text{op}}$ sending $\langle n \rangle \mapsto (n, \langle n \rangle)$. It induces canonical maps $\text{Hom}_{\text{Cat}_\infty}([n] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}_\times^{\text{op}}, \mathcal{C}) \rightarrow \text{Hom}_{\text{Cat}_\infty}([n] \times_{\mathbb{I}^{\text{op}}} \mathbb{I}^{\text{op}}, \mathcal{C}) \simeq \text{Hom}_{\text{Cat}_\infty}([n], \mathcal{C})$, hence a map $F_{\mathcal{C}^\times} \rightarrow N^r(\mathcal{C})$ of simplicial anima and therefore a map $\mathcal{C}^\times \rightarrow \mathcal{C}$.

II.43. Theorem. — If \mathcal{C} has finite products, then the map $\mathcal{C}^\times \rightarrow \mathcal{C}$ constructed above induces an equivalence

$$\text{Fun}^{\text{Op}\infty}(\mathcal{O}, \mathcal{C}^\times) \xrightarrow{\sim} \mathcal{O}\text{Mon}(\mathcal{C})$$

is an equivalence. If \mathcal{C} has finite coproducts and $\text{Hom}_{\mathcal{O}}^{\text{act}}(\emptyset, x) \simeq *$ for all $x \in \mathcal{O}$ (so that \mathcal{O} is unital, see Lemma/Definition II.45a below) then the restriction

$$\text{Fun}^{\text{Op}\infty}(\mathcal{O}, \mathcal{C}^\sqcup) \xrightarrow{\sim} \text{Fun}(\mathcal{O}_1, \mathcal{C})$$

is an equivalence as well.

Proof. See [HA]: The first part follows from Proposition 2.4.2.5, the second part from Propositions 2.4.3.16 and 2.3.1.11. \square

Theorem II.43 is supposed to motivate the following general definition.

II.44. Definition. — (a) If \mathcal{O} is an ∞ -operad and \mathcal{C}^\otimes is a symmetric monoidal structure on an ∞ -category \mathcal{C} , then we put

$$\text{Alg}_{\mathcal{O}}(\mathcal{C}^\otimes) := \text{Fun}^{\text{Op}\infty}(\mathcal{O}, \mathcal{C}^\otimes)$$

and call this the ∞ -category of \mathcal{O} -algebras in \mathcal{C} . If the symmetric monoidal structure on \mathcal{C} is clear from the context, we'll often write $\text{Alg}_{\mathcal{O}}(\mathcal{C})$ instead.

- (b) More generally, let \mathcal{O}' be another ∞ -operad with a map $\alpha: \mathcal{O}' \rightarrow \mathcal{O}$, and let \mathcal{C} be an \mathcal{O} -monoidal ∞ -category, i.e. an object of $\mathcal{O}\text{Mon}(\text{Cat}_\infty)$. Again, we denote by

$(-)^{\otimes}: \mathcal{O}\text{Mon}(\text{Cat}_{\infty}) \rightarrow \text{Op}_{\infty}/\mathcal{O}$ the functor induced by cocartesian unstraightening. Then $\text{Alg}_{\mathcal{O}'/\mathcal{O}}(\mathcal{C}^{\otimes})$ is defined as the pullback

$$\begin{array}{ccc} \text{Alg}_{\mathcal{O}'/\mathcal{O}}(\mathcal{C}^{\otimes}) & \longrightarrow & \text{Fun}^{\text{Op}_{\infty}}(\mathcal{O}', \mathcal{C}^{\otimes}) \\ \downarrow & \lrcorner & \downarrow \\ * & \xrightarrow{\alpha} & \text{Fun}^{\text{Op}_{\infty}}(\mathcal{O}', \mathcal{O}) \end{array}$$

This recovers part (a) as the special case $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes}) \simeq \text{Alg}_{\mathcal{O}/\text{Comm}}(\mathcal{C}^{\otimes})$, where we denote $\text{Comm} = \text{id}: \mathbb{I}^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$ (more on that in Example II.45(c)).

II.45. Example. — “My first algebras over ∞ -operads”:

- (a) If \mathcal{O} is the ∞ -operad $* \xrightarrow{\langle 0 \rangle} \mathbb{I}^{\text{op}}$, then $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})$ is easily identified with the fibre of \mathcal{C}^{\otimes} over $\langle 0 \rangle$, hence $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes}) \simeq *$.
- (b) If \mathcal{O} is $\mathbb{I}_{\text{int}}^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$, then $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes}) \simeq \mathcal{C}$. Hence this ∞ -operad will be denoted $\mathcal{O} = \text{Triv}$. We’ll give a sketch of why this is true. One easily finds

$$\text{Fun}_{\mathbb{I}^{\text{op}}}(\mathbb{I}_{\text{int}}^{\text{op}}, \mathcal{C}^{\otimes}) \simeq \text{Fun}_{\mathbb{I}_{\text{int}}^{\text{op}}}(\mathbb{I}_{\text{int}}^{\text{op}}, \mathcal{C}^{\otimes} \times_{\mathbb{I}^{\text{op}}} \mathbb{I}_{\text{int}}^{\text{op}}),$$

and this restricts to an equivalence

$$\text{Fun}^{\text{Op}_{\infty}}(\text{Triv}, \mathcal{C}^{\otimes}) \simeq \Gamma_{\text{cocart}}(\mathcal{C}^{\otimes} \times_{\mathbb{I}^{\text{op}}} \mathbb{I}_{\text{int}}^{\text{op}}),$$

where Γ_{cocart} is defined as in Proposition I.36. In particular, the right hand side is equivalent to $\lim(\text{St}^{\text{cocart}}(\mathcal{C}^{\otimes} \times_{\mathbb{I}^{\text{op}}} \mathbb{I}_{\text{int}}^{\text{op}}): \mathbb{I}_{\text{int}}^{\text{op}} \rightarrow \text{Cat}_{\infty})$, so we are to show that this limit is \mathcal{C} .

To see this, note that by definition of $\mathbb{I}_{\text{int}}^{\text{op}}$, its opposite category \mathbb{I}_{int} can be viewed as the category of finite sets $\langle n \rangle$ with injective maps between them. Hence there is a $\mathbb{I}_{\text{int}} \rightarrow \text{Cat}_{\infty}$ sending $\langle n \rangle$ to itself, considered as a discrete ∞ -category. Let $U \rightarrow \mathbb{I}_{\text{int}}^{\text{op}}$ be its cartesian unstraightening, so that the fibre $U_n := U \times_{\mathbb{I}_{\text{int}}^{\text{op}}} \{\langle n \rangle\}$ is isomorphic to $\langle n \rangle$. Then $\mathcal{C}_n^{\otimes} \simeq \mathcal{C}^n \simeq \lim_{U_n} \mathcal{C}$. Hence we may apply the dual of Proposition I.42 to see that the limit we’re looking for is given as $\lim_U \mathcal{C}$. But now it’s straightforward to check that the unique point of the fibre U_1 is an initial object of U , hence $\lim_U \mathcal{C} \simeq \mathcal{C}$, as required.

- (c) If \mathcal{O} is $\text{id}: \mathbb{I}^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$, then we define $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes}) =: \text{CMon}(\mathcal{C}^{\otimes})$. If \mathcal{C} has finite products and $\mathcal{C}^{\otimes} \simeq \mathcal{C}^{\times}$ is the ∞ -operad giving the cartesian monoidal structure, then this notation is consistent with our previous definition of $\text{CMon}(\mathcal{C})$ by Theorem II.43. Consequently, the ∞ -operad $\text{id}: \mathbb{I}^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$ will be denoted $\mathcal{O} = \text{Comm}$ (or sometimes \mathbb{E}_{∞}).
- (d) In view of (c), a natural question is whether there exists an ∞ -operad Assoc satisfying $\text{Alg}_{\text{Assoc}}(\mathcal{C}^{\times}) \simeq \text{Mon}(\mathcal{C})$. This is indeed true and we will construct Assoc soon (to be precise, in a future version of Example II.45(d)).
- (e) Similarly, there is an ∞ -operad LMod encoding pairs (A, M) of associative monoids A and an object M on which A acts from the left. LMod will also be constructed in a future version of II.45(e).
- (f) If $p: \mathcal{C}^{\otimes} \rightarrow \mathbb{I}^{\text{op}}$ is any symmetric monoidal ∞ -category, then we would expect that the tensor unit $1_{\mathcal{C}} \in \mathcal{C}$ gives rise to an object in $\text{CMon}(\mathcal{C}^{\otimes})$. This is indeed true and works in greater generality. To give this result the space it deserves, we outsource it to Lemma/Definition II.45a below.

II.45a. Lemma/Definition. — For any ∞ -operad \mathcal{O} , the unique object $*_{\mathcal{O}} \in \mathcal{O}_0$ is terminal in \mathcal{O} . Moreover, the following conditions are equivalent:

- (a) $\mathrm{Hom}_{\mathcal{O}}^{\mathrm{act}}(\emptyset, x) \simeq *$ for all $x \in \mathcal{O}_1$.
- (b) The unique object $*_{\mathcal{O}} \in \mathcal{O}_0$ is initial in \mathcal{O} .

Such ∞ -Operads are called *unital*. If \mathcal{O} is unital and $p: \mathcal{C}^{\otimes} \rightarrow \mathcal{O}$ an \mathcal{O} -monoidal ∞ -category (see Definition II.44(b)), then the tensor unit of \mathcal{C}^{\otimes} , i.e. the unique element of $\mathrm{St}^{\mathrm{cocart}}(p)(*_{\mathcal{O}})$, canonically gives rise to an initial object $\mathbb{1} \in \mathrm{Alg}_{\mathcal{O}/\mathcal{O}}(\mathcal{C}^{\otimes})$. In the case $\mathcal{O} \simeq \mathrm{Comm}$, the evaluation of $\mathbb{1}$ at $\langle 1 \rangle \in \mathrm{Comm}$ is indeed the usual tensor unit $1_{\mathcal{C}} \in \mathcal{C}$.

Proof. The construction of $\mathbb{1} \in \mathrm{Alg}_{\mathcal{O}/\mathcal{O}}(\mathcal{C}^{\otimes})$ was done in the lecture (in the special case $\mathcal{O} \simeq \mathrm{Comm}$), but the rest wasn't.

The fact that $*_{\mathcal{O}}$ is terminal follows immediately from Definition II.36(c). For the equivalence, observe that there is only one map $\langle 0 \rangle \rightarrow \langle 1 \rangle$ in \mathbb{T}^{op} , hence $\mathrm{Hom}_{\mathcal{O}}^{\mathrm{act}}(\emptyset, x) \simeq \mathrm{Hom}_{\mathcal{O}}(*_{\mathcal{O}}, x)$ for all $x \in \mathcal{O}_1$. This immediately implies (b) \Rightarrow (a). To see (a) \Rightarrow (b), note that $\mathrm{Hom}_{\mathcal{O}}(*_{\mathcal{O}}, x) \simeq *$ for all $x \in \mathcal{O}_1$ already suffices to show the same for all $x \in \mathcal{O}$, by Definition II.36(c).

Now let \mathcal{O} unital ∞ -operad and $p: \mathcal{C}^{\otimes} \rightarrow \mathcal{O}$ an \mathcal{O} -monoidal ∞ -category. To construct $\mathbb{1} \in \mathrm{Alg}_{\mathcal{O}/\mathcal{O}}(\mathcal{C}^{\otimes}) \subseteq \mathrm{Fun}_{\mathcal{O}}(\mathcal{O}, \mathcal{C}^{\otimes})$, our goal is to produce a canonical ∞ -operad map

$$\begin{array}{ccc} \mathcal{O} & \xrightarrow{\quad} & \mathcal{C}^{\otimes} \\ \mathrm{id} \searrow & & \swarrow p \\ & \mathcal{O} & \end{array}$$

We will choose it to be a map between cocartesian fibrations even. After straightening, a map between cocartesian fibrations over \mathcal{O} corresponds to a natural transformation in $\mathrm{Fun}(\mathcal{O}, \mathrm{Cat}_{\infty})$ between $\mathrm{const} *$ and the functor $\mathrm{St}^{\mathrm{cocart}}(p): \mathcal{O} \rightarrow \mathrm{Cat}_{\infty}$. Since $*_{\mathcal{O}} \in \mathcal{O}$ is initial, we get a natural transformation $\mathrm{const} *_{\mathcal{O}} \Rightarrow \mathrm{id}_{\mathcal{O}}$ in $\mathrm{Fun}(\mathcal{O}, \mathcal{O})$, hence a natural transformation

$$\mathrm{const} * \simeq \mathrm{const} \mathrm{St}^{\mathrm{cocart}}(p)(*_{\mathcal{O}}) \Rightarrow \mathrm{St}^{\mathrm{cocart}}(p)$$

in $\mathrm{Fun}(\mathcal{O}, \mathrm{Cat}_{\infty})$, as required. In the special case $\mathcal{O} \simeq \mathrm{Comm}$ (to which our considerations apply since $\langle 0 \rangle \in \mathrm{Comm}_0$ is initial in Comm), the evaluation of $\mathbb{1}$ at $\langle 1 \rangle$ is the endpoint of a cocartesian lift of $\langle 0 \rangle \rightarrow \langle 1 \rangle$, hence indeed $1_{\mathcal{C}} \in \mathcal{C}$.

For simplicity, let me only prove that $\mathbb{1}$ is initial the special case $\mathcal{O} \simeq \mathrm{Comm}$, since the general case works just the same, but would require some new notation. So consider another commutative algebra $A \in \mathrm{Alg}_{\mathrm{Comm}}(\mathcal{C}^{\otimes})$, corresponding to a functor $A: \mathbb{T}^{\mathrm{op}} \rightarrow \mathcal{C}^{\otimes}$ that preserves inerts and satisfies $p \circ A \simeq \mathrm{id}_{\mathbb{T}^{\mathrm{op}}}$. Let's also write $A_n = A(\langle n \rangle)$. Since A preserves inerts, we have $A_n \simeq (A_1, \dots, A_1)$ via the Segal maps. Also $\mathbb{1}_n \simeq (1_{\mathcal{C}}, \dots, 1_{\mathcal{C}})$ by construction. We compute

$$\begin{aligned} \mathrm{Hom}_{\mathrm{Alg}_{\mathrm{Comm}}(\mathcal{C}^{\otimes})}(\mathbb{1}, A) &\simeq \mathrm{Hom}_{\mathrm{Fun}_{\mathbb{T}^{\mathrm{op}}}(\mathrm{Comm}, \mathcal{C}^{\otimes})}(\mathbb{1}, A) \\ &\simeq \mathrm{Nat}(\mathbb{1}, A) \times_{\mathrm{Nat}(\mathrm{id}_{\mathbb{T}^{\mathrm{op}}}, \mathrm{id}_{\mathbb{T}^{\mathrm{op}}})} \{\mathrm{id}\}, \end{aligned}$$

using that $\mathrm{Alg}_{\mathrm{Comm}}(\mathcal{C}^{\otimes}) \subseteq \mathrm{Fun}_{\mathbb{T}^{\mathrm{op}}}(\mathrm{Comm}, \mathcal{C}^{\otimes})$ is a full sub- ∞ -category and that taking Hom anima commutes with limits of the ∞ -categories they are taken in. Now plug Corollary I.49

into the right-hand side and use the fact that limits commute (i.e. the dual of Proposition I.42) to obtain

$$\begin{aligned}
\mathrm{Hom}_{\mathrm{Alg}_{\mathrm{Comm}}(\mathcal{C}^\otimes)}(\mathbb{1}, A) &\simeq \lim_{(\alpha: \langle m \rangle \rightarrow \langle n \rangle) \in \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}})} \mathrm{Hom}_{\mathcal{C}^\otimes}(\mathbb{1}_m, A_n) \times_{\mathrm{Hom}_{\mathbb{I}^{\mathrm{op}}}(\langle m \rangle, \langle n \rangle)} \{\alpha\} \\
&\simeq \lim_{(\alpha: \langle m \rangle \rightarrow \langle n \rangle) \in \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}})} \mathrm{Hom}_{\mathcal{C}^\otimes}^\alpha(\mathbb{1}_m, A_n) \\
&\simeq \lim_{(\alpha: \langle m \rangle \rightarrow \langle n \rangle) \in \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}})} \mathrm{Hom}_{\mathcal{C}^\otimes}^\alpha(\mathbb{1}_m, A_n) \\
&\simeq \lim_{(\alpha: \langle m \rangle \rightarrow \langle n \rangle) \in \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}})} \mathrm{Hom}_{\mathcal{C}^n}^\alpha(\mathbb{1}_n, A_n) \\
&\simeq \lim_{(\alpha: \langle m \rangle \rightarrow \langle n \rangle) \in \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}})} \mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, A_1)^n.
\end{aligned}$$

In the second-last step we used that $\mathbb{1}_m \rightarrow \alpha_* \mathbb{1}_m \simeq \mathbb{1}_n$ is a p -cocartesian lift of α . Now the target projection $t: \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}}) \rightarrow \mathbb{I}^{\mathrm{op}}$ is final. Indeed, this can be shown as in the proof of Corollary I.50 (where we considered the source projection $s: \mathrm{TwAr}(\mathcal{I}) \rightarrow \mathcal{I}^{\mathrm{op}}$, but we'll see that things dualize in the right way): t is cocartesian, since $(s, t): \mathrm{TwAr}(\mathbb{I}^{\mathrm{op}}) \rightarrow \mathbb{I} \times \mathbb{I}^{\mathrm{op}}$ is a left fibration and $\mathrm{pr}_2: \mathbb{I} \times \mathbb{I}^{\mathrm{op}} \rightarrow \mathbb{I}^{\mathrm{op}}$ is cocartesian. Hence we may apply Exercise I.50a to get $|t/\langle n \rangle| \simeq |t^{-1}\{\langle n \rangle\}|$. But the fibres of t are given by $t^{-1}\{\langle n \rangle\} \simeq \mathbb{I}/\langle n \rangle$, hence weakly contractible, as required by Definition I.44. So t is indeed final.

The upshot is that we may now write

$$\mathrm{Hom}_{\mathrm{Alg}_{\mathrm{Comm}}(\mathcal{C}^\otimes)}(\mathbb{1}, A) \simeq \lim_{\langle n \rangle \in \mathbb{I}^{\mathrm{op}}} \mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, A_1)^n \simeq \mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, A_1)^0 \simeq *,$$

since $\langle 0 \rangle \in \mathbb{I}^{\mathrm{op}}$ is initial. We're done. \square

II.46. Construction. — Let \mathcal{C}^\otimes be a symmetric monoidal ∞ -category and \mathcal{O} an ∞ -operad. Define $F(\mathcal{C}^\otimes, \mathcal{O}) \rightarrow \mathbb{I}^{\mathrm{op}}$ via II.38a applied to the presheaf $F: \Delta/\mathbb{I}^{\mathrm{op}} \rightarrow \mathbf{An}$ which is given by

$$F([n] \rightarrow \mathbb{I}^{\mathrm{op}}) \simeq \mathrm{Hom}_{\mathrm{Cat}_{\infty}/\mathbb{I}^{\mathrm{op}}}([n] \times_{\mathbb{I}^{\mathrm{op}}} \mathcal{C}^\otimes, \mathcal{O})$$

Then F defines a complete Segal anima over $\mathbb{N}^r(\mathbb{I}^{\mathrm{op}})$. Indeed, this can be shown as in the proof of Proposition II.42, the only difference being that one can no longer show by hand that $I^n \times_{\mathbb{N}^r(\mathbb{I}^{\mathrm{op}})} \mathcal{C}^\otimes \rightarrow \Delta^n \times_{\mathbb{N}^r(\mathbb{I}^{\mathrm{op}})} \mathcal{C}^\otimes$ is a Joyal equivalence; instead, one can for example use that pullback along cocartesian fibrations preserves Joyal equivalences by [HTT, Proposition 3.3.1.3] or a stronger version of [HCII, Theorem IX.17].

Once we know F is complete Segal, we may apply the same method as in the proof of Proposition II.40 to compute

$$F(\mathcal{C}^\otimes, \mathcal{O})_n \simeq \mathrm{Fun}(\mathcal{C}_n^\otimes, \mathcal{O}_n) \simeq \mathrm{Fun}(\mathcal{C}^n, \mathcal{O}_1^n).$$

Let the *day convolution* $\mathrm{Day}(\mathcal{C}^\otimes, \mathcal{O}) \subseteq F(\mathcal{C}^\otimes, \mathcal{O})$ be the (non-full!) sub- ∞ -category spanned by those functors $F: \mathcal{C}_n^\otimes \rightarrow \mathcal{O}_n$ which split as products $F \simeq F_1 \times \cdots \times F_n$ and those morphisms that “split similarly”. To make it more precise which morphisms we allow, let $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ be a morphism in \mathbb{I}^{op} . Regarding α as a map $\alpha: [1] \rightarrow \mathbb{I}^{\mathrm{op}}$, we form the pullbacks $\mathcal{C}_\alpha^\otimes := [1] \times_{\mathbb{I}^{\mathrm{op}}} \mathcal{C}^\otimes$ and $\mathcal{O}_\alpha := [1] \times_{\mathbb{I}^{\mathrm{op}}} \mathcal{O}$. Let, finally, α_i denote the active map $\alpha^{-1}(i) \rightarrow [1]$ for all $i \in \langle n \rangle$. Then there are decompositions

$$\mathcal{C}_\alpha^\otimes \simeq \mathcal{C}^S \times_{[1]} \mathcal{C}_{\alpha_1}^\otimes \times_{[1]} \cdots \times_{[1]} \mathcal{C}_{\alpha_n}^\otimes \quad \text{and} \quad \mathcal{O}_\alpha \simeq \mathcal{O}_1^S \times_{[1]} \mathcal{O}_{\alpha_1} \times_{[1]} \cdots \times_{[1]} \mathcal{O}_{\alpha_n},$$

where $S \subseteq \langle m \rangle$ is the subset where α is undefined. By definition, a morphism in $F(\mathcal{C}^\otimes, \mathcal{O})$ covering α is a map $\mathcal{C}_\alpha^\otimes \rightarrow \mathcal{O}$ in $\text{Cat}_\infty/\mathbb{T}^{\text{op}}$, or equivalently, a map $\mathcal{C}_\alpha^\otimes \rightarrow \mathcal{O}_\alpha$ in $\text{Cat}_\infty/[1]$. Such a morphism is permitted into $\text{Day}(\mathcal{C}^\otimes, \mathcal{O})$ iff it respects the above decompositions. Note that this condition is always satisfied if α is active.

II.47. Proposition. — *With notation as in Construction II.46:*

- (a) $\text{Day}(\mathcal{C}^\otimes, \mathcal{O}) \rightarrow \mathbb{T}^{\text{op}}$ is an ∞ -operad with underlying ∞ -category $\text{Fun}(\mathcal{C}, \mathcal{O}_1)$.
- (b) If $q: \mathcal{C}^\otimes \rightarrow \mathcal{C}'^\otimes$ is an ∞ -operad map, or $p: \mathcal{O} \rightarrow \mathcal{O}'$ is one, so are

$$q^*: \text{Day}(\mathcal{C}'^\otimes, \mathcal{O}) \longrightarrow \text{Day}(\mathcal{C}^\otimes, \mathcal{O}) \quad \text{and} \quad p_*: \text{Day}(\mathcal{C}^\otimes, \mathcal{O}) \longrightarrow \text{Day}(\mathcal{C}^\otimes, \mathcal{O}').$$

- (c) If $\mathcal{O} \simeq \mathcal{D}^\otimes$, where \mathcal{D} is a cocomplete symmetric monoidal category such that $- \otimes_{\mathcal{D}} -$ commutes with colimits in each variable, then $\text{Day}(\mathcal{C}^\otimes, \mathcal{D}^\otimes)$ represents a symmetric monoidal structure on $\text{Fun}(\mathcal{C}, \mathcal{D})$ with tensor product

$$(F_1 \otimes_{\text{Day}} F_2)(c) \simeq \text{colim}_{d \otimes_{\mathcal{C}} d' \rightarrow c} (F_1(d) \otimes_{\mathcal{D}} F_2(d')).$$

To be really precise, the colimit on the right-hand side is taken over the slice category $(- \otimes_{\mathcal{C}} -)/c$, where $- \otimes_{\mathcal{C}} -: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ denotes the tensor product on \mathcal{C} . Moreover, the underlying ∞ -category of $\text{Day}(\mathcal{C}^\otimes, \mathcal{D}^\otimes)$ is again cocomplete and $- \otimes_{\text{Day}} -$ commutes with colimits in either variable.

- (d) In the situation of (c), let $q: \mathcal{C} \rightarrow \mathcal{C}'$ be a strongly symmetric monoidal functor. Then q^* has a left adjoint (left Kan extension) which is strongly symmetric monoidal.

Proof sketch.* Let $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ and define $\mathcal{C}_\alpha^\otimes, \mathcal{O}_\alpha$ as above. Then $\text{Hom}_{\text{Cat}_\infty/\mathbb{T}^{\text{op}}}(\mathcal{C}_\alpha^\otimes, \mathcal{O}) \simeq \text{Hom}_{\text{Cat}_\infty/[1]}(\mathcal{C}_\alpha^\otimes, \mathcal{O})$, and likewise $\text{Hom}_{\text{Cat}_\infty/\mathbb{T}^{\text{op}}}(\mathcal{C}_m^\otimes, \mathcal{O}) \simeq \text{Hom}_{\text{Cat}_\infty}(\mathcal{C}_m^\otimes, \mathcal{O}_m)$. Using this and adapting the arguments from the proof sketch of Proposition II.40, one shows that we have pullback squares

$$\begin{array}{ccc} \text{Hom}_{F(\mathcal{C}^\otimes, \mathcal{O})}^\alpha(F, G) & \xrightarrow{\quad} & \text{Hom}_{\text{Cat}_\infty/[1]}(\mathcal{C}_\alpha^\otimes, \mathcal{O}_\alpha) \\ \downarrow & \lrcorner & \downarrow (d_1, d_0) \\ * & \xrightarrow{(F, G)} & \text{Hom}_{\text{Cat}_\infty}(\mathcal{C}_m^\otimes, \mathcal{O}_m) \times \text{Hom}_{\text{Cat}_\infty}(\mathcal{C}_n^\otimes, \mathcal{O}_n) \end{array}$$

for all $F, G \in F(\mathcal{C}^\otimes, \mathcal{O})$ with images $\langle m \rangle$ and $\langle n \rangle$ respectively (but we do not claim that $F(\mathcal{C}^\otimes, \mathcal{O})$ is an ∞ -operad). If we restrict each factor to the full subanima spanned by those functors that meet the requirements from Construction II.46, we obtain a similar pullback diagram for $\text{Hom}_{\text{Day}(\mathcal{C}^\otimes, \mathcal{O})}^\alpha(F, G)$. Also note that we can plug in and pull out the decompositions of $\mathcal{O}_\alpha, \mathcal{O}_m$ and \mathcal{O}_n to make this pullback more confusing.

To prove (a), we must provide cocartesian lifts of inert morphisms. If α is inert, then lifting α to $F(\mathcal{C}^\otimes, \mathcal{O})$ means (by unravelling the pullback diagram above) we need to solve a lifting problem

$$\begin{array}{ccc} \mathcal{C}_m^\otimes & \longrightarrow & \mathcal{O}_m \\ d_1 \downarrow & & \downarrow \\ \mathcal{C}_\alpha^\otimes & \dashrightarrow & \mathcal{O}_\alpha \end{array}$$

Additionally, we want that the solution lies in $\text{Day}(\mathcal{C}^\otimes, \mathcal{O})$ and is cocartesian. Since α is inert, we have $\mathcal{C}_\alpha^\otimes \simeq \mathcal{C}^S \times \mathcal{C}^n \times [1]$ and $\mathcal{O}_\alpha \simeq \mathcal{O}_1^S \times \mathcal{O}_1^n \times [1]$, where again $S \subseteq \langle m \rangle$ is the subset

where α isn't defined. Moreover, the top arrow in the diagram above represents an element of $\text{Day}(\mathcal{C}^\otimes, \mathcal{O})_m$, hence a functor $F: \mathcal{C}^m \rightarrow \mathcal{O}_1^m$ which decomposes as $F \simeq F_1 \times \cdots \times F_m$. Thus we can choose the dashed arrow in the diagram above to be F_i on those factors with $i \in S$ and to be $F_i \times \text{id}_{[1]}$ on those factors with $i \notin S$. This clearly defines a map in $\text{Day}(\mathcal{C}^\otimes, \mathcal{O})$, and to show that it's cocartesian one can adapt the arguments from the proof sketch of Proposition II.40 and use the description of $\text{Hom}_{\text{Day}(\mathcal{C}^\otimes, \mathcal{O})}^\alpha(-, -)$ given above. This shows that $\text{Day}(\mathcal{C}^\otimes, \mathcal{O}) \rightarrow \mathbb{I}^{\text{op}}$ satisfies Definition II.36(a). The Segal condition from Definition II.36(b) holds pretty much by definition—or does it? Frankly, I don't see why that should be, but I hope I can present you a proper argument at some point. Finally, the condition from Definition II.36(c) can again be checked using our description of $\text{Hom}_{\text{Day}(\mathcal{C}^\otimes, \mathcal{O})}^\alpha(-, -)$. This finishes part (a).

For (b), we get functors $q^*: F(\mathcal{C}'^\otimes, \mathcal{O}) \rightarrow F(\mathcal{C}^\otimes, \mathcal{O})$ and $p_*: F(\mathcal{C}^\otimes, \mathcal{O}) \rightarrow F(\mathcal{C}^\otimes, \mathcal{O}')$ since the formation of the presheaf F in Construction II.46 is clearly functorial in \mathcal{C}^\otimes and \mathcal{O} . So all we need to check is that q^* and p_* preserve the decomposition conditions from Construction II.46 as well as cocartesian lifts of inert morphisms. But both things are clear from our constructions.

Now let's prove (c). Let $F, G \in F(\mathcal{C}^\otimes, \mathcal{O})$ lie over $\langle m \rangle$ and $\langle n \rangle$. Then F and G define functors $F: \mathcal{C}^m \rightarrow \mathcal{D}^m$ and $G: \mathcal{C}^n \rightarrow \mathcal{D}^n$. The following claim will be used several times during the proof:

(\boxtimes) *Let $\alpha: \langle m \rangle \rightarrow \langle n \rangle$ be an active morphism. By cocartesian unstraightening, α induces morphisms $\alpha_{\mathcal{C}}: \mathcal{C}^m \rightarrow \mathcal{C}^n$ and $\alpha_{\mathcal{D}}: \mathcal{D}^m \rightarrow \mathcal{D}^n$. Then*

$$\text{Hom}_{F(\mathcal{C}^\otimes, \mathcal{O})}^\alpha(F, G) \simeq \text{Nat}(\alpha_{\mathcal{D}} \circ F, G \circ \alpha_{\mathcal{C}})$$

Proving (\boxtimes) should be a matter of playing around with the pullback square above and the straightening equivalence, but I have yet to come up with a good argument so the proof is “left to the reader” for now. Also note that if F and G belong to $\text{Day}(\mathcal{C}^\otimes, \mathcal{D}^\otimes)$, then $\text{Hom}_{F(\mathcal{C}^\otimes, \mathcal{D}^\otimes)}^\alpha(F, G) \simeq \text{Hom}_{\text{Day}(\mathcal{C}^\otimes, \mathcal{D}^\otimes)}^\alpha(F, G)$ since α is supposed to be active, so (\boxtimes) gives a way of computing the right-hand side.

We'll use the recognition criterion from Example II.37(a) to show that $\text{Day}(\mathcal{C}^\otimes, \mathcal{D}^\otimes)$ is symmetric monoidal. As usual, let $f_n: \langle n \rangle \rightarrow \langle 1 \rangle$ denote the unique active morphism. Let $F: \mathcal{C}^n \rightarrow \mathcal{D}^n$ such that $F \simeq F_1 \times \cdots \times F_n$ and let $G: \mathcal{C} \rightarrow \mathcal{D}$. The maps $(f_n)_{\mathcal{C}}: \mathcal{C}^n \rightarrow \mathcal{C}$ and $(f_n)_{\mathcal{D}}: \mathcal{D}^n \rightarrow \mathcal{D}$ are given by the n -fold tensor products $\otimes_{\mathcal{C}}^n: \mathcal{C}^n \rightarrow \mathcal{C}$ and $\otimes_{\mathcal{D}}^n: \mathcal{D}^n \rightarrow \mathcal{D}$. Thus (\boxtimes) shows

$$\text{Hom}_{\text{Day}(\mathcal{C}^\otimes, \mathcal{O})}^{\text{act}}((F_1, \dots, F_n), G) \simeq \text{Hom}_{\text{Day}(\mathcal{C}^\otimes, \mathcal{O})}^{f_n}(F, G) \simeq \text{Nat}(\otimes_{\mathcal{D}}^n \circ F, G \circ \otimes_{\mathcal{C}}^n),$$

Hence a lift $F \rightarrow F'$ of f_n is locally cocartesian iff $F': \mathcal{C} \rightarrow \mathcal{D}$ is the left Kan extension of $\otimes_{\mathcal{D}}^n \circ F: \mathcal{C}^n \rightarrow \mathcal{D}$ along $\otimes_{\mathcal{C}}^n: \mathcal{C}^n \rightarrow \mathcal{C}$. Since \mathcal{D} is cocomplete, these Kan extensions exist by Theorem I.52. In particular, for $F_1, F_2: \mathcal{C} \rightarrow \mathcal{D}$ we have that $F_1 \otimes_{\text{Day}} F_2$ is the left Kan extension of $F_1 \otimes_{\mathcal{D}} F_2: \mathcal{C}^2 \rightarrow \mathcal{D}$ along $\otimes_{\mathcal{C}}^2: \mathcal{C}^2 \rightarrow \mathcal{C}$, which shows that the desired formula holds true (but we don't know yet that $-\otimes_{\text{Day}}-$ defines a symmetric monoidal structure).

So far we haven't used that $-\otimes_{\mathcal{D}}-$ commutes with colimits in either variable, but we'll need it to show that the locally cocartesian edges compose, so that $\text{Day}(\mathcal{C}^\otimes, \mathcal{O}) \rightarrow \mathbb{I}^{\text{op}}$ is not only locally cocartesian, but honestly cocartesian by [HCII, Proposition IX.13]. To show this, it will suffice to check that $-\otimes_{\text{Day}}-$ is associative and unital. For associativity, it suffices to check that $F_1 \otimes_{\mathcal{D}} (F_2 \otimes_{\text{Day}} F_3): \mathcal{C}^3 \rightarrow \mathcal{D}$ is the left Kan extension of $F_1 \otimes_{\mathcal{D}} F_2 \otimes_{\mathcal{D}} F_3: \mathcal{C}^3 \rightarrow \mathcal{D}$ along $\text{id}_{\mathcal{C}} \times \otimes_{\mathcal{C}}^2: \mathcal{C}^3 \rightarrow \mathcal{C}^2$ (and likewise for $(F_1 \otimes_{\text{Day}} F_2) \otimes_{\mathcal{D}} F_3$), since then the fact that Kan extension

composes will do the rest. Given $(c, c') \in \mathcal{C}^2$, we have $(\text{id}_{\mathcal{C}} \times \otimes_{\mathcal{C}}^2)/(c, c') \simeq \text{id}_{\mathcal{C}}/c \times \otimes_{\mathcal{C}}^2/c'$. Now $\{c\}$ is cofinal in $\text{id}_{\mathcal{C}}/c$, hence $\{c\} \times \otimes_{\mathcal{C}}^2/c'$ is cofinal in $\text{id}_{\mathcal{C}}/c \times \otimes_{\mathcal{C}}^2/c'$. The value at (c, c') of the left Kan extension in question can therefore be computed as

$$\begin{aligned} \text{colim}_{(c, d \otimes_{\mathcal{C}} d') \rightarrow (c, c')} (F_1(c) \otimes_{\mathcal{D}} F_2(d) \otimes_{\mathcal{D}} F_3(d')) &\simeq F_1(c) \otimes_{\mathcal{D}} \text{colim}_{d \otimes_{\mathcal{C}} d' \rightarrow c'} (F_2(d) \otimes_{\mathcal{D}} F_3(d')) \\ &\simeq F_1(c) \otimes_{\mathcal{D}} (F_2 \otimes_{\text{Day}} F_3)(c'), \end{aligned}$$

as desired. Here where we finally used that $- \otimes_{\mathcal{D}} -$ commutes with colimits. In the exact same way one shows $F \otimes_{\text{Day}} 1_{\text{Day}} \simeq F$, where $1_{\text{Day}}: \mathcal{C} \rightarrow \mathcal{D}$ is given by the left Kan extension of $1_{\mathcal{D}}: * \rightarrow \mathcal{D}$ along $1_{\mathcal{C}}: * \rightarrow \mathcal{C}$. This concludes the proof that $\text{Day}(\mathcal{C}^{\otimes}, \mathcal{D}^{\otimes})$ is symmetric monoidal. The additional assertion is clear since left Kan extension preserves colimits and colimits in functor categories are pointwise. This proves (c).

For (d), note that $q^*: \text{Fun}(\mathcal{C}', \mathcal{D}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$ has a left adjoint $q_!$ given by left Kan extension. Using the definition of $- \otimes_{\text{Day}} -$ via left Kan extension and the fact that left Kan extension composes, it's straightforward to check $q_!(F_1 \otimes_{\text{Day}} F_2) \simeq q_!(F_1) \otimes_{\text{Day}} q_!(F_2)$ and $q_!(1_{\text{Day}}) \simeq 1_{\text{Day}}$. Hence Lemma* II.37b can be applied to finish the proof. \square

II.48. Definition. — Let \mathcal{O} be an ∞ -operad and $F: \mathcal{I} \rightarrow \mathcal{O}_1$ be a diagram. Then a colimit/limit of F is called *operadic* if for every $x_1, \dots, x_n, y \in \mathcal{O}_1$ we have

$$\begin{aligned} \text{Hom}_{\mathcal{O}}^{\text{act}} \left(\left(\text{colim}_{\mathcal{I}} F, x_2, \dots, x_n \right), y \right) &\simeq \lim_{\mathcal{I}^{\text{op}}} \text{Hom}_{\mathcal{O}}^{\text{act}} \left((F(-), x_2, \dots, x_n), y \right) \\ \text{Hom}_{\mathcal{O}}^{\text{act}} \left((x_1, \dots, x_n), \lim_{\mathcal{I}} F \right) &\simeq \lim_{\mathcal{I}} \text{Hom}_{\mathcal{O}}^{\text{act}} \left((x_1, \dots, x_n), F(-) \right) \end{aligned}$$

An ∞ -operad is *pointed/semi-additive/additive/stable* if \mathcal{O}_1 is and the required limits and colimits are operadic. Spelling this out explicitly, we obtain:

- (a) \mathcal{O} is *pointed* if \mathcal{O}_1 has a zero object which is also both operadic initial and operadic terminal. We define $\text{Op}_{\infty}^* \subseteq \text{Op}_{\infty}$ as the (non-full) sub- ∞ -category spanned by ∞ -operads with an operadic terminal object and maps preserving these. We denote by $\text{Op}_{\infty}^{\text{pt}} \subseteq \text{Op}_{\infty}^*$ the full sub- ∞ -category spanned by pointed ∞ -operads.
- (b) \mathcal{O} is *semi-additive/additive* if \mathcal{O}_1 is and all finite products and coproducts are operadic. We define $\text{Op}_{\infty}^{\times} \subseteq \text{Op}_{\infty}$ as the (non-full) sub- ∞ -category spanned by ∞ -operads with finite operadic products and maps preserving these. We denote by $\text{Op}_{\infty}^{\text{semi-add}}, \text{Op}_{\infty}^{\text{add}} \subseteq \text{Op}_{\infty}^*$ the full sub- ∞ -categories spanned by semi-additive and additive ∞ -operads, respectively.
- (c) \mathcal{O} is *stable* if \mathcal{O}_1 is and all finite limits and colimits are operadic. We define $\text{Op}_{\infty}^{\text{lex}} \subseteq \text{Op}_{\infty}$ as the (non-full) sub- ∞ -category spanned by ∞ -operads with finite operadic limits and maps preserving these. We denote by $\text{Op}_{\infty}^{\text{st}} \subseteq \text{Op}_{\infty}^*$ the full sub- ∞ -category spanned by stable ∞ -operads.

II.48a. Remark*. — If $\mathcal{O} \simeq \mathcal{C}^{\otimes}$ is a symmetric monoidal ∞ -category, then all limits are operadic because $\text{Hom}_{\mathcal{C}^{\otimes}}^{\text{act}}((x_1, \dots, x_n), -) \simeq \text{Hom}_{\mathcal{C}}(x_1 \otimes \dots \otimes x_n, -)$, and a colimit is operadic iff $\text{colim}_{\mathcal{I}}(F(-) \otimes x) \simeq (\text{colim}_{\mathcal{I}} F) \otimes x$ for all $x \in \mathcal{C}$.

In general, a limit in \mathcal{O}_1 is operadic iff it is also a limit in \mathcal{O} . Indeed, the limit condition in \mathcal{O} means that

$$\lim_{\mathcal{I}^{\text{op}}} \text{Hom}_{\mathcal{O}}^{\alpha} \left((x_1, \dots, x_n), \lim_{\mathcal{I}} F \right) \simeq \lim_{\mathcal{I}} \text{Hom}_{\mathcal{O}}^{\alpha} \left((x_1, \dots, x_n), F(-) \right)$$

for every $\alpha: \langle n \rangle \rightarrow \langle 1 \rangle$ in \mathbb{I}^{op} . The special case $\alpha = f_n$ shows that if the limit over $F: \mathcal{I} \rightarrow \mathcal{O}_1 \subseteq \mathcal{O}$ happens to lie inside \mathcal{O}_1 , then it is operadic. Conversely, any α can be factored into $\alpha = f_m \circ \iota$, where $\iota: \langle n \rangle \rightarrow \langle m \rangle$ is inert. If $x \rightarrow \iota_* x$ is a cocartesian lift of ι starting at $x = (x_1, \dots, x_n)$, then $\text{Hom}_{\mathcal{O}}^{\alpha}(x, -) \simeq \text{Hom}_{\mathcal{O}}^{\text{act}}(\iota_* x, -)$, hence operadic limits are also limits in \mathcal{O} .

Given the limit case, one might expect that a colimit is operadic iff $(\text{colim}_{\mathcal{I}} F, x_2, \dots, x_n)$ is a colimit in \mathcal{O} over $(F(-), x_2, \dots, x_n): \mathcal{I} \rightarrow \mathcal{O}$ for all $x_2, \dots, x_n \in \mathcal{O}_1$. But this is false unless \mathcal{I} is weakly contractible! The reason why this fails is rather stupid: Similar to above, we must check the colimit condition on $\text{Hom}_{\mathcal{O}}^{\alpha}$ for any $\alpha: \langle n \rangle \rightarrow \langle 1 \rangle$. If we factor $\alpha = f_m \circ \iota$, then it may happen that ι isn't defined at $1 \in \langle n \rangle$, and in this case the colimit condition reads $\text{Hom}_{\mathcal{O}}^{\text{act}}(\iota_* x, y) \simeq \lim_{\mathcal{I}^{\text{op}}} \text{Hom}_{\mathcal{O}}^{\text{act}}(\iota_* x, y)$, which only holds in general if \mathcal{I} is weakly contractible.

Given an ∞ -operad \mathcal{O} with sufficient operadic limits and colimits, our goal is to construct new ∞ -operads

$$\mathcal{O}_*^{\text{Op}\infty}, \quad \text{CMon}^{\text{Op}\infty}(\mathcal{O}), \quad \text{CGrp}^{\text{Op}\infty}(\mathcal{O}), \quad \text{and} \quad \text{Sp}^{\text{Op}\infty}(\mathcal{O})$$

with underlying ∞ -categories $*/\mathcal{O}_1$, $\text{CMon}(\mathcal{O}_1)$, $\text{CGrp}(\mathcal{O}_1)$, and $\text{Sp}(\mathcal{O}_1)$ respectively.

II.49. Construction. — We will tackle all four constructions at once! Consider the following ∞ -operads:

$$[1]^{\min} \longrightarrow \mathbb{I}^{\text{op}}, \quad (\mathbb{I}^{\text{op}})^{\times} \longrightarrow \mathbb{I}^{\text{op}}, \quad \text{and} \quad (\mathbb{F}^{\text{op}})^{\wedge} \longrightarrow \mathbb{I}^{\text{op}}.$$

The first one is given by the symmetric monoidal structure on $[1]$ obtained by taking the minimum (in particular, $1 \in [1]$ is the tensor unit). The second one is the cartesian symmetric monoidal structure (as introduced in Proposition II.42) on \mathbb{I}^{op} . The third one is given by the smash product. We didn't construct this yet, but it will arise as follows: Starting from the cartesian symmetric monoidal structure An^{\times} on An , the smash product will be defined as its “pointification” $(*/\text{An})^{\wedge} := (\text{An}^{\times})_*^{\text{Op}\infty}$. Then $(\mathbb{F}^{\text{op}})^{\wedge} \subseteq (*/\text{An})^{\wedge}$ can be defined via Lemma II.37a(a). So to be super precise, we would need to prove the assertions about $(-)_*^{\text{Op}\infty}$ in Theorem II.50 below before we could even formulate anything about $\text{Sp}^{\text{Op}\infty}(-)$, but obviously that's not what we're going to do. We will also check below Remark* II.51a that the smash product coincides with the one you know from topology.

If $\mathcal{O} \in \text{Op}_{\infty}^*$, we let

$$\mathcal{O}_*^{\text{Op}\infty} \subseteq \text{Day}([1]^{\min}, \mathcal{O})$$

denote the ∞ -operad obtained via Example II.37(d) from the full sub- ∞ -category $(*/\mathcal{O}_1) \subseteq \text{Fun}([1], \mathcal{O}_1)$ spanned by those functors $F: [1] \rightarrow \mathcal{O}_1$ with $F(0) = *$. Similarly, if $\mathcal{O} \in \text{Op}_{\infty}^{\times}$, we let

$$\text{CGrp}^{\text{Op}\infty}(\mathcal{O}) \subseteq \text{CMon}^{\text{Op}\infty}(\mathcal{O}) \subseteq \text{Day}((\mathbb{I}^{\text{op}})^{\times}, \mathcal{O})$$

be induced by $\text{CGrp}(\mathcal{O}_1) \subseteq \text{CMon}(\mathcal{O}_1) \subseteq \text{Fun}(\mathbb{I}^{\text{op}}, \mathcal{O}_1)$, i.e. the full sub- ∞ -categories of functors satisfying the Segal condition and—in the case of $\text{CGrp}(\mathcal{O}_1)$ —additionally the condition from Definition II.4. Finally, for $\mathcal{O} \in \text{Op}_{\infty}^{\text{lex}}$, we let

$$\text{Sp}^{\text{Op}\infty}(\mathcal{O}) \subseteq \text{Day}((\mathbb{F}^{\text{op}})^{\wedge}, \mathcal{O})$$

be given by the full sub- ∞ -category $\text{Sp}(\mathcal{O}_1) \subseteq \text{Fun}(\mathbb{F}^{\text{op}}, \mathcal{O}_1)$ spanned by reduced and excisive functors (see Definition II.33).

These ∞ -operads come equipped with canonical maps

$$\mathcal{O} \longleftarrow \mathcal{O}_*^{\mathrm{Op}\infty} \longleftarrow \mathrm{CMon}^{\mathrm{Op}\infty}(\mathcal{O}) \longleftarrow \mathrm{CGrp}^{\mathrm{Op}\infty}(\mathcal{O}) \xleftarrow{\Omega^\infty} \mathrm{Sp}^{\mathrm{Op}\infty}(\mathcal{O}).$$

Except for the fourth one, which comes from the inclusion $\mathrm{CMon}(\mathcal{O}_1) \supseteq \mathrm{CGrp}(\mathcal{O}_1)$, these maps are induced via functoriality of Day convolution (see Proposition II.47(b)) by the ∞ -operad maps

$$\mathrm{Comm} \longrightarrow [1]^{\mathrm{min}} \longrightarrow (\mathbb{I}^{\mathrm{op}})^\times \longrightarrow (\mathbb{F}^{\mathrm{op}})^\wedge.$$

All of these are induced by strongly monoidal functors between the underlying categories: The first one comes from $1: * \rightarrow [1]$, the second one from $[1] \rightarrow \mathbb{I}^{\mathrm{op}}$ sending $i \mapsto \langle i \rangle$ for $i = 0, 1$, and the third one comes from the functor $(-)_+: \mathbb{I}^{\mathrm{op}} \rightarrow \mathbb{F}^{\mathrm{op}}$ introduced in Definition II.33.

With these constructions we can now formulate a striking theorem of Thomas Nikolaus!

II.50. Theorem ([Nik16, Theorem 4.11 and Theorem 5.7]). — *The ∞ -operads $\mathcal{O}_*^{\mathrm{Op}\infty}$, $\mathrm{CMon}^{\mathrm{Op}\infty}(\mathcal{O})$, $\mathrm{CGrp}^{\mathrm{Op}\infty}(\mathcal{O})$, and $\mathrm{Sp}^{\mathrm{Op}\infty}(\mathcal{O})$ from Construction II.49 are pointed, semi-additive, additive and stable respectively. Moreover, the respective constructions assemble into functors*

$$\begin{aligned} (-)_*^{\mathrm{Op}\infty}: \mathrm{Op}_\infty^* &\longrightarrow \mathrm{Op}_\infty^{\mathrm{pt}}, & \mathrm{CMon}^{\mathrm{Op}\infty}(-): \mathrm{Op}_\infty^\times &\longrightarrow \mathrm{Op}_\infty^{\mathrm{semi-add}}, \\ \mathrm{CGrp}^{\mathrm{Op}\infty}(-): \mathrm{Op}_\infty^\times &\longrightarrow \mathrm{Op}_\infty^{\mathrm{add}}, & \mathrm{Sp}^{\mathrm{Op}\infty}(-): \mathrm{Op}_\infty^{\mathrm{lex}} &\longrightarrow \mathrm{Op}_\infty^{\mathrm{st}}, \end{aligned}$$

each of which is left-adjoint to the respective fully faithful inclusion in the other direction (i.e., a left Bousfield localisation).

*Proof sketch**. Once we know that our four functors take values in the correct ∞ -categories, we can proceed as in Theorem II.19 and Theorem II.30 to show that they are left Bousfield localisations. So we only need to prove the first part of the theorem.

The most subtle part is to show that the required colimits in the ∞ -operads $\mathcal{O}_*^{\mathrm{Op}\infty}$, $\mathrm{CMon}^{\mathrm{Op}\infty}(\mathcal{O})$, $\mathrm{CGrp}^{\mathrm{Op}\infty}(\mathcal{O})$, and $\mathrm{Sp}^{\mathrm{Op}\infty}(\mathcal{O})$ are actually operadic. Thomas Nikolaus does this with a beautiful trick: Let's first assume $\mathcal{O} \simeq \mathcal{C}^\otimes$, where \mathcal{C} is a symmetric monoidal ∞ -category for which Proposition II.51 below is applicable. Then this proposition says that in fact all small colimits are operadic and we're happy. In general, one can find a fully faithful map of ∞ -operads $i: \mathcal{O} \rightarrow \mathcal{C}^\otimes$ such that Proposition II.51 is applicable to \mathcal{C}^\otimes and $i_1: \mathcal{O}_1 \rightarrow \mathcal{C}$ preserves the required finite limits, see [Nik16, Proposition 2.7] and Remark* II.51 a. The required finite colimits in $*/\mathcal{O}_1$, $\mathrm{CMon}(\mathcal{O}_1)$, $\mathrm{CGrp}(\mathcal{O}_1)$, and $\mathrm{Sp}(\mathcal{O}_1)$ are also finite colimits in $*/\mathcal{C}$, $\mathrm{CMon}(\mathcal{C})$, $\mathrm{CGrp}(\mathcal{C})$, and $\mathrm{Sp}(\mathcal{C})$ since each of them (zero objects, finite coproducts in semi-additive ∞ -categories, finite colimits in stable ∞ -categories) can be built from finite limits which $i_1: \mathcal{O}_1 \rightarrow \mathcal{C}$ preserves. Hence they are operadic and we win. \square

In general, neither of the constructions from Construction II.49 gives a symmetric monoidal category, even when \mathcal{O} itself is symmetric monoidal. The following proposition gives a criterion for the symmetric monoidal structure to be inherited.

II.51. Proposition. — *Suppose $\mathcal{O} \simeq \mathcal{C}^\otimes$ is a symmetric monoidal ∞ -categories with all small operadic colimits (this is a short way of saying that \mathcal{C} is cocomplete and $- \otimes_{\mathcal{C}} -$ commutes with colimits in either variable, see Remark* II.48a). Suppose furthermore that the inclusions*

$$\begin{aligned} */\mathcal{C} &\subseteq \mathrm{Ar}(\mathcal{C}), & \mathrm{CMon}(\mathcal{C}) &\subseteq \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathcal{C}), \\ \mathrm{CGrp}(\mathcal{C}) &\subseteq \mathrm{Fun}(\mathbb{I}^{\mathrm{op}}, \mathcal{C}), & \mathrm{Sp}(\mathcal{C}) &\subseteq \mathrm{Fun}(\mathbb{F}^{\mathrm{op}}, \mathcal{C}) \end{aligned}$$

have left adjoints. Then $(\mathcal{C}^\otimes)_*^{\text{Op}\infty}$, $\text{CMon}^{\text{Op}\infty}(\mathcal{C}^\otimes)$, $\text{CGrp}^{\text{Op}\infty}(\mathcal{C}^\otimes)$, and $\text{Sp}^{\text{Op}\infty}(\mathcal{C}^\otimes)$ are symmetric monoidal ∞ -categories again and have all small operadic colimits. Moreover, there are maps ∞ -operads

$$\begin{aligned} (-)_+ : \mathcal{C}^\otimes &\longrightarrow (\mathcal{C}^\otimes)_*^{\text{Op}\infty}, & \text{Free}^{\text{CMon}} : \mathcal{C}^\otimes &\longrightarrow \text{CMon}^{\text{Op}\infty}(\mathcal{C}^\otimes), \\ \text{Free}^{\text{CGrp}}(-) : \mathcal{C}^\otimes &\longrightarrow \text{CGrp}^{\text{Op}\infty}(\mathcal{C}^\otimes), & \Sigma^\infty : \mathcal{C}^\otimes &\longrightarrow \text{Sp}^{\text{Op}\infty}(\mathcal{C}^\otimes) \end{aligned}$$

which are strongly monoidal and left-adjoint to the respective maps in the other direction constructed at the end of Construction II.49. Similar assertions hold for other left adjoints among these. For instance, in the case $\mathcal{C}^\otimes = \text{An}^\times$ we get a fully faithful strongly monoidal left adjoint $B^\infty : \text{CGrp}^{\text{Op}\infty}(\text{An}^\times) \rightarrow \text{Sp}^{\text{Op}\infty}(\text{An}^\times)$ of Ω^∞ .

Proof.* We only prove the assertions about Sp , since the arguments can be copied verbatim for the other cases. By Proposition II.47(c), $\text{Day}((\mathbb{F}^{\text{op}})^\wedge, \mathcal{C}^\otimes)$ is symmetric monoidal and has all small operadic colimits. Since $\text{Sp}(\mathcal{C}) \subseteq \text{Fun}(\mathbb{F}^{\text{op}}, \mathcal{C})$ has a left adjoint, we may apply Lemma II.37a(b) to see that $\text{Sp}^{\text{Op}\infty}(\mathcal{C}^\otimes)$ is again symmetric monoidal and its inclusion into $\text{Day}((\mathbb{F}^{\text{op}})^\wedge, \mathcal{C}^\otimes)$ has a left adjoint $\text{Day}((\mathbb{F}^{\text{op}})^\wedge, \mathcal{C}^\otimes) \rightarrow \text{Sp}^{\text{Op}\infty}(\mathcal{C}^\otimes)$ which is strongly monoidal. Together with our computation of colimits in Bousfield localisations (Corollary I.61) this shows that $\text{Sp}^{\text{Op}\infty}(\mathcal{C}^\otimes)$ has all small operadic colimits.

Moreover, to produce the strongly monoidal left adjoint $\Sigma^\infty : \mathcal{C}^\otimes \rightarrow \text{Sp}^{\text{Op}\infty}(\mathcal{C}^\otimes)$, it suffices to construct a strongly monoidal left adjoint of the canonical map

$$\text{Day}((\mathbb{F}^{\text{op}})^\wedge, \mathcal{C}) \longrightarrow \text{Day}(\text{Comm}, \mathcal{C}^\otimes) \simeq \mathcal{C}^\otimes$$

induced by the strongly monoidal map $\text{Comm} \rightarrow (\mathbb{F}^{\text{op}})^\wedge$. But this is precisely what Proposition II.47(d) does! The additional assertion about B^∞ can be shown in the same way. \square

II.51a. Remark*. — Proposition II.51 is always applicable when $\mathcal{C}^\otimes \simeq \text{Day}(\mathcal{D}^\otimes, \text{An}^\times)$ for some small symmetric monoidal ∞ -category \mathcal{D} . That is, whenever $\mathcal{C} \simeq \mathcal{P}(\mathcal{D})$, equipped with the Day convolution structure. Since \mathcal{C}^\otimes in the proof of Theorem II.50 can be always chosen in this way (see the proof of [Nik16, Proposition 2.7]), this is all we need.

So let's prove that the assumptions of Proposition II.51 are really satisfied for $\mathcal{C} \simeq \mathcal{P}(\mathcal{D})$! First of all, $\mathcal{P}(\mathcal{D})$ has all small operadic colimits by Proposition II.47(c), so only the required left adjoints have to be constructed. Let's first do the case of Sp . By Proposition II.34a, the inclusion $\text{Sp}(\mathcal{C}) \subseteq \text{Fun}_*(\mathbb{F}^{\text{op}}, */\mathcal{C})$ into the reduced functors has a left adjoint whenever $*/\mathcal{C}$ has sequential colimits and $\Omega : */\mathcal{C} \rightarrow */\mathcal{C}$ commutes with them. This is satisfied for $*/\text{An}$, as we checked after Proposition II.31, hence also for $\text{const } */\mathcal{P}(\mathcal{D}) \simeq \text{Fun}(\mathcal{D}^{\text{op}}, */\text{An})$ since everything is pointwise. Now $\text{Fun}_*(\mathbb{F}^{\text{op}}, */\mathcal{C}) \simeq \text{Fun}_*(\mathbb{F}^{\text{op}}, \mathcal{C})$, since every reduced functor $F : \mathbb{F}^{\text{op}} \rightarrow \mathcal{C}$ canonically lifts to $*/\mathcal{C}$, so it suffices to construct a left adjoint of $\text{Fun}_*(\mathbb{F}^{\text{op}}, \mathcal{C}) \subseteq \text{Fun}(\mathbb{F}^{\text{op}}, \mathcal{C})$. If \mathcal{C} has pushouts, we can send an arbitrary $F : \mathbb{F}^{\text{op}} \rightarrow \mathcal{C}$ to $\text{cofib}(F(*) \rightarrow F(-)) : \mathbb{F}^{\text{op}} \rightarrow \mathcal{C}$ to obtain the desired left adjoint. Again, $\mathcal{P}(\mathcal{D})$ clearly has pushouts, whence we are done.

Next, we treat CMon . Denote $i : \mathbb{I}^{\text{op}}_{\leq 1} \hookrightarrow \mathbb{I}^{\text{op}}$. If \mathcal{C} has small limits, then restriction and right Kan extension along i gives a functor $i_*i^* : \text{Fun}_*(\mathbb{I}^{\text{op}}, \mathcal{C}) \rightarrow \text{CMon}(\mathcal{C})$. This is a left Bousfield localisation, which can be easily checked using Proposition I.61a and an assertion similar to claim (\boxtimes) from the proof of Proposition II.16. Composing this with a left Bousfield localisation $\text{Fun}(\mathbb{I}^{\text{op}}, \mathcal{C}) \rightarrow \text{Fun}_*(\mathbb{I}^{\text{op}}, \mathcal{C})$ as above gives the required localisation $\text{Fun}(\mathbb{I}^{\text{op}}, \mathcal{C}) \rightarrow \text{CMon}(\mathcal{C})$ in the CMon case. This immediately implies the CGrp case, since

$\mathrm{CGrp}(\mathrm{An}) \subseteq \mathrm{CMon}(\mathrm{An})$ has a left adjoint by Corollary II.21, hence the same is true for $\mathrm{CGrp}(\mathcal{P}(\mathcal{D})) \simeq \mathrm{Fun}(\mathcal{D}, \mathrm{CGrp}(\mathrm{An})) \subseteq \mathrm{Fun}(\mathcal{D}, \mathrm{CMon}(\mathrm{An})) \simeq \mathrm{CMon}(\mathcal{P}(\mathcal{D}))$.

I'll leave the final case, i.e. showing that $*/\mathcal{C} \subseteq \mathrm{Ar}(\mathcal{C})$ has a left adjoint in the special case $\mathcal{C} \simeq \mathcal{P}(\mathcal{D})$, to you.

II.51b. Example. — Let's discuss two special cases of Proposition II.51. The first one wasn't in the lecture, but it will show up later.

- (a) We claim that $\mathrm{CGrp}^{\mathrm{Op}\infty}(\mathrm{Set}^\times) \simeq \mathrm{Ab}^\otimes$ is the symmetric monoidal structure on the 1-category of abelian groups given by the usual tensor product. Indeed, let $A, B \in \mathrm{Ab}$, thought of as functors $A, B: \mathbb{I}^{\mathrm{op}} \rightarrow \mathrm{Set}$. Let $A \otimes B$ denote their Day convolution $\mathrm{Day}((\mathbb{I}^{\mathrm{op}})^\times, \mathrm{Set}^\times)$. By Proposition II.47(c), we can compute it as the left Kan extension of

$$\begin{array}{ccc} \mathbb{I}^{\mathrm{op}} \times \mathbb{I}^{\mathrm{op}} & \xrightarrow{A \times B} & \mathrm{Set} \times \mathrm{Set} \xrightarrow{\times_{\mathrm{Set}}} \mathrm{Set} \\ \times_{\mathbb{I}^{\mathrm{op}}} \downarrow & & \nearrow A \otimes B \\ \mathbb{I}^{\mathrm{op}} & \xrightarrow{\quad} & \end{array}$$

By the universal property of left Kan extension, this means that

$$\mathrm{Hom}_{\mathrm{Ab}}(A \otimes B, C) \simeq \mathrm{Nat}(A \otimes B, C) \simeq \mathrm{Nat}(\times_{\mathrm{Set}} \circ (A \times B), C \circ \times_{\mathbb{I}^{\mathrm{op}}})$$

for all abelian groups C . Unravelling, we find that the right-hand side is given by \mathbb{Z} -bilinear maps $A \times B \rightarrow C$, hence $A \otimes B$ is indeed the usual tensor product.

The inclusion $\mathrm{Set} \subseteq \mathrm{An}$ induces inclusions $\mathrm{Ab} \simeq \mathrm{CGrp}(\mathrm{Set}) \subseteq \mathrm{CGrp}(\mathrm{An})$ and $\mathrm{Ab}^\otimes \simeq \mathrm{CGrp}^{\mathrm{Op}\infty}(\mathrm{Set}^\times) \subseteq \mathrm{CGrp}^{\mathrm{Op}\infty}(\mathrm{An})$, which are fully faithful again. To see this, note that the left adjoint $\pi_0: \mathrm{An} \rightarrow \mathrm{Set}$ induces left adjoints for each of the inclusions, and the condition that the counit transformation is an equivalence is inherited, hence so is fully faithfulness by Proposition/Definition I.58(a).

- (b) Proposition II.51 shows that $*/\mathrm{An}$ carries a pointed symmetric monoidal structure, called the *smash product*, via $(*/\mathrm{An})^\wedge := (\mathrm{An}^\times)_*^{\mathrm{Op}\infty}$. We wish to compute $- \wedge -$ explicitly to show that it coincides with the smash product we know from topology.

So let $X, Y \in */\mathrm{An} \subseteq \mathrm{Ar}(\mathrm{An})$. Let's first compute the Day convolution $X \times_{\mathrm{Day}} Y$: Using the explicit formula from Proposition II.47(c), we obtain

$$\begin{aligned} (X \times_{\mathrm{Day}} Y)(0) &\simeq \mathrm{colim} \left(\begin{array}{ccc} * \times * & \longrightarrow & * \times Y \\ \downarrow & & \\ X \times * & & \end{array} \right) \simeq X \vee Y, \\ (X \times_{\mathrm{Day}} Y)(1) &\simeq \mathrm{colim} \left(\begin{array}{ccc} * \times * & \longrightarrow & * \times Y \\ \downarrow & \searrow & \downarrow \\ X \times * & \longrightarrow & X \times Y \end{array} \right) \simeq X \times Y. \end{aligned}$$

Since the left adjoint of $*/\mathrm{An} \subseteq \mathrm{Ar}(\mathrm{An})$ sends any $T: [1] \rightarrow \mathrm{An}$ to $\mathrm{cofib}(T(0) \rightarrow T(1))$, we obtain $X \wedge Y \simeq \mathrm{cofib}(X \wedge Y \rightarrow X \times Y)$ after unravelling what the proofs* of Proposition II.51 and Lemma II.37a(b) actually do. Hence $X \wedge Y$ is what we would expect.

II.52. Definition. — The *tensor product* (or traditionally *smash product*) of spectra is defined by the ∞ -operad $\mathrm{Sp}^\otimes := \mathrm{Sp}^{\mathrm{Op}\infty}(\mathrm{An}^\times)$, which is a symmetric monoidal ∞ -category by Proposition II.51.

Some Brave New Algebra

LECTURE 17
7th Jan, 2021

The term “brave new algebra” roughly refers to the idea that Sp should behave like the category of abelian groups, and more generally to notion of \mathbb{E}_∞ -ring spectra and modules over them, which should behave very much like ordinary commutative rings and modules. Making this precise and proving it rigorously takes Lurie about 400 pages in [HA]! Of course there’s no way to even remotely cover this in this course. So be warned that proofs will become more and more sparse towards the end of this final subsection of Chapter II. Nevertheless, Fabian made it his goal to introduce all relevant definitions and theorems in the lecture, and even more material in the lecture notes [A&HK, Chapter II pp. 104–132], to give us at least a guide for how to read Lurie. Starting from Chapter III, we will then just assume that homological algebra works with \mathbb{E}_∞ -ring spectra just as it does for ordinary rings.

II.53. First Steps with the Tensor Product on Sp . — Recall the symmetric monoidal structure on Sp from Definition II.52.

The tensor unit 1_{Sp} is given by the sphere spectrum \mathbb{S} ! The quickest way to see this is that $\mathbb{S}[-]: \mathrm{An}^\times \rightarrow \mathrm{Sp}^\otimes$ is strongly symmetric monoidal because so are both its constituents $(-)_+: \mathrm{An}^\times \rightarrow (*/\mathrm{An})^\wedge$ and $\Sigma^\infty: (*/\mathrm{An})^\wedge \rightarrow \mathrm{Sp}^\otimes$ by Proposition II.51, and the tensor unit $* \in \mathrm{An}$ is sent to \mathbb{S} .

As an easy consequence, we obtain

$$X \otimes E \simeq \mathbb{S}[X] \otimes E \quad \text{and} \quad (X, x) \otimes E \simeq \Sigma^\infty(X, x) \otimes E,$$

for all $(X, x) \in */\mathrm{An}$ and $E \in \mathrm{Sp}$, where $X \otimes E$ and $(X, x) \otimes E$ are defined as in the proof of Proposition II.34. Indeed, for the left-hand equivalence it’s enough to show that both $- \otimes E: \mathrm{An} \rightarrow \mathrm{Sp}$ and $\mathbb{S}[-] \otimes E: \mathrm{An} \rightarrow \mathrm{Sp}$ preserve colimits and that they agree on $* \in \mathrm{An}$, for then they must agree by Theorem I.51. That both sides agree on $*$ precisely says that \mathbb{S} is the tensor unit, which we just checked. That $- \otimes E$ preserves colimits is clear by inspection, whereas for $\mathbb{S}[-] \otimes E$ it follows from the fact that the tensor product on spectra commutes with colimits in either variable by Proposition II.51. Hence the left-hand equivalence holds. The right-hand one follows, as $(X, x) \otimes E \simeq \mathrm{cofib}(\{x\} \otimes E \rightarrow X \otimes E)$ and $\Sigma^\infty(X, x) \simeq \mathrm{cofib}(\mathbb{S}[\{x\}] \rightarrow \mathbb{S}[X])$.

This can be used to derive a general formula for the tensor product of spectra. Any spectrum E can be written as

$$E \simeq \mathrm{colim}_{n \in \mathbb{N}} (\Sigma^\infty(\Omega^{\infty-n} E))[-n].$$

Indeed, using the $(\Sigma^\infty, \Omega^\infty)$ -adjunction, we can calculate

$$\begin{aligned} \mathrm{Hom}_{\mathrm{Sp}} \left(\mathrm{colim}_{n \in \mathbb{N}} (\Sigma^\infty(\Omega^{\infty-n} E))[-n], E' \right) &\simeq \lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Hom}_{\mathrm{Sp}} (\Sigma^\infty(\Omega^{\infty-n} E), E'[n]) \\ &\simeq \lim_{n \in \mathbb{N}^{\mathrm{op}}} \mathrm{Hom}_{*/\mathrm{An}} (\Omega^{\infty-n} E, \Omega^{\infty-n} E'), \end{aligned}$$

and the right-hand side is $\mathrm{Hom}_{\mathrm{Sp}}(E, E')$ since Hom anima in limits are limits of Hom anima. Hence the claimed equivalence follows from Yoneda’s lemma.

Together with the fact that the tensor product of spectra commutes with arbitrary colimits, this shows the general formula

$$E \otimes E' \simeq \mathrm{colim}_{n \in \mathbb{N}} (\Sigma^\infty(\Omega^{\infty-n} E))[-n] \otimes E' \simeq \mathrm{colim}_{n \in \mathbb{N}} (\Omega^{\infty-n} E \otimes E')[-n],$$

as was already announced after Definition II.35. Note that the tensor products on the right-hand side are no longer those in Sp , but defined as in the proof of Proposition II.34, so this formula gives—at least in theory—a way to compute tensor products of spectra.

In real life though, the colimits in question are seldom computable. To give an example of how complicated even simple tensor products can become, write

$$\mathrm{Tor}_i^{\mathbb{S}}(E, E') := \pi_*(E \otimes E') \quad \text{and} \quad \mathrm{Ext}_{\mathbb{S}}^i(E, E') \simeq \pi_i \mathrm{hom}_{\mathrm{Sp}}(E, E'),$$

and consider $\mathrm{Tor}_*(H\mathbb{F}_p, H\mathbb{F}_p) = \mathcal{A}_p^\vee$ and $\mathrm{Ext}_{\mathbb{S}}^*(H\mathbb{F}_p, H\mathbb{F}_p) = \mathcal{A}_p$, called the (*dual* and *non-dual*) *Steenrod algebra*. These are infinite-dimensional graded \mathbb{F}_p -algebras, but completely computed (albeit not by the formula above). For example, $\mathcal{A}_2^\vee = \mathbb{F}_2[x_i \mid i \in \mathbb{N}]$ with $\deg(x_i) = 2^i - 1$ is a polynomial ring in infinitely many variables.

To show that our formula isn't completely useless, I'd like to use it to prove the tensor-Hom adjunction. This wasn't discussed in the lecture, but Fabian has also written out a proof in the official lecture notes, [A&HK, Remarks II.53(iv)].

II.53a. Lemma*. — *For all $E, E', E'' \in \mathrm{Sp}$, we have an equivalence*

$$\mathrm{hom}_{\mathrm{Sp}}(E \otimes E', E'') \simeq \mathrm{hom}_{\mathrm{Sp}}(E, \mathrm{hom}_{\mathrm{Sp}}(E', E''))$$

It is functorial in all three variables, i.e. an equivalence in $\mathrm{Fun}(\mathrm{Sp}^{\mathrm{op}} \times \mathrm{Sp}^{\mathrm{op}} \times \mathrm{Sp}, \mathrm{Sp})$.

Proof.* Let's first check it in the special case $E' \simeq \mathbb{S}[X]$. Note that $\mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}, -) : \mathrm{Sp} \rightarrow \mathrm{Sp}$ is the identity functor. Indeed, we've already verified $\mathrm{Hom}_{\mathrm{Sp}}(\mathbb{S}, -) \simeq \Omega^\infty$ (see before Lemma II.31c), and $\Omega^\infty : \mathrm{Sp} \rightarrow \mathrm{An}$ clearly lifts to $\mathrm{id}_{\mathrm{Sp}}$ via Theorem II.30. Hence the functors

$$\mathrm{hom}_{\mathrm{Sp}}(E \otimes \mathbb{S}[-], E'')^{\mathrm{op}}, \mathrm{hom}_{\mathrm{Sp}}(E, \mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}[-], E''))^{\mathrm{op}} : \mathrm{An} \longrightarrow \mathrm{Sp}^{\mathrm{op}}$$

agree on $* \in \mathrm{An}$ and both clearly preserve colimits, hence they must agree by Theorem I.51. This settles the special case $E' \simeq \mathbb{S}[X]$. More generally, this shows

$$\mathrm{hom}_{\mathrm{Sp}}(- \otimes \mathbb{S}[-], -)^{\mathrm{op}} \simeq \mathrm{hom}_{\mathrm{Sp}}(-, \mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}[-], -))^{\mathrm{op}}.$$

Indeed, we can interpret both sides as functors $\mathrm{Sp}^{\mathrm{op}} \times \mathrm{Sp} \rightarrow \mathrm{Fun}(\mathrm{An}, \mathrm{Sp}^{\mathrm{op}})$. Upon inspection, they land in the full sub- ∞ -category $\mathrm{Fun}^{\mathrm{colim}}(\mathrm{An}, \mathrm{Sp}^{\mathrm{op}})$ of colimit-preserving functors. But $\mathrm{Fun}^{\mathrm{colim}}(\mathrm{An}, \mathrm{Sp}^{\mathrm{op}}) \simeq \mathrm{Fun}(*, \mathrm{Sp}^{\mathrm{op}})$, and if we interpret the two functors above as functors $\mathrm{Sp}^{\mathrm{op}} \times \mathrm{Sp} \rightarrow \mathrm{Fun}(*, \mathrm{Sp}^{\mathrm{op}})$ then they clearly agree since both $- \otimes \mathbb{S}$ and $\mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}, -)$ are equivalent to the identity. The case $E' \simeq \Sigma^\infty(X, x)$, and more generally the equivalence of functors

$$\mathrm{hom}_{\mathrm{Sp}}(- \otimes \Sigma^\infty -, -) \simeq \mathrm{hom}_{\mathrm{Sp}}(-, \mathrm{hom}_{\mathrm{Sp}}(\Sigma^\infty -, -))$$

in $\mathrm{Fun}(\mathrm{Sp}^{\mathrm{op}} \times (*/\mathrm{An})^{\mathrm{op}} \times \mathrm{Sp}, \mathrm{Sp})$ immediately follows.

Finally, we can write $E' \simeq \mathrm{colim}_{n \in \mathbb{N}}(\Sigma^\infty(\Omega^{\infty-n} E'))[-n]$, and then the formula from II.53 reduces the general case to the case $E' \simeq \Sigma^\infty(X, x)$. This also proves the full functoriality statement, since we can similarly write $\mathrm{id}_{\mathrm{Sp}} \simeq \mathrm{colim}_{n \in \mathbb{N}}(\Sigma^\infty(\Omega^{\infty-n} -))[-n]$. \square

\mathbb{E}_∞ -Ring Spectra

II.54. Definition and Examples. — For any symmetric monoidal ∞ -operad, we define $\mathrm{CAlg}(\mathcal{C}^\otimes) := \mathrm{Alg}_{\mathrm{Comm}}(\mathcal{C}^\otimes)$. We already introduced the notation $\mathrm{CMon}(\mathcal{C}^\otimes)$ for this in Example II.45(c), but nevermind. In the special case $\mathcal{C}^\otimes \simeq \mathrm{Sp}^\otimes$, we put

$$\mathrm{CAlg} := \mathrm{CAlg}(\mathrm{Sp}^\otimes) = \mathrm{Alg}_{\mathrm{Comm}}(\mathrm{Sp}^\otimes)$$

and call this the ∞ -category of \mathbb{E}_∞ -ring spectra. Any \mathbb{E}_∞ -ring spectrum has an underlying spectrum, given by its image under $\mathrm{ev}_{\langle 1 \rangle}: \mathrm{CAlg} \simeq \mathrm{Fun}^{\mathrm{Op}\infty}(\mathrm{Comm}, \mathrm{Sp}^\otimes) \rightarrow \mathrm{Sp}$. Occasionally (and abusively) we won't distinguish between an \mathbb{E}_∞ -ring spectrum and its underlying spectrum.

In our picture where Sp corresponds to the 1-category Ab of abelian groups, CAlg takes the place of the 1-category CRing of commutative rings (which makes a lot of sense, as $\mathrm{CRing} \simeq \mathrm{CAlg}(\mathrm{Ab}^\otimes)$ by unwinding of definitions). Now come “my first \mathbb{E}_∞ -ring spectra”:

- (a) Recall from II.53 that $\mathbb{S}[-]: \mathrm{An}^\times \rightarrow \mathrm{Sp}^\otimes$ is strongly monoidal, and that $\mathrm{CAlg}(\mathrm{An}^\times) \simeq \mathrm{CMon}(\mathrm{An})$ by Theorem II.43. Hence $\mathbb{S}[-]$ upgrades to a functor

$$\mathbb{S}[-]: \mathrm{CMon}(\mathrm{An}) \longrightarrow \mathrm{CAlg}.$$

For $M \in \mathrm{CMon}(\mathrm{An})$, we think of $\mathbb{S}[M] \in \mathrm{CAlg}$ as the “spherical monoid ring on M ”.

- (b) Every ordinary commutative ring defines an \mathbb{E}_∞ -ring spectrum, and more generally so does every commutative differential graded algebra (CGDA in the following). We will discuss this in detail in Lemma II.55, Corollary II.55a, and the space in between.

Let me also mention another example, that came only up at a later point in the course, but fits here very well.

- (c*) Let R be a commutative ring and $k(R)$ its algebraic K -theory space from Definition II.12a. We know from Example II.21a(a) that it is an \mathbb{E}_∞ -group. But $B^\infty k(R)$ even has a canonical refinement as an \mathbb{E}_∞ -ring spectrum! To see this, put

$$\mathrm{CMon}(\mathrm{Grpd})^\otimes := \mathrm{CMon}^{\mathrm{Op}\infty}(\mathrm{Grpd}^\times) \quad \text{and} \quad \mathrm{CMon}(\mathrm{An})^\otimes := \mathrm{CMon}^{\mathrm{Op}\infty}(\mathrm{An}^\times).$$

The tensor product $-\otimes_R-$ turns $\mathrm{Proj}(R)$, which is already an object of $\mathrm{CMon}(\mathrm{Grpd})$ via $-\oplus-$, into an object of $\mathrm{CAlg}(\mathrm{CMon}(\mathrm{Grpd})^\otimes) \subseteq \mathrm{CAlg}(\mathrm{CMon}(\mathrm{An})^\otimes)$. The functors $(-)^\infty\text{-grp}: \mathrm{CMon}(\mathrm{An})^\otimes \rightarrow \mathrm{CGrp}(\mathrm{An})^\otimes$ and $B^\infty: \mathrm{CGrp}(\mathrm{An})^\otimes \rightarrow \mathrm{Sp}^\otimes$ are strongly monoidal by Proposition II.51 (the former is hidden under “similar assertions hold for other left adjoints among these”), hence $B^\infty k(R) \simeq B^\infty \mathrm{Proj}(R)^\infty\text{-grp}$ is indeed an element of CAlg .

Let's discuss now how to turn ordinary commutative rings and CDGAs into \mathbb{E}_∞ -ring spectra, as promised in (b):

II.55. Lemma. — *The Eilenberg–MacLane functor $H: \mathcal{D}(\mathbb{Z}) \rightarrow \mathrm{Sp}$ refines to a map*

$$H: \mathcal{D}(\mathbb{Z})^{\otimes_{\mathbb{Z}}^L} \longrightarrow \mathrm{Sp}^\otimes$$

of ∞ -operad, i.e. it is a lax symmetric monoidal functor. More generally, for arbitrary commutative rings R , the functor $C_\bullet(-, R) = C_\bullet(-) \otimes_{\mathbb{Z}}^L R: \mathrm{Sp} \rightarrow \mathcal{D}(R)$ has a left adjoint $H: \mathcal{D}(R) \rightarrow \mathrm{Sp}$, which refines to a lax symmetric functor

$$H: \mathcal{D}(R)^{\otimes_R^L} \longrightarrow \mathrm{Sp}^\otimes.$$

Proof. This isn't anything special about $\mathcal{D}(R)$, but an instance of the following more general principle:

- (*) *If \mathcal{C}^\otimes is a symmetric monoidal ∞ -category, then $\mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, -): \mathcal{C} \rightarrow \mathrm{An}$ has a canonical lax symmetric monoidal refinement. If \mathcal{C}^\otimes is stable as an ∞ -operad, then the same is true for the enriched Hom functor $\mathrm{hom}_{\mathcal{C}}(1_{\mathcal{C}}, -): \mathcal{C} \rightarrow \mathrm{Sp}$.*

The derived base change $-\otimes_{\mathbb{Z}}^L R: \mathcal{D}(\mathbb{Z}) \rightarrow \mathcal{D}(R)$ has the forgetful functor $\mathcal{D}(R) \rightarrow \mathcal{D}(\mathbb{Z})$ as a left adjoint, hence $C_{\bullet}(-, R)$ has indeed a right adjoint, which deserves to be named H as well since it really does nothing but forgetting to $\mathcal{D}(\mathbb{Z})$ and then applying $H: \mathcal{D}(\mathbb{Z}) \rightarrow \mathbf{Sp}$. Corollary II.30a implies $H \simeq \mathrm{hom}_{\mathcal{D}(R)}(R[0], -)$. Since $R[0] \in \mathcal{D}(R)$ is the tensor unit, we see that $(*)$ indeed implies the assertion of the lemma.

To prove $(*)$, let's first consider an arbitrary symmetric monoidal ∞ -category \mathcal{D} with all small operadic colimits. Then the Day convolution $\mathrm{Day}(\mathcal{C}^{\otimes}, \mathcal{D}^{\otimes})$ is symmetric monoidal and its tensor unit is given by

$$\mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, -) \otimes 1_{\mathcal{D}}: \mathcal{C} \longrightarrow \mathcal{D}.$$

Note that the tensor product here is neither $\otimes_{\mathcal{C}}$, nor $\otimes_{\mathcal{D}}$, but the one defined in the proof of Proposition II.34! That is, the functor under consideration is actually given by $\mathrm{colim}_{\mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, -)} \mathrm{const} 1_{\mathcal{D}}$, and one immediately checks that this coincides with the left Kan extension of $1_{\mathcal{D}}: * \rightarrow \mathcal{D}$ along $1_{\mathcal{C}}: * \rightarrow \mathcal{C}$, which is how we constructed the tensor unit in the proof of Proposition II.47(c). Now, per construction (just take a sharp look at Construction II.46 again), there is a map

$$\mathrm{Alg}_{\mathrm{Comm}}(\mathrm{Day}(\mathcal{C}^{\otimes}, \mathcal{D}^{\otimes})) \longrightarrow \mathrm{Fun}^{\mathrm{Op}\infty}(\mathcal{C}^{\otimes}, \mathcal{D}^{\otimes})$$

(it is even an equivalence, but we won't need that). By Lemma/Definition II.45a, the tensor unit of $\mathrm{Day}(\mathcal{C}^{\otimes}, \mathcal{D}^{\otimes})$ defines canonically an element of the left-hand side, hence also an element of the right-hand side. This shows that $\mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, -) \otimes 1_{\mathcal{D}}$ has a canonical lax symmetric monoidal refinement.

Now plug in $\mathcal{D}^{\otimes} \simeq \mathbf{An}^{\times}$. Then $1_{\mathcal{D}} \simeq *$ and $X \otimes * \simeq \mathrm{colim}_X \mathrm{const} * \simeq X$ for all $X \in \mathbf{An}$, hence we obtain that $\mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, -): \mathcal{C} \rightarrow \mathbf{An}$ is the Day convolution tensor unit and thus has a lax symmetric monoidal refinement $\mathrm{Hom}_{\mathcal{C}}(1_{\mathcal{C}}, -): \mathcal{C}^{\otimes} \rightarrow \mathbf{An}^{\times}$, as required. Moreover, it preserves limits (which are automatically operadic), hence the refinement lifts to $\mathrm{hom}_{\mathcal{C}}(1_{\mathcal{C}}, -): \mathcal{C}^{\otimes} \rightarrow \mathbf{Sp}^{\otimes}$ by Theorem II.50 if \mathcal{C}^{\otimes} is a stable ∞ -operad. \square

Now we can resume our discussion on how to include commutative rings and CDGAs into CAlg. Observe that we have two more lax symmetric monoidal functors

$$\mathrm{Ch}(\mathbb{Z})^{\otimes_{\mathbb{Z}}} \longrightarrow \mathcal{K}(\mathbb{Z})^{\otimes_{\mathbb{Z}}} \longrightarrow \mathcal{D}(\mathbb{Z})^{\otimes_{\mathbb{Z}}^L},$$

where $\mathrm{Ch}(\mathbb{Z})$ is the 1-category of chain complexes over \mathbb{Z} . The left functor comes from the fact that $\mathcal{K}(\mathbb{Z}) \simeq \mathbf{N}^c(\mathrm{Ch}(\mathbb{Z}))$ is the coherent nerve of the Kan-enrichment of $\mathrm{Ch}(\mathbb{Z})$. The right functor was constructed in Example II.37(e). Unwinding definitions, we see that $\mathrm{CAlg}(\mathrm{Ch}(\mathbb{Z})^{\otimes_{\mathbb{Z}}}) \simeq \mathrm{CDGAlg}$ is the 1-category of commutative differential graded algebras. Moreover, we have an inclusion $\mathbf{Ab}^{\otimes} \subseteq \mathrm{Ch}(\mathbb{Z})^{\otimes_{\mathbb{Z}}}$ of symmetric monoidal 1-categories, given by interpreting abelian groups as chain complexes concentrated in degree 0, which gives a map $\mathrm{CRing} \simeq \mathrm{CAlg}(\mathbf{Ab}^{\otimes}) \rightarrow \mathrm{CAlg}(\mathrm{Ch}(\mathbb{Z})^{\otimes_{\mathbb{Z}}}) \simeq \mathrm{CDGAlg}$. Summarizing, if we apply $\mathrm{CAlg}(-)$ everywhere, we obtain functors

$$\mathrm{CRing} \longrightarrow \mathrm{CDGAlg} \longrightarrow \mathrm{CAlg}(\mathcal{D}(\mathbb{Z})^{\otimes_{\mathbb{Z}}^L}) \longrightarrow \mathrm{CAlg}.$$

In fact, it turns out that we can completely import ordinary commutative ring theory:

II.55a. Corollary. — *The above induces a fully faithful functor $\mathrm{CRing} \rightarrow \mathrm{CAlg}$ with essential image those $R \in \mathrm{CAlg}$ for which $\pi_i(R) = 0$ for all $i \neq 0$.*

Here we define the homotopy groups of an \mathbb{E}_∞ -ring spectrum as those of its underlying spectrum. Also note that in the composition above, the middle functor $\mathrm{CDGAlg} \rightarrow \mathrm{CAlg}(\mathcal{D}(\mathbb{Z})^{\otimes \frac{L}{2}})$ from the 1-category of CDGAs to the ∞ -category of \mathbb{E}_∞ -chain complexes is not fully faithful, so this is really a thing about the composition.

Proof sketch of Corollary II.55a.* The proof wasn't discussed in the lecture, but is taken from Fabian's lecture notes, [A&HK, Remarks II.53(ii)]. Recall from Example II.51b that

$$\mathrm{Ab}^\otimes \simeq \mathrm{CGrp}^{\mathrm{Op}\infty}(\mathrm{Set}^\times) \longrightarrow \mathrm{CGrp}^{\mathrm{Op}\infty}(\mathrm{An}^\times) \simeq \mathrm{Sp}_{\geq 0}^\otimes$$

is fully faithful with left adjoint π_0 . The adjunction, together with the condition that the counit is an equivalence, persists to taking $\mathrm{CAlg}(-)$ everywhere. This almost shows what we want, except that we have to explain why the functor $i^\otimes: \mathrm{CRing} \rightarrow \mathrm{CAlg}(\mathrm{Sp}_{\geq 0}^\otimes) \subseteq \mathrm{CAlg}$ that we get out is the one induced by H , i.e. the one from the formulation of the corollary. It's more or less clear from the constructions that the underlying functor of i^\otimes is indeed $H \simeq \mathrm{hom}_{\mathrm{Ab}}(\mathbb{Z}, -): \mathrm{Ab} \rightarrow \mathrm{Sp}$, but we need to check that the lax symmetric monoidal refinement is the right one.

Fabian outlines two ways to do so. One can use the result from Example II.37(f) to see that lax monoidal refinements of right adjoints are equivalent to oplax monoidal refinements of left adjoints, so it suffices that both candidates induce the same refinement of π_0 .

Alternatively, one can argue as follows: By construction and Lemma/Definition II.45a, the lax symmetric monoidal refinement of $\mathrm{Hom}_{\mathrm{Ab}}(\mathbb{Z}, -): \mathrm{Ab} \rightarrow \mathrm{An}$ from the proof of Lemma II.55 is initial in

$$\mathrm{CAlg}(\mathrm{Day}(\mathrm{Ab}^\otimes, \mathrm{An}^\times)) \simeq \mathrm{Fun}^{\mathrm{Op}\infty}(\mathrm{Ab}^\otimes, \mathrm{An}^\times)$$

(we didn't need this to be an equivalence in the proof of Lemma II.55). Hence there is a unique map to $\Omega^\infty i^\otimes$, which must be an equivalence since it is an equivalence on underlying functor $\mathrm{Ab} \rightarrow \mathrm{An}$ (and then Segal conditions do the rest). Now use Theorem II.50 to see that our lax symmetric monoidal refinement of $\mathrm{hom}_{\mathrm{Ab}}(\mathbb{Z}, -)$ coincides with i^\otimes . \square

Before we go on and construct module categories, I'd like to go on another detour and prove that the homology and cohomology theories associated to a spectrum really do what they're supposed to do.

II.55b. Corollary*. — *The homology and cohomology associated to a spectrum E by Definition II.35 can be described as*

$$\begin{aligned} E_*(-) &\simeq \pi_*(\mathbb{S}[-] \otimes E) & \text{and} & & E^*(-) &\simeq \pi_{-*} \mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}[-], E) \\ &\simeq \mathrm{Tor}_*^{\mathbb{S}}(\mathbb{S}[-], E) & & & &\simeq \mathrm{Ext}_{\mathbb{S}}^*(\mathbb{S}[-], E) \end{aligned}$$

They satisfy the Eilenberg–Steenrod axioms. Moreover, the homology and cohomology theory associated to an Eilenberg–MacLane spectrum HA for some abelian group A coincide with singular homology and cohomology, i.e.

$$HA_*(-) \simeq H_*(-, A): \mathrm{An} \longrightarrow \mathrm{Ab} \quad \text{and} \quad HA^*(-) \simeq H^*(-, A): \mathrm{An}^{\mathrm{op}} \longrightarrow \mathrm{Ab}.$$

Finally, the homology and cohomology of a spectrum E itself, as defined in II.26, is given by

$$H_*(E, A) \simeq \pi_*(E \otimes HA) \quad \text{and} \quad H^*(E, A) \simeq \pi_{-*} \mathrm{hom}_{\mathrm{Sp}}(E, HA).$$

*Proof**. The description of $E_*(-)$ follows from our computations in [II.53](#). Along the same lines one can also prove $E^X \simeq \text{hom}_{\text{Sp}}(\mathbb{S}[X], E)$, hence the description of $E^*(-)$.

Next we check the Eilenberg–Steenrod axioms. To construct the long exact sequences and excision, first note that fibre and cofibre sequences in the stable ∞ -category Sp coincide, and that to each such sequence there is an associated long exact sequence on homotopy group (because the functors $\Omega^{\infty-i}: \text{Sp} \rightarrow */\text{An}$ preserve fibre sequences, so the long exact sequences from $*/\text{An}$ are inherited). Now both $\mathbb{S}[-] \otimes E$ and $\text{hom}_{\text{Sp}}(\mathbb{S}[-], E)$ transform cofibre sequences in An into fibre sequences in Sp , providing the long exact sequences for $E_*(-)$ and $E^*(-)$.

It remains to see that they transform disjoint unions into direct sums/products. For $E^*(-)$, this is easy: $\text{hom}_{\text{Sp}}(\mathbb{S}[-], E)$ sends disjoint unions to products, and taking homotopy groups commutes with products. For $E_*(-)$, we can argue similarly that $\mathbb{S}[-] \otimes E$ sends disjoint unions to coproducts, but now we need to show that π_* sends coproducts of spectra to products. Arbitrary coproducts can be written as filtered colimits of finite coproducts, and π_* commutes with filtered colimits (Remark* [II.31b](#)), hence it suffices to check the case of finite coproducts. But Sp is stable, hence finite coproducts agree with finite products, and π_* preserves the latter.

Finally, let’s do our reality check for Eilenberg–MacLane spectra. The cohomology case is easy: From the (C_\bullet, H) -adjunction combined with Corollary [II.30a](#), we get

$$\text{hom}_{\text{Sp}}(E, HA) \simeq \text{hom}_{\mathcal{D}(\mathbb{Z})}(C_\bullet(E), A) \simeq H(R\text{Hom}_{\mathbb{Z}}(C_\bullet(E), A))$$

By Example [II.25\(c\)](#), the homotopy groups of the right-hand side are the homology groups of $R\text{Hom}_{\mathbb{Z}}(C_\bullet(E), A)$, which are the singular cohomology groups $H^*(E, A)$ (in negative degrees) by definition. Since $H^*(\mathbb{S}[X], A) \simeq H^*(X, A)$ for all $X \in \text{An}$ by Lemma [II.31c](#), plugging in the special case $E \simeq \mathbb{S}[X]$ also shows $HA^*(X) \simeq H^*(X, A)$.

For the homology case we actually need to use fact that H is lax symmetric monoidal by Lemma [II.55](#). Combining this with the unit $E \rightarrow HC_\bullet(E)$ of the (C_\bullet, H) -adjunction gives functorial maps

$$E \otimes HA \longrightarrow HC_\bullet(E) \otimes HA \longrightarrow H(C_\bullet(E) \otimes_{\mathbb{Z}}^L A).$$

If we can show that the composition is an equivalence, taking homotopy groups on both sides will show $\pi_*(E \otimes HA) \simeq H_*(E, A)$, as desired, and then the special case $E \simeq \mathbb{S}[X]$ together with Lemma [II.31c](#) will also show $HA_*(X) \simeq H_*(X, A)$.

To show that the composition is indeed an equivalence, first note that it is an equivalence for $E \simeq \mathbb{S}$ by Lemma [II.31c](#), and that both sides commute with shifts and colimits in E . For this we need that H commutes with colimits, which is perhaps a bit surprising at first, given that H is only a right adjoint, but we give an argument in the proof sketch* of Theorem [II.57](#) below (and once you know this theorem, H preserving colimits is not that surprising any more). Now all of Sp can be generated from \mathbb{S} using shifts and colimits, hence

$$E \otimes HA \xrightarrow{\sim} H(C_\bullet(E) \otimes_{\mathbb{Z}}^L A)$$

is an equivalence everywhere. □

∞ -Categories of Modules

\mathbb{E}_∞ -ring spectra have categories of modules that behave like the derived categories of modules over ordinary rings. In particular, as we’ll see in Theorem [II.57](#) below, the category of modules over HR for some $R \in \text{CRing}$ is nothing else but $\mathcal{D}(R)$.

As a first step, we'll construct certain operads $\mathbb{A}\text{ssoc}$ and $\mathbb{L}\text{Mod}$. These have already been teased in Example II.45(d) and (e), so we'll hijack that numbering and define them properly.

II.45. Example. — What it should have been:

- (d) We would like to have an ∞ -operad that describes associative monoids. So we define $\mathbb{E}_1 = \mathbb{A}\text{ssoc} \rightarrow \mathbb{I}^{\text{op}}$ as follows: The objects of $\mathbb{A}\text{ssoc}$ are finite sets and a morphism $I \rightarrow J$ is a map in \mathbb{I}^{op} together with a total ordering on each preimage $\alpha^{-1}(j)$. Composition is given by composition in \mathbb{I}^{op} and lexicographic ordering. If we equip the 1-category $\mathbb{A}\text{ssoc}$ with the obvious functor $\mathbb{A}\text{ssoc} \rightarrow \mathbb{I}^{\text{op}}$, it becomes an ∞ -operad with underlying category $\mathbb{A}\text{ssoc}_1 = \{*\}$ and

$$\text{Hom}_{\mathbb{A}\text{ssoc}}^{\text{act}}(\underbrace{(*, \dots, *)}_{n \text{ entries}}, *) = \{\text{total orders on } \langle n \rangle\}.$$

(just as there are $n!$ ways to multiply n factors in an associative, but not necessarily commutative monoid).

- (e) We would also like to have an ∞ -operad that describes an associative monoid A together with an object M it acts upon (we think of this as an algebra together with a module over it). So we define $\mathbb{L}\text{Mod} \rightarrow \mathbb{I}^{\text{op}}$ as follows: Its objects are pairs (I, S) with $I \in \mathbb{I}^{\text{op}}$ and $S \subseteq I$ (think of I as the set of factors we'd like to multiply, and of S as the set of those factors that come from M ; of course, we can't multiply more than one element from M). A map $(I, S) \rightarrow (J, T)$ is given by a map $\alpha: I \rightarrow J$ in $\mathbb{A}\text{ssoc}$ such that $\alpha(S) = T$ and for all $t \in T$ the set $\alpha^{-1}(t) \cap S$ consists exactly of the maximal element of $\alpha^{-1}(t)$. Composition is inherited from $\mathbb{A}\text{ssoc}$. Again, if we equip the 1-category $\mathbb{L}\text{Mod}$ with the obvious map $\mathbb{L}\text{Mod} \rightarrow \mathbb{I}^{\text{op}}$, it becomes an ∞ -operad. The underlying category $\mathbb{L}\text{Mod}_1$ has precisely two objects: $a = (\langle 1 \rangle, \emptyset)$ and $m = (\langle 1 \rangle, \langle 1 \rangle)$, which we think of corresponding to A and M respectively. We have

$$\text{Hom}_{\mathbb{L}\text{Mod}}^{\text{act}}((b_1, \dots, b_n), a) = \begin{cases} \emptyset & \text{if } b_i = m \text{ for some } i \\ \{\text{total orders on } \langle n \rangle\} & \text{if } b_i = a \text{ for all } i \end{cases}$$

This makes sense since “as soon as one of the factors is from M , the result will be no longer an element of A , but one of M ”. Similarly,

$$\text{Hom}_{\mathbb{L}\text{Mod}}^{\text{act}}((b_1, \dots, b_n), m) = \begin{cases} \emptyset & \text{unless exactly one } b_i = m \\ \left\{ \begin{array}{l} \text{total orders on } \langle n \rangle \\ \text{with max. element } i \end{array} \right\} & \text{if } b_i = m \end{cases}$$

since there can be only one factor from M in any product, and if there is, it must be the right-most factor.

Sending the unique element $* \in \mathbb{A}\text{ssoc}_1$ to $a \in \mathbb{L}\text{Mod}_1$ defines a map $\mathbb{A}\text{ssoc} \rightarrow \mathbb{L}\text{Mod}$ of ∞ -operads, which induces a map $a^*: \text{Alg}_{\mathbb{L}\text{Mod}}(\mathcal{C}^{\otimes}) \rightarrow \text{Alg}_{\mathbb{A}\text{ssoc}}(\mathcal{C}^{\otimes})$ (“we forget M and only retain A ”) for any symmetric monoidal category \mathcal{C}^{\otimes} . For any $A \in \text{Alg}_{\mathbb{A}\text{ssoc}}(\mathcal{C}^{\otimes})$, the pullback

$$\begin{array}{ccc} \text{LMod}_A(\mathcal{C}^{\otimes}) & \longrightarrow & \text{Alg}_{\mathbb{L}\text{Mod}}(\mathcal{C}^{\otimes}) \\ \downarrow & \lrcorner & \downarrow a^* \\ * & \xrightarrow{A} & \text{Alg}_{\mathbb{A}\text{ssoc}}(\mathcal{C}^{\otimes}) \end{array}$$

is called the ∞ -category of left modules in \mathcal{C} over A . We can convince ourselves in examples that this terminology really makes sense: If $\mathcal{C}^\otimes = \text{Set}^\times$, then an algebra $A \in \text{Alg}_{\text{Assoc}}(\text{Set}^\times) \simeq \text{Mon}(\text{Set})$ is a monoid and elements of $\text{LMod}_A(\text{Set}^\times)$ are really left A -sets. Similarly, if $\mathcal{C}^\otimes = \text{Ab}^\otimes$, then A is an associative (not necessarily commutative) ring and $\text{LMod}_A(\text{Ab}^\otimes)$ is really the 1-category of left A -modules.

II.56. General Properties of Module ∞ -Categories. — We will now discuss (and partially prove) a series of results to show that Assoc and LMod really do what they are supposed to do, and to see how one works with the left module ∞ -categories $\text{LMod}_A(\mathcal{C}^\otimes)$. After that, we are ready for modules over \mathbb{E}_∞ -ring spectra!

First off, recall the functor $\text{Cut}: \Delta^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$ from Definition II.15, which sends a totally ordered set I to its set of Dedekind cuts. Since the set of Dedekind cuts inherits a total order, Cut lifts to a functor $\text{Cut}: \Delta^{\text{op}} \rightarrow \text{Assoc}$ over \mathbb{I}^{op} . Also recall the notation introduced before Theorem II.43: For any ∞ -operad \mathcal{O} and any ∞ -category \mathcal{C} with finite products, we denote by $\mathcal{O}\text{Mon}(\mathcal{C}) \subseteq \text{Fun}(\mathcal{O}, \mathcal{C})$ the full sub- ∞ -category of functors satisfying the Segal condition.

II.56a. Theorem (Lurie, [HA, Proposition 4.1.2.10]). — *If \mathcal{C} has finite products, then the induced functor*

$$\text{Cut}^*: \text{AssocMon}(\mathcal{C}) \simeq \text{Alg}_{\text{Assoc}}(\mathcal{C}^\times) \xrightarrow{\sim} \text{Mon}(\mathcal{C})$$

is an equivalence. More generally, if \mathcal{C}^\otimes is any symmetric monoidal ∞ -category, then

$$\text{Cut}^*: \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes) \longrightarrow \text{Fun}(\Delta^{\text{op}}, \mathcal{C}^\otimes) \times_{\text{Fun}(\Delta^{\text{op}}, \mathbb{I}^{\text{op}})} \{\text{Cut}\}$$

is fully faithful, with essential image those functors $F: \Delta^{\text{op}} \rightarrow \mathcal{C}^\otimes$ that send inerts to inerts. That is, whenever $\alpha: [n] \rightarrow [m]$ is inert in Δ (without $^{\text{op}}$), the induced morphism $F(\alpha^{\text{op}}): F([m]) \rightarrow F([n])$ is inert in \mathcal{C}^\otimes .

II.56b. Remark*. — We won't prove Theorem II.56a, but let me at least give some additional motivation. The condition that F preserves inerts is just a somewhat unusual formulation of the Segal condition. Indeed, if we denote $A = F([1]) \in \mathcal{C}_1^\otimes \simeq \mathcal{C}$, then the fact that the Segal maps $e_i: [1] \rightarrow [n]$ induce inert maps in \mathcal{C}^\otimes shows

$$F([n]) \simeq (A, \dots, A) \in \mathcal{C}_n^\otimes \simeq \mathcal{C}^n$$

(in particular, $F([0]) \simeq 1_{\mathcal{C}}$). For general $\alpha: [n] \rightarrow [m]$ in Δ , we can think of the induced map $F(\alpha^{\text{op}}): F([m]) \rightarrow F([n])$ as follows: Let $\text{Cut}(\alpha^{\text{op}})_*: (A, \dots, A) \rightarrow (B_1, \dots, B_n)$ be a cocartesian lift of $\text{Cut}(\alpha^{\text{op}}): \langle m \rangle \rightarrow \langle n \rangle$. A quick unravelling shows

$$B_i \simeq \bigotimes_{\alpha(i-1) < j \leq \alpha(i)} A$$

for $i = 1, \dots, n$. Then $\text{Cut}(\alpha^{\text{op}})_*$ and $F(\alpha^{\text{op}})$ induce a map $\Lambda_0^2 \rightarrow \mathcal{C}^\otimes$, which has an extension to Δ^2 since $\mathcal{C}^\otimes \rightarrow \mathbb{I}^{\text{op}}$ is a cocartesian fibration. Restricting the extension to $\Delta^{\{1,2\}}$ gives maps $B_i \rightarrow A$ in \mathcal{C} , which we think of as the “multiplication maps in the algebra A ”.

LECTURE 18
12th Jan, 2021

There is a similar description of $\text{Alg}_{\text{LMod}}(\mathcal{C}^\otimes)$. To this end, we can extend Cut to a functor $\text{LCut}: \Delta^{\text{op}} \times [1] \rightarrow \text{LMod}$ that fits into a diagram

$$\begin{array}{ccc} \Delta^{\text{op}} & \xrightarrow{\text{Cut}} & \text{Assoc} \\ \downarrow - \times \{1\} & & \downarrow a \\ \Delta^{\text{op}} \times [1] & \xrightarrow{\text{LCut}} & \text{LMod} \end{array} \quad \begin{array}{c} \nearrow p \\ \searrow \end{array} \quad \mathbb{I}^{\text{op}}$$

LCut sends elements of the form $([n], 1)$ to the “algebra-like” element $(a, \dots, a) \in \mathbb{LMod}_n$ (this guy has n repetitions of a), and elements of the form $([n], 0)$ to the “module-like” element $(a, \dots, a, m) \in \mathbb{LMod}_{n+1}$ (this guy also has n repetitions of a). In particular, and this might be confusing at first, the structure map $p \circ \text{LCut}: \Delta^{\text{op}} \times [1] \rightarrow \Gamma^{\text{op}}$ is *not* given by projecting to the first factor and applying $\text{Cut}: \Delta^{\text{op}} \rightarrow \Gamma^{\text{op}}$. But if you think about this, it makes sense to have it that way: The morphisms $\text{LCut}([n], 0) \rightarrow \text{LCut}([n], 1)$ should be given by forgetting the “module part” of (a, \dots, a, m) and only retaining the algebra part (a, \dots, a) —which only works if the morphism in question runs from \mathbb{LMod}_{n+1} to \mathbb{LMod}_n .

II.56c. Theorem (Lurie, [HA, Proposition 4.2.2.12]). — *The functor*

$$\text{LCut}^*: \text{Alg}_{\mathbb{LMod}}(\mathcal{C}^{\otimes}) \longrightarrow \text{Fun}(\Delta^{\text{op}} \times [1], \mathcal{C}^{\otimes}) \times_{\text{Fun}(\Delta^{\text{op}} \times [1], \Gamma^{\text{op}})} \{p \circ \text{LCut}\}$$

is fully faithful, with essential image those functors $F: \Delta^{\text{op}} \times [1] \rightarrow \mathcal{C}^{\otimes}$ that fulfill the following conditions:

- (a) The “algebra-like” part $F|_{\Delta^{\text{op}} \times \{1\}}$ of F satisfies the condition from Theorem II.56a. That is, whenever $\alpha: [n] \rightarrow [m]$ is inert in Δ (without $^{\text{op}}$), the induced morphism $F(\alpha^{\text{op}}, \text{id}_1): F([m], 1) \rightarrow F([n], 1)$ is inert in \mathcal{C}^{\otimes} .
- (b) $F([n], 0) \rightarrow F([n], 1)$ is always inert in \mathcal{C}^{\otimes} .
- (c) If $\alpha^{\text{op}}: [n] \rightarrow [m]$ is an inert morphism in Δ satisfying $\alpha(n) = m$, then the induced morphism $F(\alpha^{\text{op}}, \text{id}_0): F([m], 0) \rightarrow F([n], 0)$ is inert in \mathcal{C}^{\otimes} .

II.56d. Remark*. — Again, we won’t prove Theorem II.56c, but at least explain how the structure of an algebra and a module over it are encoded in the right-hand side. Let $M = F([0], 0)$ and $A = F([1], 1)$; both are elements of $\mathcal{C}_1^{\otimes} \simeq \mathcal{C}$ by construction. By condition (a), the part $F|_{\Delta^{\text{op}} \times \{1\}}$ encodes the algebra structure on A , so in particular, $F([n], 1) \simeq (A, \dots, A) \in \mathcal{C}^n$ as in Remark* II.56b. Combining this with conditions (b) and (c) shows

$$F([n], 0) \simeq (A, \dots, A, M) \in \mathcal{C}^{n+1}.$$

Given a general map $\alpha: [n] \rightarrow [m]$ in Δ , we can think of $F(\alpha^{\text{op}}, \text{id}_0): F([m], 0) \rightarrow F([n], 0)$ as follows: Choose a cocartesian lift $p(\text{LCut}(\alpha^{\text{op}}))_*: (A, \dots, A, M) \rightarrow (B_1, \dots, B_n, N)$ of $p(\text{LCut}(\alpha^{\text{op}})): \langle m+1 \rangle \rightarrow \langle n+1 \rangle$. Then

$$B_i \simeq \bigotimes_{\alpha(i-1) < j \leq \alpha(i)} A, \quad \text{and} \quad N \simeq \left(\bigotimes_{\alpha(n) < j \leq m} A \right) \otimes M.$$

As in Remark* II.56b, we get maps $B_i \rightarrow A$ and $N \rightarrow M$, which encode the multiplication in A and the left action of A on M .

In particular Theorem II.56a says that any algebra $A \in \text{Alg}_{\text{Assoc}}(\mathcal{C}^{\otimes})$ defines an element $\text{Cut}^*(A) \in \text{Fun}_{\Gamma^{\text{op}}}(\Delta^{\text{op}}, \mathcal{C}^{\otimes})$ which satisfies the Segal condition. And then by Theorem II.56c, we find that $\mathbb{LMod}_A(\mathcal{C}^{\otimes})$ is a full sub- ∞ -category of the pullback

$$\begin{array}{ccc} P & \longrightarrow & \text{Fun}_{\Gamma^{\text{op}}}(\Delta^{\text{op}} \times [1], \mathcal{C}^{\otimes}) \\ \downarrow & \lrcorner & \downarrow (- \times \{1\})^* \\ * & \xrightarrow{\text{Cut}^*(A)} & \text{Fun}_{\Gamma^{\text{op}}}(\Delta^{\text{op}}, \mathcal{C}^{\otimes}) \end{array}$$

Note that besides the operad map $a: \text{Assoc} \rightarrow \mathbb{L}\text{Mod}$ of extracting algebras, there is also a map $m: \text{Triv} \rightarrow \mathbb{L}\text{Mod}$ that extracts the module object. Also recall that $\text{Alg}_{\text{Triv}}(\mathcal{C}^\otimes) \simeq \mathcal{C}$ as shown in Example II.45(b).

II.56e. Corollary. — *If \mathcal{C}^\otimes has small/finite operadic colimits, then $\text{LMod}_A(\mathcal{C}^\otimes)$ has small/finite colimits, and these are computed “underlyingly”. That is, the functor*

$$m^*: \text{LMod}_A(\mathcal{C}^\otimes) \longrightarrow \text{Alg}_{\text{Triv}}(\mathcal{C}^\otimes) \simeq \mathcal{C}$$

preserves small/finite colimits. Likewise, if \mathcal{C} has small/finite limits (which are automatically operadic), then so has $\text{LMod}_A(\mathcal{C}^\otimes)$, and m^ preserves small/finite limits. In particular, if \mathcal{C}^\otimes is a stable ∞ -operad, then $\text{LMod}_A(\mathcal{C}^\otimes)$ is a stable ∞ -category for all $A \in \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)$.*

Proof sketch. Let $F: \mathcal{I} \rightarrow \text{LMod}_A(\mathcal{C}^\otimes)$ be a diagram. Then Theorem II.56c induces a diagram $F: \mathcal{I} \rightarrow \text{Fun}(\Delta^{\text{op}} \times [1], \mathcal{C}^\otimes)$ satisfying

$$F(i)([n], 1) \simeq (A, \dots, A) \quad \text{and} \quad F([n], 0) \simeq (A, \dots, A, M(i)),$$

where $M: \mathcal{I} \rightarrow \mathcal{C}$ is another diagram (the “module part” of F). We have to check that the there are functors

$$\begin{aligned} \overline{F}_{\text{lim}}, \overline{F}_{\text{colim}}: \mathcal{I} \times \Delta^{\text{op}} \times [1] &\longrightarrow \mathcal{C}^\otimes \\ (i, [n], 0) &\longmapsto (A, \dots, A) \\ (i, [n], 1) &\longmapsto \begin{cases} (A, \dots, A, \lim_{i \in \mathcal{I}} M(i)) & \text{for } \overline{F}_{\text{lim}} \\ (A, \dots, A, \text{colim}_{i \in \mathcal{I}} M(i)) & \text{for } \overline{F}_{\text{colim}} \end{cases} \end{aligned}$$

Moreover, we need to verify that $\overline{F}_{\text{lim}}, \overline{F}_{\text{colim}}$ constitute the limit and colimit over F , taken inside the pullback P above, and that both satisfy the conditions from Theorem II.56c so that they lie inside the full subcategory $\text{LMod}_A(\mathcal{C}^\otimes) \subseteq P$. We didn’t work this out in the lecture, but in these notes I would like to give a rough sketch of what I think happens.

Naively, I would expect that the limit and colimit over F , taken in P , should be given by some kind of pointwise limit in $\text{Fun}(\Delta^{\text{op}} \times [1], \mathcal{C}^\otimes)$. There are two problems though: First, we see from Remark* II.48a that $(A, \dots, A, \text{colim}_{i \in \mathcal{I}} M(i))$ doesn’t necessarily coincide with the colimit $\text{colim}_{i \in \mathcal{I}} (A, \dots, A, M(i))$ in \mathcal{C}^\otimes , even though colimits in \mathcal{C} are operadic, unless \mathcal{I} is weakly contractible. This problem doesn’t occur in the limit case, but there is a second one, which affects both cases: We aren’t taking limits/colimits in $\text{Fun}(\Delta^{\text{op}} \times [1], \mathcal{C}^\otimes)$, but in some pullback P of it. And if \mathcal{I} isn’t weakly contractible, then the limit/colimit over the constant A -diagram in $\text{Fun}_{\Gamma^{\text{op}}}(\Delta^{\text{op}}, \mathcal{C}^\otimes)$ might not be A any more. Hence we can’t expect that limits/colimits in the pullback P can be computed in each factor separately.

Fortunately, there is a trick that solves both problems at once. Since \mathcal{C} is cocomplete by assumption, it has an initial object, hence the diagram $M: \mathcal{I} \rightarrow \mathcal{C}$ extends canonically to a diagram $M^\triangleleft: \mathcal{I}^\triangleleft \rightarrow \mathcal{C}$ that sends the tip of the cone to the initial object. Moreover,

$$\text{colim}_{i \in \mathcal{I}} M(i) \simeq \text{colim}_{i \in \mathcal{I}^\triangleleft} M^\triangleleft(i),$$

and the right-hand side is also a colimit in \mathcal{C}^\otimes by Remark* II.48a, since $\mathcal{I}^\triangleleft$ is weakly contractible. Similarly, taking the colimit over an $\mathcal{I}^\triangleleft$ -shaped in $\text{Fun}_{\Gamma^{\text{op}}}(\Delta^{\text{op}}, \mathcal{C}^\otimes)$ which is constant on A gives A again. After some unravelling, we see (I think) that this gives a way of computing \mathcal{I} -shaped colimits in P by factorwise $\mathcal{I}^\triangleleft$ -shaped colimits. A similar trick works for limits. I’ll leave it to you to make this argument precise and also to check that the conditions from Theorem II.56c are satisfied. \square

The structure map $\text{Assoc} \rightarrow \mathbb{I}^{\text{op}}$ defines an ∞ -operad map $\text{Assoc} \rightarrow \text{Comm}$, which in turn induces a map $\mathcal{C}\text{Alg}(\mathcal{C}^\otimes) \simeq \text{Alg}_{\text{Comm}}(\mathcal{C}^\otimes) \rightarrow \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)$, as one would expect. In particular, the tensor unit $1_{\mathcal{C}} \in \mathcal{C}$ also defines an associative algebra. In fact, one can show similar to Lemma/Definition II.45a that $1_{\mathcal{C}}$ is also initial in $\text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)$.

II.56f. Corollary. — *For any symmetric monoidal ∞ -category $q: \mathcal{C}^\otimes \rightarrow \mathbb{I}^{\text{op}}$, the forgetful functor*

$$m^*: \text{LMod}_{1_{\mathcal{C}}}(\mathcal{C}^\otimes) \xrightarrow{\sim} \text{Alg}_{\text{Triv}}(\mathcal{C}^\otimes) \simeq \mathcal{C}$$

is an equivalence.

Proof sketch. Since the map $\emptyset \rightarrow (1_{\mathcal{C}})$ in \mathcal{C}^\otimes is a q -cocartesian lift of $\langle 0 \rangle \rightarrow \langle 1 \rangle$, we see that a functor $F: \Delta^{\text{op}} \times [1] \rightarrow \mathcal{C}^\otimes$ fulfills the conditions from Theorem II.56c iff all induced maps in \mathcal{C}^\otimes are q -cocartesian. But since $([0], 0)$ is initial in $\Delta^{\text{op}} \times [1]$, a functor $F: \Delta^{\text{op}} \times [1] \rightarrow \mathcal{C}^\otimes$ with $F([1], 1) \simeq 1_{\mathcal{C}}$ is the same thing as a natural transformation

$$\eta_F: \text{const } F([0], 0) \Longrightarrow F|_{\Delta^{\text{op}} \times [1] \setminus \{([0], 0)\}}$$

in $\text{Fun}(\Delta^{\text{op}} \times [1] \setminus \{([0], 0)\}, \mathcal{C}^\otimes)$. In other words, F corresponds to a lift in the diagram

$$\begin{array}{ccc} [0] & \xrightarrow{\text{const } F([0], 0)} & \text{Fun}(\Delta^{\text{op}} \times [1] \setminus \{([0], 0)\}, \mathcal{C}^\otimes) \\ \downarrow & \nearrow \eta_F & \downarrow q_* \\ [1] & \xrightarrow{\text{const } \langle 0 \rangle \Rightarrow p \circ \text{LCut}} & \text{Fun}(\Delta^{\text{op}} \times [1] \setminus \{([0], 0)\}, \mathbb{I}^{\text{op}}) \end{array}$$

(in the bottom arrow, $p: \mathbb{L}\text{Mod} \rightarrow \mathbb{I}^{\text{op}}$ denotes the structure map of $\mathbb{L}\text{Mod}$).

As we've seen above, F defines an element of $\text{LMod}_{1_{\mathcal{C}}}(\mathcal{C}^\otimes)$ iff it takes all maps in $\Delta^{\text{op}} \times [1]$ to q -cocartesian ones. Using [HCII, Corollary IX.25], this is equivalent to η_F being a q_* -cocartesian edge. Hence the space of lifts η_F , and thus the space of functors F with given $F([0], 0)$, is the space of q_* -cocartesian lifts in the above diagram. Now the functor $m^*: \text{LMod}_{1_{\mathcal{C}}}(\mathcal{C}^\otimes) \rightarrow \mathcal{C}$ corresponds to the choice of $F([0], 0)$, i.e. to the choice of starting point. Since q_* is a cocartesian fibration, we see that m^* is essentially surjective, and since a map between starting points defines a contractible space of maps between q_* -cocartesian lifts (you can make this precise using [HCII, Proposition IX.24]), we see that m^* is fully faithful. \square

Fabian's script mentions a (stronger version of) another corollary, which we'll need to prove Theorem II.57 below. I thought about whether there is an easy proof using what we know so far, but didn't succeed, so for now I'll just refer to Lurie.

II.56g. Corollary* (Lurie, [HA, Proposition 4.2.4.2]). — *Let \mathcal{C}^\otimes be a symmetric monoidal ∞ -category and $A \in \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)$ an algebra in it (whose underlying object we will also denote $A \in \mathcal{C}$). Then the functor $A \otimes -: \mathcal{C} \rightarrow \mathcal{C}$ lifts to a functor*

$$\begin{array}{ccc} & & \text{LMod}_A(\mathcal{C}^\otimes) \\ & \nearrow A \otimes - & \downarrow m^* \\ \mathcal{C} & \xrightarrow{A \otimes -} & \mathcal{C} \end{array}$$

which is right-adjoint to the forgetful functor $m^: \text{LMod}_A(\mathcal{C}^\otimes) \rightarrow \mathcal{C}$. In particular, we see that $A \in \text{LMod}_A(\mathcal{C}^\otimes)$.*

With that out of the way, let's discuss a cool application. Any \mathbb{E}_∞ -ring spectrum R defines an element of $\mathrm{Alg}_{\mathrm{Assoc}}(\mathrm{Sp}^\otimes)$ via the canonical map $\mathrm{CAlg} \rightarrow \mathrm{Alg}_{\mathrm{Assoc}}(\mathrm{Sp}^\otimes)$. Consequently, we may define the ∞ -category of modules over R as

$$\mathrm{Mod}_R := \mathrm{LMod}_R(\mathrm{Sp}^\otimes).$$

We know from Corollary II.56e that this is a stable ∞ -category.

II.57. Theorem. — *Let R be an ordinary commutative ring. Then the canonical functor*

$$\mathcal{D}(R) \simeq \mathrm{LMod}_{R[0]}(\mathcal{D}(R)^{\otimes_R^L}) \xrightarrow{H} \mathrm{LMod}_{HR}(\mathrm{Sp}^\otimes) \simeq \mathrm{Mod}_{HR}$$

induced by Corollary II.56f and Lemma II.55 is an equivalence.

We can interpret this theorem as follows: Since \mathbb{S} is the tensor unit in Sp (by II.53), it is the initial element in CAlg (Lemma/Definition II.45a) and we have $\mathrm{Sp} \simeq \mathrm{Mod}_{\mathbb{S}}$ (Corollary II.56f). Hence, what Theorem II.57 says is that the passage from $\mathcal{D}(R)$ to Sp is nothing else but the forgetful functor $\mathrm{Mod}_{HR} \rightarrow \mathrm{Mod}_{\mathbb{S}}$ with respect to the unique map $\mathbb{S} \rightarrow HR$ in CAlg .

As an easy consequence, we obtain that also

$$K: \mathcal{D}_{\geq 0}(R) \xrightarrow{\sim} \mathrm{LMod}_R(\mathrm{CGrp}(\mathrm{An})^\otimes)$$

is an equivalence, where $\mathrm{CGrp}(\mathrm{An})^\otimes \simeq \mathrm{CGrp}^{\mathrm{Op}\infty}(\mathrm{An}^\times)$ denotes the symmetric monoidal structure on $\mathrm{CGrp}(\mathrm{An})$ obtained via Proposition II.51.

Proof sketch of Theorem II.57.* We didn't prove it in the lecture, but Fabian has sketched out a proof in the lecture notes [A&HK, Theorem II.57], which I include here.

Recall that an object c in an ∞ -category \mathcal{C} is called *compact* if $\mathrm{Hom}_{\mathcal{C}}(c, -)$ commutes with filtered colimits (which were introduced in Remark* II.31b). The proof is based on the following general fact:

(\boxtimes) *Let \mathcal{C} be an ∞ -category with a set S of compact objects such that the functors*

$$\mathrm{Hom}_{\mathcal{C}}(s, -): \mathcal{C} \longrightarrow \mathrm{An}$$

are jointly conservative. Let $\mathcal{C}^\omega \subseteq \mathcal{C}$ denote the full sub- ∞ -category of compact objects. Then the restricted Yoneda embedding $\mathcal{C} \rightarrow \mathcal{P}(\mathcal{C}^\omega)$ is fully faithful with essential image described equivalently as

- (a) *those functors $(\mathcal{C}^\omega)^{\mathrm{op}} \rightarrow \mathrm{An}$ that preserve finite limits, or*
- (b) *the smallest subcategory of $\mathcal{P}(\mathcal{C}^\omega)$ that contains \mathcal{C}^ω and is closed under filtered colimits (also known as the ind-completion $\mathrm{Ind}(\mathcal{C}^\omega)$).*

Such ∞ -categories are called *compactly generated*. Note that $\mathcal{D}(R)$ is compactly generated, with $S = \{R[i] \mid i \in \mathbb{Z}\}$. Indeed, to show that $\mathrm{Hom}_{\mathcal{D}(R)}(R[i], -) \simeq \Omega^\infty \mathrm{hom}_{\mathcal{D}(R)}(R[i], -)$ are jointly conservative and commute with filtered colimits, it suffices to show the same for $\pi_j \mathrm{hom}_{\mathcal{D}(R)}(R[i], -)$, using that Ω^∞ and π_j preserve filtered colimits (Remark* II.31b) and that equivalences of spectra can be detected on homotopy groups (Lemma* II.23c). We've seen in the proof of Lemma II.55 that $\mathrm{hom}_{\mathcal{D}(R)}(R[0], -) \simeq H: \mathcal{D}(R) \rightarrow \mathrm{Sp}$. Hence

$$\pi_j \mathrm{hom}_{\mathcal{D}(R)}(R[i], -) \simeq \pi_j H(-[-i]) \simeq H_{i+j}: \mathcal{D}(R) \longrightarrow \mathrm{Ab}$$

is the functor taking homology in degree $i + j$. These functors preserve filtered colimits and are clearly jointly conservative, as claimed.

We remark for future use that this argument also proves that $H: \mathcal{D}(R) \rightarrow \mathrm{Sp}$ is conservative and commutes with filtered colimits. In fact, it even shows that H commutes with arbitrary colimits! Indeed, H is an exact functor between stable ∞ -categories, hence it also preserves finite colimits. But every colimit can be obtained from coproducts (which are filtered colimits of finite coproducts) and pushouts (which are finite). The upshot is that also the functor $H: \mathcal{D}(R) \rightarrow \mathrm{Mod}_{HR}$ we're interested in is conservative and colimit-preserving, since colimits and equivalences in Mod_{HR} can be detected on underlying spectra by Corollary II.56e, and the Segal condition from Theorem II.56c, respectively.

Similarly, Mod_T is a compactly generated ∞ -category for any \mathbb{E}_∞ -ring spectrum T , with $S = \{T[i] \mid i \in \mathbb{Z}\}$ a set of compact generators. Indeed, Corollary* II.56g implies

$$\mathrm{Hom}_{\mathrm{Mod}_T}(T[i], -) \simeq \mathrm{Hom}_{\mathrm{Sp}}(\mathbb{S}[i], -) \simeq \Omega^{\infty+i},$$

and the right-hand sides are clearly jointly conservative and preserve filtered colimits by Remark* II.31b.

Now we can begin with the actual proof. It's a general fact that a conservative functor is an equivalence iff it has a left adjoint and the counit is an equivalence (use the triangle relations to show that the unit must be an equivalence as well). By Lurie's presentability magic (more precisely, [HTT, Corollary 5.5.2.9]), a functor between compactly generated ∞ -categories has a left adjoint if it preserves limits and filtered colimits, and a right adjoint if it preserves colimits. This is certainly true for $H: \mathcal{D}(R) \rightarrow \mathrm{Mod}_{HR}$: It preserves limits since these can be computed on underlying spectra by Corollary II.56e and the underlying functor $H: \mathcal{D}(R) \rightarrow \mathrm{Sp}$ is a right adjoint. Preservation of colimits was checked above. Hence we get a left adjoint (and a right adjoint, which we won't need) for free and all we need to check is that the counit is an equivalence.

If $L: \mathrm{Mod}_{HR} \rightarrow \mathcal{D}(R)$ denotes the left adjoint, then

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}(R)}(L(HR), C) &\simeq \mathrm{Hom}_{\mathrm{Mod}_{HR}}(HR, HC) \\ &\simeq \mathrm{Hom}_{\mathrm{Sp}}(\mathbb{S}, HC) \\ &\simeq \mathrm{Hom}_{\mathcal{D}(R)}(R[0], C), \end{aligned}$$

using the adjunctions from Corollary* II.56g and Lemma II.55. Hence the counit $LH \Rightarrow \mathrm{id}_{\mathcal{D}(R)}$ is an equivalence on $R[0]$. Since Mod_{HR} is stable by Corollary II.56e, L is a colimit-preserving functor between stable ∞ -categories, hence exact by Corollary II.28. Thus both LH and $\mathrm{id}_{\mathcal{D}(R)}$ are exact and preserve colimits, and therefore the sub- ∞ -category of $\mathcal{D}(R)$ where the counit is an equivalence must be stable under shifts and colimits. But then it must be all of $\mathcal{D}(R)$: It contains $R[0]$, hence all $R[i]$, hence everything as $\mathcal{D}(R)$ is compactly generated. \square

Fabian also mentioned in the lecture (and elaborates in the script, see [A&HK, Theorem II.58]) that one can more or less the same argument as in the proof of Theorem II.57 to show a vast generalisation: The *Schwede–Shipley theorem*.

II.58. Theorem (Schwede–Shipley). — *If \mathcal{C} is a complete stable ∞ -category containing an object $x \in \mathcal{C}$ such that $\mathrm{hom}_{\mathcal{C}}(x, -): \mathcal{C} \rightarrow \mathrm{Sp}$ is conservative and commutes with colimits, then this functor lifts to an equivalence*

$$\mathrm{hom}_{\mathcal{C}}(x, -): \mathcal{C} \xrightarrow{\sim} \mathrm{Mod}_{\mathrm{end}(x)^{\mathrm{op}}},$$

where $\text{end}(x)^{\text{op}}$ denotes the opposite algebra of the canonical refinement of $\text{end}(x) := \text{hom}_{\mathcal{C}}(x, x) \in \text{Sp}$ to an object of $\text{Alg}_{\text{Assoc}}(\text{Sp}^{\otimes})$.

Tensor Products over Arbitrary Bases

By Theorem II.57, the symmetric monoidal structure on $\mathcal{D}(R)$ can be carried over to Mod_{HR} . More generally, we would like to define symmetric monoidal structures Mod_T^{\otimes} on Mod_T for any \mathbb{E}_{∞} -ring spectrum T . In the special case $T \simeq HR$ it should be given by

$$HM \otimes_{HR} HN \simeq H(M \otimes_R^L N).$$

for any $M, N \in \mathcal{D}(R)$. In fact, constructing symmetric monoidal structures on ∞ -categories of algebras works ridiculous generality: Given a symmetric monoidal ∞ -category $q: \mathcal{C}^{\otimes} \rightarrow \mathbb{T}^{\text{op}}$ and an ∞ -operad $p: \mathcal{O} \rightarrow \mathbb{T}^{\text{op}}$, the ∞ -category of algebras $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})$ inherits a symmetric monoidal structure $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})^{\otimes}$,

II.59. Construction. — With notation as above, define $\overline{\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})}^{\otimes} \rightarrow \mathbb{T}^{\text{op}}$ via II.38a applied to the presheaf $F: \Delta/\mathbb{T}^{\text{op}} \rightarrow \text{An}$ which is given by

$$F(f: [n] \rightarrow \mathbb{T}^{\text{op}}) \simeq \text{Hom}_{\text{Cat}_{\infty}}([n] \times \mathcal{O}, \mathcal{C}^{\otimes}) \times_{\text{Hom}_{\text{Cat}_{\infty}}([n] \times \mathcal{O}, \mathbb{T}^{\text{op}})} \{f \times p\},$$

where $f \times p: [n] \times \mathcal{O} \rightarrow \mathbb{T}^{\text{op}}$ denotes the composition $[n] \times \mathcal{O} \rightarrow \mathbb{T}^{\text{op}} \times \mathbb{T}^{\text{op}} \xrightarrow{\times} \mathbb{T}^{\text{op}}$. Let's compute the fibres

$$\overline{\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})}_n^{\otimes} \simeq \overline{\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})}^{\otimes} \times_{\mathbb{T}^{\text{op}}} \{\langle n \rangle\}.$$

Using the method from the proof sketch of Proposition II.40, we find that the Rezk nerve of the fibre over $\langle n \rangle$ is a simplicial anima Y given by $Y_k \simeq F(\text{const } \langle n \rangle: [k] \rightarrow \mathbb{T}^{\text{op}})$. Plugging in the definitions, we get

$$\begin{aligned} Y_k &\simeq \text{Hom}_{\text{Cat}_{\infty}}([k] \times \mathcal{O}, \mathcal{C}^{\otimes}) \times_{\text{Hom}_{\text{Cat}_{\infty}}([k] \times \mathcal{O}, \mathbb{T}^{\text{op}})} \{\text{const } \langle n \rangle \times p\} \\ &\simeq \text{Hom}_{\text{Cat}_{\infty}}([k], \text{Fun}(\mathcal{O}, \mathcal{C}^{\otimes})) \times_{\text{Hom}_{\text{Cat}_{\infty}}([k], \text{Fun}(\mathcal{O}, \mathbb{T}^{\text{op}}))} \{\text{const } \langle n \rangle \times p\} \\ &\simeq \text{Hom}_{\text{Cat}_{\infty}}([k], \text{Fun}(\mathcal{O}, \mathcal{C}^{\otimes}) \times_{\text{Fun}(\mathcal{O}, \mathbb{T}^{\text{op}})} \{\langle n \rangle \times p\}) \\ &\simeq \text{Hom}_{\text{Cat}_{\infty}}([k], \text{Fun}_{\mathbb{T}^{\text{op}}}(\mathcal{O}, \underbrace{\mathcal{C}^{\otimes} \times_{\mathbb{T}^{\text{op}}} \cdots \times_{\mathbb{T}^{\text{op}}} \mathcal{C}^{\otimes}}_{n \text{ times}})). \end{aligned}$$

For the second and third equivalence, we use currying and the fact that $\text{Hom}_{\text{Cat}_{\infty}}([k], -)$ commutes with pullbacks. For the fourth equivalence, we use that maps $\mathcal{O} \rightarrow \mathcal{C}^{\otimes}$ over \mathbb{T}^{op} , where \mathcal{O} is equipped with the structure map $\langle n \rangle \times p: \mathcal{O} \rightarrow \mathbb{T}^{\text{op}}$, are the same as maps from \mathcal{O} with its usual structure map p to the pullback of \mathcal{C}^{\otimes} along $\langle n \rangle \times -: \mathbb{T}^{\text{op}} \rightarrow \mathbb{T}^{\text{op}}$. Via the Segal maps, said pullback is easily identified with the n -fold fibre product of \mathcal{C}^{\otimes} over \mathbb{T}^{op} .

The calculation above shows that $Y \simeq N^r(\text{Fun}_{\mathbb{T}^{\text{op}}}(\mathcal{O}, \mathcal{C}^{\otimes})^n)$, hence the fibre we're interested in is given by

$$\overline{\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})}_n^{\otimes} \simeq \text{Fun}_{\mathbb{T}^{\text{op}}}(\mathcal{O}, \mathcal{C}^{\otimes})^n.$$

We let $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})^{\otimes} \subseteq \overline{\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})}^{\otimes}$ be the full sub- ∞ -category spanned fibrewise by those (F_1, \dots, F_n) such that each F_i is a map of ∞ -operads.

II.60. Proposition (Lurie, [HA, Proposition 3.2.4.3]). — *The map $\text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})^{\otimes} \rightarrow \mathbb{T}^{\text{op}}$ is always a symmetric monoidal monoidal ∞ -category, and for every $x \in \mathcal{O}_1$, evaluation at x refines to a strongly monoidal functor*

$$\text{ev}_x: \text{Alg}_{\mathcal{O}}(\mathcal{C}^{\otimes})^{\otimes} \longrightarrow \mathcal{C}^{\otimes}.$$

II.61. Observation. — *In the special case $\mathcal{O} \simeq \mathbb{C}\text{omm}$, the symmetric monoidal structure on $\text{Alg}_{\mathbb{C}\text{omm}}(\mathcal{C}^\otimes) \simeq \text{CAlg}(\mathcal{C}^\otimes)$ is the cocartesian symmetric monoidal structure as defined in Proposition II.40. In particular, the tensor product of two algebras is also their coproduct in $\text{CAlg}(\mathcal{C}^\otimes)$.*

Proof sketch of Observation II.61. Use a generalisation of Lemma II.20, which we used to show that $\text{CMon}(\mathcal{C})$ is semi-additive. In fact, Lurie’s version ([HA, Proposition 2.4.3.19]) of that lemma gives a criterion for a symmetric monoidal structure to be cocartesian. \square

By Observation II.61 the second half of Theorem II.43, we get

$$\begin{aligned} \text{CAlg}(\text{CAlg}(\mathcal{C}^\otimes)^\otimes) &\simeq \text{Fun}^{\text{Op}\infty}(\mathbb{C}\text{omm}, \text{CAlg}(\mathcal{C}^\otimes)^\otimes) \\ &\simeq \text{Fun}(\mathbb{C}\text{omm}_1, \text{CAlg}(\mathcal{C}^\otimes)) \\ &\simeq \text{CAlg}(\mathcal{C}^\otimes). \end{aligned}$$

Hence every commutative algebra $A \in \text{CAlg}(\mathcal{C}^\otimes)$ canonically refines to a map of ∞ -operads $A: \mathbb{C}\text{omm} \rightarrow \text{CAlg}(\mathcal{C}^\otimes)^\otimes$. Now consider the following pullback diagram:

$$\begin{array}{ccccccc} \text{LMod}_A(\mathcal{C}^\otimes) & \longrightarrow & \text{LMod}_A(\mathcal{C}^\otimes)^{\otimes_A} & \xrightarrow{\quad} & \text{Alg}_{\text{LMod}}(\mathcal{C}^\otimes)^\otimes \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow a^* \\ * & \xrightarrow{A} & \mathbb{C}\text{omm} & \xrightarrow{A} & \text{Alg}_{\mathbb{C}\text{omm}}(\mathcal{C}^\otimes)^\otimes \longrightarrow \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)^\otimes \end{array}$$

II.62. Theorem. — *Suppose \mathcal{C}^\otimes has all small operadic colimits. Then*

$$a^*: \text{Alg}_{\text{LMod}}(\mathcal{C}^\otimes)^\otimes \longrightarrow \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)^\otimes$$

is a cocartesian fibration. In particular, $\text{LMod}_A(\mathcal{C}^\otimes)^{\otimes_A} \rightarrow \mathbb{C}\text{omm}$ defines a symmetric monoidal structure on $\text{LMod}_A(\mathcal{C}^\otimes)$ for all $A \in \text{CAlg}(\mathcal{C}^\otimes)$. Moreover, $\text{LMod}_A(\mathcal{C}^\otimes)^{\otimes_A}$ has again all operadic colimits.

II.62a. The Bar Construction. — The tensor product in $\text{LMod}_A(\mathcal{C}^\otimes)$ can be computed by the *bar construction*: For $M, N \in \text{LMod}_A(\mathcal{C}^\otimes)$, the underlying \mathcal{C} -object of $M \otimes_A N$ (which we’ll abusively also denote that way) is given by

$$M \otimes_A N \simeq \text{colim}_{\Delta^{\text{op}}} B_\bullet(M, A, N).$$

Informally, the functor $B_\bullet(M, A, N): \Delta^{\text{op}} \rightarrow \mathcal{C}$ sends $[n] \in \Delta^{\text{op}}$ to $M \otimes A^{\otimes n} \otimes N$, with the tensor product being taken in \mathcal{C} .

In the script [A&HK, Chapter II pp. 123–124], Fabian explains how to define the bar construction formally. Let $\text{rev}: \Delta^{\text{op}} \rightarrow \Delta^{\text{op}}$ denote the functor that reverses the order of a totally ordered set. Using Theorem II.56a, it induces a functor

$$\text{rev}^*: \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes) \longrightarrow \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes).$$

But one easily checks that $\text{Cut}: \Delta^{\text{op}} \rightarrow \mathbb{I}^{\text{op}}$ and $\text{Cut} \circ \text{rev}: \Delta^{\text{op}} \rightarrow \Delta^{\text{op}}$ are naturally equivalent, hence rev^* becomes equivalent to the identity upon composition with the canonical map $\text{CAlg}(\mathcal{C}^\otimes) \rightarrow \text{Alg}_{\text{Assoc}}(\mathcal{C}^\otimes)$. Of course, this is nothing but the ∞ -categorical way of saying that a commutative algebra is naturally equivalent to its opposite.

Using Theorem II.56c, we may now consider the functors

$$M \circ (\text{rev} \times \text{id}_{[1]}), N: \Delta^{\text{op}} \times [1] \longrightarrow \mathcal{C}^{\otimes}.$$

Let's also write $M \circ (\text{rev} \times \text{id}_{[1]}) =: M_A$ for simplicity (indicating that we now consider the right- A module structure on M , although we never defined anything of that sort). Then $M_{A|\Delta^{\text{op}} \times \{1\}} \simeq A^{\text{rev}} \simeq A \simeq N_{|\Delta^{\text{op}} \times \{1\}}$ by what we just showed. In particular, if we regard the functors M_A and N as natural transformations $M_{A|\Delta^{\text{op}} \times \{0\}} \Rightarrow M_{A|\Delta^{\text{op}} \times \{1\}}$ and $N_{|\Delta^{\text{op}} \times \{0\}} \Rightarrow N_{|\Delta^{\text{op}} \times \{1\}}$ in $\text{Fun}(\Delta^{\text{op}}, \mathcal{C}^{\otimes})$, then their endpoints coincide, and we can form the pullback $P \in \text{Fun}(\Delta^{\text{op}}, \mathcal{C}^{\otimes})$. Pullbacks in functor categories are formed pointwise (Lemma I.39), hence $P([n])$ fits into a pullback diagram

$$\begin{array}{ccc} P([n]) & \longrightarrow & N([n], 0) \\ \downarrow & \lrcorner & \downarrow \\ M_A([n], 0) & \longrightarrow & A([n]) \end{array}$$

in \mathcal{C}^{\otimes} . By construction, we have $M_A([n], 0) \simeq (M, A, \dots, A)$, $N([n], 0) \simeq (A, \dots, A, N)$, and $A([n]) \simeq (A, \dots, A)$, where we abusively identify algebras and modules with their underlying \mathcal{C} -objects, and the maps between them are the evident projections. Thus, using Definition II.36(c) one easily checks

$$P([n]) \simeq (M, A, \dots, A, N) \in \mathcal{C}_{n+2}^{\otimes}.$$

Now observe that $P: \Delta^{\text{op}} \rightarrow \mathcal{C}^{\otimes}$ factors over $\mathcal{C}_{\text{act}}^{\otimes} := \mathcal{C}^{\otimes} \times_{\Gamma^{\text{op}}} \Gamma_{\text{act}}^{\text{op}}$. Indeed, for a general $\alpha: [n] \rightarrow [m]$ in Δ , the morphism $\text{Cut}(\alpha^{\text{op}}): \langle m \rangle \rightarrow \langle n \rangle$ is only undefined on those $i \in \langle m \rangle$ for which $i \leq \alpha(0)$ or $\alpha(n) < i$, see Definition II.15. But those i with $\alpha(n) < i$ are used to encode the left-action of A on N (compare this to Remark* II.56d), and those i with $i \leq \alpha(0)$ are used to encode the right action of A on M_A . Hence the image of $P(\alpha^{\text{op}})$ in Γ^{op} must be active, as claimed.

The unique active maps $f_n: \langle n \rangle \rightarrow \langle 1 \rangle$ define a natural transformation $\text{id}_{\Gamma_{\text{act}}^{\text{op}}} \Rightarrow \text{const} \langle 1 \rangle$ in $\text{Fun}(\Gamma_{\text{act}}^{\text{op}}, \Gamma_{\text{act}}^{\text{op}})$. By unstraightening magic, it defines a map of cocartesian fibrations from $\mathcal{C}_{\text{act}}^{\otimes}$ to its pullback along $\text{const} \langle 1 \rangle$, which is $\mathcal{C} \times \Gamma_{\text{act}}^{\text{op}}$. Composing P with that map gives a functor

$$B_{\bullet}(M, A, N): \Delta^{\text{op}} \longrightarrow \mathcal{C},$$

which is the one we're looking for. Since cocartesian lifts of the active maps $f_{n+2}: \langle n+2 \rangle \rightarrow \langle 1 \rangle$ just tensor stuff together, we see that $B(M, A, N)$ does indeed take $[n]$ to $M \otimes A^{\otimes n} \otimes N$.

We won't explain why the bar construction computes the tensor product in $\text{LMod}_A(\mathcal{C}^{\otimes})$, and refer instead to [HA] again: In Proposition 4.4.2.8, Lurie proves that the tensor product in *bimodules* is given by the bar construction, and in Theorem 4.5.2.1 he shows that the tensor product on $\text{LMod}_A(\mathcal{C}^{\otimes})$ for $A \in \text{CAlg}(\mathcal{C}^{\otimes})$ factors over the specialisation from bimodules to left modules.

II.62b. Relative Tensor Products of Spectra*. — We can now define symmetric monoidal structures on the stable ∞ -categories $\text{Mod}_R^{\otimes R}$ for all \mathbb{E}_{∞} -ring spectra R . We'd hope the following properties to hold:

- (a) *The functor $\text{hom}_{\text{Mod}_R}: \text{Mod}_R^{\text{op}} \times \text{Mod}_R \rightarrow \text{Sp}$ has a natural refinement to a functor $\text{hom}_R: \text{Mod}_R^{\text{op}} \times \text{Mod}_R \rightarrow \text{Mod}_R$ satisfying the tensor-Hom adjunction*

$$\text{hom}_R(M \otimes_R N, L) \simeq \text{hom}_R(M, \text{hom}_R(N, L)).$$

(b) If $R \rightarrow S$ is a map in \mathbf{CAlg} , we get a symmetric monoidal functor

$$S \otimes_R - : \mathbf{Mod}_R^{\otimes_R} \longrightarrow \mathbf{Mod}_S^{\otimes_S},$$

which does what it says on underlying spectra. If moreover $S \rightarrow T$ is another map in \mathbf{CAlg} , then $T \otimes_S (S \otimes_R -) \simeq T \otimes_R -$.

(c) The functor $S \otimes_R : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_S$ has a right adjoint, the forgetful functor

$$F_{S/R} : \mathbf{Mod}_S \longrightarrow \mathbf{Mod}_R,$$

which is the identity on underlying spectra. Moreover, $F_{S/R}$ has a right adjoint itself, which is a refinement of $\mathrm{hom}_R(S, -) : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_R$.

*Proof**. Part (b) works in fact in much greater generality, see [HA, Theorem 4.5.3.1]. We will see that the rest are formal consequences.

The existence of a functor $\mathrm{hom}_R : \mathbf{Mod}_R^{\mathrm{op}} \times \mathbf{Mod}_R \rightarrow \mathbf{Mod}_R$ in part (a) follows from adjoint functor theorem magic (initially, we only get a tensor-Hom adjunction with the outer hom_R replaced by the actual Hom functor $\mathrm{Hom}_{\mathbf{Mod}_R} : \mathbf{Mod}_R^{\mathrm{op}} \times \mathbf{Mod}_R \rightarrow \mathbf{An}$, but then it formally follows for hom_R as well). To see that hom_R equals $\mathrm{hom}_{\mathbf{Mod}_R}$ on underlying spectra, we have to prove (c) first.

The existence of $F_{S/R}$ follows again from Lurie's adjoint functor theorem: We know that $S \otimes_R -$, as an endofunctor of \mathbf{Mod}_R , preserves colimits by Theorem II.62, hence the same is true for $S \otimes_R - : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_S$, since both equivalences and colimits in \mathbf{Mod}_S can be detected on underlying spectra by the Segal condition from Theorem II.56c, and Corollary II.56e, respectively. Hence $F_{S/R}$ exists. To see that it is the identity on underlying spectra, note that $S \otimes_R (R \otimes_S -) \simeq S \otimes_S -$ implies $F_{R/S} \circ F_{S/R} \simeq F_{S/S}$, and both $F_{R/S}$ and $F_{S/S}$ are just extraction of underlying spectra by Corollary* II.56g. This also implies that $F_{S/R}$ preserves colimits, since these can be detected on underlying spectra. Hence $F_{S/R}$ has itself a right adjoint $h_{S/R} : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_S$. Since right adjoints compose, $F_{S/R} \circ h_{S/R}$ is a right adjoint of $S \otimes_R - : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_R$, hence $h_{S/R}$ is indeed a refinement of $\mathrm{hom}_R(S, -) : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_R$. This proves (c).

Finally, to show that the underlying spectrum of $\mathrm{hom}_R(M, N)$ is $\mathrm{hom}_{\mathbf{Mod}_R}(M, N)$, it suffices to consider the case $M = R$, since then the general case follows by writing M as a colimit of shifts of R (see the proof of Theorem II.57 for why that's possible). But $\mathrm{hom}_R(R, N) \simeq N$ by the tensor-Hom adjunction, and $\mathrm{hom}_{\mathbf{Mod}_R}(R, N) \simeq \mathrm{hom}_{\mathbf{Sp}}(\mathbb{S}, N) \simeq N$ by the adjunction from Corollary* II.56g. \square

It remains to see why in the case of an ordinary ring R , the derived tensor product on $\mathcal{D}(R)$ and the tensor product on \mathbf{Mod}_{HR} coincide. To this end, observe that for arbitrary symmetric monoidal ∞ -categories, there's a strongly monoidal functor

$$\mathcal{C}^{\otimes} \longrightarrow \mathrm{Alg}_{\mathbf{LMod}}(\mathcal{C}^{\otimes})^{\otimes},$$

which informally sends objects (c_1, \dots, c_n) to $((1_{\mathcal{C}}, c_1), \dots, (1_{\mathcal{C}}, c_n))$, considering each c_i as a module over the trivial algebra $1_{\mathcal{C}}$ via Corollary II.56f.

II.63. Proposition. — *The composite*

$$\mathcal{D}(R)^{\otimes_R^L} \longrightarrow \mathrm{Alg}_{\mathbf{LMod}}(\mathcal{D}(R)^{\otimes_R^L})^{\otimes_R^L} \xrightarrow{H} \mathrm{Alg}_{\mathbf{LMod}}(\mathbf{Sp}^{\otimes})^{\otimes}$$

induces an equivalence of symmetric monoidal ∞ -categories

$$H : \mathcal{D}(R)^{\otimes_R^L} \xrightarrow{\sim} \mathbf{Mod}_{HR}^{\otimes_{HR}}.$$

*Proof sketch**. We already know that H is an equivalence of underlying ∞ -categories. Hence it suffices to show

$$H(R[0]) \simeq HR \quad \text{and} \quad H(M \otimes_R^L N) \simeq HM \otimes_{HR} HN$$

(since then it formally follows that H preserves cocartesian lifts of active morphisms, hence all cocartesian lifts). There's nothing to show for the first equivalence. For the second, we'll use the bar construction from II.62a. Since H , being an equivalence, preserves colimits, it suffices to consider the case $M = R[0]$. Hence we must show

$$HN \simeq \operatorname{colim}_{\Delta^{\text{op}}} B_{\bullet}(HR, HR, HN).$$

This is, in fact, a special case of a general principle. Observe that the bar construction $B_{\bullet} := B_{\bullet}(HR, HR, HN)$ is the *décalage* of another simplicial HR -module. That is, there is another functor $B'_{\bullet}: \Delta^{\text{op}} \rightarrow \operatorname{Mod}_{HR}$ such that the following diagram commutes:

$$\begin{array}{ccc} \Delta^{\text{op}} & \xrightarrow{\sigma^{\text{op}}} & \Delta^{\text{op}} \\ & \searrow B_{\bullet} & \swarrow B'_{\bullet} \\ & \operatorname{Mod}_{HR} & \end{array}$$

Here $\sigma: \Delta \rightarrow \Delta$ denotes the functor taking objects $[n]$ to $[n+1]$ and morphisms $\alpha: [n] \rightarrow [m]$ to $\sigma(\alpha): [n+1] \rightarrow [m+1]$ given by $\sigma(\alpha)(i) = \alpha(i)$ for $i \leq n$ and $\sigma(\alpha)(n+1) = m+1$.

It's not hard to construct B'_{\bullet} in our example. Basically, one has to place another HN in degree 0 and add those maps $HR^{\otimes n} \otimes HN \rightarrow HR^{\otimes n+1} \otimes HN$ that are given by $\mathbb{S} \rightarrow HR$ on the first factor and the identity on the rest, as well as those maps $HR^{\otimes n+1} \otimes HN \rightarrow HR^{\otimes n} \otimes HN$ that multiply the first factor with HN (so we are forming some kind of cyclic bar construction). I'll leave the formal construction to you.

In general, if $B_{\bullet}, B'_{\bullet}: \Delta^{\text{op}} \rightarrow \mathcal{C}$ are simplicial objects in some ∞ -category \mathcal{C} , and if $B_{\bullet} \simeq \operatorname{dec}(B'_{\bullet})$ is the *décalage* of B'_{\bullet} , then

$$\operatorname{colim}_{\Delta^{\text{op}}} B_{\bullet} \simeq \operatorname{colim}_{\Delta^{\text{op}}} \operatorname{const} B'_0 \simeq B'_0.$$

The right equivalence follows from the fact that Δ^{op} is weakly contractible (with $[0]$ an initial object). For the left-hand side, observe that there are canonical transformations $B_{\bullet} \Rightarrow \operatorname{const} B'_0$ and $\operatorname{const} B'_0 \Rightarrow B_{\bullet}$ in $\operatorname{Fun}(\Delta^{\text{op}}, \mathcal{C})$, induced by suitable natural transformations $\eta: \sigma^{\text{op}} \Rightarrow \operatorname{const} [0]$ and $\tau: \operatorname{const} [0] \Rightarrow \sigma^{\text{op}}$ in $\operatorname{Fun}(\Delta^{\text{op}}, \Delta^{\text{op}})$. Using higher order transformations between η and σ , one shows that

$$\tau^*: \operatorname{Nat}(B_{\bullet}, \operatorname{const} x) \xrightarrow[\sim]{\sim} \operatorname{Nat}(\operatorname{const} B'_0, \operatorname{const} x) : \eta^*$$

are mutually inverse homotopy equivalences, which is what we need. \square

Miscellanea

The following stuff was only implicitly mentioned in the lecture (if at all), but Fabian gave a detailed account in his notes, [A&HK, Remarks II.53].

II.64. Graded Structures on Homotopy Groups*. — The functor $\pi_0: \mathbf{Sp} \rightarrow \mathbf{Ab}$ is lax symmetric monoidal, since it is the underlying functor of the composition

$$\mathbf{Sp}^{\otimes} \xrightarrow{\Omega^{\infty}} \mathbf{CGrp}(\mathbf{An})^{\otimes} \xrightarrow{\pi_0} \mathbf{Ab}^{\otimes},$$

where we denote $\mathbf{CGrp}(\mathbf{An})^{\otimes} \simeq \mathbf{CGrp}^{\mathrm{Op}\infty}(\mathbf{An}^{\times})$ (which makes sense, since the right-hand side is symmetric monoidal by Proposition II.51), and use $\mathbf{CGrp}^{\mathrm{Op}\infty}(\mathbf{Set}^{\times}) \simeq \mathbf{Ab}^{\otimes}$ by Example II.51b(a).

This implies that π_0 induces functors

$$\begin{aligned} \pi_0: \mathbf{CAlg} &\simeq \mathbf{CAlg}(\mathbf{Sp}^{\otimes}) \longrightarrow \mathbf{CAlg}(\mathbf{Ab}^{\otimes}) \simeq \mathbf{CRing}, \\ \pi_0: \mathbf{Mod}_R &\simeq \mathbf{LMod}_R(\mathbf{Sp}^{\otimes}) \longrightarrow \mathbf{LMod}_{\pi_0(R)}(\mathbf{Ab}^{\otimes}) \simeq \mathbf{Mod}_{\pi_0(R)} \end{aligned}$$

for every \mathbb{E}_{∞} -ring spectrum R . In particular, $\pi_0(R)$ is an ordinary ring, and if M is an R -module, then $\pi_0(M)$ is an ordinary $\pi_0(R)$ -module. But we can do better! For all $i, j \in \mathbb{Z}$, the multiplication $\mu: R \otimes M \rightarrow M$ induces morphisms

$$\pi_0(\mathbb{S}[-i] \otimes R) \otimes \pi_0(\mathbb{S}[-j] \otimes M) \longrightarrow \pi_0(\mathbb{S}[-(i+j)] \otimes R \otimes M) \xrightarrow{\mu} \pi_0(\mathbb{S}[-(i+j)] \otimes M),$$

which provide a multiplication $\pi_i(R) \otimes \pi_j(M) \rightarrow \pi_{i+j}(M)$. In the special case $R = M$, it turns $\pi_*(R)$ into a graded ring, and for general M we get a graded $\pi_*(R)$ -module structure on $\pi_*(M)$!

In fact, $\pi_*(R)$ is graded commutative. To see this, we need to investigate the effect of the canonical equivalence $\mathbb{S}[i] \otimes \mathbb{S}[j] \simeq \mathbb{S}[j] \otimes \mathbb{S}[i]$ (we drop the minus signs for convenience) on homotopy groups. This equivalence corresponds to an element of

$$\pi_0 \mathrm{Hom}_{\mathbf{Sp}}(\mathbb{S}[i+j], \mathbb{S}[i+j]) \simeq \pi_{i+j}(\mathbb{S}[i+j]) \simeq \mathbb{Z}.$$

Now $\mathbb{S}[i] \simeq \Sigma^{\infty} \mathbb{S}^i$ and $\mathbb{S}[j] \simeq \Sigma^{\infty} \mathbb{S}^j$. Since Σ^{∞} is strongly monoidal (Proposition II.51), we see that the equivalence $\mathbb{S}[i] \otimes \mathbb{S}[j] \simeq \mathbb{S}[j] \otimes \mathbb{S}[i]$ corresponds to

$$\mathbb{S}^i \wedge \mathbb{S}^j \simeq (\mathbb{S}^1)^{\wedge i} \wedge (\mathbb{S}^1)^{\wedge j} \simeq (\mathbb{S}^1)^{\wedge j} \wedge (\mathbb{S}^1)^{\wedge i} \simeq \mathbb{S}^j \wedge \mathbb{S}^i,$$

given by permutation of factors. The permutation in question can be written as ij transpositions, hence its sign is $(-1)^{ij}$. Hence its action on $\pi_{i+j}(\mathbb{S}^{i+j}) \simeq \mathbb{Z}$ is given by $(-1)^{ij}$ as well. This proves graded commutativity.

Combining these considerations with II.15c(c*) for example, we see that the algebraic K -theory $K_*(R) = \pi_*(B^{\infty}k(R))$ of any commutative ring R carries a graded commutative ring structure, called the *cup product* on K -theory. In the next paragraph we'll see that also the cup product on singular cohomology arises in this way.

II.65. The Cup Product and the Pontryagin Product*. — If $M \in \mathbf{CGrp}(\mathbf{An})$, then $\mathbb{S}[M]$ is an \mathbb{E}_{∞} -ring spectrum, as noted way back in II.54. Using Proposition II.60, this implies that $\mathbb{S}[M] \otimes HR$ is an \mathbb{E}_{∞} -ring spectrum too for any ordinary ring R . Hence, by II.64 and Corollary* II.55b there is a graded commutative ring structure on

$$\pi_*(\mathbb{S}[M] \otimes HR) \simeq H_*(M, R).$$

This is (our definition of) the *Pontryagin product*.

A natural question is whether the cup product in cohomology arises in a similar way. It does! The main ingredient is a construction due to Glasman, which upgrades the Yoneda embedding $Y^{\mathcal{C}}: \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \text{An})$ to a lax symmetric monoidal functor

$$\mathcal{C}^{\otimes} \longrightarrow \text{Day}(\mathcal{C}_{\otimes}^{\text{op}}, \text{An}^{\times}),$$

where $\mathcal{C}_{\otimes}^{\text{op}}$ denotes the induced symmetric monoidal structure on \mathcal{C}^{op} (i.e., compose the functor $\text{St}^{\text{cocart}}(\mathcal{C}^{\otimes}): \mathbb{I}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ with $(-)^{\text{op}}$ and observe that the Segal condition survives). In particular, if $A \in \text{CAlg}(\mathcal{C}^{\otimes})$ is a commutative algebra, which we abusively identify with its underlying \mathcal{C} -object, then

$$\text{Hom}_{\mathcal{C}}(-, A): \mathcal{C}^{\text{op}} \longrightarrow \text{An}$$

is an element of $\text{CAlg}(\text{Day}(\mathcal{C}_{\otimes}^{\text{op}}, \text{An}^{\times})) \simeq \text{Fun}^{\text{Op}\infty}(\mathcal{C}_{\otimes}^{\text{op}}, \text{An}^{\times})$ (as in the proof of Lemma II.55, we don't even need that this is an equivalence), hence a lax symmetric monoidal functor. Thus it defines a functor

$$\text{Hom}_{\mathcal{C}}(-, A): \text{CAlg}(\mathcal{C}_{\otimes}^{\text{op}}) \longrightarrow \text{CAlg}(\text{An}^{\times}).$$

The right-hand side equals $\text{CMon}(\text{An})$ by Theorem II.43. In particular, if $B \in \text{CAlg}(\mathcal{C}_{\otimes}^{\text{op}})$ is a coalgebra in \mathcal{C} , then $\text{Hom}_{\mathcal{C}}(B, A)$ has a canonical refinement in $\text{CMon}(\text{An})$.

Moreover one can check that $\text{Sp}^{\text{Op}\infty}(\text{Day}(\mathcal{C}_{\otimes}, \mathcal{O})) \simeq \text{Day}(\mathcal{C}_{\otimes}^{\text{op}}, \text{Sp}^{\text{Op}\infty}(\mathcal{O}))$ for all ∞ -operads $\mathcal{O} \in \text{Op}_{\infty}^{\text{lex}}$. Hence, if \mathcal{C}^{\otimes} is a stable ∞ -operad, we can apply Theorem II.50 to the functors $\text{Hom}_{\mathcal{C}}(-, A)$ above and obtain upgraded lax symmetric monoidal functors

$$\begin{aligned} \text{hom}_{\mathcal{C}}(-, A): \mathcal{C}_{\otimes}^{\text{op}} &\longrightarrow \text{Sp}^{\otimes}, \\ \text{hom}_{\mathcal{C}}(-, A): \text{CAlg}(\mathcal{C}_{\otimes}^{\text{op}}) &\longrightarrow \text{CAlg}(\text{Sp}^{\otimes}) \simeq \text{CAlg}. \end{aligned}$$

If $\mathcal{C}^{\otimes} \simeq \mathcal{C}^{\times}$ is the cartesian monoidal structure from Proposition II.42, then $\mathcal{C}_{\times}^{\text{op}} \simeq (\mathcal{C}^{\text{op}})^{\sqcup}$ is the cocartesian monoidal structure, and $\text{CAlg}(\mathcal{C}_{\times}^{\text{op}}) \simeq \text{CAlg}((\mathcal{C}^{\text{op}})^{\sqcup}) \simeq \mathcal{C}^{\text{op}}$ by the second half of Theorem II.43. In particular, every anima $X \in \text{An}$ is a coalgebra in the cartesian monoidal structure. Hence $\mathbb{S}[X] \in \text{CAlg}(\text{Sp}_{\otimes}^{\text{op}})$, and then we see that $\text{hom}_{\text{Sp}}(\mathbb{S}[X], E)$ is canonically an \mathbb{E}_{∞} -ring spectrum for every \mathbb{E}_{∞} -ring spectrum E . Now Corollary* II.55b implies

$$\pi_{-*} \text{hom}_{\text{Sp}}(\mathbb{S}[X], HR) \simeq H^*(X, R),$$

whence II.64 endows the cohomology of X with a graded commutative ring structure. This is (our definition of) the *cup product*.

II.66. \mathbb{E}_1 -Ring Spectra*. — Elements of $\text{Alg}_{\text{Assoc}}(\text{Sp}^{\otimes})$ are called *\mathbb{E}_1 -ring spectra*. Many of our results about \mathbb{E}_{∞} -ring spectra can be generalised to \mathbb{E}_1 -ring spectra. But let me only mention those that we'll need in Chapter III.

(a) Since $\mathbb{S}[-]: \text{An}^{\times} \rightarrow \text{Sp}^{\otimes}$ is symmetric monoidal by Proposition II.51, it induces a functor

$$\mathbb{S}[-]: \text{Mon}(\text{An}) \simeq \text{Alg}_{\text{Assoc}}(\text{An}^{\times}) \longrightarrow \text{Alg}_{\text{Assoc}}(\text{Sp}^{\otimes}).$$

In particular, if $M \in \text{Mon}(\text{An})$ is an \mathbb{E}_1 -monoid, then its spherical group ring $\mathbb{S}[M]$ is an \mathbb{E}_1 -ring spectrum.

- (b) Tensor products of \mathbb{E}_1 -ring spectra inherit a \mathbb{E}_1 -ring structure by Proposition II.60.
- (c) If A is an \mathbb{E}_1 -ring spectrum and M a left module over it, then $\pi_*(A)$ is a graded (but of course not necessarily commutative) ring and $\pi_*(M)$ a graded left $\pi_*(A)$ -module.

CHAPTER III.

III

The Group Completion Theorem and the K -Theory of Finite Fields

Our first goal in this chapter, and the content of the group completion theorem, is to understand $\mathbb{S}[M^{\infty\text{-grp}}]$ in terms of $\mathbb{S}[M]$ for $M \in \mathbf{CMon}(\mathbf{An})$. This turns out to be an easy to understand localisation. So let's understand localisations first.

Localisations of \mathbb{E}_∞ -Ring Spectra

Recall that $\pi_0(E) \simeq \pi_0 \operatorname{hom}_{\mathbb{S}\mathbf{p}}(\mathbb{S}, E) \simeq \pi_0 \operatorname{Hom}_{\mathbb{S}\mathbf{p}}(\mathbb{S}, E)$ for every spectrum E (this follows for example from Lemma* II.53a). Hence every element $e \in \pi_0(E)$ defines, up to homotopy, a map $e: \mathbb{S} \rightarrow E$.

III.1. Definition. — Let $R \in \mathbf{CAlg}$ is an \mathbb{E}_∞ -ring spectrum, $M \in \mathbf{Mod}_R$ some module over it, and $S \subseteq \pi_0(R)$ some subset. Then M is called *S -local* if for every $s \in S$ the map

$$M \simeq \mathbb{S} \otimes M \xrightarrow{s \otimes \operatorname{id}_M} R \otimes M \xrightarrow{\mu} M$$

(“multiplication by s ”) is an equivalence.

Our goal is to produce a localisation functor. Let's do the case of a single element $s \in \pi_0(R)$ first. We put

$$M[s^{-1}] := \operatorname{colim}_{\mathbb{N}} \left(M \xrightarrow{\cdot s} M \xrightarrow{\cdot s} M \xrightarrow{\cdot s} \dots \right),$$

the colimit being taken in \mathbf{Mod}_R (which is cocomplete by Corollary II.56e; this corollary also shows that we get a colimit on underlying spectra too). Since the tensor product on \mathbf{Mod}_R commutes with colimits (Theorem II.62), we have $M[s^{-1}] \simeq R[s^{-1}] \otimes_R M$, and since taking homotopy groups commutes with sequential colimits (Lemma II.31a(a^*)), we get $\pi_*(M[s^{-1}]) \simeq \pi_*(M)[s^{-1}]$. Also there is a functorial map $M \rightarrow M[s^{-1}]$, given by including M as the first element of the colimit.

III.2. Proposition. — *The functor $-[s^{-1}]: \mathbf{Mod}_R \rightarrow \mathbf{Mod}_R$ is a Bousfield localisation onto the $\{s\}$ -local R -modules.*

Proof. It's clear that $M[s^{-1}]$ is s -local. Indeed, whether $s: M[s^{-1}] \rightarrow M[s^{-1}]$ is an equivalence can be checked on homotopy groups of underlying spectra (by the Segal condition from Theorem II.56c plus Lemma* II.23c), where it is evidently true as $\pi_*(M[s^{-1}]) \simeq \pi_*(M)[s^{-1}]$.

Therefore, it suffices to check that the functorial maps $\eta_M: M \rightarrow M[s^{-1}]$ satisfy the condition from Proposition I.61a. That is, we must show that

$$\eta_{M[s^{-1}]}, \eta_M[s^{-1}]: M[s^{-1}] \xrightarrow{\sim} (M[s^{-1}])[s^{-1}]$$

are equivalences. This is clearly true for $\eta_{M[s^{-1}]}$ because $M[s^{-1}]$ is $\{s\}$ -local, hence it follows for $\eta_M[s^{-1}]$ by the same “coordinate flip” trick as in the proof of Theorem II.30. \square

This now allows us to define localisations for arbitrary $S \subseteq \pi_0(R)$: If $T \subseteq S$ is a finite subset, say, $T = \{s_1, \dots, s_n\}$, put $M[T^{-1}] := M[(s_1 \dots s_n)^{-1}]$. Since being T -local can be detected on homotopy groups, one readily checks that the T -local R -modules are precisely the $\{s_1 \dots s_n\}$ -local R -modules, hence $-[T^{-1}]: \text{Mod}_R \rightarrow \text{Mod}_R$ is a Bousfield localisation onto the T -local R -modules by Proposition III.2. In general, we put

$$M[S^{-1}] := \text{colim}_{T \subseteq S \text{ finite}} M[T^{-1}].$$

Note that a priori there is a coherence issue in this colimit: Namely, the s_i in $T = \{s_1, \dots, s_n\}$ are only defined up to homotopy. However, we have described $M[T^{-1}]$ in terms of a universal property independent of any choice, so we don’t have to worry about coherence.

III.3. Corollary. — *The functor $-[S^{-1}]: \text{Mod}_R \rightarrow \text{Mod}_R$ is a Bousfield localisation onto the S -local objects.*

Proof. Analogous to Proposition III.2. \square

Now what’s still missing is an argument why $R[S^{-1}]$ is an \mathbb{E}_∞ -ring spectrum again, and why the forgetful functor $\text{Mod}_{R[S^{-1}]} \rightarrow \text{Mod}_R$ (induced by II.62b and the map $R \rightarrow R[S^{-1}]$, which we are to show is a map of \mathbb{E}_∞ -ring spectra) is an equivalence onto the S -local objects. As we will just see, all of these follow from general principles, since $R[S^{-1}] \in \text{Mod}_R^{\otimes_R}$ is a typical example of a \otimes -idempotent object.

III.4. Proposition/Definition. — *Let \mathcal{C}^\otimes be a symmetric monoidal ∞ -category with tensor unit $1_{\mathcal{C}} \in \mathcal{C}$. A map $f: 1_{\mathcal{C}} \rightarrow x$ in \mathcal{C} is called \otimes -idempotent if*

$$\text{id}_x \otimes f: x \otimes 1_{\mathcal{C}} \xrightarrow{\sim} x \otimes x$$

is an equivalence. In this case, the following assertions hold:

- (a) *The functor $- \otimes x = L_f: \mathcal{C} \rightarrow \mathcal{C}$ is a localisation onto the full sub- ∞ -category $\mathcal{C}_f \subseteq \mathcal{C}$ spanned by all $y \in \mathcal{C}$ that absorb f , i.e. those y for which $\text{id}_y \otimes f: y \otimes 1_{\mathcal{C}} \xrightarrow{\sim} y \otimes x$ is an equivalence.*
- (b) *\mathcal{C}_f^\otimes is a symmetric monoidal ∞ -category, $L_f: \mathcal{C} \rightarrow \mathcal{C}_f$ refines to a strongly monoidal functor. Moreover, for any $y \in \mathcal{C}_f$, $z \in \mathcal{C}$ we have $y \otimes z \in \mathcal{C}_f$, and $\mathcal{C}_f^\otimes \subseteq \mathcal{C}^\otimes$ only fails to be strongly monoidal because it doesn’t preserve tensor units.*
- (c) *The strongly monoidal functor $L_f: \mathcal{C}^\otimes \rightarrow \mathcal{C}_f^\otimes$ from (b) induces a Bousfield localisation $L_f: \text{CAlg}(\mathcal{C}^\otimes) \rightarrow \text{CAlg}(\mathcal{C}_f^\otimes) \subseteq \text{CAlg}(\mathcal{C}^\otimes)$ onto those algebras whose underlying \mathcal{C} -object absorbs f . In particular, $x \in \text{CAlg}(\mathcal{C}^\otimes)$, and the induced functor*

$$\mathcal{C}_f \simeq \text{LMod}_x(\mathcal{C}_f^\otimes) \xrightarrow{\sim} \text{LMod}_x(\mathcal{C}^\otimes)$$

is an equivalence.

- (d) If \mathcal{C}^\otimes has all small operadic colimits, then the equivalence from (c) refines to a strongly symmetric monoidal equivalence

$$\mathcal{C}_f^\otimes \xrightarrow{\sim} \mathrm{LMod}_x(\mathcal{C}^\otimes)^{\otimes x}.$$

Proof. For (a), we'll apply Proposition I.61a, as usual. The map $f: 1_{\mathcal{C}} \rightarrow x$ induces a natural transformation $\eta: \mathrm{id}_{\mathcal{C}} \simeq - \otimes 1_{\mathcal{C}} \Rightarrow - \otimes x \simeq L_f$, so we must show that ηL_f and $L_f \eta$ are equivalences. Up to reordering tensor factors, both amount to showing that

$$\mathrm{id}_y \otimes \mathrm{id}_x \otimes f: y \otimes x \otimes 1_{\mathcal{C}} \xrightarrow{\sim} y \otimes x \otimes x$$

is an equivalence for all $y \in \mathcal{C}$, which is clear from the condition on x . Hence L_f is indeed a Bousfield localisation, with unit η . By definition, an element $y \in \mathcal{C}$ absorbs f iff $\eta_y: y \rightarrow L_f(y)$ is an equivalence, hence \mathcal{C}_f is indeed the essential image of L_f .

For (b), we use the criterion from Lemma II.37a(b). Unravelling the statement, we must show that whenever $g: y \rightarrow y'$ is a morphism in \mathcal{C} such that $L_f(g) \simeq \mathrm{id}_x \otimes g: x \otimes y \xrightarrow{\sim} x \otimes y'$ is an equivalence, then so is $L_f(g \otimes \mathrm{id}_z) \simeq \mathrm{id}_x \otimes g \otimes \mathrm{id}_z: x \otimes y \otimes z \xrightarrow{\sim} x \otimes y' \otimes z$ for all $z \in \mathcal{C}$. But that's just obvious, and so are the additional two assertions from (b).

For (c), note that the adjunction $L_f: \mathcal{C} \rightleftarrows \mathcal{C}_f: i$ descends to adjunctions

$$\begin{aligned} L_f: \mathrm{CAlg}(\mathcal{C}^\otimes) &\rightleftarrows \mathrm{CAlg}(\mathcal{C}_f^\otimes) : i \\ L_f: \mathrm{LMod}_x(\mathcal{C}^\otimes) &\rightleftarrows \mathrm{LMod}_x(\mathcal{C}_f^\otimes) : i, \end{aligned}$$

because unit and counit as well as the triangle identities are inherited (and for the lower adjunction we also need $L_f(x) \simeq x$). Moreover, the unit being an equivalence is inherited as well, hence $\mathrm{CAlg}(\mathcal{C}_f^\otimes) \rightarrow \mathrm{CAlg}(\mathcal{C}^\otimes)$ is fully faithful, and the characterisation of its essential image follows by inspection. The same argument shows that $\mathrm{LMod}_x(\mathcal{C}_f^\otimes) \rightarrow \mathrm{LMod}_x(\mathcal{C}^\otimes)$ is fully faithful. To see that it's essentially surjective as well, we must show that the underlying \mathcal{C} -object of every left x -module already absorbs f . So let $m \in \mathrm{LMod}_x(\mathcal{C}^\otimes)$. By abuse of notation, we identify m with its underlying \mathcal{C} -object. Then there is a multiplication map $\mu: x \otimes m \rightarrow m$ fitting into a diagram

$$\begin{array}{ccccc} m & \xrightarrow{f \otimes \mathrm{id}_m} & x \otimes m & \xrightarrow{\mu} & m \\ f \otimes \mathrm{id}_m \downarrow & & \downarrow f \otimes \mathrm{id}_x \otimes \mathrm{id}_m & & \downarrow f \otimes \mathrm{id}_m \\ x \otimes m & \xrightarrow{f \otimes \mathrm{id}_x \otimes \mathrm{id}_m} & x \otimes x \otimes m & \xrightarrow{\mathrm{id}_x \otimes \mu} & x \otimes m \end{array}$$

which shows that $f \otimes \mathrm{id}_m: m \rightarrow x \otimes m$ is a retract of an equivalence, hence an equivalence itself. Hence $\mathrm{LMod}_x(\mathcal{C}_f^\otimes) \xrightarrow{\sim} \mathrm{LMod}_x(\mathcal{C}^\otimes)$ is an equivalence. Finally, $\mathcal{C}_f^\otimes \simeq \mathrm{LMod}_x(\mathcal{C}_f^\otimes)$ follows from Corollary II.56f, since x is the tensor unit of \mathcal{C}_f^\otimes .

For (d), first note that by functoriality of the relative tensor product construction from Theorem II.62, we get a lax monoidal functor

$$i: \mathrm{LMod}_x(\mathcal{C}_f^\otimes)^{\otimes x} \longrightarrow \mathrm{LMod}_x(\mathcal{C}^\otimes)^{\otimes x}.$$

We would like to show that it's a strongly monoidal equivalence. It certainly is an equivalence on underlying ∞ -categories by (c). Hence it suffices to check $i(x) \simeq x$ and $i(m \otimes_x n) \simeq i(m) \otimes_x i(n)$. The former is obvious. For the latter, recall that i only fails to be strongly monoidal by not preserving the tensor unit, and hence both sides can be computed by the exact same bar construction (see II.62a).

Now, by construction and Proposition II.60, there are ∞ -operad maps

$$u: \mathrm{LMod}_x(\mathcal{C}_f^\otimes)^{\otimes x} \longrightarrow \mathrm{Alg}_{\mathrm{LMod}}(\mathcal{C}_f^\otimes)^\otimes \xrightarrow{\mathrm{ev}_m} \mathcal{C}_f^\otimes.$$

We are done if we can show that their composition u is a strongly monoidal equivalence. It is an equivalence on underlying ∞ -categories by (c), hence it suffices to show $u(x) \simeq x$ and $u(m \otimes_x n) \simeq u(m) \otimes u(n)$. The former is obvious again and the latter follows from the fact that the bar construction computing $u(m \otimes_x n)$ is constant on $u(m) \otimes u(n)$, as $B_n(u(m), x, u(n)) \simeq u(m) \otimes x^{\otimes n} \otimes u(n)$. \square

III.4a. Corollary. — *The R -module $R[S^{-1}]$ is naturally an \mathbb{E}_∞ -ring spectrum and the map $R \rightarrow R[S^{-1}]$ is a map of \mathbb{E}_∞ -ring spectra. The corresponding base change functor (see II.62b)*

$$R[S^{-1}] \otimes_R -: \mathrm{Mod}_R \longrightarrow \mathrm{Mod}_{R[S^{-1}]}$$

is a Bousfield localisation onto the S -local objects $\mathrm{Mod}_{R,S} \subseteq \mathrm{Mod}_R$. Under this equivalence, the symmetric monoidal structures on $\mathrm{Mod}_{R[S^{-1}]}$ (via Theorem II.62) and $\mathrm{Mod}_{R,S}$ (via Proposition/Definition III.4(b)) get identified. Finally, for any \mathbb{E}_∞ -ring spectrum T ,

$$\mathrm{Hom}_{\mathrm{CAlg}}(R[S^{-1}], T) \subseteq \mathrm{Hom}_{\mathrm{CAlg}}(R, T)$$

is the collection of path components of maps $R \rightarrow T$ that take $S \subseteq \pi_0(R)$ to units in $\pi_0(T)$.

*Proof**. This may seem tautological after Proposition/Definition III.4, but I don't think it is. For the first part, we know from Proposition/Definition III.4(c) that $R \rightarrow R[S^{-1}]$ is a map in $\mathrm{CAlg}(\mathrm{Mod}_R^{\otimes_R})$, hence it's also a map of \mathbb{E}_∞ -ring spectra via the forgetful functor $\mathrm{Mod}_R^{\otimes_R} \rightarrow \mathrm{Mod}_S^{\otimes} \simeq \mathrm{Sp}^{\otimes}$. Moreover, II.62b provides an adjunction

$$R[S^{-1}] \otimes_R -: \mathrm{Mod}_R \rightleftarrows \mathrm{Mod}_{R[S^{-1}]} : F_{R[S^{-1}]/R}.$$

The forgetful functor $F_{R[S^{-1}]/R}$ takes values in the full sub- ∞ -category $\mathrm{Mod}_{R,S} \subseteq \mathrm{Mod}_R$ of S -local objects. We claim the counit $R[S^{-1}] \otimes_R F_{R[S^{-1}]/R} \Rightarrow \mathrm{id}$ is an equivalence. But equivalences of $R[S^{-1}]$ -modules can be detected on underlying spectra (even on homotopy groups), so it definitely suffices that $R[S^{-1}] \otimes_R N \xrightarrow{\sim} N$ is an equivalence in Mod_R (rather than $\mathrm{Mod}_{R[S^{-1}]}$) for every $R[S^{-1}]$ -module N . But this follows from Corollary III.3 and the fact that N is S -local, as was noted above.

In particular, $\mathrm{Mod}_{R[S^{-1}]}$ is a full subcategory of Mod_R . To show that it coincides with $\mathrm{Mod}_{R,S}$, consider the diagram

$$\begin{array}{ccc} \mathrm{LMod}_{R[S^{-1}]}(\mathrm{Mod}_R^{\otimes_R}) & \longrightarrow & \mathrm{LMod}_R(\mathrm{Mod}_R^{\otimes_R}) \\ \downarrow & & \downarrow \wr \\ \mathrm{LMod}_{R[S^{-1}]}(\mathrm{Sp}^{\otimes}) & \longrightarrow & \mathrm{LMod}_R(\mathrm{Sp}^{\otimes}) \end{array}$$

induced by the lax symmetric monoidal functor $\mathrm{Mod}_R^{\otimes_R} \rightarrow \mathrm{Sp}^{\otimes}$. The right vertical arrow is an equivalence by Corollary II.56f, and by Proposition/Definition III.4(c), the top arrow can be identified with the inclusion $\mathrm{Mod}_{R,S} \subseteq \mathrm{Mod}_R$. Hence the bottom arrow, which is fully faithful with essential image contained in $\mathrm{Mod}_{R,S}$, also hits of $\mathrm{Mod}_{R,S}$. In particular, the left vertical arrow is an equivalence as well, and one can even show that it is strongly monoidal: It clearly preserves tensor units, hence all we need to check is

$$\mathrm{colim}_{\Delta^{\mathrm{op}}} B_\bullet(M, R, N) \simeq \mathrm{colim}_{\Delta^{\mathrm{op}}} B_\bullet(M, R[S^{-1}], N)$$

for all S -local R -modules M and N . But we can permute $\operatorname{colim}_{\Delta^{\operatorname{op}}}$ with the colimits defining $R[S^{-1}]$, and then M and N being S -local does the rest.

The final assertion about $\operatorname{Hom}_{\operatorname{CAlg}}(R[S^{-1}], T)$ can be checked on (derived) fibres over $\operatorname{Hom}_{\operatorname{CAlg}}(R, T)$. That is, it suffices to check that once we choose a map $f: R \rightarrow T$,

$$\operatorname{Hom}_{R/\operatorname{CAlg}}(R[S^{-1}], T)$$

is either contractible or empty, depending on whether f maps $S \subseteq \pi_0(R)$ to units in $\pi_0(T)$ or not. If it doesn't, then $\operatorname{Hom}_{\operatorname{CAlg}}(R[S^{-1}], T) \rightarrow \operatorname{Hom}_{\operatorname{CAlg}}(R, T)$ doesn't hit the path component of f , hence the fibre—derived or not—is empty. So let's assume f maps $S \subseteq \pi_0(R)$ to units in $\pi_0(T)$. To handle this case, we need to bring out a big gun: By a theorem of Lurie,

$$R/\operatorname{CAlg} \simeq \operatorname{CAlg}(\operatorname{Mod}_R^{\otimes_R}).$$

The proof is in [HA]: See Corollary 3.4.1.7 for the statement, Example 3.3.1.12 for why $\mathbb{C}\operatorname{om}$ is *coherent*, and 4.5.1.5 for a proof that our definition of module ∞ -categories coincides with Lurie's.

In any case, we see that $f: R \rightarrow T$ turns T into an element of $\operatorname{CAlg}(\operatorname{Mod}_R^{\otimes_R})$. By our assumption on f , T is even an element of the full sub- ∞ -category $\operatorname{CAlg}(\operatorname{Mod}_{R,S}^{\otimes_R})$, hence

$$\operatorname{Hom}_{\operatorname{CAlg}(\operatorname{Mod}_R^{\otimes_R})}(R[S^{-1}], T) \simeq \operatorname{Hom}_{\operatorname{CAlg}(\operatorname{Mod}_{R,S}^{\otimes_R})}(R, T)$$

by Proposition/Definition III.4(c). The right-hand side is contractible since R is initial in $\operatorname{CAlg}(\operatorname{Mod}_R^{\otimes_R}) \simeq R/\operatorname{CAlg}$, whence we are done. \square

LECTURE 19
14th Jan, 2021

As an example of this machinery, consider the unique \mathbb{E}_∞ ring map $\mathbb{S} \rightarrow H\mathbb{Z}$ (using that \mathbb{S} is initial in CAlg by Lemma/Definition II.45a). This map is sometimes called the “Hurewicz map”—if you tensor with $\mathbb{S}[X]$ for some $X \in \operatorname{An}$ and take homotopy groups, you get the stable Hurewicz morphism $\pi_*^s(X) \rightarrow H_*(X, \mathbb{Z})$. Localising at $\mathbb{Z} \setminus \{0\} \subseteq \mathbb{Z} = \pi_0\mathbb{S}$ gives an equivalence

$$\mathbb{S}[p^{-1} \mid p \text{ prime}] \xrightarrow{\sim} H\mathbb{Q}.$$

Indeed, first of all, the natural map $H\mathbb{Z} \rightarrow H\mathbb{Q}$ factors over $H\mathbb{Z}[p^{-1} \mid p \text{ prime}] \rightarrow H\mathbb{Q}$ by the second part of Corollary III.4a, and this map must be an equivalence as one immediately sees on homotopy groups. Moreover, the $\pi_i(\mathbb{S})$ are finite for $i > 0$ by a theorem of Serre, hence they die in the localisation, so $\mathbb{S}[p^{-1} \mid p \text{ prime}] \xrightarrow{\sim} H\mathbb{Q}$ is an equivalence on homotopy groups as well.

III.5. Corollary. — *The Eilenberg–MacLane functor*

$$H: \mathcal{D}(\mathbb{Q}) \longrightarrow \operatorname{Sp}$$

defines an equivalence onto all spectra whose homotopy groups are rational (i.e., \mathbb{Q} -vector spaces).

Proof. Having rational homotopy groups is equivalent to being $(\mathbb{Z} \setminus \{0\})$ -local, hence these spectra are precisely $\operatorname{Mod}_{\mathbb{S}[p^{-1} \mid p \text{ prime}]} \simeq \operatorname{Mod}_{H\mathbb{Q}}$. On the other hand, $H: \mathcal{D}(\mathbb{Q}) \xrightarrow{\sim} \operatorname{Mod}_{H\mathbb{Q}}$ is an equivalence by Theorem II.57. \square

So in particular, $H(M \otimes_{\mathbb{Q}}^L N) \simeq HM \otimes HN$ holds for all $M, N \in \mathcal{D}(\mathbb{Q})$ by Proposition/Definition III.4(b). We would already know this for $-\otimes_{H\mathbb{Q}}-$ instead of $-\otimes-$ (see Proposition II.63), but as it is, it's really new information. We also get that the “rational stable Hurewicz map”

$$\pi_*^s(X) \otimes \mathbb{Q} \simeq \pi_*(\mathbb{S}[X]) \otimes \mathbb{Q} \xrightarrow{\sim} H_*(X, \mathbb{Q})$$

is an equivalence for all $X \in \text{An}$.

The Group Completion Theorem and $K_1(R)$

From Proposition II.51 we get an adjunction $\mathbb{S}[-]: \text{An}^\times \rightleftarrows \text{Sp}^\otimes: \Omega^\infty$, in which the right adjoint is lax monoidal and the left adjoint even strongly so. As usual, this adjunction persists after taking $\text{CAlg}(-)$ on both sides. Since $\text{CAlg}(\text{An}^\times) \simeq \text{CMon}(\text{An})$ by Theorem II.43, the new adjunction appears as

$$\mathbb{S}[-]: \text{CMon}(\text{An}) \rightleftarrows \text{CAlg}: \Omega^\infty.$$

In particular, if R is an \mathbb{E}_∞ -ring spectrum, then $\Omega^\infty R$ carries a \mathbb{E}_∞ -monoid structure. In fact, it carries another one, due to the fact that $\Omega^\infty: \text{Sp} \rightarrow \text{An}$ factors over $\text{CGrp}(\text{An})$. But these two structures are different! The former comes from the *multiplicative* structure on R , whereas the latter comes from the *additive structure*: Since $\Omega^\infty: \text{Sp} \rightarrow \text{CGrp}(\text{An})$ is an exact functor between additive categories, it induces a strongly monoidal functor $\text{Sp}^\oplus \rightarrow \text{CGrp}(\text{An})^\oplus$ between the cartesian monoidal structures from Proposition II.42 (and also $\text{Sp} \simeq \text{CMon}(\text{Sp}^\otimes)$, $\text{CGrp}(\text{An}) \simeq \text{CMon}(\text{CGrp}(\text{An})^\otimes)$ by Theorem II.19), but these are obviously different from Sp^\otimes and $\text{CGrp}(\text{An})^\otimes$. For example, on discrete abelian groups, $\text{CGrp}(\text{An})^\otimes$ encodes the usual tensor product (Example II.51b(a)), whereas $\text{CGrp}(\text{An})^\oplus$ encodes their product.

Back to the matters at hand: Let $R \in \text{CAlg}$, and take $\Omega^\infty R$ with its multiplicative \mathbb{E}_∞ -monoid structure. Let moreover $M \in \text{CMon}(\text{An})$. We compute

$$\begin{aligned} \text{Hom}_{\text{CAlg}}(\mathbb{S}[M^{\infty\text{-grp}}], R) &\simeq \text{Hom}_{\text{CMon}(\text{An})}(M^{\infty\text{-grp}}, \Omega^\infty R) \\ &\subseteq \text{Hom}_{\text{CMon}(\text{An})}(M, \Omega^\infty R) \\ &\simeq \text{Hom}_{\text{CAlg}}(\mathbb{S}[M], R), \end{aligned}$$

and the image of the inclusion on the second line consists of all path components of maps $M \rightarrow \Omega^\infty R$ that take $\pi_0(M)$ to units in $\pi_0(\Omega^\infty R)$. Now regard $\pi_0(M)$ as sitting inside $\pi_0(\mathbb{S}[M])$ via the isomorphism

$$\pi_0(\mathbb{S}[M]) \simeq H_0(M, \mathbb{Z}) \simeq \mathbb{Z}[\pi_0(M)]$$

from Corollary* II.55b. Then we have more or less just shown:

III.6. Theorem (Group completion theorem, McDuff–Segal). — *For all $E \in \text{Sp}$ and $M \in \text{CMon}(\text{An})$ there is a canonical equivalence*

$$(E[M])[\pi_0(M)^{-1}] \xrightarrow{\sim} E[M^{\infty\text{-grp}}]$$

as $\mathbb{S}[M]$ -module spectra, and even $\mathbb{S}[M]$ -algebra spectra if $E \in \text{CAlg}$. In particular,

$$H_*(M^{\infty\text{-grp}}, \mathbb{Z}) \simeq H_*(M, \mathbb{Z})[\pi_0(M)^{-1}].$$

In Theorem III.6, and henceforth, we use the notation $E[X] := E \otimes \mathbb{S}[X]$ for $X \in \mathbf{An}$. Also recall that $M^{\infty\text{-grp}} \simeq \Omega BM$ by Corollary II.21, which is how you'll probably find this result in the literature.

Proof of Theorem III.6. It follows from our calculation above that

$$\mathrm{Hom}_{\mathrm{CAlg}}(\mathbb{S}[M^{\infty\text{-grp}}], R) \subseteq \mathrm{Hom}_{\mathrm{CAlg}}(\mathbb{S}[M], R)$$

is the collection of path components of those $\mathbb{S}[M] \rightarrow R$ that send $\pi_0(M) \subseteq \pi_0(\mathbb{S}[M]) \simeq \mathbb{Z}[\pi_0(M)]$ into units in $\pi_0(R)$. But this shows

$$(\mathbb{S}[M])[\pi_0(M)^{-1}] \simeq \mathbb{S}[M^{\infty\text{-grp}}]$$

by Corollary III.4a. The general case follows by tensoring with E . For the statement about homology, take $E \simeq H\mathbb{Z}$ and apply Corollary* II.55b. \square

To warm up for our calculation of $K_1(R)$, let's compute the homology of $(\Omega^\infty \mathbb{S})_0$, the 0-component of the \mathbb{E}_∞ -group $\Omega^\infty \mathbb{S} \simeq \mathrm{colim}_{n \in \mathbb{N}} \Omega^n \mathbb{S}^n$. To really warm up, Fabian decided to give both corollaries the number III.7, but I'm not going to do that as well.

III.6a. Corollary. — *There is a canonical isomorphism*

$$H_*((\Omega^\infty \mathbb{S})_0, \mathbb{Z}) \simeq \mathrm{colim}_{n \in \mathbb{N}} H_*^{\mathrm{grp}}(\mathfrak{S}_n, \mathbb{Z}).$$

Here $H_*^{\mathrm{grp}}(\mathfrak{S}_n, \mathbb{Z})$ denotes the group homology of the n^{th} symmetric group, and the transition maps $\mathfrak{S}_n \rightarrow \mathfrak{S}_{n+1}$ are given by mapping \mathfrak{S}_n to the permutations that fix the element $n+1$.

Proof. First up, recall the standard fact that $H_*^{\mathrm{grp}}(\mathfrak{S}_n, \mathbb{Z}) \simeq H_*(B\mathfrak{S}_n, \mathbb{Z})$, where $B\mathfrak{S}_n$ is the image of \mathfrak{S}_n under

$$\mathrm{Grp}(\mathrm{Set}) \subseteq \mathrm{Grp}(\mathbf{An}) \xrightarrow{\perp\!\!\!\perp} \mathbf{An}.$$

(see II.6 for notation). Also recall that

$$\pi_i(B\mathfrak{S}_n, *) \simeq \begin{cases} 0 & \text{if } i \neq 1 \\ \mathfrak{S}_n & \text{if } i = 1 \end{cases},$$

since \mathfrak{S}_n is a discrete anima and B shifts homotopy groups up (because its inverse Ω shifts them down).

Now let $\mathfrak{S} \simeq \coprod_{n \geq 0} B\mathfrak{S}_n$ denote the free commutative monoid on a point, or equivalently, the symmetric monoidal groupoid $(\{\text{finite sets, bijections}\}, \sqcup)$; see page 86. In particular, $\pi_0(\mathfrak{S}) \simeq \mathbb{N}$. So in order to invert $\pi_0(\mathfrak{S})$ in $H_*(\mathfrak{S}, \mathbb{Z}) \simeq \mathbb{Z}[\pi_0(\mathfrak{S})]$, we only need to invert the generator $[1] \in \pi_0(\mathfrak{S})$. Fabian remarks that this notation is a bit awkward: $[1]$ doesn't correspond to the unit in the ring $\mathbb{Z}[\pi_0(\mathfrak{S})]$, which is instead given by the unit $1 = [0] \in \pi_0(\mathfrak{S})$ of the monoid $\pi_0(\mathfrak{S}) \simeq \mathbb{N}$. Also, by the Baratt–Priddy–Quillen theorem (Corollary II.32) we have $\mathfrak{S}^{\infty\text{-grp}} \simeq \Omega^\infty \mathbb{S}$. Hence Theorem III.6 shows

$$H_*(\Omega^\infty \mathbb{S}, \mathbb{Z}) \simeq \mathrm{colim}_{\mathbb{N}} \left(H_*(\mathfrak{S}, \mathbb{Z}) \xrightarrow{\cdot[1]} H_*(\mathfrak{S}, \mathbb{Z}) \xrightarrow{\cdot[1]} \dots \right).$$

Now isolate the components corresponding to $(\Omega^\infty \mathbb{S})_0 \simeq (\mathfrak{S}^{\infty\text{-grp}})_0$ on both sides: In the colimit on the right-hand side, $\cdot[1]$ maps the component $B\mathfrak{S}_0$ into $B\mathfrak{S}_1$, then into $B\mathfrak{S}_2$, then into $B\mathfrak{S}_3$ etc., and these maps $B\mathfrak{S}_n \rightarrow B\mathfrak{S}_{n+1}$ are precisely the transition specified in the formulation of the corollary. Hence we really obtain

$$H_*((\Omega^\infty \mathbb{S})_0, \mathbb{Z}) \simeq \mathrm{colim}_{n \in \mathbb{N}} H_*(B\mathfrak{S}_n, \mathbb{Z}) \simeq \mathrm{colim}_{n \in \mathbb{N}} H_*^{\mathrm{grp}}(\mathfrak{S}_n, \mathbb{Z}). \quad \square$$

III.6b. — There's another way of phrasing the result from Corollary III.6a. Consider the “group of finitely supported permutations” $\mathfrak{S}_\infty := \operatorname{colim}_{n \in \mathbb{N}} \mathfrak{S}_n$. The canonical map $\mathfrak{S} \rightarrow \mathfrak{S}^{\infty\text{-grp}} \simeq \Omega^\infty \mathbb{S}$ induces a map

$$\mathbb{Z} \times B\mathfrak{S}_\infty \simeq \operatorname{colim}_{n \in \mathbb{N}} \left(\mathfrak{S} \xrightarrow{[1]} \mathfrak{S} \xrightarrow{[1]} \dots \right) \longrightarrow \operatorname{colim}_{n \in \mathbb{N}} \left(\Omega^\infty \mathbb{S} \xrightarrow{[1]} \Omega^\infty \mathbb{S} \xrightarrow{[1]} \dots \right) \simeq \Omega^\infty \mathbb{S}$$

(where we use that $\Omega^\infty \mathbb{S}$ is already an \mathbb{E}_∞ -group, so all transition maps in the second colimit are equivalences). As all components of an \mathbb{E}_∞ -group are equivalent, Corollary III.6a implies that the map $\mathbb{Z} \times B\mathfrak{S}_\infty \rightarrow \Omega^\infty \mathbb{S}$ induces isomorphisms

$$H_*(\mathbb{Z} \times B\mathfrak{S}_\infty, \mathbb{Z}) \xrightarrow{\sim} H_*(\Omega^\infty \mathbb{S}, \mathbb{Z})$$

on homology. But it's far from being a homotopy equivalence! Amusingly, this already follows from Corollary III.6a itself: As seen in its proof,

$$\pi_1(\mathbb{Z} \times B\mathfrak{S}_\infty, (0, \operatorname{id})) \simeq \operatorname{colim}_{n \in \mathbb{N}} \pi_1(B\mathfrak{S}_n, \operatorname{id}) \simeq \mathfrak{S}_\infty,$$

whereas

$$\pi_1(\Omega^\infty \mathbb{S}, 0) \simeq H_1((\Omega^\infty \mathbb{S})_0, \mathbb{Z}) \simeq \operatorname{colim}_{n \in \mathbb{N}} H_1(\mathfrak{S}_n, \mathbb{Z}) \simeq \operatorname{colim}_{n \in \mathbb{N}} \mathfrak{S}_n^{\text{ab}} \simeq \mathbb{Z}/2\mathbb{Z}.$$

The first isomorphism follows from the Poincaré lemma, as $\pi_1(\Omega^\infty \mathbb{S}, 0) \simeq \pi_1(\mathbb{S})$ is already abelian, the third isomorphism follows from the Poincaré lemma in group homology, and the fourth from the fact that the commutator $[\mathfrak{S}_n, \mathfrak{S}_n] \simeq \mathfrak{A}_n$ is the alternating group for $n \geq 2$, which has index 2.

So $\mathbb{Z} \times B\mathfrak{S}_\infty \rightarrow \Omega^\infty \mathbb{S}$ is an example where the homology Whitehead theorem fails for anima which aren't simple. And also it shows that the group completion theorem fails rather drastically before taking suspension spectra. The exact nature of this failure will be thoroughly investigated in III.11a.

III.7. Corollary. — *Let R be a ring. Then there's a canonical isomorphism*

$$H_*(k(R)_0, \mathbb{Z}) \xrightarrow{\sim} \operatorname{colim}_{n \in \mathbb{N}} H_*^{\text{grp}}(\operatorname{GL}_n(R), \mathbb{Z}).$$

In particular,

$$K_1(R) = \operatorname{colim}_{n \in \mathbb{N}} \operatorname{GL}_n(R)^{\text{ab}} \simeq \operatorname{GL}_\infty(R)^{\text{ab}}$$

Proof. While we don't understand $\pi_0 \operatorname{Proj}(R)$, to invert it in $H_*(\operatorname{Proj}(R), \mathbb{Z})$ it suffices to invert the element $[R] \in \pi_0 \operatorname{Proj}(R)$. Indeed, for every finite projective R -module P , the class $[P]$ “divides” some power of $[R]$ in $\pi_0 \operatorname{Proj}(R)$ since there is some Q such that $P \oplus Q \simeq R^n$. Thus from Theorem III.6 applied to $k(R) \simeq \operatorname{Proj}(R)^{\infty\text{-grp}}$ we find

$$H_*(k(R), \mathbb{Z}) \simeq \operatorname{colim}_{n \in \mathbb{N}} \left(H_*(\operatorname{Proj}(R), \mathbb{Z}) \xrightarrow{[R]} H_*(\operatorname{Proj}(R), \mathbb{Z}) \xrightarrow{[R]} \dots \right)$$

We have $\operatorname{Proj}(R) \simeq \coprod_{[P] \in \pi_0 \operatorname{Proj}(R)} B\operatorname{GL}(P)$ by direct inspection. Indeed, by II.6, constructing an equivalence from $B\operatorname{GL}_n(P)$ onto the component of P is the same as constructing an equivalence $B\operatorname{GL}_n(P) \xrightarrow{\sim} \Omega_P \operatorname{Proj}(R) \simeq \operatorname{Hom}_{\operatorname{Proj}(R)}(P, P)$. But since $\operatorname{Proj}(R)$ is the

1-groupoid of finite projective R -modules, the right-hand side just *is* the discrete anima $\mathrm{GL}_n(P)$, as required.

As in the proof of Corollary III.6a, the assertion now follows by isolating the component of 0 on both sides: In the colimit right-hand side, the component of 0 is mapped to $B\mathrm{GL}_1(R)$, then to $B\mathrm{GL}_2(R)$, then to $B\mathrm{GL}_3(R)$ etc., so we really get

$$H_*(k(R)_0, \mathbb{Z}) \simeq \operatorname{colim}_{n \in \mathbb{N}} H_*(B\mathrm{GL}_n(R), \mathbb{Z}) \simeq \operatorname{colim}_{n \in \mathbb{N}} H_*^{\mathrm{grp}}(\mathrm{GL}_n(R), \mathbb{Z}).$$

In the special case $*$ = 1, we apply the Poincaré lemma in singular homology (using that $\pi_1(k(R), 0)$ is already abelian) and group homology to get the desired formula for $K_1(R)$. \square

III.7a. — As for Corollary III.6a, we could also reformulate Corollary III.7 as follows: There exists a map $K_0(R) \times B\mathrm{GL}_\infty(R) \rightarrow k(R)$, which induces an isomorphism on homology.

Fabian also mentioned in the lecture, and elaborates in his notes [A&HK, Chapter III pp. 48–49], that for $r \in \{\text{quad, even}\}$, one can compute $H_*(\mathrm{gw}^r(\mathbb{Z}), \mathbb{Z})$ via a similar calculation, but this requires non-trivial input about $\pi_0 \mathrm{Unimod}^r(\mathbb{Z})$.

III.7b. $K_1(R)$ and Determinants. — Note that for $P \in \mathrm{Proj}(R)$ we get a map

$$\mathrm{Aut}(P) \simeq \pi_1(\mathrm{Proj}(R), P) \longrightarrow \pi_1(k(R), P) \simeq K_1(R)$$

(the last equivalence follows since all path components of the \mathbb{E}_∞ -group $k(R)$ must be equivalent). If R is commutative then the determinant gives a map $\det: K_1(R) \simeq \mathrm{GL}_\infty(R)^{\mathrm{ab}} \rightarrow R^\times$. Composing it with the morphism $\mathrm{Aut}(P) \rightarrow K_1(R)$ from above gives a way to extend determinants to automorphisms of arbitrary finite projective modules. Of course, such an extension could also be obtained by hand, but it pops out for free here. In general, for not necessarily commutative rings R , we may think of the morphism $\mathrm{Aut}(P) \rightarrow K_1(R)$ as a good “generalised determinant”.

If R is commutative, the kernel of $\det: K_1(R) \rightarrow R^\times$ is usually denoted $SK_1(R)$ (the “special” K -group). Since \det admits a splitting via $R^\times \simeq \mathrm{GL}_1(R) \rightarrow \mathrm{GL}_\infty^{\mathrm{ab}}$, we obtain

$$K_1(R) \simeq R^\times \oplus SK_1(R).$$

III.8. Proposition (Whitehead’s lemma). — *The commutator subgroup of $\mathrm{GL}_\infty(R)$ is generated by the elementary matrices*

$$\begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & \ddots & r & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \cdots & 0 & 1 \end{pmatrix} = \mathbb{1}_n + re_{i,j}$$

for $i, j \in \{1, \dots, n\}$, $i \neq j$, and $r \in R$; in other words, by those (finite) matrices with all diagonal entries equal to 1 and one off-diagonal entry equal to some $r \in R$.

So $SK_1(R)$ is the obstruction group for doing row operations to get any matrix into diagonal form. In particular, $SK_1(R) = 0$ if R is a euclidean domain.

Proof. Omitted. But Fabian’s notes [A&HK, Proposition III.8] have some details. \square

LECTURE 20
19th Jan, 2021

Note that $SK_1(R) = 0$ is not necessarily true when R is a PID, so we really need R to have a euclidean algorithm. However, counterexamples are rather obscure (due to the fact that $SK_1(\mathcal{O}_F) = 0$ whenever \mathcal{O}_F is the ring of integers in a number field): One can take $R = \mathbb{Z}[X][\Phi_n^{-1} \mid n \in \mathbb{N}]$ for example, where Φ_n denotes the n^{th} cyclotomic polynomial.

Towards the K -Theory of Finite Fields

Quillen's computation of $K_*(\mathbb{F}_q)$ proceeds in a rather roundabout way, with the essential step being a comparison with complex K -theory. This necessitates a detour into the land of (various versions of) topological K -theory. But before that, we'll take another detour and compute the rational K -groups $K_*(R) \otimes \mathbb{Q}$, which will already have a heavy topological flavour.

Rational K -Theory

Recall that the rational cohomology of any anima $X \in \mathbf{An}$ has a comultiplication

$$\Delta: H_*(X, \mathbb{Q}) \xrightarrow{\Delta_*} H_*(X \times X, \mathbb{Q}) \xleftarrow{\sim} H_*(X, \mathbb{Q}) \otimes H_*(X, \mathbb{Q}),$$

where the right arrow is the Künneth isomorphism. It is coassociative and counital (the counit being $\varepsilon: H_*(X, \mathbb{Q}) \rightarrow H_*(*, \mathbb{Q}) \simeq \mathbb{Q}$), hence makes $H_*(X, \mathbb{Q})$ into a *coalgebra*—in fancy words, an element of $\mathrm{Alg}_{\mathbf{Assoc}}((\mathbf{Ab}^{\otimes})^{\mathrm{op}})$. As it turns out, $H_*(X, \mathbb{Q})$ is moreover graded commutative.

If X is connected, we have a canonical element $1 \in H_0(X, \mathbb{Q})$, given by the image of $1 \in H_0(X, \mathbb{Z}) \simeq \mathbb{Z}$. An arbitrary element $\alpha \in H_*(X, \mathbb{Q})$ is then called *primitive* if

$$\Delta(\alpha) = 1 \otimes \alpha + \alpha \otimes 1.$$

To acquaint myself with the comultiplication a little more, I decided to put some of its properties (which are probably clear to you) into a lemma.

III.8a. Lemma*. — *Let X be connected.*

- (a) *The primitive elements form a sub- \mathbb{Q} -vector space of $H_*(X, \mathbb{Q})$.*
- (b) *For general $\alpha \in H_*(X, \mathbb{Q})$, we have*

$$\Delta(\alpha) = 1 \otimes \alpha + \alpha \otimes 1 + \sum_i \alpha'_i \otimes \alpha''_i,$$

where the α'_i and α''_i are elements sitting in positive degrees.

- (c) *The rational Hurewicz map $h: \pi_*(X, x) \otimes \mathbb{Q} \rightarrow H_*(X, \mathbb{Q})$ lands in the primitives.*

Proof.* Part (a) is clear since Δ is \mathbb{Q} -linear. For (b), we consider the following diagram:

$$\begin{array}{ccccc} & & H_*(\ast \times X, \mathbb{Q}) & \xleftarrow{\sim} & \mathbb{Q} \otimes H_*(X, \mathbb{Q}) \\ & \nearrow 1 \otimes - & \uparrow \mathrm{pr}_{2,*} & & \uparrow \varepsilon \otimes \mathrm{id} \\ H_*(X, \mathbb{Q}) & \xrightarrow{\Delta_*} & H_*(X \times X, \mathbb{Q}) & \xleftarrow{\sim} & H_*(X, \mathbb{Q}) \otimes H_*(X, \mathbb{Q}) \\ & \searrow - \otimes 1 & \downarrow \mathrm{pr}_{1,*} & & \downarrow \mathrm{id} \otimes \varepsilon \\ & & H_*(X \times \ast, \mathbb{Q}) & \xleftarrow{\sim} & H_*(X, \mathbb{Q}) \otimes \mathbb{Q} \end{array}$$

The diagram shows that the part of $\Delta(\alpha)$ not contained in $\bigoplus_{i,j \geq 1} H_i(X, \mathbb{Q}) \otimes H_j(X, \mathbb{Q})$ must be $1 \otimes \alpha + \alpha \otimes 1$, as claimed.

Finally, for (c) take some element $[f] \in \pi_n(X, x)$, represented by a map $f: \mathbb{S}^n \rightarrow X$. Then $h[f]$ is in the image of $f_*: H_*(\mathbb{S}^n, \mathbb{Q}) \rightarrow H_*(X, \mathbb{Q})$ and hence it suffices to check that all elements of $H_*(\mathbb{S}^n, \mathbb{Q})$ are primitive. But $H_*(\mathbb{S}^n, \mathbb{Q})$ is only non-zero in degrees 0 and n , and Δ respects the degree of an element, hence the formula from (b) shows that all elements must indeed be primitive. \square

III.9. Proposition (Cartan–Serre). — *Let X be a connected simple anima. That is, the action of $\pi_1(X)$ on $\pi_n(X)$ is trivial for all n (in particular, $\pi_1(X)$ acts trivially on itself, it must be abelian since it's a well-known fact that $\pi_1(X)$ acts on itself via conjugation). Suppose furthermore that the rational cohomology of X is a free graded-commutative \mathbb{Q} -algebra of finite type. Then the rational Hurewicz map*

$$\pi_*(X) \otimes \mathbb{Q} \longrightarrow H_*(X, \mathbb{Q})$$

is an isomorphism onto the primitives for all $ > 0$.*

Proof sketch. Choose free algebra generators $x_i \in H^{n_i}(X, \mathbb{Q})$, $i = 1, \dots, r$. By Theorem I.56a, these determine (up to homotopy) a map

$$f: X \longrightarrow \prod_{i=1}^r K(\mathbb{Q}, n_i).$$

We claim:

(\boxtimes) *The map f induces an isomorphism on rational homotopy groups $\pi_* \otimes \mathbb{Q}$.*

Once we have (\boxtimes), it suffices to check the assertion of the proposition in the special case where X is a rational *gem* (“generalised Eilenberg–MacLane space”). This can be done by a direct computation, using that the rational cohomology of Eilenberg–MacLane spaces is known.

Since we also need it to show (\boxtimes), let’s recall the result (a reference is [Dic08, Theorem (20.7.1)]):

$$H^*(K(\mathbb{Z}, n), \mathbb{Q}) = H^*(K(\mathbb{Q}, n), \mathbb{Q}) = \begin{cases} \mathbb{Q}[t_{2i}] & \text{if } n = 2i \\ \Lambda_{\mathbb{Q}}[t_{2i+1}] & \text{if } n = 2i + 1 \end{cases}$$

(since $K(\mathbb{Z}, n) \rightarrow K(\mathbb{Q}, n)$ is an isomorphism on rational homotopy groups, it’s also an isomorphism on rational cohomology by Serre’s rational Hurewicz theorem, which is in turn an application of the Leray–Serre spectral sequence). So if $n = 2i$ is even, we get a polynomial ring on a single generator in degree $2i$, and if $n = 2i + 1$ is odd, we get an alternating algebra on a single generator in degree $2i + 1$. Note that $\Lambda_{\mathbb{Q}}[t_{2i+1}] = H^*(\mathbb{S}^{2i+1}, \mathbb{Q})$ has only two non-zero degrees.

By assumption on X and our choice of the x_i , this implies that the map

$$f^*: \bigotimes_{i=1}^r H^*(K(\mathbb{Q}, n_i), \mathbb{Q}) \simeq H^*\left(\prod_{i=1}^r K(\mathbb{Q}, n_i), \mathbb{Q}\right) \xrightarrow{\sim} H^*(X, \mathbb{Q})$$

is an isomorphism. In other words, f is an isomorphism in rational cohomology, hence also in rational homology (all cohomology groups are finite-dimensional \mathbb{Q} -vector spaces, thus isomorphic to their duals). Now (\boxtimes) follows from Serre’s rational Hurewicz theorem. \square

III.9a. Remark. — Fabian remarks that there are versions of Proposition III.9 that also work without the finiteness assumptions. But then one has to be a bit careful to put the right topology on our graded rings, and also to replace “free” by “topologically free”.

III.9b. Hopf Algebras. — If M is a connected \mathbb{E}_1 -group whose rational homology groups $H_*(M, \mathbb{Q})$ are finite-dimensional \mathbb{Q} -vector spaces for all $n \geq 0$, then Proposition III.9 applies to M . Indeed, the fact that M is simple holds more generally for all H -spaces, see [Hat02, Example 4A.3] (but mind that Hatcher uses “abelian” rather than “simple”). To see that $H^*(M, \mathbb{Q})$ is a free graded commutative \mathbb{Q} -algebra, we need to use that $A = H_*(M, \mathbb{Q})$ is a *Hopf algebra* over \mathbb{Q} . This means the following:

- (a) A has a multiplication $\mu: A \otimes A \rightarrow A$ and a unit map $u: \mathbb{Q} \rightarrow A$, which turn A into an associative and unital (but not necessarily commutative) \mathbb{Q} -algebra. In our case $A = H_*(M, \mathbb{Q}) = \pi_*(H\mathbb{Q}[M])$, we use that $H\mathbb{Q}[M]$ is an \mathbb{E}_1 -ring spectrum, hence its homotopy groups form a graded ring (see II.66).
- (b) A has a comultiplication $\Delta: A \rightarrow A \otimes A$ and a counit $\varepsilon: A \rightarrow \mathbb{Q}$, turning A into a coassociative and counital \mathbb{Q} -coalgebra. In our case $A = H_*(M, \mathbb{Q})$ we use the structure from the beginning of the subsection.
- (c) μ and u are morphisms of \mathbb{Q} -coalgebras, or equivalently, Δ and ε are morphism of \mathbb{Q} -algebras. I’ll leave it to you to verify that this is satisfied in our case.
- (d) A has an *antipode*, i.e. a \mathbb{Q} -linear map $i: A \rightarrow A$ fitting into a commutative diagram

$$\begin{array}{ccccc}
 & A \otimes A & \xrightarrow{(\text{id}_A, i)} & A \otimes A & \\
 \Delta \nearrow & & & & \searrow \mu \\
 A & \xrightarrow{\varepsilon} & \mathbb{Q} & \xrightarrow{u} & A \\
 \Delta \searrow & & & & \nearrow \mu \\
 & A \otimes A & \xrightarrow{(i, \text{id}_A)} & A \otimes A &
 \end{array}$$

In our case, we can i to be induced by $(-)^{-1}: M \rightarrow M$.

Any graded commutative and degree-wise finite-dimensional Hopf algebra A over \mathbb{Q} has a free graded commutative underlying algebra (see [Hat02, Theorem 3C.4] for example). In particular, if the rational homology of our connected \mathbb{E}_1 -group M is degree-wise finite dimensional, we may apply this result to the dual Hopf algebra $H^*(M, \mathbb{Q}) \simeq H_*(M, \mathbb{Q})^\vee$ (which is graded commutative since $H_*(M, \mathbb{Q})$ is graded cocommutative) and obtain that $H^*(M, \mathbb{Q})$ is a free graded commutative \mathbb{Q} -algebra. Hence Proposition III.9 applies to M and we obtain that

$$\pi_*(M) \otimes \mathbb{Q} \longrightarrow H_*(M, \mathbb{Q})$$

is an isomorphism onto the primitives.

This is still true in the case where $H_*(M, \mathbb{Q})$ isn’t necessarily finite-dimensional in every degree, since Proposition III.9 can also be extended to this case (see Remark III.9a above). Following [MM65, Appendix], we can say even more: Let A be a nonnegatively graded Hopf algebra over \mathbb{Q} with $A_0 = \mathbb{Q}$ (i.e. A is connected). Then the primitive elements $P(A) \subseteq A$ form a Lie algebra under the commutator, and if A is graded cocommutative, then $U(P(A)) \simeq A$ by a theorem of Leray (where $U(\mathfrak{g})$ denotes the universal enveloping algebra of a Lie algebra \mathfrak{g}). Applying this to the graded cocommutative connected Hopf algebra $H_*(M, \mathbb{Q})$, we get

$$U(\pi_*(M) \otimes \mathbb{Q}) \simeq H_*(M, \mathbb{Q}).$$

The Lie algebra structure on $\pi_*(M) \otimes \mathbb{Q}$ can be described using the *Samelson product* $[-, -]: M \wedge M \rightarrow M$.

We can still say more: Let A be a nonnegatively graded Hopf algebra over \mathbb{Q} with $A_0 = \mathbb{Q}$. If A is both graded commutative and graded cocommutative, then it must be free on its primitive elements by another general result. A reference for this result is [MM65, Corollary 4.18], but this needs a short explanation: What the reference says is that in this case the primitive elements of A coincide with its *indecomposables*, i.e. with

$$\text{indec}_* A := \text{coker} \left(A_{>0} \otimes A_{>0} \xrightarrow{\mu} A_{>0} \right).$$

If you think about this for a moment, this precisely means that A is free on its primitives, as desired. Observe that if M is a connected \mathbb{E}_∞ -monoid, then the Hopf algebra $H_*(M, \mathbb{Q})$ is both graded commutative and cocommutative, hence the result can be applied and $H_*(M, \mathbb{Q})$ is free on its primitive elements $\pi_*(M) \otimes \mathbb{Q}$. In the special case $M \simeq \Omega^\infty$ for some $E \in \text{Sp}$, the discussion around Corollary III.5 shows $\pi_*(E) \otimes \mathbb{Q} \simeq \pi_*(E \otimes H\mathbb{Q})$. Put the right-hand side is $H_*(E, \mathbb{Q})$ by Corollary* II.55b. Hence the primitive elements $\pi_* \Omega^\infty E \otimes \mathbb{Q} \simeq H_{*\geq 0}(E, \mathbb{Q})$ are given by the nonnegative part of the homology of E . This shows that

$$H_*(\Omega^\infty E, \mathbb{Q}) \simeq \text{Sym}_{\mathbb{Q}}^{\text{gr}}(H_{*\geq 0}(E, \mathbb{Q}))$$

is the free graded commutative \mathbb{Q} -algebra on $H_{*\geq 0}(E, \mathbb{Q})$.

III.10. Corollary. — *For any ring R and all $i \geq 1$, there are a canonical isomorphisms*

$$K_i(R) \otimes \mathbb{Q} \simeq \text{indec}_i H_*^{\text{grp}}(\text{GL}_\infty(R), \mathbb{Q}).$$

Here $H_*^{\text{grp}}(\text{GL}_\infty(R), -)$ is understood to denote $\text{colim}_{n \in \mathbb{N}} H_*^{\text{grp}}(\text{GL}_n(R), -)$.

Proof.* By definition, we have $K_i(R) \otimes \mathbb{Q} \simeq \pi_i(k(R)_0) \otimes \mathbb{Q}$, which is isomorphic to the degree- i primitive elements of $H_*(k(R)_0, \mathbb{Q})$ by the discussion above, since $k(R)_0$ is a connected \mathbb{E}_∞ -group. But from Corollary III.7 and the universal coefficient theorems for singular homology and group homology, we get $H_*(k(R)_0, \mathbb{Q}) \simeq H_*^{\text{grp}}(\text{GL}_\infty(R), \mathbb{Q})$. Since this is a graded commutative and graded cocommutative Hopf algebra, its primitive elements coincide with its indecomposables by [MM65, Corollary 4.18], whence we are done. \square

Corollary III.10 reduces the computation of rational K -theory to a problem purely from group homology. This still isn't easy by any means, but known in many cases. For example, we have the following theorem (whose proof, as Fabian explained, is pretty crazy and way beyond the scope of this lecture):

III.11. Theorem (Borel). — *If \mathcal{O}_F is the ring of integers in a number field F (i.e. a finite extension F/\mathbb{Q}), then the K -groups of \mathcal{O}_F are given by*

$$K_i(\mathcal{O}_F) \otimes \mathbb{Q} = \begin{cases} \mathbb{Q} & \text{if } i = 0 \\ 0 & \text{if } i = 1 \text{ or } i > 0 \text{ even} \\ \mathbb{Q}^{r+s} & \text{if } i \equiv 1 \pmod{4} \text{ and } i > 1 \\ \mathbb{Q}^s & \text{if } i \equiv 3 \pmod{4} \end{cases}.$$

Here r and s are the numbers of real and complex embeddings of F , respectively.

The Cyclic Invariance Condition and Quillen's Plus Construction

Let's analyse what goes wrong with the group completion theorem before taking $\mathbb{S}[-]$, as promised after Corollary III.6a.

III.11a. A Subtlety. — For simplicity, we assume that $M \in \mathbf{CMon}(\mathbf{An})$ is an \mathbb{E}_∞ -monoid for which there is an $s \in \pi_0(M)$ with $(\pi_0(M))[s^{-1}] \simeq \pi_0(M)^{\text{grp}}$. This is always true in our cases of interest; for example, we’ve seen it for $M = \mathfrak{S}$ and $M = k(R)$ in the proofs of Corollaries III.6a and III.7. If you’re not willing to make this assumption, you’ll have to iterate the constructions to come.

Since $\mathbf{CMon}(\mathbf{An}) \simeq \mathbf{CAlg}(\mathbf{An}^\times)$ by Theorem II.43, we can form the module category $\mathbf{LMod}_M(\mathbf{An}^\times)$ as in Example II.45(e). The adjunction from Corollary* II.56g shows $\pi_0 \mathbf{Hom}_{\mathbf{LMod}_M(\mathbf{An}^\times)}(M, M) \simeq \pi_0 \mathbf{Hom}_{\mathbf{An}}(*, M) \simeq \pi_0(M)$, hence multiplication with s is an M -module map (as we would expect). Now put

$$T(M, s) \simeq \operatorname{colim}_{\mathbb{N}} \left(M \xrightarrow{s} M \xrightarrow{s} M \xrightarrow{s} \dots \right).$$

By Corollary II.56e, $T(M, s)$ is naturally an M -module again. But $T(M, s)$ is usually *not* s -local, i.e. the map $s: T(M, s) \rightarrow T(M, s)$ fails to be an equivalence!

For example, take $M = \mathfrak{S}$ and $s = [1]$, then $T(\mathfrak{S}, [1]) \simeq \mathbb{Z} \times B\mathfrak{S}_\infty$, as seen after Corollary III.6a. But multiplication with $[1]$ does not correspond to shifting components! It does take $\{n\} \times B\mathfrak{S}_\infty \rightarrow \{n+1\} \times B\mathfrak{S}_\infty$, but this map is not the identity on $B\mathfrak{S}_\infty$, and in fact not even an equivalence. Instead, it can be described as follows: Let’s identify permutations with their corresponding permutation matrices and consider the map

$$\begin{aligned} \varphi: \mathfrak{S}_\infty &\longrightarrow \mathfrak{S}_\infty \\ A &\longmapsto \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

Then $[1]: \{n\} \times B\mathfrak{S}_\infty \rightarrow \{n+1\} \times B\mathfrak{S}_\infty$ is given by $B\varphi: B\mathfrak{S}_\infty \rightarrow B\mathfrak{S}_\infty$. This can’t be an equivalence since it’s not even the induced morphism on fundamental groups, which is φ , is an isomorphism.

To unwind what’s going on, let temporarily $r \in \pi_0(M)$ be another element (we’ll soon take $r = s$, but doing it right away would be confusing rather than illuminating). Then the multiplication $r: T(M, s) \rightarrow T(M, s)$ can be described by a big diagram

$$\begin{array}{ccc} T(M, s) & & \\ \downarrow r & \simeq \operatorname{colim}_{\mathbb{N}} & \left(\begin{array}{ccccc} M & \xrightarrow{s} & M & \xrightarrow{s} & M & \xrightarrow{s} & \dots \\ \downarrow r & \text{///} \tau & \downarrow r & \text{///} \tau & \downarrow r & & \\ M & \xrightarrow{s} & M & \xrightarrow{s} & M & \xrightarrow{s} & \dots \end{array} \right) \\ T(M, s) & & \end{array}$$

The homotopy τ occurring in every square comes from the *symmetric* monoidal structure on M , so in particular, τ is an equivalence and the squares commute. However, in the special case $r = s$, τ doesn’t need to be the constant homotopy, i.e. the identity in $\mathbf{Hom}_{\mathbf{Fun}(M, M)}(s^2, s^2)$, but rather whatever equivalence the symmetric monoidal structure gives us. After all, commutativity is a *structure*, not a *property*!

In the example of $M = \mathfrak{S}$, the symmetry isomorphism between $[n] \cdot [m]: \mathfrak{S} \rightarrow \mathfrak{S}$ and $[m] \cdot [n]: \mathfrak{S} \rightarrow \mathfrak{S}$ is not the identity. Indeed, on matrix representations of permutations we may visualise $[n] \cdot [m]$ and $[m] \cdot [n]$ as

$$[n] \cdot [m]: A \longmapsto \begin{pmatrix} A & 0 & 0 \\ 0 & \mathbb{1}_m & 0 \\ 0 & 0 & \mathbb{1}_n \end{pmatrix} \quad \text{and} \quad [m] \cdot [n]: A \longmapsto \begin{pmatrix} A & 0 & 0 \\ 0 & \mathbb{1}_n & 0 \\ 0 & 0 & \mathbb{1}_m \end{pmatrix}$$

and in this picture, the symmetry isomorphism “flips the lower two blocks”. This looks super tautological and you really have to look twice to see why this isn’t the identity. But it really isn’t: If $m = n = 1$, then $[1] \cdot [1]: \mathfrak{S} \rightarrow \mathfrak{S}$ adds two elements to each finite set $S \in \mathfrak{S}$, and the symmetry isomorphism *swaps these two new elements*!

Back to the general situation: Since we’ve seen that $s: T(M, s) \rightarrow T(M, s)$ might not be an equivalence, the argument we would normally use cannot apply. So what is this “usual argument”? If $\tilde{s}: T(M, s) \rightarrow T(M, s)$ denotes the map induced by the constant homotopies, we can construct an inverse $t: T(M, s) \rightarrow T(M, s)$ by means of the diagram

$$\begin{array}{ccc} T(M, s) & & \\ \uparrow t & \simeq \operatorname{colim}_{\mathbb{N}} & \left(\begin{array}{ccccc} M & \xrightarrow{s} & M & \xrightarrow{s} & M & \xrightarrow{s} & \dots \\ & \nearrow s & // \operatorname{id} & \nearrow s & // \operatorname{id} & \nearrow s & \\ M & \xrightarrow{s} & M & \xrightarrow{s} & M & \xrightarrow{s} & \dots \end{array} \right) \\ T(M, s) & & \end{array}$$

(use cofinality to see that t is really an inverse). But since $s: T(M, s) \rightarrow T(M, s)$ doesn’t necessarily coincide with \tilde{s} , the map t doesn’t necessarily provide an inverse.

III.11b. Why Does It Work for Spectra Though? — For an \mathbb{E}_∞ -ring spectrum R , a module M over it, and an element $s \in \pi_0(R)$, we have defined the localisation $M[s^{-1}]$ as $T(M, s)$ in Proposition III.2, and have had no problems with M not being s -local. This was because we can detect on homotopy groups whether $s: T(M, s) \rightarrow T(M, s)$ is an equivalence. The reason why this fails in general, even though equivalences can still be detected on homotopy groups of underlying anima, are basepoint issues: In general, we must consider all possible basepoints, and there might be non-equivalent connected components, whereas for spectra the canonical choice of basepoints always suffices (which was something we had to argue in the proof* of Lemma* II.23c).

III.11c. When Does It Work in General? — We’ve seen in III.11a that the map $s: T(M, s) \rightarrow T(M, s)$ is an equivalence if $\tau = \operatorname{id}$. But we can do better: Actually, it suffices to have $\tau^n = \operatorname{id} \in \operatorname{Hom}_{\operatorname{Fun}(M, M)}(s^{n+1}, s^{n+1})$ for some $n \geq 1$, since we can also write

$$T(M, s) \simeq \operatorname{colim}_{\mathbb{N}} \left(M \xrightarrow{s^n} M \xrightarrow{s^n} M \xrightarrow{s^n} \dots \right)$$

by cofinality. Moreover, we don’t need the 2-cell τ (or τ^n for that matter) to be trivial right away, it suffices when it’s trivial after mapping to $T(M, s)$.

To formalise these considerations, let’s construct natural maps $\mathfrak{S}_n \rightarrow \pi_1(M, m^n)$ for all $m \in M$ and $n \geq 1$. I’m still a bit confused about how we did this in the lecture, so I’ll give an alternative description of (hopefully) the same construction: Since $M \in \operatorname{CMon}(\operatorname{An})$, it defines a functor $M^{(-)}: \mathbb{I}^{\operatorname{op}} \rightarrow \operatorname{An}$ sending $\langle n \rangle$ to M^n . From functoriality of $M^{(-)}$ we get a homotopy-commutative diagram

$$\begin{array}{ccccccc} \mathfrak{S}_n & \longrightarrow & \operatorname{Hom}_{\mathbb{I}^{\operatorname{op}}}(\langle n \rangle, \langle n \rangle) & \longrightarrow & \operatorname{Hom}_{\operatorname{An}}(M^n, M^n) & \longrightarrow & \operatorname{Hom}_{\operatorname{An}}(\{(m, \dots, m)\}, M^n) \simeq M^n \\ \downarrow & & \downarrow (f_n)_* & & \downarrow \mu_* & & \downarrow \mu_* \\ * & \xrightarrow{f_n} & \operatorname{Hom}_{\mathbb{I}^{\operatorname{op}}}(\langle n \rangle, \langle 1 \rangle) & \longrightarrow & \operatorname{Hom}_{\operatorname{An}}(M^n, M) & \longrightarrow & \operatorname{Hom}_{\operatorname{An}}(\{(m, \dots, m)\}, M) \simeq M \end{array}$$

It induces a map $\mathfrak{S}_n \rightarrow M^n \times_M \{\mu\}$, unique up to contractible choice. But upon inspection, the composition $\mathfrak{S}_n \rightarrow M^n$ from the top row sends everyone of the $n!$ points of the discrete anima \mathfrak{S}_n to the point $(m, \dots, m) \in M^n$, hence we even get a map

$$\mathfrak{S}_n \longrightarrow \{(m, \dots, m)\} \times_{M^n} M^n \times_M \{\mu\} \simeq \{m^n\} \times_M \{m^n\} \simeq \Omega_{m^n} M,$$

again unique up to contractible choice. This finally provides an honestly unique map $\mathfrak{S}_n \rightarrow \pi_0 \Omega_{m^n} M \simeq \pi_1(M, m^n)$. With this, we can formulate the following criterion:

III.12. Proposition (“Cyclic invariance criterion”). — *Let $M \in \mathbf{CMon}(\mathbf{An})$ be an \mathbb{E}_∞ -monoid with an element $s \in M$ such that $\pi_0(M)[s^{-1}] \simeq \pi_0(M)^{\text{grp}}$. Then the following conditions are equivalent:*

- (a) *The map $s: T(M, s) \rightarrow T(M, s)$ is an equivalence.*
- (b) *The fundamental groups of all components of $T(M, s)$ are abelian.*
- (c) *The fundamental groups of all components of $T(M, s)$ are hypoabelian, i.e. have no perfect subgroups except the trivial group $\{e\}$.*
- (d) *The map $\mathfrak{S}_3 \rightarrow \pi_1(M, m^3) \rightarrow \pi_1(T(M, s), m^3)$ kills the permutation $(123) \in \mathfrak{S}_3$ for all $m \in M$.*
- (e) *There is an $n \geq 2$ such that $\mathfrak{S}_n \rightarrow \pi_1(M, m^n) \rightarrow \pi_1(T(M, s), m^n)$ kills the permutation $(12 \dots n) \in \mathfrak{S}_n$ for all $m \in M$.*

In this case, $T(M, s)$ inherits a canonical \mathbb{E}_∞ -monoid structure, and

$$M^{\infty\text{-grp}} \simeq T(M, s).$$

LECTURE 21
21st Jan, 2021

Proof. Let’s prove first that (a) implies the addendum. This will take us a bit longer than in the lecture or in Fabian’s notes, since I’m trying to be super precise when dealing with M -modules and their symmetric monoidal structure, to unconfuse myself. The reason Fabian has to deal with this technical stuff at all is an inconspicuous but nasty problem: It’s not at all clear why $T(M, s)$ is an \mathbb{E}_∞ -monoid again. And it doesn’t help us to know that $\mathbf{CMon}(\mathbf{An})$ has colimits, since the maps $s: M \rightarrow M$ the colimit is taken over are not even maps of \mathbb{E}_∞ -monoids!

Enough of that and let’s get going. From Theorem II.62 we get a symmetric monoidal ∞ -category $\mathcal{M}^{\otimes_M} := \mathbf{LMod}_M(\mathbf{An}^\times)^{\otimes_M}$ whose tensor unit is M and whose tensor product commutes with colimits in either variable. We’ve already seen in III.11a that $T(M, s)$ is canonically an element of \mathcal{M} . We claim that $M \rightarrow T(M, s)$ is \otimes_M -idempotent in the sense of Proposition/Definition III.4. Indeed, since M is the tensor unit and $-\otimes_M T(M, s)$ commutes with colimits, what we need to show is that

$$T(M, s) \xrightarrow{\sim} \operatorname{colim}_{\mathbb{N}} \left(T(M, s) \xrightarrow{s} T(M, s) \xrightarrow{s} T(M, s) \xrightarrow{s} \dots \right)$$

is an equivalence. But that follows from (a). Now the whole package from Proposition/Definition III.4 can be applied. By essentially the same argument as for $T(M, s)$, we see that an element $N \in \mathcal{M}$ absorbs $M \rightarrow T(M, s)$ iff $s: N \xrightarrow{\sim} N$ is an equivalence, i.e., iff N is $\{s\}$ -local. Hence $-\otimes_M T(M, s): \mathcal{M} \rightarrow \mathcal{M}$ is a Bousfield localisation onto the full sub ∞ -category $\mathcal{M}_s \subseteq \mathcal{M}$ of $\{s\}$ -local objects. Moreover, $T(M, s) \in \mathbf{CAlg}(\mathcal{M}_s^{\otimes_M}) \subseteq \mathbf{CAlg}(\mathcal{M}^{\otimes_M})$.

Now we use that

$$\mathbf{CAlg}(\mathcal{M}^{\otimes_M}) \simeq M/\mathbf{CAlg}(\mathbf{An}^\times) \simeq M/\mathbf{CMon}(\mathbf{An})$$

(see [HA, Corollary 3.4.1.7] and the references in the proof of Corollary III.4a). In particular, $T(M, s)$ is an \mathbb{E}_∞ -monoid, hence an \mathbb{E}_∞ -group since $\pi_0(T(M, s)) \simeq \pi_0(M)[s^{-1}] \simeq \pi_0(M)^{\text{grp}}$ is a group by assumption. Hence the map $M \rightarrow T(M, s)$ factors over a map $M^{\infty\text{-grp}} \rightarrow T(M, s)$, which we are to show is an equivalence. By Yoneda’s lemma, it suffices to show that

$$\operatorname{Hom}_{\mathbf{CMon}(\mathbf{An})}(T(M, s), X) \xrightarrow{\sim} \operatorname{Hom}_{\mathbf{CMon}(\mathbf{An})}(M^{\infty\text{-grp}}, X)$$

is an equivalence for all $X \in \mathbf{CGrp}(\mathbf{An})$. This can be done on fibres over $\mathrm{Hom}_{\mathbf{CMon}(\mathbf{An})}(M, X)$. That is, it suffices to show that once we choose a map $M \rightarrow X$, we get

$$\mathrm{Hom}_{M/\mathbf{CMon}(\mathbf{An})}(T(M, s), X) \xrightarrow{\sim} \mathrm{Hom}_{M/\mathbf{CGrp}(\mathbf{An})}(M^{\infty\text{-grp}}, X)$$

The right-hand side is contractible since $\mathrm{Hom}_{\mathbf{CMon}(\mathbf{An})}(M^{\infty\text{-grp}}, X) \simeq \mathrm{Hom}_{\mathbf{CMon}(\mathbf{An})}(M, X)$. Writing $R/\mathbf{CMon}(\mathbf{An}) \simeq \mathbf{CAlg}(\mathcal{M}_M^{\otimes})$, we see that $X \in \mathbf{CAlg}(\mathcal{M}_M^{\otimes})$. But X is clearly $\{s\}$ -local since it is an \mathbb{E}_∞ -group, thus it is already an element of $\mathbf{CAlg}(\mathcal{M}_s^{\otimes})$. Hence Proposition/Definition III.4(c) implies

$$\mathrm{Hom}_{\mathbf{CAlg}(\mathcal{M}_M^{\otimes})}(T(M, s), X) \simeq \mathrm{Hom}_{\mathbf{CAlg}(\mathcal{M}_M^{\otimes})}(M, X),$$

and the right-hand side is contractible as M is initial in $\mathbf{CAlg}(\mathcal{M}_M^{\otimes}) \simeq M/\mathbf{CMon}(\mathbf{An})$. This finishes our rather verbose proof of the first implication.

The addendum implies (b), since the fundamental group of the unit component of any \mathbb{E}_1 -monoid is abelian by the Eckmann–Hilton argument, and all path components of $T(M, s)$ are equivalent since it is even an \mathbb{E}_∞ -group.

The implications (b) \Rightarrow (c), (b) \Rightarrow (d), and (d) \Rightarrow (e) are all trivial. For (c) \Rightarrow (e), note that $\mathfrak{A}_6 \subseteq \mathfrak{S}_6$ is perfect, hence the image of $(12 \dots 6) \in \mathfrak{A}_6$ in the hypoabelian group $\pi_1(T(M, s), m^n)$ must vanish.

Finally, let's show (e) \Rightarrow (a). We've seen in III.11a that $s: T(M, s) \rightarrow T(M, s)$ is an equivalence if $\tau = \mathrm{id}$, which holds if the image of the flip (12) under $\mathfrak{S}_2 \rightarrow \pi_1(M, m^2)$ vanishes. But as argued in III.11c, we really only need that some power of τ vanishes in $T(M, s)$, which replaces the vanishing of (12) under $\mathfrak{S}_2 \rightarrow \pi_1(M, m^2)$ with the vanishing of $(12 \dots n)$ under $\mathfrak{S}_n \rightarrow \pi_1(M, m^n)$. \square

If the assumptions of Proposition III.12 are not satisfied, we can still describe $M^{\infty\text{-grp}}$ in terms of $T(M, s)$. To this end let $\mathbf{An}^{\mathrm{hypo}} \subseteq \mathbf{An}$ be the full sub- ∞ -category of *hypoabelian* anima, i.e. those X for which $\pi_1(X, x)$ has no perfect subgroup except the trivial group $\{e\}$ for every basepoint $x \in X$.

III.13. Proposition (Kervaire, Quillen). — *The inclusion $\mathbf{An}^{\mathrm{hypo}} \subseteq \mathbf{An}$ admits a left adjoint $(-)^+: \mathbf{An} \rightarrow \mathbf{An}^{\mathrm{hypo}}$. The natural unit map $X \rightarrow X^+$ induces an equivalence*

$$\mathbb{S}[X] \xrightarrow{\sim} \mathbb{S}[X^+],$$

hence an isomorphism in homology. Moreover, $(-)^+$ preserves products.

As you undoubtedly have guessed already, $(-)^+$ is called the *Quillen plus construction*, which is a bad name since the construction is actually due to Kervaire.

Proof of Proposition III.13. Throughout the proof we assume without restriction that X is connected (so no basepoints of fundamental groups need to be specified). Since the proof will be a bit lengthy, we divide it into four steps.

Step 1. We construct X^+ in the case where $\pi_1(X)$ is perfect itself.

Pick generators $f_i: \mathbb{S}^1 \rightarrow X$, $i \in I$, of $\pi_1(X)$ and form the pushout

$$\begin{array}{ccc} \coprod_{i \in I} \mathbb{S}^1 & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \overline{X} \end{array}$$

Then an easy excision calculation shows that $H_2(\overline{X}, X, \mathbb{Z}) = \mathbb{Z}^{\oplus I}$ is free. Also note that $H_1(X, \mathbb{Z}) = \pi_1(X)^{\text{ab}} = 0$ since the perfect group $\pi_1(X)$ equals its own commutator.

Hence the long exact homology sequence

$$\dots \longrightarrow H_2(\overline{X}, \mathbb{Z}) \longrightarrow H_2(\overline{X}, X, \mathbb{Z}) \longrightarrow H_1(X, \mathbb{Z}) = 0$$

shows that $H_2(\overline{X}, \mathbb{Z}) \twoheadrightarrow H_2(\overline{X}, X, \mathbb{Z})$ is surjective. It's also easy to check that \overline{X} is simply connected; for example, use Seifert–van Kampen ([Hat02, Theorem 1.20]) plus some fiddling to get its assumptions fulfilled. Hence $\pi_2(\overline{X}) = H_2(\overline{X}, \mathbb{Z})$ by Hurewicz's theorem. Putting everything together, we see that we may pick maps $g_i: \mathbb{S}^2 \rightarrow \overline{X}$, $i \in I$, such that their images in $H_2(\overline{X}, X, \mathbb{Z})$ form a basis. Now put

$$\begin{array}{ccc} \coprod_{i \in I} \mathbb{S}^2 & \longrightarrow & \overline{X} \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & X^+ \end{array}$$

Again, it's easy to check that X^+ is simply connected. Also $X \rightarrow X^+$ is an isomorphism in homology. To see this, one can for example use the same excision calculation as before to compute that $H_i(X^+, \overline{X}, \mathbb{Z})$ equals $\mathbb{Z}^{\oplus I}$ for $i = 3$ and vanishes in all other degrees, and then use the long exact sequence of the triple (X^+, \overline{X}, X) to deduce $H_*(X^+, X, \mathbb{Z}) = 0$.

Step 2. Still in the case where $\pi_1(X)$ is perfect, we use obstruction theory to verify that $\text{Hom}_{\text{An}}(X^+, Z) \xrightarrow{\sim} \text{Hom}_{\text{An}}(X, Z)$ is an equivalence for all hypoabelian anima Z .

We didn't have time for this step in the lecture, so here's my own argument. It suffices to check that

$$\text{Hom}_{\text{An}}(K, \text{Hom}_{\text{An}}(X^+, Z)) \longrightarrow \pi_0 \text{Hom}_{\text{An}}(K, \text{Hom}_{\text{An}}(X, Z))$$

is a bijection for all anima K . We may rewrite the left-hand side as $\pi_0 \text{Hom}_{\text{An}}(X \times K, Z)$ and the right-hand side as $\pi_0 \text{Hom}_{\text{An}}(X^+ \times K, Z)$. By the Künneth theorem and the universal coefficients formulas, we get that $X \times K \rightarrow X^+ \times K$ is an isomorphism in homology and cohomology with arbitrary coefficients. In particular,

$$H^*(X^+ \times K, X \times K, \pi_n(Z, z)) = 0$$

(as long as the homotopy group that occurs is abelian). If Z is simply connected, or at least simple, we can apply [Hat02, Corollary 4.73] (mind that Hatcher calls simple spaces “abelian” instead) to see that $\pi_0 \text{Hom}_{\text{An}}(X^+ \times K, Z) \xrightarrow{\sim} \pi_0 \text{Hom}_{\text{An}}(X \times K, Z)$. But it's also injective: The map

$$(X^+ \times K) \times \{0, 1\} \cup (X \times K) \times \Delta^1 \longrightarrow (X^+ \times K) \times \Delta^1$$

is a homology equivalence again (use Mayer–Vietoris for example), hence we may use the same argument to lift homotopies. This settles the case where Z is simply connected.

For the general case, the idea is to consider the universal covering of Z , but the details get a little technical. First, it suffices that

$$\text{Hom}_{*/\text{An}}((X^+, x), (Z, z)) \xrightarrow{\sim} \text{Hom}_{*/\text{An}}((X, x), (Z, z))$$

is an equivalence for all choices of base points, since equivalences of anima can be checked on (derived) fibres. Since $\text{ev}_x: F(X, Z) \rightarrow F(\{x\}, Z) = Z$ is a Kan fibration, we may use

$F_*((X, x), (Z, z)) = F(X, Z) \times_Z \{z\}$ and similarly $F_*((X^+, x), (Z, z)) = F(X^+, Z) \times_Z \{z\}$, with pullbacks taken in \mathbf{sSet} , as explicit simplicial models for the Hom anima in $*/\mathbf{An}$. Moreover, we may assume that $Z = \text{Sing } Y$ for some actual CW complex Y . In this situation, we claim that there is an actual equality of simplicial sets

$$F_*((X, x), \text{Sing}(Y, y)) = F_*((X, x), \text{Sing}(\tilde{Y}, \tilde{y})),$$

where $(\tilde{Y}, \tilde{y}) \rightarrow (Y, y)$ is the universal covering. Together with the analogous assertion for X^+ , this will seal the deal since we already know that $\text{Hom}_{*/\mathbf{An}}((X, x), \text{Sing}(\tilde{Y}, \tilde{y}))$ and $\text{Hom}_{*/\mathbf{An}}((X^+, x), \text{Sing}(\tilde{Y}, \tilde{y}))$ are equivalent from the simply connected case.

To see the claim, we unwind that the set of n -simplices $F_*((X, x), \text{Sing}(Y, y))_n$ is the set of all maps $X \times \Delta^n \rightarrow \text{Sing } Y$ that send $\{x\} \times \Delta^n$ to the basepoint $\{y\}$, or by adjunction, the set of all continuous maps $f: |X| \times |\Delta^n| \rightarrow Y$ with the same basepoint property. But since $\pi_1(|X| \times |\Delta^n|) = \pi_1(X)$ is perfect, whereas $\pi_1(Y, y)$ is hypoabelian, any such continuous map f induces the trivial map $f_* = \text{const } 1: \pi_1(|X| \times |\Delta^n|) \rightarrow \pi_1(Y, y)$ on fundamental groups. Hence, by covering theory (see [Hat02, Propositions 1.33 and 1.34] for example), f has a unique lift

$$\begin{array}{ccc} & \exists! \tilde{f} & \nearrow \\ & & \tilde{Y} \\ |X| \times |\Delta^n| & \xrightarrow{f} & Y \end{array}$$

such that \tilde{f} sends $\{x\} \times |\Delta^n|$ to $\{\tilde{y}\}$. And so we are done.

Step 3. We construct X^+ in general and show that it satisfies the required universal property.

Let $P \subseteq \pi_1(X)$ be the largest perfect subgroup (just consider the subgroup generated by all perfect subgroup and check it's perfect). Let $\tilde{X} \rightarrow X$ be the covering space associated to P ([Hat02, Theorem 1.38]). Note that $P \subseteq \pi_1(X)$ is normal (since its conjugates are also perfect), hence $\pi_1(\tilde{X}) = P$ by more covering theory. This means that we know how to construct \tilde{X}^+ and can put

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & \tilde{X}^+ \\ \downarrow & \lrcorner & \downarrow \\ X & \longrightarrow & X^+ \end{array}$$

in general. Since \tilde{X}^+ is simply connected by construction, a Seifert–van Kampen argument again shows $\pi_1(X^+) = \pi_1(X)/P$, which is now hypoabelian by construction. Moreover,

$$\text{Hom}_{\mathbf{An}}(X^+, Z) \simeq \text{Hom}_{\mathbf{An}}(X, Z) \times_{\text{Hom}_{\mathbf{An}}(\tilde{X}, Z)} \text{Hom}_{\mathbf{An}}(\tilde{X}^+, Z) \simeq \text{Hom}_{\mathbf{An}}(X, Z)$$

for all hypoabelian anima Z , since we already know $\text{Hom}_{\mathbf{An}}(\tilde{X}, Z) \simeq \text{Hom}_{\mathbf{An}}(\tilde{X}^+, Z)$.

Step 4. We verify that $\mathbb{S}[X] \xrightarrow{\sim} \mathbb{S}[X^+]$ is an equivalence and that $(-)^+$ commutes with finite products.

Both assertions are completely formal. We have $\text{Hom}_{\mathbf{Sp}}(\mathbb{S}[X], E) \simeq \text{Hom}_{\mathbf{An}}(X, \Omega^\infty E)$ and $\text{Hom}_{\mathbf{Sp}}(\mathbb{S}[X^+], E) \simeq \text{Hom}_{\mathbf{An}}(X^+, \Omega^\infty E)$. But the right-hand sides coincide because $\Omega^\infty E$ is hypoabelian. In fact, it is an \mathbb{E}_∞ -group and thus all its path components are equivalent and have abelian fundamental group. Thus $\mathbb{S}[X] \simeq \mathbb{S}[X^+]$ by Yoneda.

For the second assertion, $*^+ \simeq *$ holds for trivial reasons. To see $(X \times Y)^+ \simeq X^+ \times Y^+$, first note that the right-hand side is hypoabelian since $\pi_1: \mathbf{An} \rightarrow \mathbf{Grp}$ commutes with

products. Hence it suffices to show that the composite $X \times Y \rightarrow X^+ \times Y \rightarrow X^+ \times Y^+$ induces equivalences after taking $\mathrm{Hom}_{\mathrm{An}}(-, Z)$ for Z hypoabelian. But

$$\begin{aligned} \mathrm{Hom}_{\mathrm{An}}(X^+ \times Y, Z) &\simeq \mathrm{Hom}_{\mathrm{An}}(Y, \mathrm{Hom}_{\mathrm{An}}(X^+, Z)) \\ &\simeq \mathrm{Hom}_{\mathrm{An}}(Y, \mathrm{Hom}_{\mathrm{An}}(X, Z)) \\ &\simeq \mathrm{Hom}_{\mathrm{An}}(X \times Y, Z) \end{aligned}$$

if Z is hypoabelian, hence $X^+ \times Y \rightarrow X \times Y$ induces an equivalence after taking $\mathrm{Hom}_{\mathrm{An}}(-, Z)$, and for $X^+ \times Y \rightarrow X^+ \times Y^+$ we can use the same argument again. \square

III.14. Proposition. — *If $M \in \mathrm{CMon}(\mathrm{An})$ has an element $s \in \pi_0(M)$ such that $\pi_0(M)[s^{-1}] = \pi_0(M)^{\mathrm{grp}}$, then*

$$T(M, s)^+ \simeq T(M^+, s) \simeq M^{\infty\text{-grp}}.$$

Proof. First of all, Proposition III.12(b) and (e) imply that $T(M^+, s)$ has abelian fundamental group: Indeed, it suffices to check that the permutation $(12\dots 6) \in \mathfrak{S}_6$ gets killed under the map $\mathfrak{S}_6 \rightarrow \pi_1(T(M^+, s), m^n)$, but this map factors through the hypoabelian group $\pi_1(M^+, m^n)$, in which the perfect subgroup $\mathfrak{A}_6 \subseteq \mathfrak{S}_6$ containing $(12\dots 6)$ already dies. Thus, to show $T(M, s)^+ \simeq T(M^+, s)$, it suffices to check that the right-hand side satisfies the universal property of the Quillen plus construction:

$$\begin{aligned} \mathrm{Hom}_{\mathrm{An}}(T(M^+, s), Z) &\simeq \lim_{\mathrm{Nop}} \left(\dots \xrightarrow{s^*} \mathrm{Hom}_{\mathrm{An}}(M^+, Z) \xrightarrow{s^*} \mathrm{Hom}_{\mathrm{An}}(M^+, Z) \right) \\ &\simeq \lim_{\mathrm{Nop}} \left(\dots \xrightarrow{s^*} \mathrm{Hom}_{\mathrm{An}}(M, Z) \xrightarrow{s^*} \mathrm{Hom}_{\mathrm{An}}(M, Z) \right) \\ &\simeq \mathrm{Hom}_{\mathrm{An}}(T(M, s)^+, Z) \end{aligned}$$

holds for all $Z \in \mathrm{An}^{\mathrm{hypo}}$, so indeed $T(M, s)^+ \simeq T(M^+, s)$. Now consider the square

$$\begin{array}{ccc} T(M, s)^+ & \longrightarrow & M^{\infty\text{-grp}} \\ \downarrow \wr & & \downarrow \wr \\ T(M^+, s) & \xrightarrow{\sim} & (M^+)^{\infty\text{-grp}} \end{array}$$

The left vertical arrow is an equivalence as we just showed, and the bottom arrow is an equivalence by Proposition III.12. The right vertical arrow is an isomorphism on $\mathbb{S}[-]$, using $\mathbb{S}[M] \simeq \mathbb{S}[M^+]$ by Proposition III.13 and Theorem III.6, hence an isomorphism on homology by Corollary* II.55b, hence an equivalence by Whitehead's theorem, since both spaces are simple (because any H -space is; see [Hat02, Example 4A.3], but mind that Hatcher uses “abelian” instead of “simple”). \square

We obtain Quillen's first definition of algebraic K -theory.

III.14a. Corollary. — *For any ring R ,*

$$k(R) = K_0(R) \times \mathrm{BGL}_{\infty}(R)^+.$$

Proof. By Proposition III.14 and the arguments from the beginning of the proof of Corollary III.7, we have

$$k(R) \simeq \mathrm{Proj}(R)^{\infty\text{-grp}} \simeq \mathrm{colim}_{\mathbb{N}} \left(\mathrm{Proj}(R) \xrightarrow{[R]} \mathrm{Proj}(R) \xrightarrow{[R]} \dots \right)^+$$

Isolating the 0-component on both sides gives $k_0(R) \simeq (\mathrm{colim}_{n \in \mathbb{N}} BGL_n(R))^+ \simeq BGL_{\infty}(R)^+$. Now $k(R) \simeq \pi_0 k(R) \times k(R)_0 \simeq K_0(R) \times BGL_{\infty}(R)^+$ as all components in an \mathbb{E}_{∞} -group are equivalent. \square

Topological K -Theory

Our next goal is to get some understanding of ku and the Adams operations on it, which will play a prominent role in Quillen's comparison with the K -theory of finite fields. The first thing to do is to get another description of ku as well as the other \mathbb{E}_{∞} -groups ko , $k\mathrm{top}$, and $k\mathrm{sph}$ from Definition II.14a.

III.15. Corollary. — For $V \in \{\mathrm{O}, \mathrm{U}, \mathrm{Top}, \mathcal{G}\}$ put $BV := \mathrm{colim}_{n \in \mathbb{N}} BV(n)$, where in the last two cases we define $\mathrm{Top}(n) := \mathrm{Aut}_{\mathrm{Top}}(\mathbb{R}^n)$, equipped with the compact-open topology, and $\mathcal{G}(n) := \mathrm{Aut}_{*/\mathrm{An}}(\mathbb{S}^n, *)$. Then

$$ko \simeq \mathbb{Z} \times BO, \quad ku \simeq \mathbb{Z} \times BU, \quad k\mathrm{top} \simeq \mathbb{Z} \times B\mathrm{Top}, \quad \text{and} \quad k\mathrm{sph} \simeq \mathbb{Z} \times B\mathcal{G}.$$

Moreover if X is a finitely dominated CW complex, i.e. a retract of a finite one, then the comparison maps from II.14 are equivalences

$$\begin{aligned} \mathrm{Vect}_{\mathbb{R}}(X)^{\infty\text{-grp}} &\xrightarrow{\sim} \mathrm{Hom}_{\mathrm{An}}(X, ko), & \mathcal{E}\mathrm{ucl}_{\mathbb{R}}(X)^{\infty\text{-grp}} &\longrightarrow \mathrm{Hom}_{\mathrm{An}}(X, k\mathrm{top}), \\ \mathrm{Vect}_{\mathbb{C}}(X)^{\infty\text{-grp}} &\xrightarrow{\sim} \mathrm{Hom}_{\mathrm{An}}(X, ku), & \mathrm{Sph}_{\mathbb{R}}(X)^{\infty\text{-grp}} &\xrightarrow{\sim} \mathrm{Hom}_{\mathrm{An}}(X, k\mathrm{sph}). \end{aligned}$$

In particular, $ko^0(X) = \pi_0 \mathrm{Hom}(X, ko)$ and $ku^0(X) = \pi_0 \mathrm{Hom}_{\mathrm{An}}(X, ku)$ are the group completions of the (ordinary) monoids of isomorphism classes of real and complex vector bundles on X , respectively.

Proof. We'll only do the cases of ko and ku , but Fabian's script [A&HK, Chapter III pp. 21–24] also has interesting things to say about $k\mathrm{sph}$. First recall Lemma* II.14b, which says

$$\mathrm{Vect}_{\mathbb{R}} \simeq \coprod_{n \geq 0} BO(n) \quad \text{and} \quad \mathrm{Vect}_{\mathbb{C}} \simeq \coprod_{n \geq 0} BU(n).$$

Hence $\pi_0 \mathrm{Vect}_{\mathbb{R}} = \mathbb{N}$ and $\pi_0 \mathrm{Vect}_{\mathbb{C}} = \mathbb{N}$, with generators $[\mathbb{R}]$ and $[\mathbb{C}]$ respectively, and we find

$$T(\mathrm{Vect}_{\mathbb{R}}, [\mathbb{R}]) \simeq \mathbb{Z} \times BO \quad \text{and} \quad T(\mathrm{Vect}_{\mathbb{C}}, [\mathbb{C}]) \simeq \mathbb{Z} \times BU.$$

Since B shift homotopy groups up, we get $\pi_1(BO) \simeq \pi_0(\mathrm{O}) \simeq \{\pm 1\}$ (via the determinant) and likewise $\pi_1(BU) \simeq \pi_0(\mathrm{U}) \simeq \{\pm 1\}$. These groups are abelian, thus Proposition III.12(b) shows that $\mathbb{Z} \times BO$ and $\mathbb{Z} \times BU$ are already the \mathbb{E}_{∞} -group completions of $\mathrm{Vect}_{\mathbb{R}}$ and $\mathrm{Vect}_{\mathbb{C}}$. That is, they coincide with ko and ku , as claimed.

For the assertion about the comparison maps, recall $\mathrm{Vect}_{\mathbb{R}}(X) \simeq \mathrm{Hom}_{\mathrm{An}}(X, \mathrm{Vect}_{\mathbb{R}})$ (see II.14). If X is a finite anima, then $\mathrm{Hom}_{\mathrm{An}}(X, -)$ commutes with sequential (and even filtered) colimits. This clearly still holds if X is only finitely dominated. Hence

$$\mathrm{Hom}_{\mathrm{An}}(X, ko) \simeq \mathrm{Hom}_{\mathrm{An}}(X, T(\mathrm{Vect}_{\mathbb{R}}, [\mathbb{R}])) \simeq T(\mathrm{Vect}_{\mathbb{R}}(X), [X \times \mathbb{R}]).$$

Since ko is an \mathbb{E}_∞ -group, so is the left-hand side, hence so is the right-hand side. Now use the fact that every vector bundle on X is a direct summand of a trivial bundle to see that $s = [X \times \mathbb{R}]$ satisfies the condition from Proposition III.12, and then that proposition shows $T(\text{Vect}_\mathbb{R}(X), [X \times \mathbb{R}]) \simeq \text{Vect}_\mathbb{R}(X)^{\infty\text{-grp}}$. For the additional assertion, recall

$$\pi_0 \text{Hom}_{\text{An}}(X, \text{Vect}_\mathbb{R}) \simeq \pi_0 \text{Vect}_\mathbb{R}(X)$$

(the $\text{Vect}_\mathbb{R}$ on the right-hand side has no curly “ \mathcal{V} ” and denotes the groupoid of real vector bundles on X) by the classification of principal bundles and Lemma* II.14c. Hence

$$ko^0(X) = \pi_0 \text{Hom}_{\text{An}}(X, ko) = \pi_0(\text{Hom}_{\text{An}}(X, \text{Vect}_\mathbb{R})^{\infty\text{-grp}}) = (\pi_0 \text{Vect}_\mathbb{R}(X))^{\text{grp}},$$

as claimed. The complex case works completely analogous. \square

Recall that the tensor product $- \otimes_R -$ on $\text{Proj}(R)$ turns $B^\infty k(R)$ into an \mathbb{E}_∞ -ring spectrum, as explained in II.54(c*). The same argument essentially works again for the real tensor product $- \otimes_\mathbb{R} -$ on $\text{Vect}_\mathbb{R}$ and the complex tensor product $- \otimes_\mathbb{C} -$ on $\text{Vect}_\mathbb{C}$, up to some check that they are compatible with the Kan enrichment (as defined in II.14) to ensure that the coherent nerves $\text{Vect}_\mathbb{R}$ and $\text{Vect}_\mathbb{C}$ are indeed elements of $\text{CAlg}(\text{CMon}(\text{An})^\otimes)$. Up to this technicality, we see that $B^\infty ko$ and $B^\infty ku$ are \mathbb{E}_∞ -ring spectra! Also note that this doesn’t work for $k\text{top}$ or $k\text{sph}$.

Using Corollary* II.55b, we now recognize

$$ko^*(X) = \pi_{-*} \text{Hom}_{\text{An}}(X, ko) = \pi_{-*} \text{hom}_{\text{Sp}}(\mathbb{S}[X], B^\infty ko) = (B^\infty ko)^*(X)$$

as the negative part of the cohomology theory associated to $B^\infty ko$. The same works for $ku^*(X)$. It might seem weird at first that $ko^*(-)$ and $ku^*(-)$ are only negative parts of cohomology theories, but then again it’s not so weird anymore if you know a bit of topological K -theory: A priori, it can only be constructed in degrees ≤ 0 . To extend it to positive degrees, one needs *Bott periodicity*—which we’ll discuss next.

Consider the *Hopf bundles* (see [Hat02, Examples 4.45–4.47] for a construction)

$$(\gamma_1^\mathbb{R} \rightarrow \mathbb{RP}^1) \in \text{Vect}_\mathbb{R}^1(\mathbb{S}^1), \quad (\gamma_1^\mathbb{H} \rightarrow \mathbb{HP}^1) \in \text{Vect}_\mathbb{R}(\mathbb{S}^4), \quad \text{and} \quad (\gamma_1^\mathbb{O} \rightarrow \mathbb{OP}^1) \in \text{Vect}_\mathbb{R}^8(\mathbb{S}^8)$$

(where \mathbb{H} denotes the quaternions and \mathbb{O} the octonions) and also

$$(\gamma_1^\mathbb{C} \rightarrow \mathbb{CP}^1) \in \text{Vect}_\mathbb{C}^1(\mathbb{S}^2)$$

(which is 1-dimensional as a complex bundle, hence the $\text{Vect}_\mathbb{C}^1$ is not a typo). As mentioned in Corollary III.15, we have $\pi_0 \text{Vect}_\mathbb{R}(\mathbb{S}^n) = \pi_0 \text{Hom}_{\text{An}}(\mathbb{S}^n, ko)$. Hence every vector bundle $\xi \rightarrow \mathbb{S}^n$ defines a homotopy class of maps $\xi: \mathbb{S}^n \rightarrow ko$. If the vector bundle in question is d -dimensional, the map ξ has image in $\{d\} \times BO \subseteq \mathbb{Z} \times BO \simeq ko$. Similar to our considerations in III.11a, there is a map $+\mathbb{R}: ko \rightarrow ko$ (induced by the one we took the colimit over to construct $ko \simeq T(\text{Vect}_\mathbb{R}, [\mathbb{R}])$ in the first place) which sends $\{i\} \times BO \rightarrow \{i+1\} \times BO$. If $-\mathbb{R}^d$ denotes the d -fold iteration of its inverse, then we can postcompose ξ with $-\mathbb{R}^d$ to obtain a map $\xi - \mathbb{R}^d: \mathbb{S}^n \rightarrow ko$, which lands in the 0-component $ko_0 \simeq \{0\} \times BO$, hence defines an element of $\pi_n(ko_0)$.

Applying this construction to the Hopf bundles $\gamma_1^\mathbb{R}$, $\gamma_1^\mathbb{H}$, and $\gamma_1^\mathbb{O}$, and its complex analogue to the Hopf bundle $\gamma_1^\mathbb{C}$, we obtain elements

$$\begin{aligned} \eta &= \gamma_1^\mathbb{R} - [\mathbb{R}] \in \pi_1(ko_0), & \beta &= \gamma_1^\mathbb{C} - [\mathbb{C}] \in \pi_1(ku_0), \\ \sigma &= \gamma_1^\mathbb{H} - [\mathbb{R}^4] \in \pi_4(ko_0), & \nu &= \gamma_1^\mathbb{O} - [\mathbb{R}^4] \in \pi_8(ko_0). \end{aligned}$$

Since $B^\infty ko$ and $B^\infty ku$ are \mathbb{E}_∞ -ring spectra, we see that $\pi_*(ko_0) = \pi_*(B^\infty ko)$ and $\pi_*(ku_0) = \pi_*(B^\infty ku)$ are graded commutative rings by II.64 (and more generally, the same is true for $ko^*(X)$ and $ku^*(X)$ using the cup product from II.65). In fact, we'll see in a second that they are even honest commutative, since the odd degrees vanish or have characteristic 2.

III.16. Theorem (Bott periodicity). — *The elements $\beta \in \pi_*(ku_0)$ and $\eta, \sigma, \nu \in \pi_*(ko_0)$ define isomorphisms*

$$\mathbb{Z}[\beta] \xrightarrow{\sim} \pi_*(ku_0) \quad \text{and} \quad \mathbb{Z}[\eta, \sigma, \nu]/(2\eta, \eta^3, \eta\sigma, \sigma^2 - 4\nu) \xrightarrow{\sim} \pi_*(ko_0).$$

In particular, we can refine the description from II.14 as follows:

$$\pi_n(ku) \simeq \begin{cases} \mathbb{Z}\langle\beta^i\rangle & \text{if } n = 2i \\ 0 & \text{if } n = 2i + 1 \end{cases} \quad \text{and} \quad \pi_n(ko) \simeq \begin{cases} \mathbb{Z}\langle\nu^i\rangle & \text{if } n = 8i \\ \mathbb{Z}/2\langle\eta\nu^i\rangle & \text{if } n = 8i + 1 \\ \mathbb{Z}/2\langle\eta^2\nu^i\rangle & \text{if } n = 8i + 2 \\ 0 & \text{if } n = 8i + 3 \\ \mathbb{Z}\langle\sigma\nu^i\rangle & \text{if } n = 8i + 4 \\ 0 & \text{if } n = 8i + 5 \\ 0 & \text{if } n = 8i + 6 \\ 0 & \text{if } n = 8i + 7 \end{cases}.$$

We won't prove Theorem III.16. If you want to read up on this, you'll find proofs in any book on topological K -theory but Fabian particularly recommends Bott's original proof in [Bot59].

As a consequence, we can now define spectra KU and KO with $\Omega^{\infty-2i}KU = ku$ and $\Omega^{\infty-8i}KO \simeq ko$ for all $i \in \mathbb{Z}$, which are thus 2-periodic and 8-periodic, respectively. Another description is

$$KU \simeq (B^\infty ku)[\beta^{-1}] \quad \text{and} \quad KO \simeq (B^\infty ko)[\nu^{-1}]$$

(in Proposition III.2, we only saw how to localise elements from π_0 , but the construction is straightforward to generalise to elements from arbitrary degrees). In this formulation we immediately see $KO, KU \in \text{CAlg}$. The associated cohomology theories were studied by Atiyah and Hirzebruch, and are known as (*periodic*) *topological K -theory*.

III.17. Adams Operations on K -Theory. — The Adams operations are certain maps $\psi^i: ku \rightarrow ku$. We'll construct them in two steps:

Step 1. We'll massage the exterior power maps $\Lambda^i: \text{Vect}_{\mathbb{C}}(X) \rightarrow \text{Vect}_{\mathbb{C}}(X)$ on complex vector bundles long enough to turn them into maps $\lambda^i: ku \rightarrow ku$ of anima (not respecting any \mathbb{E}_∞ -structures though).

Let's suppose X is finitely dominated at first. Then $ku^0(X) \simeq (\pi_0 \text{Vect}_{\mathbb{C}}(X))^{\text{grp}}$ by Corollary III.15. The main obstacle later on will be to bypass the finiteness assumptions on X , but right now, there's another problem: We would like to extend

$$\Lambda^i: \pi_0 \text{Vect}_{\mathbb{C}}(X) \longrightarrow ku^0(X)$$

to all of $ku^0(X)$. However, knowing that $ku^0(X) \simeq (\pi_0 \text{Vect}_{\mathbb{C}}(X))^{\text{grp}}$ doesn't help us as Λ^i is not even a map of monoids! This can be fixed as follows: Instead of considering each Λ^i on its own, we combine them into a single map

$$\begin{aligned} \Lambda: \pi_0 \text{Vect}_{\mathbb{C}}(X) &\longrightarrow ku^0(X)[[t]] \\ [V] &\longmapsto \sum_{i \geq 0} [\Lambda^i V] t^i, \end{aligned}$$

where $ku^0(X)[[t]]$ denotes the ring of power series over $ku^0(X)$, which is a ring itself as denoted before Theorem III.16. From the formula

$$\Lambda^n(V \oplus W) = \bigoplus_{i+j=n} \Lambda^i(V) \otimes \Lambda^j(W),$$

we get that Λ is a homomorphism onto the subgroup $1 + t \cdot ku^0(X)[[t]] \subseteq ku^0(X)[[t]]^\times$ of the group of units of the ring $ku^0(X)[[t]]$. In the case where X is finitely dominated, hence $ku^0(X) \simeq (\pi_0 \text{Vect}_{\mathbb{C}}(X))^{\text{grp}}$, we may thus extend Λ to a map

$$\Lambda: ku^0(X) \longrightarrow ku^0(X)[[t]],$$

which is a group homomorphism into $ku^0(X)[[t]]^\times$. Writing $\Lambda = \sum_{i \geq 0} t^i \Lambda^i$, we obtain natural maps $\Lambda^i: ku^0(X) \rightarrow ku^0(X)$ for all finitely dominated X (but note that the Λ^i neither preserve the additive, nor the multiplicative structure).

We would like to extend the Λ^i to natural maps $\Lambda^i: ku^0(X) \rightarrow ku^0(X)$ for *all* anima X , i.e., to a natural transformation from the functor $ku^0(-): (\pi \text{An})^{\text{op}} \rightarrow \text{Set}$ to itself. By definition, we have

$$ku^0(X) = \pi_0 \text{Hom}_{\text{An}}(X, ku) = \text{Hom}_{\pi \text{An}}(X, ku),$$

hence $ku^0(-) = \text{Hom}_{\pi \text{An}}(-, ku)$ is represented by ku . By the 1-categorical Yoneda lemma, any endotransformation of $ku^0(-)$ thus corresponds to a map $ku \rightarrow ku$ in πAn , or put differently, to an element of $ku^0(ku)$. The idea how to construct the element $\lambda^i \in ku^0(ku)$ we're looking for is simple: We "approximate" ku by suitable finite anima, on which we already know what Λ^i does, and show that all finite stages together combine into a unique element of $ku^0(ku)$. Let's give a sketch of how to make this precise!

First off, we need to compute $ku^0(ku)$. Since $\Omega^\infty KU \simeq ku$, we may equivalently compute $KU^0(ku)$. In the lecture we worked with ku^* right away, but I think this might get us into trouble (please don't take my word for it, I'm way out of my mathematical comfort zone here). Now KU is an example of a *multiplicative complex oriented cohomology theory*. These are \mathbb{E}_∞ -ring spectra E for which the natural inclusion $\mathbb{S}^2 \simeq \mathbb{CP}^1 \subseteq \mathbb{CP}^\infty$ induces a surjection $E^2(\mathbb{CP}^\infty) \twoheadrightarrow E^2(\mathbb{S}^2)$. This is by the way the reason why I don't think we could take ku^* instead of KU^* : The former is not a cohomology theory, only the negative part of one (and we can't just extend it by 0 to positive degrees since that would violate the suspension isomorphism). A general fact about multiplicative complex oriented cohomology theories is the formula

$$E^*(BU(n)) \simeq \pi_*(E)[[c_1, \dots, c_n]],$$

where the $c_i \in E^{2i}(BU(n))$ are the *Chern classes* in E -cohomology. You can find a proof in [Lur10, Lecture 4]. If you are confused like me about how a power series ring is supposed to

be graded: It appears to be a convention to regard graded cohomology rings as products $E^*(X) = \prod_{n \in \mathbb{Z}} E^n(\mathbb{Z})$ rather than direct sums $\bigoplus_{n \in \mathbb{N}} E^n(X)$.

Applying the above formula to $E \simeq KU$, which has $\pi_*(KU) = \mathbb{Z}[\beta, \beta^{-1}]$, where β is the Bott element from Theorem III.16, and using $ku \simeq \mathbb{Z} \times BU$, we find

$$ku^0(ku) = KU^0(ku) = \prod_{n \in \mathbb{Z}} \mathbb{Z} [\beta^{-i} c_i \mid i \geq 1] .$$

Now that we have computed $ku^0(ku)$, we enter phase 2 of our plan and try to “approximate” ku by finite anima. Let $\mathrm{Gr}_k(\mathbb{C}^m)$ be the *Grassmannian* parametrising k -dimensional sub- \mathbb{C} -vector spaces of \mathbb{C}^m , and let $\gamma_{k,m}^{\mathbb{C}} \rightarrow \mathrm{Gr}_k(\mathbb{C}^m)$ be the tautological bundle. By the classification of vector bundles, it defines a homotopy class of maps $\mathrm{Gr}_k(\mathbb{C}^m) \rightarrow BU(k)$. Furthermore, we have a maps $BU(k) \rightarrow \{n\} \times BU \subseteq ku$ for all $n \in \mathbb{Z}$. Letting k, m , and n vary thus provides a single map

$$ku^0(ku) \longrightarrow \prod_{n \in \mathbb{Z}} \prod_{0 \leq k \leq m} ku^0(\mathrm{Gr}_k(\mathbb{C}^m)) .$$

This map is injective, which follows essentially from the formula for $ku^0(ku)$ above and from the definition of Chern classes. Now the Grassmannians $\mathrm{Gr}_k(\mathbb{C}^m)$ are finite CW complexes, hence we already know what $\Lambda^i: ku^0(\mathrm{Gr}_k(\mathbb{C}^m)) \rightarrow ku^0(\mathrm{Gr}_k(\mathbb{C}^m))$ is supposed to do. Using the injectivity above, one checks that the $[\Lambda^i \gamma_{k,n}^{\mathbb{C}}]$ combine into a unique element $\lambda^i \in ku^0(ku)$, which is what we’re looking for.

The upshot is that we get natural maps $\lambda^i: ku \rightarrow ku$ (these are maps of anima, but don’t preserve any \mathbb{E}_∞ -structures). For any anima X , they induce maps $\lambda^i: ku^0(X) \rightarrow ku^0(X)$ which satisfy

$$\lambda^n(x + y) = \sum_{i+j=n} \lambda^i(x) \cdot \lambda^j(y) \in ku^0(X)$$

for all $x, y \in ku^0(X)$. We’ve thereby reached our goal of Step 1.

Step 2. We show how one can extract ring homomorphisms $\psi^i: ku^0(X) \rightarrow ku^0(X)$ from the λ^i by a purely algebraic procedure.

The buzzword here is that $\{\lambda^i \mid i \geq 0\}$ turn $ku^0(X)$ into a λ -ring. For a precise definition, consult the [nLab](#); in a nutshell, it means one has a ring R together with maps $\lambda^i: R \rightarrow R$ of sets that satisfy the relation above, as well as a bunch of other relations for $\lambda^n(xy)$ and $\lambda^n(\lambda^m(x))$, which are all satisfied for exterior powers of vector bundles, and thus also in our case. And now extracting the ψ^i from the λ^i is a purely algebraic procedure in that it can be done for any λ -ring.

Just as the Λ^i in Step 1 behave nicer once combined into a single power series, the same happens for the λ^i . For better normalisation behaviour later one, we put

$$\tilde{\lambda}(x) := \sum_{i \geq 0} (-1)^i \lambda^i(x) t^i \in ku^0(X)[[t]]$$

this time. As for Λ , one checks that $\tilde{\lambda}: ku^0(X) \rightarrow 1 + t \cdot ku^0(X)[[t]] \subseteq ku^0(X)[[t]]^\times$ is a group homomorphism into the unit group of $ku^0(X)[[t]]$. To get a homomorphism into the additive group of $ku^0(X)[[t]]$ instead, we would like to take “logarithms”. To this end, let

$$\ell(t) := \sum_{i \geq 1} \frac{(-1)^{i-1} t^i}{i} \in \mathbb{Q}[[t]]$$

be the formal power series of the Taylor expansion of $\log(1+x): (-1,1) \rightarrow \mathbb{R}$. Then the coefficients $\varphi^i(x)$ of $\ell(\tilde{\lambda}(x) - 1) \in (ku^0(X) \otimes \mathbb{Q})[[t]]$ are additive functions of x (note that $\tilde{\lambda} - 1$ has constant coefficient 0, so we may plug it into any other formal power series). To get integer coefficients instead of rational ones, we take formal derivatives (which preserves additivity of the coefficients), and therefore we arrive at the following definition:

III.17a. Definition. — For $i \geq 1$, the i^{th} Adams operation $\psi^i: ku^0(X) \rightarrow ku^0(X)$ is defined by

$$-\ell(\tilde{\lambda}(x) - 1)' = \sum_{i \geq 0} \psi^{i+1}(x)t^i \in ku^0(X)[[t]],$$

and for $i = 0$ we put $\psi^0(x) = \text{rk}(x) \cdot [C] \in ku^0(X)$, where $\text{rk}: ku \simeq \mathbb{Z} \times BU \rightarrow \mathbb{Z}$ is the projection to path components.

LECTURE 22
26th Jan, 2021

For later use, we record that the derivative in Definition III.17a evaluates to

$$\ell(\tilde{\lambda}(x) - 1)' = -\frac{\tilde{\lambda}(x)'}{\tilde{\lambda}(x)} = -\sum_{i \geq 0} (1 - \tilde{\lambda}(x))^i \cdot \tilde{\lambda}(x)'.$$

III.18. Proposition. — For all $X \in \text{An}$ we have:

- (a) For all $i \geq 0$, $\psi^i: ku^0(X) \rightarrow ku^0(X)$ is a ring homomorphism.
- (b) For all $i, j \geq 0$, $\psi^i \circ \psi^j = \psi^{ij}$.
- (c) If $L \in \text{Vect}_{\mathbb{C}}^1(X)$ is a line bundle on X , then $\psi^i([L]) = [L]^i = [L^{\otimes i}]$ for all $i \geq 0$.
- (d) If p is a prime, then $\psi^p(x) \equiv x^p \pmod{p}$. That is, the ψ^p are Frobenius lifts!
- (e) For all $i, n \geq 0$, $\psi^i(\beta^n) = i^n \beta^n \in ku^0(\mathbb{S}^{2n}, *)$.

We should perhaps explain the notation from Proposition III.18(e), as well as why β^n is an element of $ku^0(\mathbb{S}^{2n}, *)$. The relative ku -group $ku^0(\mathbb{S}^{2n}, *)$ is defined as

$$\begin{aligned} ku^0(\mathbb{S}^{2n}, *) &:= \ker(ku^0(\mathbb{S}^{2n}) \rightarrow ku^0(*)) \\ &= \ker(\pi_0 \text{Hom}_{\text{An}}(\mathbb{S}^{2n}, ku) \rightarrow \pi_0 \text{Hom}_{\text{An}}(*, ku)). \end{aligned}$$

Using $\pi_1(\text{Hom}_{\text{An}}(*, ku), 0) = \pi_1(ku_0) = 0$ by Theorem III.16, we see that $ku^0(\mathbb{S}^{2n}, *)$ equals $\pi_{2n}(ku_0) \simeq \mathbb{Z}\langle \beta^n \rangle$. Hence (e) makes indeed sense.

Combining Proposition III.18(a), (b), and (d), we see that the $\{\psi^i \mid i \geq 0\}$ are uniquely determined by the ψ^p for p a prime, and these guys are commuting Frobenius lifts. This holds in fact for all λ -rings! Moreover, it is a theorem of Wilkerson that conversely any choice of commuting Frobenius lifts on a \mathbb{Z} -torsion free ring defines a unique λ -structure. This makes λ -rings interesting to number theorists; in particular, to people working in finite or mixed characteristic—and especially, I'd like to add, to people working in “characteristic 1”.

Proof sketch of Proposition III.18. Parts (a), (b), and (d) work for arbitrary λ -rings and can thus be proved by horrible manipulations of power series (for (a) and (b) you'll need some additional properties of the λ^i that we didn't specify). We'll immediately see that in the case of K -theory, there's a dirty shortcut, but before we do that, let me prove (d) in general because it isn't so horrible at all and I like Frobenii.

By Definition III.17a, $\psi^p(x)$ is the coefficient of t^{p-1} in $\ell(\tilde{\lambda}(x) - 1)'$. In our formula above, only those summands $(1 - \tilde{\lambda}(x))^i \cdot \tilde{\lambda}(x)'$ with $i \leq p - 1$ will contribute to the coefficient of t^{p-1} . Using this together with

$$(1 - \tilde{\lambda}(x))^p \equiv 1 - \tilde{\lambda}(x)^p \pmod{p},$$

we get that $\psi^p(x)$ modulo p is the coefficient of t^{p-1} in

$$\begin{aligned} - \sum_{i=0}^{p-1} (1 - \tilde{\lambda}(x))^i \cdot \tilde{\lambda}(x)' &\equiv - \frac{1 - (1 - \tilde{\lambda}(x))^p}{1 - (1 - \tilde{\lambda}(x))} \cdot \tilde{\lambda}(x)' \\ &\equiv - \frac{\tilde{\lambda}(x)^p}{\tilde{\lambda}(x)} \cdot \tilde{\lambda}(x)' \\ &\equiv -\tilde{\lambda}(x)^{p-1} \cdot \tilde{\lambda}(x)' \pmod{p} \end{aligned}$$

(dividing by $\tilde{\lambda}(x)$ is fine since it is an element of $ku^0(X)[[t]]^\times$ as seen above). Now observe that $-\tilde{\lambda}(x)^{p-1} \cdot \tilde{\lambda}(x)' = -\frac{1}{p}(\tilde{\lambda}(x)^p)'$ and that

$$\tilde{\lambda}(x)^p \equiv \sum_{i \geq 0} (-1)^{ip} \lambda^i(x)^{ip} t^{ip} \pmod{p}.$$

Hence the coefficient of t^p in $\tilde{\lambda}(x)^p$ is $(-1)^p \lambda^1(x)^p + py = -x^p + py$ for some y . Hence the coefficient of t^{p-1} in $-\frac{1}{p}(\tilde{\lambda}(x)^p)'$ is $-\frac{1}{p}(-px^p + p^2y) \equiv x^p \pmod{p}$ and we're done.

Next, we prove (c) and explain how it can be used to give a quick and dirty proof of (a), (b), and (d), which only works for $ku^0(X)$, but not for general λ -rings. For (c), we find that $\lambda([L]) = 1 - [L]t$, hence

$$-\ell(\tilde{\lambda}([L]) - 1)' = \sum_{i \geq 0} [L]^{i+1} t^i,$$

and thus $\psi^i([L]) = [L]^i$ for all $i \geq 1$. For $i = 0$ it holds by Definition III.17a.

To show (a), (b), and (d), we first verify that all $\psi^i: ku^0(X) \rightarrow ku^0(X)$ are additive. This is easy and holds more or less by construction (we took the logarithm of a homomorphism into the multiplicative unit group of $ku^0(X)[[t]]$). To show multiplicativity as well as (b) and (d), note that these follow from additivity and (c) whenever our vector bundles $[V] \in ku^0(X)$ can be decomposed into a direct sum $V = L_1 \oplus \cdots \oplus L_n$ of line bundles. This clearly doesn't hold in general. However, by the *splitting principle*, there's always a map $f: F \rightarrow X$ such that $f^*: ku^*(X) \rightarrow ku^*(F)$ is injective and $f^*(V)$ decomposes into a sum of line bundles. By naturality of the Adams operations, this allows us to reduce everything to the decomposable case, whence we are done.

Finally, we prove (e). From the 1-categorical Yoneda lemma we know that the ψ^i also exist as (homotopy classes of) maps $\psi^i: ku \rightarrow ku$. Then $\psi^i(\beta^n) \in ku^0(\mathbb{S}^{2n}, *) = \pi_{2n}(ku_0)$ corresponds to the map

$$\mathbb{S}^{2n} \xrightarrow{\beta^n} ku \xrightarrow{\psi^i} ku.$$

Using (a), we see that $\psi^i(\beta^n) = \psi^i(\beta)^n$ also holds in the homotopy ring $\pi_*(ku_0)$. Hence it suffices to consider the case $n = 1$. In this case, recall $\beta = [\gamma_1^{\mathbb{C}}] - [\mathbb{C}]$, where $[\mathbb{C}] \in ku^0(\mathbb{S}^{2n})$ acts as the multiplicative unit, and compute

$$\psi^i(\beta) = \psi^k([\gamma_1^{\mathbb{C}}] - [\mathbb{C}]) = [\gamma_1^{\mathbb{C}}]^i - [\mathbb{C}] = \beta \sum_{j=0}^{i-1} [\gamma_1^{\mathbb{C}}]^j = i\beta.$$

Here we used (c) as well as the general fact that cup products in generalised cohomology theories vanish on suspensions, so $\beta^2 = 0 \in ku^0(\mathbb{S}^2, *)$ and thus $\beta[\gamma_1^{\mathbb{C}}]^j = \beta$ holds for all $j = 0, \dots, i-1$. \square

A Sketch of Quillen's Computation

Throughout this subsection, p is a prime number and q some power of p . By Proposition III.18(a) and the 1-categorical Yoneda lemma, the Adams operations induce (homotopy classes of) maps $\psi^i: ku \rightarrow ku$, which preserve the 0-component $ku_0 \simeq \{0\} \times BU$.

III.19. Theorem (Quillen [Qui72]). — *Any embedding $\rho: \overline{\mathbb{F}}_p^\times \hookrightarrow \mathbb{C}^\times$ as the group $\bigoplus_{\ell \neq p} \mu_{\ell^\infty}$ of all roots of unity of order coprime to p gives an equivalence*

$$\mathrm{Br}^\rho: k(\mathbb{F}_q)_0 \xrightarrow{\sim} \mathrm{fib}(\psi^q - \mathrm{id}: BU \longrightarrow BU).$$

Consequently,

$$K_i(\mathbb{F}_q) = \begin{cases} \mathbb{Z} & \text{if } i = 0 \\ \mathbb{Z}/(q^n - 1) & \text{if } i = 2n - 1 \\ 0 & \text{else} \end{cases}.$$

The group μ_{ℓ^∞} of ℓ -power roots of unity is called the *Prüfer ℓ -group*. There are many more ways to write it down, such as $\mathbb{Z}[\ell^{-1}]/\mathbb{Z}$ or $\mathbb{Q}_\ell/\mathbb{Z}_\ell$. Using that the group of units in any finite field is cyclic, one easily finds $\overline{\mathbb{F}}_p^\times \simeq \bigoplus_{\ell \neq p} \mu_{\ell^\infty}$. But there is no canonical embedding of this into \mathbb{C}^\times , hence we remember ρ in the notation Br^ρ . It should be noted that Br^ρ also depends on the embedding $\mathbb{F}_q \hookrightarrow \overline{\mathbb{F}}_p$ into an algebraic closure, which we chose to suppress in the notation

Proof sketch of Theorem III.19. The computation of $K_*(\mathbb{F}_q) = \pi_*(k(\mathbb{F}_q)_0)$ is a consequence of the long exact sequence of homotopy groups associated to the fibre sequence

$$\mathrm{fib}(\psi^q - \mathrm{id}) \longrightarrow BU \xrightarrow{\psi^q - \mathrm{id}} BU,$$

using that $\pi_*(BU)$ vanishes in odd degrees by Theorem III.16, and that in even degrees the map $\psi^q - \mathrm{id}: \pi_{2n}(BU) \rightarrow \pi_{2n}(BU)$ acts as $(q^n - 1): \mathbb{Z}\langle \beta^n \rangle \rightarrow \mathbb{Z}\langle \beta^n \rangle$ by Proposition III.18(e).

Now we'll sketch a proof of the first part, up to the extensive group homology calculations that go into it. To get a map

$$k(R)_0 \longrightarrow \mathrm{fib}(\psi^q - \mathrm{id}),$$

we will first construct “compatible” maps $BGL_n(R) \rightarrow \mathrm{fib}(\psi^q - \mathrm{id})$ for all $n \geq 1$ to get a map $BGL_\infty(\mathbb{F}_q) \rightarrow \mathrm{fib}(\psi^q - \mathrm{id})$. Then observe that the right-hand side is simple: Indeed, it's a general fact about fibre sequences $F \rightarrow E \rightarrow B$ that the action of $\pi_1(F, f)$ on $\pi_*(F, f)$ factors through an action of $\pi_1(E, f)$ on $\pi_*(F, f)$. Since no one (including Quillen) ever seems to spell that out, I'll give the details in Lemma* III.19b below. Anyway, this shows that $\mathrm{fib}(\psi^q - \mathrm{id})$ is indeed simple since $\pi_1(BU) = 0$. The fundamental group of any simple space is abelian, hence the universal property from Proposition III.13 together with Corollary III.14a provide an extension

$$k(R)_0 \simeq BGL_\infty(\mathbb{F}_q)^+ \longrightarrow \mathrm{fib}(\psi^q - \mathrm{id}).$$

If we can show that this is an isomorphism on homology, then we're done by Whitehead's theorem, since both sides are simple: $k(R)_0$ because it is an H -space, and $\mathrm{fib}(\psi^q - \mathrm{id})$ because

we just checked this. By Proposition III.13, we may replace $BGL_\infty(\mathbb{F}_q)^+$ by $BGL_\infty(\mathbb{F}_q)$, and by “standard arguments”, it suffices to have isomorphisms on homology with coefficients in \mathbb{Q} , \mathbb{Z}/p , and \mathbb{Z}/ℓ for all $\ell \neq p$. The latter is again something no one ever spells out, so we’ll give a proper argument in our brief discussion of completions of spectra, see [TODO].

Since $\pi_* \text{fib}(\psi^q - \text{id})$ are p -torsionfree torsion abelian groups, the “Hurewicz theorem modulo a Serre class” (see for example [Hat02, Theorem 5.7] in Hatcher’s additional chapter on spectral sequences) shows that the same must be true for $H_*(\text{fib}(\psi^q - \text{id}), \mathbb{Z})$, hence

$$H_*(\text{fib}(\psi^q - \text{id}), \mathbb{Q}) = 0 \quad \text{and} \quad H_*(\text{fib}(\psi^q - \text{id}), \mathbb{Z}/p) = 0.$$

Furthermore, if G is a finite group, then the group homology $H_*^{\text{grp}}(G, A)$ for any coefficients A is annihilated by $\#G$. In particular, it vanishes for $A = \mathbb{Q}$, hence

$$H_*(BGL_\infty(\mathbb{F}_q), \mathbb{Q}) = \text{colim}_{n \in \mathbb{N}} H_*^{\text{grp}}(GL_n(\mathbb{F}_q), \mathbb{Q}) = 0$$

The bulk of Quillen’s proof goes into comparing the homology $H_*(BGL_\infty(\mathbb{F}_q), \mathbb{Z}/\ell)$ and $H_*(\text{fib}(\psi^q - \text{id}), \mathbb{Z}/\ell)$ for $\ell \neq p$, and another significant portion into $H_*(BGL_\infty(\mathbb{F}_q), \mathbb{Z}/p) = 0$. Both problems are essentially about group homology and would take us too much time, so we’ll refer to Quillen’s original paper [Qui72] instead.

What we will do, however, is to construct the desired map $BGL_\infty(\mathbb{F}_q) \rightarrow \text{fib}(\psi^q - \text{id})$. To give this construction some structure, I decided to divide it into four steps.

Step 0. We recall some facts from the representation theory of finite groups.

If you are someone like me who dodged representation theory for their entire mathematical life, Fabian recommends you put a copy of [Ser77] on your nightstand and read a page per day; it will make you a better mathematician within a semester. So let G be a finite group and $R_k(G)$ its *representation ring* over some field k (that is, $R_k(G) = K_0(k[G])$). We’ll need the following facts:

- (a) There is a map

$$\text{rep}: R_{\mathbb{C}}(G) \longrightarrow ku^0(BG),$$

sending a representation $V \in R_{\mathbb{C}}(G)$, i.e. a \mathbb{C} -vector space with a G -action, to the vector bundle $EG \times_G V \rightarrow BG$. Alternatively, it can be described as sending a group homomorphism $r: G \rightarrow U(n)$ to the map $BG \rightarrow BU(n) \rightarrow \{n\} \times BU \subseteq ku$. It suffices to consider homomorphisms $r: G \rightarrow U(n) \subseteq GL_n(\mathbb{C})$, since we can always find a G -equivariant scalar product on any G -representation V (start with any scalar product $\langle \cdot, \cdot \rangle'$ on V and put $\langle u, v \rangle = \#G^{-1} \sum_{g \in G} \langle r(g)u, r(g)v \rangle'$ to make it G -equivariant).

- (b) For any field k , there are exterior power maps $\Lambda^i: \text{LMod}_{k[G]} \rightarrow \text{LMod}_{k[G]}$. Just as in Step 1 of III.17 (minus the technical mumbo-jumbo), these induce maps of sets $\lambda^i: R_k(G) \rightarrow R_k(G)$, turning the representation ring $R_k(G) = K_0(k[G])$ into a λ -ring. And just as in Step 2 of III.17, we can then extract Adams operations $\psi^i: R_k(G) \rightarrow R_k(G)$ from them. In the case $k = \mathbb{C}$, it follows from the constructions that $\text{rep}: R_{\mathbb{C}}(G) \rightarrow ku^0(BG)$ from (a) commutes with the Adams operations on both sides.
- (c) Recall that a *class function* on G is a function $G \rightarrow \mathbb{C}$ which is constant on conjugacy classes. One of the first results about complex representations is that the function

$$\chi: R_{\mathbb{C}}(G) \hookrightarrow \{\text{class functions } G \rightarrow \mathbb{C}\}$$

is injective (and an isomorphism on $\text{Rep}_{\mathbb{C}}(G) \otimes \mathbb{C}$). Here χ sends a representation V to the class function $\chi_V: G \rightarrow \mathbb{C}$ given by $\chi_V(g) = \text{tr}(g: V \rightarrow V)$.

(d) The map χ from (c) interacts with the Adams operations ψ^n from (b) as follows:

$$\chi(\psi^n(V))(g) = \chi(V)(g^n).$$

To see this, fix V and g , and let $\lambda_1, \dots, \lambda_d$ be the eigenvalues of $g: V \rightarrow V$, counted with algebraic multiplicity (so that $d = \dim V$). Then the eigenvalues of $\Lambda^m g: \Lambda^m V \rightarrow \Lambda^m V$ are given by the products $\lambda_{i_1} \cdots \lambda_{i_m}$ for $1 \leq i_1 < \cdots < i_m \leq m$. In particular, $\text{tr}(\Lambda^m g)$ is the m^{th} elementary symmetric polynomial in the λ_i . Hence

$$\chi(\tilde{\lambda}(V))(g) = \sum_{m=0}^d \chi(\Lambda^m V)(g) t^m = \prod_{i=1}^d (1 - \lambda_i t).$$

The derivative of the right-hand side is $-\sum_{i=1}^d \lambda_i \prod_{j \neq i} (1 - \lambda_j t)$. Plugging in our definitions, together with the formula after Definition III.17a, thus shows

$$\sum_{n \geq 0} \chi(\psi^{n+1}(V))(g) t^n = \sum_{i=1}^d \frac{\lambda_i}{1 - \lambda_i t} = \sum_{n \geq 0} \sum_{i=1}^d \lambda_i^{n+1} t^n.$$

Comparing coefficients, plus the fact that the eigenvalues of g^n are precisely $\lambda_1^n, \dots, \lambda_d^n$ (for example by Jordan decomposition), proves the claim.

With that out of the way, let's start with the actual construction.

Step 1. For any embedding $\rho: \overline{\mathbb{F}}_p^\times \hookrightarrow \mathbb{C}^\times$ we construct the famous Brauer lift

$$\text{Br}^\rho: R_{\overline{\mathbb{F}}_p}(\text{GL}_n(\mathbb{F}_q)) \longrightarrow R_{\mathbb{C}}(\text{GL}_n(\mathbb{F}_q)).$$

For any $\overline{\mathbb{F}}_p$ -representation V of $\text{GL}_n(\mathbb{F}_q)$, we form its *Brauer character*

$$\chi_V^\rho: \text{GL}_n(\mathbb{F}_q) \longrightarrow \mathbb{C},$$

sending an element g to $\chi_V^\rho(g) = \sum_{\lambda} \text{mult}(\lambda) \rho(\lambda)$, where the sum runs over the $\overline{\mathbb{F}}_p$ -eigenvalues of $g: V \rightarrow V$, with their algebraic multiplicities $\text{mult}(\lambda)$. Since eigenvalues are preserved under conjugation, χ_V^ρ is a class function on $\text{GL}_n(\mathbb{F}_q)$. By a classical result of Green [Gre55, Theorem 1], χ_V^ρ is in the image of the injection

$$\chi: R_{\mathbb{C}}(\text{GL}_n(\mathbb{F}_q)) \hookrightarrow \{\text{class functions } \text{GL}_n(\mathbb{F}_q) \longrightarrow \mathbb{C}\}$$

from (c) above. This gives rise to the *Brauer lift* $\text{Br}^\rho: R_{\overline{\mathbb{F}}_p}(\text{GL}_n(\mathbb{F}_q)) \rightarrow R_{\mathbb{C}}(\text{GL}_n(\mathbb{F}_q))$. We can now form a diagram

$$\begin{array}{ccc} R_{\overline{\mathbb{F}}_p}(\text{GL}_n(\mathbb{F}_q)) & \xrightarrow{\text{Br}^\rho} & R_{\mathbb{C}}(\text{GL}_n(\mathbb{F}_q)) \\ \uparrow \scriptstyle \overline{\mathbb{F}}_p \otimes_{\mathbb{F}_q} - & & \downarrow \scriptstyle \text{rep} \\ R_{\mathbb{F}_q}(\text{GL}_n(\mathbb{F}_q)) & \dashrightarrow & ku^0(B\text{GL}_n(\mathbb{F}_q)) \end{array}$$

Also recall that $ku^0(B\text{GL}_n(\mathbb{F}_q)) = \pi_0 \text{Hom}_{\text{An}}(B\text{GL}_n(\mathbb{F}_q), ku)$. Hence the canonical representation $\text{GL}_n(\mathbb{F}_q) \curvearrowright \mathbb{F}_q^n$ defines a homotopy class of maps $B\text{GL}_n(\mathbb{F}_q) \rightarrow ku$. We have yet to show that these maps lift to $\text{fib}(\psi^q - \text{id})$, and that the lifts are compatible for varying n .

Step 2. As a first step in that direction, we verify that the composition

$$R_{\mathbb{F}_q}(\mathrm{GL}_n(\mathbb{F}_q)) \longrightarrow R_{\overline{\mathbb{F}_p}}(\mathrm{GL}_n(\mathbb{F}_q)) \xrightarrow{\mathrm{Br}^\rho} R_{\mathbb{C}}(\mathrm{GL}_n(\mathbb{F}_q))$$

takes values in the fixed points of ψ^q .

Since the map χ from (c) above is injective, it's enough to verify this after composition with χ . Recall that $\mathbb{F}_q \subseteq \overline{\mathbb{F}_p}$ is the fixed field of the Frobenius $(-)^q: \overline{\mathbb{F}_p} \rightarrow \overline{\mathbb{F}_p}$. Hence, if λ is an eigenvalue of $g: \overline{\mathbb{F}_p} \otimes_{\mathbb{F}_q} V \rightarrow \overline{\mathbb{F}_p} \otimes_{\mathbb{F}_q} V$, then so are $\lambda^q, \lambda^{q^2}, \lambda^{q^3}, \dots$, each of them with the same multiplicity as λ . Hence

$$\chi_{\overline{\mathbb{F}_p} \otimes_{\mathbb{F}_q} V}^\rho(g) = \sum_{[\mu]} \mathrm{mult}(\mu) \sum_{\lambda \in [\mu]} \rho(\lambda),$$

where the first sum ranges over all Frobenius orbits $[\mu]$. Using assertion (d) from Step 0 shows that applying ψ^q only replaces g by g^q . But the eigenvalues of g^q are the q^{th} powers of the eigenvalues of g (for example by Jordan decomposition), and replacing each λ by λ^q only permutes the summands in $\sum_{\lambda \in [\mu]} \rho(\lambda)$, hence $\chi_{\overline{\mathbb{F}_p} \otimes_{\mathbb{F}_q} V}^\rho$ is indeed invariant under the action of ψ^q .

The upshot is that we obtain a compatible system of maps

$$R_{\mathbb{F}_q}(\mathrm{GL}_n(\mathbb{F}_q)) \longrightarrow \mathrm{ku}^0(B\mathrm{GL}_n(\mathbb{F}_q))^{\psi^q} = \pi_0 \mathrm{Hom}_{\mathrm{An}}(B\mathrm{GL}_n(\mathbb{F}_q), \mathrm{ku})^{\psi^q},$$

where $(-)^{\psi^q}$ denotes fixed points under ψ^q . Here we also use that the canonical map $\mathrm{rep}: R_{\mathbb{C}}(\mathrm{GL}_n(\mathbb{F}_q)) \rightarrow \mathrm{ku}^0(B\mathrm{GL}_n(\mathbb{F}_q))$ from assertion (c) above is compatible with the Adams operations on both sides.

Step 3. To get a compatible system of maps $B\mathrm{GL}_n(\mathbb{F}_q) \rightarrow \mathrm{fib}(\psi^q - \mathrm{id})$, we show that there are bijections

$$\begin{aligned} \pi_0 \mathrm{Hom}_{\mathrm{An}}(B\mathrm{GL}_n(\mathbb{F}_q), \mathrm{fib}(\psi^q - \mathrm{id})) &\xrightarrow{\sim} \pi_0 \mathrm{Hom}_{\mathrm{An}}(B\mathrm{GL}_n(\mathbb{F}_q), \mathrm{ku})^{\psi^q}, \\ \pi_0 \mathrm{Hom}_{\mathrm{An}}(B\mathrm{GL}_\infty(\mathbb{F}_q), \mathrm{fib}(\psi^q - \mathrm{id})) &\xrightarrow{\sim} \lim_{n \in \mathbb{N}} \pi_0 \mathrm{Hom}_{\mathrm{An}}(B\mathrm{GL}_n(\mathbb{F}_q), \mathrm{fib}(\psi^q - \mathrm{id})). \end{aligned}$$

The first map is automatically surjective. Indeed, the left-hand side consists of maps to the homotopy fibre $\mathrm{fib}(\psi^q - \mathrm{id})$, which given by maps to ku plus a homotopy to the identity after composition with $\psi^q: \mathrm{ku} \rightarrow \mathrm{ku}$. On the right-hand side, we forget the choice of homotopy and only remember that there is some. To show injectivity, we use that $\mathrm{Hom}_{\mathrm{An}}(B\mathrm{GL}_n(\mathbb{F}_q), -)$ preserves fibre sequences. Hence, by the long exact sequence of a fibration, there is a surjection from $\pi_1(B\mathrm{GL}_n(\mathbb{F}_q), \mathrm{ku}) = \mathrm{ku}^{-1}(B\mathrm{GL}_n(\mathbb{F}_q))$ onto the kernel. But we have the following general result:

III.19a. Theorem (Atiyah–Segal completion theorem). — *If G is a finite group (or a compact Lie group), then*

$$KU^i(BG) = \begin{cases} R_{\mathbb{C}}(G)_I^\wedge & \text{if } i \text{ is even} \\ 0 & \text{if } i \text{ is odd} \end{cases},$$

where I is the kernel of $\mathrm{rk}: R_{\mathbb{C}}(G) \rightarrow \mathbb{Z}$, and $(-)_I^\wedge$ denotes I -adic completion.

This shows $ku^{-1}(BGL_n(\mathbb{F}_q)) = 0$, and thus the first map is indeed a bijection. I suspect that there should be a less overkill proof that ku^{-1} vanishes, but right now I'm too lazy to think about this.

The second map is part of a *Milnor exact sequence* in general (see for example the corresponding [nLab](#) article). This shows that it is surjective and its kernel is given by

$$R^1 \lim_{n \in \mathbb{N}} \pi_1 \operatorname{Hom}_{\operatorname{An}}(BGL_n(\mathbb{F}_q), ku).$$

But this vanishes by the Atiyah–Segal completion theorem again.

Therefore, we have constructed a homotopy class of maps $BGL_\infty(\mathbb{F}_q) \rightarrow \operatorname{fib}(\psi^q - \operatorname{id})$. Now extend it over the Quillen plus construction $BGL_\infty(\mathbb{F}_q)^+$ and the computation of group homology may begin ... \square

We have yet to show the fibration lemma that was used to prove that $\operatorname{fib}(\psi^q - \operatorname{id})$ is simple. The topology gang will probably find it trivial ...

III.19b. Lemma*. — *Let $F \rightarrow E \xrightarrow{p} B$ be a fibre sequence of anima. Then for all $f \in F$, the action of $\pi_1(F, f)$ on $\pi_*(F, f)$ factors through an action of $\pi_1(E, f)$ on $\pi_*(F, f)$.*

Proof.* Let's first recall how the action of $\pi_1(F, f)$ on $\pi_*(F, f)$ works. So let \square^n denote the n -cube and $\alpha: (\square^n, \partial\square^n) \rightarrow (F, f)$ a map of pairs representing an element of $\pi_n(F, f)$, and let $\gamma: \Delta^1 \rightarrow F$ represent a loop in F with basepoint f . Consider

$$\gamma \cdot \alpha := (\alpha \times \{0\}) \cup (\operatorname{id}_{\partial\square^n} \times \gamma): (\square^n \times \{0\}) \cup (\partial\square^n \times \Delta^1) \longrightarrow F.$$

Once we remember $(\square^n \times \{0\}) \cup (\partial\square^n \times \Delta^1) \simeq \square^n$, it provides a well-defined element $[\gamma] \cdot [\alpha] \in \pi_n(F, f)$, which is the action we're looking for.

Now suppose $\gamma: \Delta^1 \rightarrow E$ only represents an element of $\pi_1(E, f)$. We can still apply the construction $\gamma \cdot \alpha$ above and obtain a lifting diagram

$$\begin{array}{ccc} (\square^n \times \{0\}) \cup (\partial\square^n \times \Delta^1) & \xrightarrow{\gamma \cdot \alpha} & E \\ \downarrow & \nearrow \text{dashed} & \downarrow p \\ \square^n \times \Delta^1 & \xrightarrow{\operatorname{id}_{\square^n} \times p(\gamma)} & B \end{array}$$

Restricting any lift to $\square^n \times \{1\}$ gives a map $\square^n \rightarrow E$, whose restriction to $\partial\square^n$ is constant on f and whose composition with p is constant on $p(f)$, hence an element $[\gamma] \cdot [\alpha] \in \pi_n(F, f)$. It's straightforward to check that the action of $\pi_1(F, f)$ on $\pi_*(F, f)$ factors through this new action. \square

Next, we investigate how the equivalence Br^ρ depends on the choices of embeddings of \mathbb{F}_q into an algebraic closure $\overline{\mathbb{F}}_p$, and $\rho: \overline{\mathbb{F}}_p^\times \hookrightarrow \mathbb{C}^\times$ as the group $\prod_{\mu_\ell}^\infty$.

III.20. Proposition. — *The computation of K -groups from Theorem III.19 via Br^ρ satisfies the following functoriality behaviour:*

(a) *If $f: \mathbb{F}_q \rightarrow \mathbb{F}_{q^n}$ is a field homomorphism compatible with the chosen embeddings into an algebraic closure $\overline{\mathbb{F}}_p$, then the following diagram commutes:*

$$\begin{array}{ccc} K_{2i-1}(\mathbb{F}_q) & \xrightarrow{f_*} & K_{2i-1}(\mathbb{F}_{q^n}) \\ \operatorname{Br}^\rho \downarrow \wr & & \operatorname{Br}^\rho \downarrow \wr \\ \mathbb{Z}/(q^i - 1) & \xrightarrow{\sum_{k=0}^{n-1} q^{ki}} & \mathbb{Z}/(q^{ni} - 1) \end{array}$$

(b) The Frobenius $\text{Frob} = (-)^p: \mathbb{F}_q \rightarrow \mathbb{F}_q$ induces the maps

$$p^i: K_{2i-1}(\mathbb{F}_q) \longrightarrow K_{2i-1}(\mathbb{F}_q)$$

on K -theory.

Proof. For (a), note first that $\psi^{q^n} - \text{id} = (\psi^q)^n - \text{id} = (\psi^q - \text{id}) \sum_{k=0}^{n-1} \psi^{q^k}$ by Proposition III.18(b). Now per construction of the Brauer lift also the left square in

$$\begin{array}{ccccc} k(\mathbb{F}_q)_0 & \xrightarrow{\text{Br}^\rho} & ku & \xrightarrow{\psi^q - \text{id}} & ku \\ f_* \downarrow & & \parallel & & \downarrow \sum_{k=0}^{n-1} \psi^{q^k} \\ k(\mathbb{F}_q)_0 & \xrightarrow{\text{Br}^\rho} & ku & \xrightarrow{\psi^{q^n} - \text{id}} & ku \end{array}$$

commutes. From the bijectivity of

$$\pi_0 \text{Hom}_{\text{An}}(B\text{GL}_n(\mathbb{F}_q), \text{fib}(\psi^q - \text{id})) \xrightarrow{\sim} \pi_0 \text{Hom}_{\text{An}}(B\text{GL}_n(\mathbb{F}_q), ku)^{\psi^q}$$

(Step 3 in the proof of Theorem III.19), we then obtain that f_* agrees with the map induced by the right-hand square on fibres. Hence the claim follows from Proposition III.18(e) and the way we used it in the proof of Theorem III.19.

For (b), one similarly computes from the definition of the Brauer lift and Proposition III.18(b) that

$$\begin{array}{ccccc} k(\mathbb{F}_q)_0 & \xrightarrow{\text{Br}^\rho} & ku & \xrightarrow{\psi^q - \text{id}} & ku \\ \text{Frob}_* \downarrow & & \downarrow \psi^p & & \downarrow \psi^p \\ k(\mathbb{F}_q)_0 & \xrightarrow{\text{Br}^\rho} & ku & \xrightarrow{\psi^q - \text{id}} & ku \end{array}$$

commutes. Then the same argument works. \square

III.21. Corollary. — Any embedding $\rho: \overline{\mathbb{F}}_p^\times \hookrightarrow \mathbb{C}^\times$ induces isomorphisms

$$K_i(\overline{\mathbb{F}}_p) \simeq \begin{cases} \mathbb{Z} & \text{if } i = 0 \\ \bigoplus_{\ell \neq p} \mu_{\ell^\infty} & \text{if } i \text{ is odd} \\ 0 & \text{else} \end{cases}$$

Moreover, if $\mathbb{F}_q \hookrightarrow \overline{\mathbb{F}}_p$ is any inclusion into an algebraic closure, then

$$K_*(\mathbb{F}_q) \xrightarrow{\sim} K_*(\overline{\mathbb{F}}_p)^{\text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_q)}.$$

One can actually make the computation of $K_*(\overline{\mathbb{F}}_p)$ canonical. We already know from Corollary III.7 and Proposition III.8 that $K_1(\overline{\mathbb{F}}_p) \simeq \overline{\mathbb{F}}_p^\times$ canonically. In general, using that $\overline{\mathbb{F}}_p^\times \otimes_{\mathbb{Z}}^L \overline{\mathbb{F}}_p^\times \simeq \overline{\mathbb{F}}_p^\times[1]$, we get

$$K_{2i-1}(\overline{\mathbb{F}}_p) = H_{i-1} \left(\underbrace{\overline{\mathbb{F}}_p^\times \otimes_{\mathbb{Z}}^L \cdots \otimes_{\mathbb{Z}}^L \overline{\mathbb{F}}_p^\times}_{i \text{ times}} \right),$$

and hence an $\text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_p) = \widehat{\mathbb{Z}}$ -equivariant isomorphism

$$K_*(\overline{\mathbb{F}}_p) = H_* \left(\text{Free}_{\mathcal{D}(\mathbb{Z}) \otimes_{\mathbb{Z}}^L}^{\text{Alg}}(\overline{\mathbb{F}}_p^\times[1]) \right).$$

Be warned, however, that this isomorphism only exists at the level of homology, but not at the level of \mathbb{E}_∞ -ring spectra.

LECTURE 23
28th Jan, 2021

Proof of Corollary III.21. Note that K -theory commutes with filtered colimits. One way to see this is that $\mathrm{GL}_n(-): \mathrm{Ring} \rightarrow \mathrm{Grp}(\mathrm{Set})$ commutes with filtered colimits by inspection, hence so does $B\mathrm{GL}_n(-): \mathrm{Ring} \rightarrow \mathrm{An}$, as one can check on homotopy groups (by Remark* II.31b). Now $(-)^+: \mathrm{An} \rightarrow \mathrm{An}$ doesn't commute with colimits in general (it only does if the target is replaced with $\mathrm{An}^{\mathrm{hypo}}$). But in a filtered colimit $\mathrm{colim}_{i \in \mathcal{I}} B\mathrm{GL}_\infty(R_i)^+$, all $B\mathrm{GL}_\infty(R_i)^+$ have abelian fundamental groups rather than just hypoabelian ones (because they are \mathbb{E}_∞ -groups), and filtered colimits of abelian groups are abelian again, hence $\mathrm{colim}_{i \in \mathcal{I}} B\mathrm{GL}_\infty(R_i)^+$ is an element of $\mathrm{An}^{\mathrm{hypo}}$ and thus coincides with $B\mathrm{GL}_\infty(\mathrm{colim}_{i \in \mathcal{I}} R_i)$.

With that out of the way, we may write $\overline{\mathbb{F}}_p = \mathrm{colim}_{n \in \mathbb{N}} \overline{\mathbb{F}}_{p^{n!}}$ (instead of $n!$, we could have used any sequence converging to 0 in $\widehat{\mathbb{Z}} = \lim_{m \in \mathbb{N}} \mathbb{Z}/m$) and get

$$K_{2i}(\overline{\mathbb{F}}_p) = 0 \quad \text{and} \quad K_{2i-1}(\overline{\mathbb{F}}_p) = \mathrm{colim}_{n \in \mathbb{N}} \mathbb{Z}/((p^{n!})^i - 1).$$

The transition maps in the colimit on the right-hand side are those from Proposition III.20(a). Clearly, the formula shows that $K_{2i-1}(\overline{\mathbb{F}}_p)$ is p -torsion free. Let's determine its ℓ -power torsion part for primes $\ell \neq p$. Note that for every $r \geq 1$ there is an $m \geq 1$ such that $\ell^r \mid p^m - 1$. To see this, one doesn't need to invoke splitting fields and cyclotomic polynomials, as Fabian decided to do; it suffices to note that p , being coprime to ℓ , is an element of the multiplicative group $(\mathbb{Z}/\ell^r)^\times$, which has finite order.

In particular, we obtain $\ell^r \mid (p^{n!})^i - 1$ for all $n \geq m$. This shows that the colimit above is cofinal in the colimit

$$\bigoplus_{\ell \neq p} \mu_{\ell^\infty} = \mathrm{colim}_{p \nmid s} \mathbb{Z}/s,$$

in which there are transition maps $\mathbb{Z}/s \rightarrow \mathbb{Z}/t$ given by multiplication with $\frac{t}{s}$, whenever $s \mid t$. This proves $K_{2i-1}(\overline{\mathbb{F}}_p) = \bigoplus_{\ell \neq p} \mu_{\ell^\infty}$. The additional assertion follows by inspection. \square

III.22. Completions of Spectra. — Fabian would like to end the chapter with some remarks on the Suslin rigidity theorem. But before we can do that, we need to introduce completions of spectra. This took us on a short detour in the lecture. I decided to make it a slightly longer detour, to elaborate more on the connection to derived completion in ordinary algebra and to explain one small technical step in the proof sketch of Theorem III.19.

Let's do the spectra case first. Given a spectrum $E \in \mathrm{Sp}$ and a prime p , we define the p -completion of E as

$$\widehat{E}_p := \lim_{n \in \mathbb{N}} E/p^n E,$$

where we define $E/p^n E$ as the pushout

$$\begin{array}{ccc} E & \xrightarrow{p^n} & E \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & E/p^n E \end{array}$$

Since $\mathrm{hom}_{\mathrm{Sp}}(-, E)$ preserves (but flips) fibre sequences, we see that $E/p^n E$ can equivalently be written as $\mathrm{hom}_{\mathrm{Sp}}((\mathbb{S}/p^n \mathbb{S})[-1], E)$. This also shows

$$\widehat{E}_p \simeq \mathrm{hom}_{\mathrm{Sp}}((\mathbb{S}[p^{-1}]/\mathbb{S})[-1], E),$$

which motivates the following lemma/definition:

III.22a. Lemma/Definition*. — For a spectrum E , the following are equivalent:

- (a) $\mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}[p^{-1}], E) \simeq 0$.
- (b) $\mathrm{hom}_{\mathrm{Sp}}(M, E) \simeq 0$ for all p -local spectra M .
- (c) E is equivalent to its p -completion \widehat{E}_p .

Such spectra are called p -complete. The p -completion functor

$$(-)^\wedge_p : \mathrm{Sp} \longrightarrow \mathrm{Sp}_{p\text{-comp}}$$

is a Bousfield localisation onto the full sub- ∞ -category $\mathrm{Sp}_{p\text{-comp}} \subseteq \mathrm{Sp}$ of p -complete spectra.

Proof.* The implication (b) \Rightarrow (a) is clear, and since $M \simeq M \otimes \mathbb{S}[p^{-1}]$ by Corollary III.4a, the reverse implication follows from the tensor-Hom adjunction (Lemma* II.53a).

Using the fact that $\mathbb{S}[p^{-1}] \otimes \mathbb{S}[p^{-1}]/\mathbb{S} \simeq \mathbb{S}[p^{-1}]/\mathbb{S}[p^{-1}] \simeq 0$ together with the tensor-Hom adjunction, we see that the p -completion $\widehat{E}_p \simeq \mathrm{hom}_{\mathrm{Sp}}((\mathbb{S}[p^{-1}]/\mathbb{S})[-1], E)$ is indeed p -complete in the sense of (a), which shows (c) \Rightarrow (a). Conversely, from the fibre sequence $(\mathbb{S}[p^{-1}]/\mathbb{S})[-1] \rightarrow \mathbb{S} \rightarrow \mathbb{S}[p^{-1}]$ we get that E is p -complete in the sense of (a) iff it equals its own p -completion. The final assertion about $(-)^\wedge_p$ being a Bousfield localisation follows from Proposition I.61a, as usual. \square

III.22b. Derived Completion in General*. — Let's leave the land of spectra for the moment and consider an ordinary commutative ring R with an element $s \in R$. Following [Stacks, Tag 091N], we call a complex $K \in \mathcal{D}(R)$ *derived s -complete* if

$$R\mathrm{Hom}_R(R[s^{-1}], K) \simeq 0.$$

In complete analogy with Lemma/Definition* III.22a, one shows that this is equivalent to $R\mathrm{Hom}_R(M, K) \simeq 0$ for all $M \in \mathcal{D}(R[s^{-1}])$. Moreover, if C denotes the chain complex $(R \rightarrow R[s^{-1}])$ concentrated in degrees 0 and -1 , then the *derived s -completion* functor

$$(-)^\wedge_s := R\mathrm{Hom}_R(C, -) : \mathcal{D}(R) \longrightarrow \mathcal{D}(R)$$

is a Bousfield localisation onto the full sub- ∞ -category $\mathcal{D}_{s\text{-comp}}(R) \subseteq \mathcal{D}(R)$ of derived s -complete complexes. One might wonder perhaps why we use the complex C rather than $(R[s^{-1}]/R)[-1]$, as was the case for spectra. This is because we have to take the derived quotient. In fact, one easily checks that $C[1]$ is the mapping cone of $R \rightarrow R[s^{-1}]$, hence it sits inside a pushout diagram

$$\begin{array}{ccc} R & \longrightarrow & R[s^{-1}] \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & C[1] \end{array}$$

Thus $C \simeq (R[s^{-1}]/^L R)[-1]$ and therefore our formula for derived completion looks exactly like in III.22.

This can be generalised as follows: If $I = (s_1, \dots, s_n) \subseteq R$ is a finitely generated ideal, then a complex $K \in \mathcal{D}(R)$ is called *derived I -complete* if it is derived s -complete for all $s \in I$. It turns out that this is equivalent to K being derived s_i -complete for $i = 1, \dots, n$. Again, there is a Bousfield localisation (*derived I -completion*)

$$(-)^\wedge_I := R\mathrm{Hom}_R(C, -) : \mathcal{D}(R) \longrightarrow \mathcal{D}(R)$$

onto the full sub- ∞ -category $\mathcal{D}_{I\text{-comp}}(R) \subseteq \mathcal{D}(R)$ of derived I -complete complexes. Here C can be chosen as the “Čech complex”

$$C \simeq \left(R \longrightarrow \bigoplus_i R[s_i^{-1}] \longrightarrow \bigoplus_{i < j} R[(s_i s_j)^{-1}] \longrightarrow \dots \longrightarrow R[(s_1 \cdots s_n)^{-1}] \right)$$

in degrees $0, -1, \dots, -n$. Proofs can be found in [Stacks, Tag 091N]. Mind that The Stacks Project only proves things for the derived 1-category $D(R) \simeq \pi\mathcal{D}(R)$, but the arguments can be carried over (potentially with the help of Corollary II.30a).

Here are two pretty useful facts about derived completions:

III.22c. Lemma*. — *Let R be a commutative ring, $s \in R$ an element and $I \subseteq R$ a finitely generated ideal.*

- (a) *The functors $(-)_s^\wedge: \mathcal{D}(R) \rightarrow \mathcal{D}(R)$ and $-[s^{-1}]: \mathcal{D}(R) \rightarrow \mathcal{D}(R)$ are jointly conservative.*
- (b) *(“Derived Nakayama lemma”) If K is derived I -complete, then $K \otimes_R^L R/I \simeq 0$ implies $K \simeq 0$.*

Proof.* Part (a). Since a morphism of complexes is a quasi-isomorphism iff its cone is acyclic, we only need to show that $\widehat{K}_s \simeq 0$ and $K[s^{-1}] \simeq 0$ imply $K \simeq 0$. Suppose that $0 \simeq \widehat{K}_s \simeq R\mathrm{Hom}_R(C, K)$. Since $C \rightarrow R \rightarrow R[s^{-1}]$ is a fibre sequence, this implies

$$R\mathrm{Hom}_R(R[s^{-1}], K) \simeq R\mathrm{Hom}_R(R, K) \simeq K.$$

Hence K is already s -local, which implies $K \simeq K[s^{-1}]$. Hence $K[s^{-1}] \simeq 0$ iff $K \simeq 0$, as claimed. This shows (a). Part (b) needs more preparation and we refer to [Stacks, Tag 0G1U]. \square

III.22d. Derived Completion in the Noetherian Case*. — It is not true, in general, that derived I -completion is given by

$$\widehat{K}_I \simeq R\mathrm{lim}_{n \in \mathbb{N}} (K \otimes_R^L R/I^n)$$

(here $R\mathrm{lim}$ denotes the limit in the ∞ -category $\mathcal{D}(R)$). And neither is it true (I think, but don’t take my word for it) that derived completion is the left-derived functor of

$$\Lambda_I = \lim_{n \in \mathbb{N}} -/I^n -: \mathrm{Mod}_R \longrightarrow \mathrm{Mod}_R$$

in general. Fabian remarks that Λ_I is neither left- nor right-exact in general¹ ... but we need a right-exact functor to form left derivatives, right? Wrong! Grothendieck formulated his theory for arbitrary additive functors between abelian categories without any exactness requirements; it just won’t be true anymore that $L_0\Lambda_I \simeq \Lambda_I$.

Both of the problems above go away if we assume R is a noetherian ring. Indeed, in this case $\widehat{K}_i \simeq R\mathrm{lim}_{n \in \mathbb{N}} (K \otimes_R^L R/I^n)$ holds by [Stacks, Tag 0BKH], which also implies that $(-)_I^\wedge \simeq L\Lambda_I$, since it takes the correct values on free R -modules.

¹ ... which confused me a bit, since $R^1\mathrm{lim}_{n \in \mathbb{N}} M/I^n M = 0$ holds by the Mittag-Leffler condition. However, this isn’t enough to ensure right-exactness. For an inclusion $N \subseteq M$, the kernel of $N/I^n N \rightarrow M/I^n M$ is $(I^n M \cap N)/I^n N$, so we would need $R^1\mathrm{lim}_{n \in \mathbb{N}} (I^n M \cap N)/I^n N = 0$ instead—which isn’t true in general.

III.22e. Derived Completion over \mathbb{Z} vs. Completion of Spectra. — Now let $p \in \mathbb{Z}$ be a prime. Since $\mathbb{Z} \rightarrow \mathbb{Z}[p^{-1}]$ is injective, the complex $C = (\mathbb{Z} \rightarrow \mathbb{Z}[p^{-1}])$ concentrated in degrees 0 and -1 is quasi-isomorphic to $(\mathbb{Z}[p^{-1}]/\mathbb{Z})[-1] \simeq \mu_{p^\infty}[-1]$. Hence derived p -adic completion in $\mathcal{D}(\mathbb{Z})$ is given by

$$L\Lambda_p \simeq R\mathrm{Hom}_{\mathbb{Z}}(\mu_{p^\infty}[-1], -) : \mathcal{D}(\mathbb{Z}) \longrightarrow \mathcal{D}_{p\text{-comp}}(\mathbb{Z}).$$

In particular, if A is an abelian group, then

$$L_0\Lambda_p(A) = \mathrm{Ext}_{\mathbb{Z}}(\mu_{p^\infty}, A), \quad L_1\Lambda_p(A) = \mathrm{Hom}_{\mathbb{Z}}(\mu_{p^\infty}, A) \quad \text{and} \quad L_i\Lambda_p(A) = 0 \text{ for } i \geq 2.$$

It's easy to determine $L_1\Lambda_p(A)$: Writing $\mu_{p^\infty} = \mathrm{colim}_{n \in \mathbb{N}} \mathbb{Z}/p^n$, with transition maps given by multiplication with p , we obtain

$$L_1\Lambda_p(A) = \lim_{n \in \mathbb{N}} \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}/p^n, A) = \lim_{n \in \mathbb{N}} A[p^n],$$

where $A[p^n]$ denotes the p^n -torsion part of A and the transition morphisms are again given by multiplication with p . Moreover, there is a short exact sequence

$$0 \longrightarrow R^1 \lim_{n \in \mathbb{N}} A[p^n] \longrightarrow L^0\Lambda_p(A) \longrightarrow \Lambda_p(A) \longrightarrow 0,$$

To see where this sequence comes from, choose a projective resolution $0 \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0$ (which we can do since \mathbb{Z} has global dimension 1), note that $A[p^n] = \ker(P_1/p^n P_1 \rightarrow P_0/p^n P_0)$ (since the right-hand side computes $\mathrm{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/p^n, A)$) and play a bit around with exact sequences.

The upshot is that if A has bounded p -torsion (for example, if A is finitely generated), then there is an $n \gg 0$ such that $p^n : A[p^{2n}] \rightarrow A[p^n]$ is the zero morphism. Hence

$$R\lim_{n \in \mathbb{N}} A[p^n] \simeq 0$$

in this case and we see that $L\Lambda_p(A)$ coincides with the usual underived p -adic completion $\Lambda_p(A)[0]$ placed in degree 0. These considerations can be exploited to compute homotopy groups of p -completions of spectra:

III.22f. Lemma. — *For every spectrum E and all $i \in \mathbb{Z}$, there are short exact sequences*

$$0 \longrightarrow L^0\Lambda_p(\pi_i(E)) \longrightarrow \pi_i(\widehat{E}_p) \longrightarrow L^1\Lambda_p(\pi_{i-1}(E)) \longrightarrow 0.$$

In particular, if $\pi_(E)$ has degreewise bounded p -torsion, then its degreewise p -completion is $\pi_*(\widehat{E}_p)$. Moreover, the homotopy groups of any p -complete spectrum are derived p -complete.*

*Proof**. The homotopy groups of \widehat{E}_p sit inside the Milnor exact sequence

$$0 \longrightarrow R^1 \lim_{n \in \mathbb{N}} \pi_{i+1}(E/p^n E) \longrightarrow \pi_i(\widehat{E}_p) \longrightarrow \lim_{n \in \mathbb{N}} \pi_i(E/p^n E) \longrightarrow 0$$

(see the [nLab](#) article and note that $\Omega^{\infty+i} : \mathrm{Sp} \rightarrow \mathrm{An}$ commutes with limits, hence the Milnor sequence works for spectra as well). The morphism $p^n : E \rightarrow E$ clearly also induces $p^n : \pi_*(E) \rightarrow \pi_*(E)$ on homotopy groups. Hence the long exact sequence associated to the fibre sequence $E \rightarrow E \rightarrow E/p^n E$ provides short exact sequences

$$0 \longrightarrow \pi_i(E)/p^n \pi_i(E) \longrightarrow \pi_i(E/p^n E) \longrightarrow \pi_{i-1}(E)[p^n] \longrightarrow 0.$$

Now take limits over n . We have $R^1 \lim_{n \in \mathbb{N}} \pi_i(E)/p^n \pi_i(E) = 0$ by Mittag-Leffler, hence the six-term exact sequence splits into a short exact sequence

$$0 \longrightarrow \Lambda_p(\pi_i(E)) \longrightarrow \lim_{n \in \mathbb{N}} \pi_i(E/p^n E) \longrightarrow L_1 \Lambda_p(\pi_{i-1}(E)) \longrightarrow 0$$

and an isomorphism $R^1 \lim_{n \in \mathbb{N}} \pi_i(E/p^n E) \simeq R^1 \lim_{n \in \mathbb{N}} \pi_{i-1}(E)[p^n]$. Summarising the information we've got so far and plugging in the short exact sequence that computes $L_0 \Lambda_p(\pi_i(E))$, we obtain a solid commutative diagram

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \downarrow & & \\
 0 & \longrightarrow & R^1 \lim_{n \in \mathbb{N}} \pi_{i-1}(E)[p^n] & \longrightarrow & L_0 \Lambda_p(\pi_i(E)) & \longrightarrow & \Lambda_p(\pi_i(E)) \longrightarrow 0 \\
 & & \downarrow \wr & & \vdots & & \downarrow \\
 0 & \longrightarrow & R^1 \lim_{n \in \mathbb{N}} \pi_i(E/p^n E) & \longrightarrow & \pi_i(\widehat{E}_p) & \longrightarrow & \lim_{n \in \mathbb{N}} \pi_i(E/p^n E) \longrightarrow 0 \\
 & & & & & & \downarrow \\
 & & & & & & L_1 \Lambda_p(\pi_{i-1}(E)) \\
 & & & & & & \downarrow \\
 & & & & & & 0
 \end{array}$$

If we can show that the dashed arrow exists, then we'll be done. Indeed, the snake lemma will then show that $L_0 \Lambda_p(\pi_i(E)) \rightarrow \pi_i(\widehat{E}_p)$ is injective and that its cokernel is isomorphic to $L_1 \Lambda_p(\pi_{i-1}(E))$, which is what we want to show.

To construct the dashed arrow, it suffices to show that $\pi_i(\widehat{E}_p)$ is derived p -complete, since then the canonical morphism $\pi_i(E) \rightarrow \pi_i(\widehat{E}_p)$ will extend over the derived p -completion $L\Lambda_p(\pi_i(E))$ and hence also over $L_0 \Lambda(\pi_i(E))$ since $\pi_i(\widehat{E}_p)$ is concentrated in degree 0. But the solid diagram is already enough to conclude that $\pi_i(\widehat{E}_p)$ is derived p -complete! Indeed, we know from [Stacks, Tag 091U] that if two out of three abelian groups in a short exact sequence are derived p -complete, then so is the third. Moreover, this result and [Stacks, Tag 091T] imply that the homology of a derived p -complete complex is derived p -complete, and that p -complete abelian groups are also derived p -complete. hence $L_0 \Lambda_p(\pi_i(E))$, $\Lambda_p(\pi_i(E))$, and $L_1 \Lambda_p(\pi_{i-1}(E))$ are all derived p -complete, which suffices to show the same for the rest of the diagram. \square

Only the third part of the following lemma was mentioned in the lecture, but the other two fit nicely and provide a proof of a small technical step in the proof sketch of Theorem III.19, as promised at the beginning of III.22.

III.22g. Lemma. — (a*) *The functors $- \otimes H\mathbb{Q}: \mathrm{Sp} \rightarrow \mathrm{Sp}$ and $(-)^{\wedge}_p: \mathrm{Sp} \rightarrow \mathrm{Sp}$ for all primes p are jointly conservative.*

(b*) *The functors $- \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow \mathcal{D}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$ and $- \otimes_{\mathbb{Z}}^L \mathbb{Z}/p: \mathcal{D}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$ for all primes p are jointly conservative. In particular, if $f: X \rightarrow Y$ is a map of anima such that*

$$f_*: H_*(X, \mathbb{Q}) \xrightarrow{\sim} H_*(Y, \mathbb{Q}) \quad \text{and} \quad f_*: H_*(X, \mathbb{Z}/p) \xrightarrow{\sim} H_*(Y, \mathbb{Z}/p)$$

are isomorphisms, then f is also an isomorphism on homology with \mathbb{Z} -coefficients.

(c) For all spectra E , there is a canonical pullback square (the “arithmetic fracture square”)

$$\begin{array}{ccc} E & \longrightarrow & \prod_p \widehat{E}_p \\ \downarrow & \lrcorner & \downarrow \\ E \otimes H\mathbb{Q} & \longrightarrow & \left(\prod_p \widehat{E}_p \right) \otimes H\mathbb{Q} \end{array}$$

Observe that Lemma III.22g(b*) is wildly false for underived tensor products. For example, \mathbb{Q}/\mathbb{Z} vanishes upon tensoring with \mathbb{Q} or any \mathbb{Z}/p .

Proof of Lemma III.22g.* For (a*), it suffices to show that $E \otimes H\mathbb{Q} \simeq 0$ and $\widehat{E}_p \simeq 0$ for all p imply $E \simeq 0$. By the same argument as in Lemma* III.22c(a), $\widehat{E}_p \simeq 0$ implies that E is p -local. If that’s the case for all p , then $E \simeq E \otimes H\mathbb{Q}$ by Corollary III.5, hence the right-hand side vanishes iff E vanishes.

For part (b*), we can use the same argument to show that $- \otimes_{\mathbb{Z}} \mathbb{Q}: \mathcal{D}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$ and $(-)_p^{\wedge}: \mathcal{D}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$ are jointly conservative. But $- \otimes_{\mathbb{Z}}^L \mathbb{Z}/p: \mathcal{D}_{p\text{-comp}}(\mathbb{Z}) \rightarrow \mathcal{D}(\mathbb{Z})$ is conservative by the derived Nakayama lemma (Lemma* III.22c(b)), hence the claim. The “in particular” follows as $H_*(-, \mathbb{Q}) = H_*(C_{\bullet}(-) \otimes_{\mathbb{Z}} \mathbb{Q})$ and $H_*(-, \mathbb{Z}/p) = H_*(C_{\bullet}(-) \otimes_{\mathbb{Z}}^L \mathbb{Z}/p)$.

For (c), we may apply (a) to see that it suffices to check the pullback property after p -completion and after tensoring with $H\mathbb{Q}$. It’s easy to check that p -completion kills all rational spectra and all ℓ -complete spectra for $\ell \neq p$. Hence after p -completion the lower row becomes zero, whereas the upper row becomes the identity on \widehat{E}_p . Similarly, the pullback property becomes trivial after tensoring with $H\mathbb{Q}$. \square

As a consequence of Lemma III.22g, to understand the *algebraic K -theory spectrum* $K(F) = B^{\infty}k(F)$ of a field F , it suffices to understand its p -completions and its rationalisation. Surprisingly, it is the latter which causes problems in praxis, even though we have a formula for its homotopy groups: $K_i(F) \otimes \mathbb{Q} = \text{indec}_i H_*^{\text{grp}}(\text{GL}_{\infty}(F), \mathbb{Q})$ by Corollary III.10. But computing the right-hand side is an insanely hard problem and very much depends on F , as one can already see in the special case $K_1(F) = F^{\times}$ (by Proposition III.8).

In contrast to that, the ℓ -adic completions $K(F)_{\ell}^{\wedge}$ are well understood and don’t really depend on F :

III.23. Theorem (Suslin’s rigidity theorem). — *For any morphism $F \rightarrow F'$ of separably closed fields and all primes ℓ ,*

$$K(F)_{\ell}^{\wedge} \xrightarrow{\sim} K(F')_{\ell}^{\wedge}$$

is an equivalence. The same is true for the tautological map

$$K(\mathbb{C})_{\ell}^{\wedge} \xrightarrow{\sim} (B^{\infty}ku)_{\ell}^{\wedge}.$$

Moreover, if $p \neq \ell$ is a prime, $\mathcal{O}_p \subseteq \overline{\mathbb{Q}}_p$ the absolute ring of integers over \mathbb{Z}_p , and $\mathfrak{m} \subseteq \mathcal{O}_p$ its maximal ideal, then also

$$K(\mathcal{O}_p)_{\ell}^{\wedge} \xrightarrow{\sim} K(\overline{\mathbb{Q}}_p)_{\ell}^{\wedge} \quad \text{and} \quad K(\mathcal{O}_p)_{\ell}^{\wedge} \xrightarrow{\sim} K(\mathcal{O}_p/\mathfrak{m})_{\ell}^{\wedge}$$

are equivalences.

We won't prove Theorem III.23. But note that $\mathcal{O}_p/\mathfrak{m} \simeq \overline{\mathbb{F}}_p$. Hence we see that any pair of embeddings $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$ and $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{C}}$ induces an equivalence $K(\overline{\mathbb{F}}_p)_\ell^\wedge \simeq (B^\infty ku)_\ell^\wedge$. This fits with our calculations of their homotopy groups: Since $\pi_*(B^\infty ku) = \mathbb{Z}[\beta]$ has degreewise bounded ℓ -torsion, Lemma III.22f implies

$$\pi_i((B^\infty ku)_\ell^\wedge) = \begin{cases} \mathbb{Z}_\ell & \text{if } i \text{ is even} \\ 0 & \text{if } i \text{ is odd} \end{cases}.$$

Also recall that $K_*(\overline{\mathbb{F}}_p)$ is 0 in even degrees and $\bigoplus_{q \neq p} \mu_{q^\infty}$ in odd degrees by Corollary III.21. But one can compute

$$L_0 \Lambda_\ell \left(\bigoplus_{q \neq p} \mu_{q^\infty} \right) = 0 \quad \text{and} \quad L_1 \Lambda_\ell \left(\bigoplus_{q \neq p} \mu_{q^\infty} \right) = \mathbb{Z}_\ell,$$

hence Lemma III.22f implies that ℓ -completion shifts the homotopy groups of $K(\overline{\mathbb{F}}_p)$ from odd degrees into even degrees. Thus

$$\pi_i(K(\overline{\mathbb{F}}_p)_\ell^\wedge) = \begin{cases} \mathbb{Z}_\ell & \text{if } i \text{ is even} \\ 0 & \text{if } i \text{ is odd} \end{cases},$$

as Theorem III.23 predicts.

Fabian also mentioned that even though the Adams operations $\psi^i: B^\infty ku \rightarrow B^\infty ku$ aren't maps of \mathbb{E}_∞ -ring spectra, they can be refined to an action of $\widehat{\mathbb{Z}}$ on the ℓ -completion $(B^\infty ku)_\ell^\wedge$ through \mathbb{E}_∞ -ring spectra maps.

To end this chapter, we state the analogue of Theorem III.19 for Hermitian K -theory/Grothendieck–Witt theory.

III.24. Theorem (Friedlander, Fiedorowicz–Priddy [FP78]). — *For odd q , Brauer lifting induces equivalences*

$$\begin{aligned} \mathrm{gw}^{-\mathrm{sym}}(\mathbb{F}_q)_0 &\xrightarrow{\sim} \mathrm{fib}(\psi^q - \mathrm{id}: B\mathrm{Spin} \longrightarrow B\mathrm{Spin}) \\ \mathrm{gw}^{\mathrm{sym}}(\mathbb{F}_q)_0 &\xrightarrow{\sim} \mathrm{fib}(\psi^q - \mathrm{id}: BO \longrightarrow BO), \end{aligned}$$

and in the even case we obtain

$$\begin{aligned} \mathrm{gw}^{\mathrm{sym}}(\mathbb{F}_{2^n}) &\simeq \mathrm{gw}^{\mathrm{even}}(\mathbb{F}_{2^n}) \xrightarrow{\sim} \mathrm{fib}(\psi^q - \mathrm{id}: B\mathrm{Spin} \longrightarrow B\mathrm{Spin}) \\ \mathrm{gw}^{\mathrm{quad}}(\mathbb{F}_{2^n}) &\simeq \mathrm{gw}^{\mathrm{sym}}(\mathbb{F}_{2^n}) \times B\mathbb{Z}/2. \end{aligned}$$

CHAPTER IV.

The K -Theory of Stable ∞ -Categories

IV

The main goal is to explain Quillen's fibre sequence

$$\bigoplus_{p \text{ prime}} K(\mathbb{F}_p) \longrightarrow K(\mathbb{Z}) \longrightarrow K(\mathbb{Q}).$$

The first problem is that there are no ring maps $\mathbb{F}_p \rightarrow \mathbb{Z}$.

To get started, we note that $K_0(R)$ admits more cycles than just projective modules: Let $P \in \text{Mod}_R$ be a *perfect module*, i.e., M admits a finite projective resolution

$$0 \longrightarrow P_n \longrightarrow P_{n-1} \longrightarrow \dots \longrightarrow P_0 \longrightarrow M \longrightarrow 0.$$

Then we can put

$$[M] := \sum_{i=0}^n (-1)^i [P_i] \in K_0(R).$$

IV.0a. Exercise. — Show that $[M]$ is independent of the choice of resolution!

So you don't need a projective module to specify an element in $K_0(R)$, all you need is a *perfect* one! More generally, let $\mathcal{D}^{\text{perf}}(R) \subseteq \mathcal{D}(R)$ denote the full sub- ∞ -category spanned by the finite complexes of finite projective R -modules (these are precisely the compact objects in $\mathcal{D}(R)$).

IV.1. Lemma. — We have an isomorphism

$$\begin{aligned} (\pi_0 \text{ core } \mathcal{D}^{\text{perf}}(R)) / \sim_{\text{ext}} &\xrightarrow{\sim} K_0(R) \\ [P_n \rightarrow \dots \rightarrow P_{-m}] &\mapsto \sum_{i=-m}^n (-1)^i [P_i], \end{aligned}$$

where " \sim " is the equivalence relation generated by $B \simeq A \oplus C$ if there is a fibre sequence $A \rightarrow B \rightarrow C$ in $\mathcal{D}^{\text{perf}}(R)$.

Proof. Exercise. □

IV.2. Definition. — Given a (small) stable ∞ -category \mathcal{C} , put

$$K_0(\mathcal{C}) := \pi_0 \text{ core }(\mathcal{C}) / \sim_{\text{ext}}$$

IV.3. Theorem/Definition (Thomason). — *A sub- ∞ -category $\mathcal{D} \subseteq \mathcal{C}$ is called dense if every object in \mathcal{C} can be written as a retract of some $d \in \mathcal{D}$, i.e., there are maps*

$$c \longrightarrow d \longrightarrow c$$

composing to id_c .

If \mathcal{C} and \mathcal{D} are stable ∞ -categories and $\mathcal{D} \subseteq \mathcal{C}$ is dense, then the inclusion induces an injection

$$K_0(\mathcal{D}) \hookrightarrow K_0(\mathcal{C}),$$

and in fact a bijection of partially ordered sets

$$\{\mathcal{D} \subseteq \mathcal{C} \text{ dense}\} \xrightarrow{\sim} \{\text{subgroups of } K_0(\mathcal{C})\}.$$

[TODO: Remark about eq. classes of subcategories]. For example, the smallest stable subcategory $\mathcal{D}^{\mathrm{free}}(R) \subseteq \mathcal{D}^{\mathrm{perf}}(R)$ containing $R[0]$ corresponds to $\mathrm{im}(\mathbb{Z} \rightarrow K_0(R))$.

Proof. Inhalt...

□

Bibliography

- [A&HK] Fabian Hebestreit. *Algebraic and Hermitian K-Theory. Lecture held at the University of Bonn.* Winter term 2020/21. URL: <https://www.math.uni-bonn.de/people/fhebestr/Ktheory/>.
- [BGT13] Andrew J Blumberg, David Gepner, and Gonalo Tabuada. “A universal characterization of higher algebraic K -theory”. In: *Geometry & Topology* 17.2 (Apr. 2013), pp. 733–838. DOI: [10.2140/gt.2013.17.733](https://doi.org/10.2140/gt.2013.17.733).
- [Bot59] Raoul Bott. “The stable homotopy of the classical groups”. In: *Ann. of Math.* (2) 70 (1959), pp. 313–337. URL: <https://doi.org/10.2307/1970106>.
- [Die08] Tammo tom Dieck. *Algebraic topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zrich, 2008, pp. xii+567. URL: <https://www.maths.ed.ac.uk/~v1ranick/papers/diecktop.pdf>.
- [Fab+20_I] Baptiste Calmès et al. *Hermitian K-theory for stable ∞ -categories I: Foundations*. 2020. arXiv: [2009.07223](https://arxiv.org/abs/2009.07223) [math.KT].
- [Fab+20_{II}] Baptiste Calmès et al. *Hermitian K-theory for stable ∞ -categories II: Cobordism categories and additivity*. 2020. arXiv: [2009.07224](https://arxiv.org/abs/2009.07224) [math.KT].
- [Fab+20_{III}] Baptiste Calmès et al. *Hermitian K-theory for stable ∞ -categories III: Grothendieck-Witt groups of rings*. 2020. arXiv: [2009.07225](https://arxiv.org/abs/2009.07225) [math.KT].
- [FP78] Zbigniew Fiedorowicz and Stewart Priddy. *Homology of classical groups over finite fields and their associated infinite loop spaces*. Vol. 674. Lecture Notes in Mathematics. Springer, Berlin, 1978, pp. vi+434.
- [Gre55] J. A. Green. “The characters of the finite general linear groups”. In: *Trans. Amer. Math. Soc.* 80 (1955), pp. 402–447. URL: <https://doi.org/10.2307/1992997>.
- [HA] Jacob Lurie. *Higher Algebra*. 2017. URL: <http://people.math.harvard.edu/~lurie/papers/HA.pdf>.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002, pp. xii+544. URL: <https://pi.math.cornell.edu/~hatcher/AT/AT.pdf>.
- [Hau20] Rune Haugseng. *A fibrational mate correspondence for ∞ -categories*. 2020. arXiv: [2011.08808](https://arxiv.org/abs/2011.08808) [math.CT].
- [HC_I] Fabian Hebestreit. *Higher Categories I. Lecture held at the University of Bonn.* Winter term 2019/20. URL: <https://www.math.uni-bonn.de/people/fhebestr/Skript%20HigherCatsI/>.
- [HC_{II}] Fabian Hebestreit. *Higher Categories II. Lecture held at the University of Bonn.* Summer term 2020. URL: <https://www.math.uni-bonn.de/people/fhebestr/Skript%20HigherCatsII/>.
- [HLN20] Fabian Hebestreit, Sil Linskens, and Joost Nuiten. *Orthofibrations and monoidal adjunctions*. 2020. arXiv: [2011.11042](https://arxiv.org/abs/2011.11042) [math.CT].

BIBLIOGRAPHY

- [HTT] Jacob Lurie. *Higher Topos Theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009, pp. xviii+925. URL: <http://people.math.harvard.edu/~lurie/papers/HTT.pdf>.
- [JT07] André Joyal and Myles Tierney. “Quasi-categories vs Segal spaces”. In: *Categories in algebra, geometry and mathematical physics*. Vol. 431. Contemp. Math. Amer. Math. Soc., Providence, RI, 2007, pp. 277–326. arXiv: [math/0607820](https://arxiv.org/abs/math/0607820) [[math.AT](#)].
- [LT19] Land and Tamme. “On the K -theory of pullbacks”. In: *Annals of Mathematics* 190.3 (2019), p. 877. DOI: [10.4007/annals.2019.190.3.4](https://doi.org/10.4007/annals.2019.190.3.4).
- [Lur09] Jacob Lurie. $(\infty, 2)$ -Categories and the Goodwillie Calculus I. 2009. arXiv: [0905.0462](https://arxiv.org/abs/0905.0462) [[math.CT](#)].
- [Lur10] Jacob Lurie. *Chromatic Homotopy Theory (Math 252x)*. Lecture held at Harvard University. Spring 2010. URL: <https://www.math.ias.edu/~lurie/252x.html>.
- [MM65] John W. Milnor and John C. Moore. “On the structure of Hopf algebras”. In: *Ann. of Math. (2)* 81 (1965), pp. 211–264. URL: <https://doi.org/10.2307/1970615>.
- [Nik16] Thomas Nikolaus. *Stable ∞ -Operads and the multiplicative Yoneda lemma*. 2016. arXiv: [1608.02901](https://arxiv.org/abs/1608.02901) [[math.AT](#)].
- [Qui72] Daniel Quillen. “On the cohomology and K -theory of the general linear groups over a finite field”. In: *Ann. of Math. (2)* 96 (1972), pp. 552–586. URL: <https://doi.org/10.2307/1970825>.
- [Ser77] Jean-Pierre Serre. *Linear representations of finite groups*. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42. Springer-Verlag, New York-Heidelberg, 1977, pp. x+170.
- [Spa88] Nicolas Spaltenstein. “Resolutions of unbounded complexes”. In: *Compositio Mathematica* 65.2 (1988), pp. 121–154. URL: http://www.numdam.org/item?id=CM_1988__65_2_121_0.
- [Stacks] The Stacks Project Authors. *The Stacks Project*. 2020. URL: <http://stacks.math.columbia.edu>.