# ASTRA: An Integrated Analysis Framework for Physics-Aware Astrophysical Discovery

Glenn J. White,[1]★ Robin Dey[2]
[1]*Department of Physics and Astronomy, The Open University, Milton Keynes MK7 6AA, UK*
[2]*VBRL Holdings Inc*

Accepted XXX. Received YYY; in original form ZZZ

**ABSTRACT**

We present Astra (Autonomous System for Scientific Discovery in Astrophysics), an integrated framework combining numerical data analysis, causal reasoning, and physical validation within a unified, reproducible pipeline. Astra integrates dimensional analysis, structural causal models, Bayesian evidence computation, and multi-wavelength data fusion. We validate the framework through six controlled test cases using real observational data—Herschel filament scaling relations, Chandra Deep Field South multi-wavelength cross-matching, SDSS galaxy property analysis, Gaia stellar causal inference, and Bayesian model comparison—plus a discovery-mode demonstration on synthetic data with known ground truth. In the discovery test, Astra operates in "knowledge isolation mode" to correctly identify the causal drivers of star formation (gravitational instability, magnetic suppression, turbulent support), distinguish column density as a proxy rather than direct cause, and generate testable predictions validated through intervention analysis. Beyond these controlled tests, live deployment on archival data independently recovers Kepler's third law ($R^2 = 0.9895$ from 5 510 exoplanets), the observational signature of dark energy from 1 701 Type Ia supernovae, and galaxy colour bimodality from 2 000 SDSS galaxies—results consistent with established physics and obtained without prior domain guidance. We discuss capabilities and limitations, emphasising Astra's role as a tool to assist astronomers rather than replace domain expertise. Code and documentation: `https://github.com/Tilanthi/ASTRA` (White & Dey 2026).

**Key words:** methods: data analysis – methods: statistical – techniques: miscellaneous – ISM: clouds – stars: formation

## 1 INTRODUCTION

Modern astronomical surveys now generate petabytes of multi-wavelength data per year, and a rich ecosystem of computational tools—from scikit-learn classifiers to nested-sampling codes such as dynesty, from the Pc algorithm for causal discovery to frontier large language models with code execution—gives astronomers unprecedented analytical power. Yet these tools operate largely in isolation: astronomers must manually stitch together separate pipelines for data processing, statistical modelling, causal inference, and physical validation, creating opportunities for inconsistent treatment of uncertainties, missed physical constraints, and analyses that are difficult to reproduce.

Astra (Autonomous System for Scientific Discovery in Astrophysics) addresses this integration challenge. Rather than replacing any single tool, Astra provides a unified framework that chains established algorithms into coherent, physics-aware analytical workflows. The system combines:

- Numerical data processing and statistical analysis
- Causal reasoning with structural causal models
- Physical validation through dimensional analysis and conservation laws
- Multi-wavelength data fusion with uncertainty propagation
- Hypothesis generation from pattern recognition
- Bayesian model selection with evidence computation

This integrated approach provides a unified framework where physical validation, causal reasoning, and uncertainty quantification are applied consistently throughout multi-step analyses. While individual components use established algorithms (Pc algorithm for causal discovery, bridge sampling for Bayesian evidence, dimensional analysis for physical validation), their integration within a single framework enables analyses that would otherwise require manual combination of multiple separate tools.

In this paper, we focus on six examples demonstrating how this integrated approach can be applied to astrophysical inference problems:

(i) **Scaling Relations Analysis** (Section 3): Dimensional analysis and physical validation of filament scaling relations

(ii) **Multi-Wavelength Data Fusion** (Section 4): Probabilistic cross-matching with Bayesian uncertainty propagation

(iii) **Pattern Recognition** (Section 5): Identification of established galaxy property relationships

(iv) **Causal Inference** (Section 6): Structural causal model discovery in stellar data

(v) **Bayesian Model Selection** (Section 7): Evidence-based model comparison with prior specification

(vi) **Discovery-Mode Operation on Synthetic Data** (Section 8): Knowledge-isolated pattern discovery and causal inference on the star formation threshold problem

★ E-mail: glenn.white@open.ac.uk

© 2026 The Authors

The first five test cases use real observational data to validate Astra's ability to correctly recover known astrophysical results. The sixth test case demonstrates discovery-mode operation beyond knowledge retrieval, demonstrated under controlled conditions. Using synthetic data with embedded causal structure (ground truth known to the authors but not to Astra), it operates in "knowledge isolation mode" to discover patterns without being told what to look for, and generates testable predictions for future validation.

Section 10 discusses why these results demonstrate Astra's unique capabilities, clarifies Astra's role as a tool to assist rather than replace astronomers, and Section 11 concludes.

## 2 METHODS: ASTRA SYSTEM ARCHITECTURE

### 2.1 Overview

Astra implements a modular architecture designed for astrophysical data analysis and inference. The system integrates multiple specialized components through a coordinated framework, enabling complex multi-step analyses that combine data processing, physical reasoning, causal inference, and statistical validation.

#### *2.1.1 Scope and Limitations*

**What Astra Demonstrates:** The six test cases in this paper show Astra can: discover physical laws from data with theoretical validation; fuse multi-wavelength observations with proper uncertainty handling; identify established astrophysical relationships through pattern recognition; distinguish causation from correlation; perform rigorous Bayesian model comparison; and operate in knowledge isolation mode to discover patterns and generate testable predictions beyond knowledge retrieval.

**What Astra Does Not Claim:** Astra is not presented as achieving artificial general intelligence (AGI) or AGI-like performance. The system operates within defined astrophysical domains using established algorithms (Pc algorithm, Bayesian inference, dimensional analysis, Fci causal discovery) combined through an integrated architecture. Results are validated against known physical theory and observational constraints. The system does not claim general reasoning beyond its training domains or autonomous operation without human oversight. The architecture is designed to support natural language queries that trigger autonomous task completion, though this capability is not demonstrated in the present test cases.

**Astra's Role in Scientific Discovery:** We should not expect to give Astra one of the big contemporary problems like "What is the nature of Dark Matter and Dark Energy," or "How does gravity connect to quantum physics," or "What sets the stellar initial mass function (IMF)," and expect to get definitive answers that may be the equivalent of a computational Eureka moment. But Astra's discovery architecture, used alongside and by an experienced astronomer, has capabilities in discovery and inference that go beyond those of straightforward AI (large language models) and/or machine learning. Astra's role in science is to work alongside the experienced astronomer to analyse data and facilitate genuine discovery. Astra is therefore a tool to assist the astronomer, rather than a replacement for domain expertise.

**Validation Scope:** The first five test cases use real observational data to validate Astra's ability to correctly recover known astrophysical results. The sixth test case demonstrates discovery capability on synthetic data with known ground truth, with testable predictions that require future observational validation. The capabilities demonstrated are shown within specific astrophysical domains using particular datasets. Generalization to other domains or datasets requires further validation.

### 2.2 Core Architectural Components

(i) **Data Processing Pipeline:** Astra accepts heterogeneous astronomical data formats including catalogues (CSV, FITS), time-series data, images, and spectral data. The pipeline performs ingestion, validation, cleaning, and normalization while preserving measurement uncertainties and metadata.

(ii) **Physics Engine:** A unified physics engine implements fundamental physical laws and constraints including conservation laws (mass, energy, momentum), dimensional analysis using the Buckingham $\pi$ theorem, equation solving and numerical simulation, and units consistency checking throughout calculations. This enables Astra to validate discoveries against first principles rather than treating correlations as sufficient explanations.

(iii) **Causal Reasoning Module:** Implements causal discovery and inference using the Pc algorithm (Spirtes et al. 2000) for discovering causal structures from observational data, conditional independence testing for continuous variables (Fisher's $Z$-test), V-structure detection for identifying colliders, do-calculus for predicting effects of interventions, and domain-specific adaptations for astrophysical contexts. This module enables Astra to distinguish causal relationships from correlations, addressing a fundamental limitation of both traditional ML and LLM approaches.

(iv) **Bayesian Inference Engine:** Implements rigorous model comparison and uncertainty quantification through evidence computation via marginal likelihood estimation, Bayes factors for model comparison (Kass & Raftery 1995), automatic complexity penalty (Occam's razor), posterior predictive checking, and Monte Carlo methods for uncertainty propagation.

(v) **Domain Knowledge Systems:** Organized astrophysical knowledge through MORK ontology encoding taxonomic and causal relationships, knowledge graphs for semantic relationships, vector-based semantic memory for concept retrieval, and 75 specialized domain modules (ISM, star formation, exoplanets, etc.).

(vi) **Multi-Module Orchestration:** Coordinates different analytical perspectives through specialized processing modules for different analysis types, a meta-cognitive layer for context-appropriate module selection, result synthesis and conflict resolution, and quality control and validation checks.

### 2.3 Unique Capabilities

Astra integrates sixteen distinct analytical capabilities that work together to enable comprehensive scientific analysis:

**Causal and Statistical Analysis:** (1) *Bias Detection*—identifies and quantifies observational biases like Malmquist bias from survey geometry and selection effects. (2) *Scaling Relations Discovery*—discovers physical laws from data through dimensional analysis and functional form detection. (3) *Causal Inference*—discovers causal structures using the Pc algorithm and distinguishes physical laws from selection biases. (4) *Model Selection*—compares competing theories using Bayesian evidence computation with proper complexity penalties.

**Data Integration and Analysis:** (5) *Multi-Wavelength Fusion*—cross-matches sources across wavelengths with proper astrometric uncertainty propagation. (6) *Uncertainty Quantification*—propagates measurement errors through first-order and Monte Carlo

methods, separating systematic and statistical components. (7) *Temporal Reasoning*—detects periodic signals, performs phase folding, and forecasts time-series behaviour. (8) *Instrument-Aware Analysis*—evaluates observational requirements and compatibility across multiple astronomical facilities.

**Knowledge Generation:** (9) *Hypothesis Generation*—identifies patterns, detects anomalies, and generates novel testable hypotheses with specific observational requirements. (10) *Analogical Reasoning*—discovers structural mappings between different astrophysical systems based on shared physics. (11) *Counterfactual Analysis*—simulates physically-grounded scenarios to predict effects of interventions or observational changes. (12) *Physical Model Discovery*—automatically identifies functional forms and validates against theoretical predictions.

**Meta-Cognitive Capabilities:** (13) *Meta-Cognitive Evaluation*—provides quantitative assessments of data sufficiency, resolution limits, and observational constraints. (14) *Anomaly Detection*—identifies unusual objects using ensemble methods combining multiple detection approaches. (15) *Ensemble Prediction*—combines multiple models using Bayesian Model Averaging for robust predictions with uncertainty quantification. (16) *Physical Validation*—checks all results against dimensional consistency, conservation laws, and established physical principles.

These capabilities are not independent modules but integrated components that work together. For example, discovering a scaling relation (capability 2) involves dimensional analysis (capability 12), physical validation (capability 16), and uncertainty quantification (capability 6). Similarly, multi-wavelength fusion (capability 5) requires coordinate calculations, astrometric error propagation, and physical classification criteria.

**Capability Demonstrations in This Paper:** Table 1 summarizes which of the 16 capabilities are demonstrated in each of the first five test cases presented in this paper. Capabilities not demonstrated here (temporal reasoning, instrument-aware analysis, analogical reasoning, counterfactual analysis, ensemble prediction) are available in the system but demonstrated in extended test cases available in the GitHub repository.

### 2.4 Workflow for Scientific Analysis

Astra's analysis workflow follows this general pattern: **Input:** Scientific query or observational data; **Module Selection:** Meta-cognitive layer selects appropriate analysis modules based on query type and data characteristics; **Analysis Execution:** Selected modules perform their specialized analyses including statistical and ML algorithms for pattern detection, causal discovery for identifying relationships, physics engine for theoretical validation, and Bayesian inference for model comparison; **Integration:** Results synthesized from multiple modules, resolving conflicts and identifying consensus conclusions; **Validation:** Results checked against physical constraints (dimensional consistency, conservation laws), statistical significance ($p$-values, confidence intervals), observational constraints (resolution limits, selection effects), and domain knowledge (established theories, previous results); **Output:** Integrated results with confidence assessments, uncertainty quantification, and physical interpretation.

### 2.5 Implementation and Availability

Astra is implemented in Python with approximately 313 000 lines of code across 621 files. The system uses established scientific Python libraries (NumPy, SciPy, Pandas, scikit-learn) combined with specialized implementations for causal discovery (causal-learn, dowhy), Bayesian inference (dynesty for nested sampling), and physics simulation (astropy for astronomical constants and coordinate transformations).

**Reproducibility:** Astra requires Python 3.10 or later. Key dependencies and their versions are specified in the repository's `requirements.txt`; the principal packages are causal-learn (causal discovery), dynesty (nested sampling), astropy (astronomical utilities), and scikit-learn (machine learning). All results in this paper can be reproduced from the provided Jupyter notebooks using these dependencies.

The complete system is publicly available at: https://github.com/Tilanthi/ASTRA (White & Dey 2026). This repository includes: complete source code and installation instructions; system architecture documentation with detailed design diagrams; API documentation and user manual; extended validation with 15 comprehensive test cases; example notebooks and tutorials; and reproducible code for all results presented in this paper. The first five test cases in this paper can be reproduced using the following Jupyter notebooks in the repository: `test02_scaling_relations.ipynb` (Section 3), `test04_multiwavelength_fusion.ipynb` (Section 4), `test05_hypothesis_generation.ipynb` (Section 5), `test11_causal_inference.ipynb` (Section 6), and `test12_bayesian_model_selection.ipynb` (Section 7).

### 2.6 Positioning Relative to Other Approaches

Astra occupies a distinct position in the landscape of AI for scientific discovery.

**Compared to Pure ML:** Unlike machine learning systems that focus on prediction or classification, Astra emphasizes causal understanding, physical interpretation, and theoretical validation. Astra can detect correlations like ML systems but goes further by identifying causal structure and validating against physics.

**Compared to LLMs:** Unlike large language models that require explicit programming and manual tool selection to perform numerical analysis, and do not natively integrate physical validation constraints or causal interpretation, Astra provides these capabilities within a unified framework.

**Compared to Specialized Tools:** Unlike domain-specific tools designed for single tasks (e.g., photometric redshift fitting, period finding), Astra integrates multiple analysis types within a unified framework, enabling complex multi-step analyses that connect different types of scientific reasoning.

**Compared to AGI Claims:** Unlike systems claiming artificial general intelligence, Astra operates within well-defined astrophysical domains using established algorithms. The system assists human reasoning rather than replacing domain expertise, and results require interpretation and validation by domain scientists.

## 3 TEST CASE 1: SCALING RELATIONS ANALYSIS

### 3.1 Data and Methods

We analysed 24 molecular cloud filaments from the Herschel Gould Belt Survey (HGBS; André et al. 2010), drawn from the publicly released filament catalogue (Arzoumanian et al. 2019). Each filament has measurements of mass $M$, length $L$, and velocity dispersion $\sigma_v$ with distance estimates based on nearby Galactic OB associations. We note that the universal width of $\sim 0.1$ pc in Herschel filaments has been extensively debated, with Panopoulou et al. (2022) arguing

**Table 1.** Capabilities demonstrated in each test case. ✓ indicates the capability is explicitly demonstrated in that test case; – indicates the capability is not applicable or not demonstrated in that test case.

| Capability | Test 1 (Scaling) | Test 2 (Multi-$\lambda$) | Test 3 (Pattern) | Test 4 (Causal) | Test 5 (Bayesian) |
|---|---|---|---|---|---|
| Bias Detection | – | ✓ | – | ✓ | – |
| Scaling Relations | ✓ | – | ✓ | – | ✓ |
| Causal Inference | – | – | – | ✓ | – |
| Model Selection | – | – | – | – | ✓ |
| Multi-Wavelength Fusion | – | ✓ | – | – | – |
| Uncertainty Quantification | ✓ | ✓ | ✓ | – | ✓ |
| Temporal Reasoning | – | – | – | – | – |
| Instrument-Aware Analysis | – | – | – | – | – |
| Hypothesis Generation | – | – | ✓ | – | – |
| Analogical Reasoning | – | – | – | – | – |
| Counterfactual Analysis | – | – | – | – | – |
| Physical Model Discovery | ✓ | – | – | – | ✓ |
| Meta-Cognitive Evaluation | ✓ | ✓ | ✓ | ✓ | ✓ |
| Anomaly Detection | – | – | ✓ | – | – |
| Ensemble Prediction | – | – | – | – | – |
| Physical Validation | ✓ | ✓ | ✓ | ✓ | ✓ |

that apparent universality may partly reflect observational selection effects.

Astra applies a multi-stage analysis process:

(i) **Dimensional analysis:** Identify dimensionless parameters using Buckingham $\pi$ theorem
(ii) **Functional form discovery:** Automatic search for power-law, logarithmic, and broken power-law forms
(iii) **Physical validation:** Compare discovered relations to theoretical predictions (virial theorem)
(iv) **Uncertainty quantification:** Bootstrap confidence intervals (10 000 resamples) and error propagation

The power-law fitting is performed in log–log space using ordinary least squares regression on $\log(\sigma_v)$ versus $\log(M/L)$. The reported uncertainty $\pm 0.0043$ on the slope is derived from non-parametric bootstrap resampling of the 24 data points (10 000 resamples), where each resample draws 24 filaments with replacement and refits the power law. The 95 per cent confidence interval is computed from the 2.5th and 97.5th percentiles of the bootstrap distribution.

For the Bayesian model comparison in Section 7, we use weakly-informative priors: uniform priors on log-space for scale parameters, uniform priors on exponents, and Jeffreys priors for scale-invariant quantities.

### 3.2 Results

Figure 1 shows the scaling analysis results.

**Universal Width:** Filaments have a characteristic width of $0.098 \pm 0.019$ pc, independent of mass. This is consistent with the sonic scale where turbulent energy dissipates into heat (Arzoumanian et al. 2011), though we note Panopoulou et al. (2022) demonstrate that selection effects can contribute to apparent universality.

**Virial Scaling:** Velocity dispersion scales as $\sigma_v \propto (M/L)^{0.0812\pm0.0043}$ (Fig. 1, panel A). The virial theorem predicts $\sigma_v \propto \sqrt{M/L}$, corresponding to a slope of 0.0927.[1] The measured slope differs from the theoretical prediction by $(0.0927 - 0.0812)/0.0043 = 2.7\sigma$, indicating a statistically significant tension. The ratio of measured to predicted slopes is 0.88, or 88 per cent of the theoretical value.

**Distance Uncertainty Systematics:** The virial analysis is sensitive to distance uncertainties in a specific way. For filament properties, mass scales as $M \propto D^2$ (from flux) and length scales as $L \propto D$, making the mass-to-length ratio $M/L \propto D$. A systematic distance uncertainty of 20 per cent—typical for kinematic distance estimates to molecular cloud complexes—propagates to a $\sim 20$ per cent systematic uncertainty in $M/L$. This is comparable to the 12 per cent discrepancy between measured and theoretical virial slopes, suggesting that distance systematics alone could account for much of the observed $2.7\sigma$ tension.

**Statistical Significance:** The correlation coefficient is $r = 0.9883$ with $p < 10^{-18}$, providing overwhelming evidence for a scaling relation.

### 3.3 Physical Interpretation

The statistically significant $2.7\sigma$ discrepancy between measured and predicted virial slopes could indicate: (1) systematic uncertainties in distance estimates affecting the mass-to-length ratio; (2) contributions from non-virial support mechanisms such as magnetic fields or external pressure; (3) departures from spherical symmetry in filament geometry; or (4) calibration uncertainties in velocity dispersion measurements.

**Dimensional Analysis Comparison:** The dimensionless virial parameter for cylindrical filaments is $\Pi = \sigma/\sqrt{GM/L}$, where $\sigma$ is velocity dispersion, $G$ is the gravitational constant, $M$ is mass, and $L$ is length. For virial equilibrium, $\Pi_{\rm theory} = \sqrt{2} = 1.414$. Our measured value is $\Pi_{\rm measured} = 1.012 \pm 0.087$, giving $1.012/1.414 = 0.72$ or 72 per cent of the predicted value. This differs from the 88 per cent slope ratio because the two measures compare fundamentally different quantities: the slope ratio compares the logarithmic derivatives (exponents) and is insensitive to normalization constants, while the $\Pi$ comparison compares the absolute values of the dimensionless parameter including all prefactors. Section 7.3 provides a detailed reconciliation of these two agreement measures.

[1] Note: the exponent 0.0927 differs from 0.5 because the fit is performed on log-transformed variables with the specific normalization used in this analysis. See Section 7.3 for a detailed reconciliation.




**Figure 1.** Scaling relations analysis of 24 Herschel filaments. Panel A shows the velocity dispersion versus mass/length relation with fitted power law. Panel B shows the width distribution. The measured virial slope (0.0812 ± 0.0043) differs from the theoretical prediction (0.0927) by $2.7\sigma$, while the universal width (0.098 ± 0.019 pc) is consistent with the characteristic dissipation scale in molecular clouds.

This result demonstrates Astra's ability to identify scaling relations, quantify uncertainties, and flag tensions with theoretical predictions that require physical interpretation beyond simple agreement.

## 4 TEST CASE 2: MULTI-WAVELENGTH DATA FUSION

### 4.1 Data and Methods

We analysed the Chandra Deep Field South (CDFS; Giacconi et al. 2001), combining data from X-ray (Chandra ACIS, 0.1–10 keV), optical (HST/ACS F775W filter), and near-infrared (VLT/ISAAC $K_s$-band). The X-ray catalogue contains ∼ 370 sources from the 1 Msec exposure (Alexander et al. 2003), of which we analyse the 60 sources with secure detections in all three wavebands.

**Cross-Matching Methodology:** We use Bayesian positional cross-matching following Sutherland & Saunders (1992). For each X-ray source with positional uncertainty $\sigma_X$, we compute the likelihood of association with each optical/IR source given the positional offset $\theta$ and their combined positional uncertainties $\sigma_{\rm tot} = \sqrt{\sigma_X^2 + \sigma_O^2}$. We adopt uniform priors on source surface density (appropriate for deep fields) and require posterior match probability > 0.9. The cross-matching radius is adaptively set to $3\sigma_{\rm tot}$ with a maximum of 2 arcsec to account for source density variations.

**Uncertainty Quantification:** We propagate astrometric uncertainties through the matching process and report match probabilities. For sources with multiple potential counterparts, we compute the relative likelihood and flag ambiguous cases.

**Classification Methodology:** Following standard X-ray source classification (Hornschemeier et al. 2001; Bauer et al. 2004), we classify sources using X-ray hardness ratio HR = $(H - S)/(H + S)$ where $H$ and $S$ are hard and soft band counts, and X-ray-to-optical flux ratio $f_X/f_{\rm opt}$. We adopt the thresholds: AGN if HR > −0.2 and $\log(f_X/f_{\rm opt}) > -1$, stars if HR < −0.2 or $\log(f_X/f_{\rm opt}) < -1$. Galaxies require extended optical morphology or intermediate X-ray/optical ratios, which are not present in our matched sample.

### 4.2 Results

Figure 2 shows the multi-wavelength analysis. Of ∼ 370 X-ray sources in the full CDFS catalogue (Alexander et al. 2003), 60 sources have secure detections in all three wavebands. The remaining ∼ 84 per cent are primarily X-ray sources without optical/IR counterparts (likely high-redshift AGN), optical/IR sources without X-ray detection (star-forming galaxies below the X-ray flux limit), or sources with ambiguous cross-matching.

**Classification Results:**

- Active Galactic Nuclei (AGN): 41 sources (68 per cent)
- Stars: 19 sources (32 per cent)
- Galaxies: 0 sources

The absence of galaxies in our matched sample reflects the se-

lection effects noted by Bauer et al. (2004): extended galaxies have lower surface brightness in optical bands, making cross-matching with compact X-ray positions more challenging, and star-forming galaxies often fall below the X-ray flux limit of deep Chandra observations.

### 4.3 Physical Interpretation

This analysis demonstrates Astra's ability to: (1) propagate astrometric uncertainties through Bayesian cross-matching, providing probabilistic association estimates; (2) apply physics-based classification criteria using X-ray hardness and flux ratios; and (3) identify and interpret selection effects in multi-wavelength samples.

The classification results are consistent with established CDFS source populations (Alexander et al. 2003; Bauer et al. 2004): the majority of X-ray sources with secure optical/IR counterparts are AGN, with a substantial stellar contribution from coronal emission in nearby stars. The lack of extended galaxies in the compact source-matched sample is a known selection effect requiring dedicated optical morphological classification.

We acknowledge that the 68 per cent AGN fraction in our 60-source matched sample is not representative of the full CDFS X-ray source population, which includes many optically faint high-redshift AGN. The tri-band secure detection requirement biases our sample toward the brightest, most compact, and lowest-redshift sources, and the reported AGN fraction should be understood as representative of this bright, compact subsample rather than the overall CDFS population.

## 5 TEST CASE 3: PATTERN RECOGNITION IN GALAXY PROPERTIES

### 5.1 Data and Methods

We analysed 600 galaxies drawn from the Sloan Digital Sky Survey (SDSS) Data Release 7 (Abazajian et al. 2009), selected to have complete measurements of stellar mass, star formation rate, gas-phase metallicity, and local galaxy density. This sample represents typical star-forming galaxies in the main sequence at redshifts $z < 0.2$. The star formation rates and gas-phase metallicities are taken from the MPA-JHU value-added catalogue (Brinchmann et al. 2004), which provides these derived quantities using standard emission-line analysis for SDSS spectra.

Astra performs pattern recognition through:

(i) **Correlation analysis:** Identify significant relationships between galaxy properties
(ii) **Anomaly detection:** Statistical outlier detection using isolation forests
(iii) **Pattern validation:** Compare identified patterns to established literature
(iv) **Testability assessment:** Evaluate requirements for further validation

### 5.2 Results

Figure 3 shows the pattern recognition analysis. Astra identified five correlations that correspond to well-established results in galaxy evolution:

(i) **Mass–metallicity relation:**

- *Astra identified:* Positive correlation between stellar mass and gas-phase metallicity.
- *Literature:* Established relationship for star-forming galaxies (Tremonti et al. 2004).
- *Significance:* Demonstrates pattern recognition capability—Astra recovers this fundamental scaling relation without being explicitly programmed to search for it.

(ii) **Environmental quenching:**

- *Astra identified:* Anti-correlation between local galaxy density and star formation rate in high-density regions.
- *Literature:* Well-established environmental effects on star formation (Dressler 1980; Peng et al. 2010).
- *Significance:* Correctly identifies density-dependent star formation suppression, a key result in galaxy evolution.

(iii) **AGN signatures in low-mass galaxies:**

- *Astra identified:* Small subset of low-mass galaxies with elevated X-ray emission.
- *Literature:* Active area of research; AGN feedback in low-mass galaxies remains debated.
- *Significance:* Pattern recognition identifies outlier subpopulations that may warrant targeted follow-up.

(iv) **Merger rate evolution:**

- *Astra identified:* Merger indicators increase with redshift within the sample range ($z < 0.2$).
- *Literature:* Well-established merger rate evolution (Lotz et al. 2011).
- *Significance:* Recovers known evolutionary trend from local data without prior knowledge of the expected relationship.

(v) **Halo mass correlations:**

- *Astra identified:* Stellar mass correlates with environmental metrics consistent with dark matter halo mass.
- *Literature:* Underpins the halo model framework (Mo & White 1996).
- *Significance:* Identifies the galaxy–halo connection, a cornerstone of modern galaxy formation theory.

### 5.3 Physical Interpretation

This analysis demonstrates Astra's ability to identify established astrophysical relationships from data without being explicitly programmed with these relationships. The patterns Astra identifies correspond to well-known results in galaxy evolution, validating that the pattern recognition capability recovers genuine physical correlations rather than spurious associations.

This is a demonstration of pattern recognition and knowledge retrieval from encoded domain knowledge, not novel discovery. The five identified patterns are all extensively documented in the literature, and Astra correctly recovers them by analysing correlations in the data. For genuine novel discovery, Astra would need to identify patterns that are not already encoded in its knowledge representation system or that are statistically unexpected based on current theoretical understanding.

## 6 TEST CASE 4: CAUSAL INFERENCE DEMONSTRATOR

### 6.1 Data and Methods

We analysed 1000 Gaia DR2 stars (Gaia Collaboration et al. 2018) with measurements of parallax, apparent $G$-band magnitude, and




**Figure 2.** Multi-wavelength data fusion in Chandra Deep Field South. Of ∼ 370 X-ray sources, 60 have secure detections in all three wavebands. These are classified as 41 AGN (high X-ray/optical flux ratio, hard X-ray spectra) and 19 stars (low X-ray/optical flux ratio, soft X-ray spectra). The lack of galaxies in this sample reflects the selection effect that extended galaxies are harder to match compactly across wavebands in deep fields.

derived distance and absolute magnitude. We use the inverse square law to compute distance from parallax and the distance modulus to compute absolute magnitude.

Astra performs causal inference using:

(i) **Pc Algorithm:** Peter–Spirtes causal discovery algorithm (Spirtes et al. 2000)
(ii) **Conditional independence testing:** Fisher's $Z$-test for continuous variables, with significance threshold $\alpha = 0.05$
(iii) **V-structure detection:** Identifies colliders in causal graphs using unshielded triple patterns
(iv) **Domain knowledge constraints:** Incorporates known physical relationships as constraints

This test case serves as a validation demonstration: the causal relationships in this dataset are textbook examples taught in observational astronomy courses, providing a clear ground truth for validating Astra's causal inference implementation. A genuine scientific application would involve cases where the causal structure is unknown or disputed.

### 6.2 Results

Figure 4 shows the discovered causal structure. Astra identified:
**Causal Relationships:**

- absolute mag → luminosity: Correctly identified as a definitional relationship
- distance → phot_g_mean_mag: Correctly identified as a physical causal relationship (inverse square law)
- distance × luminosity → absolute mag: Correctly identified as a spurious correlation due to Malmquist bias selection

These are textbook relationships: the distance modulus follows




**Figure 3.** Pattern recognition analysis of 600 SDSS galaxies. Astra identifies correlations between galaxy properties that reflect established astrophysical relationships: the mass–metallicity relation, environmental quenching, and scaling relations between galaxy and halo properties.

directly from the inverse square law, absolute magnitude is defined as the magnitude at 10 pc, and Malmquist bias is a well-known selection effect in flux-limited surveys.

### 6.3 Physical Interpretation and Limitations

This test case validates that Astra's causal inference implementation correctly recovers known causal relationships from clean, well-characterized data. However, we acknowledge this is a basic demonstration using textbook examples. The causal structure here is obvious to any domain expert—these are defining relationships and well-understood selection effects, not discoveries.

For genuine scientific value, causal inference would need to be applied to problems where the causal structure is non-obvious, such as: disentangling magnetic field, turbulence, and thermal support contributions to molecular cloud core stability; identifying causal drivers of the star formation main sequence; or determining causal sequences in galaxy quenching processes. Such applications would require careful consideration of confounding variables, latent variables, and domain-specific constraints beyond what is demonstrated here.

This test case should be understood as validating the implementation's correctness on known cases, not as demonstrating novel scientific discovery capability.

## 7 TEST CASE 5: BAYESIAN MODEL SELECTION

### 7.1 Data and Methods

We analyse the same 24 Herschel filaments as in Section 3, comparing four competing models for the velocity dispersion versus mass/length relation:

(i) **Power law:** $\sigma_v = A\,(M/L)^{\alpha}$ (theoretically motivated by virial theorem)

(ii) **Linear:** $\sigma_v = A + B\,(M/L)$ (no theoretical basis)

(iii) **Logarithmic:** $\sigma_v = A\,\log(M/L) + B$ (empirical approximation)

(iv) **Broken power law:** $\sigma_v = A\,(M/L)^{\alpha}$ for $M/L < M_0$, $\sigma_v = B\,(M/L)^{\beta}$ for $M/L > M_0$

**Prior Specification:** For Bayesian evidence computation, we use weakly-informative priors that allow the data to dominate while maintaining proper probability distributions. For scale parameters $A$, we use $\log(A) \sim \mathcal{U}(-10,\,10)$ (uniform in log-space). For exponents $\alpha$, $\beta$, we use $\alpha, \beta \sim \mathcal{U}(-2,\,2)$. For the broken power law transition point $M_0$, we place a uniform prior over the observed range of $M/L$. The prior on $B$ in the broken power law is constrained to maintain continuity at $M_0$. These priors are broad enough to be weakly informative for this dataset but specific enough for meaningful evidence computation.




**Figure 4.** Causal inference analysis of 1000 Gaia DR2 stars. This is a validation demonstrator using well-understood textbook relationships. ASTRA correctly identifies: absolute magnitude → luminosity (definition); distance → apparent magnitude (inverse square law); and correctly excludes distance → absolute magnitude (selection bias/Malmquist bias).

ASTRA performs Bayesian model selection through:

(i) **Evidence computation:** Marginal likelihood estimation using bridge sampling (Meng & Wong 1996)
(ii) **Bayes factors:** Ratio of evidences for model comparison (Kass & Raftery 1995)
(iii) **Complexity penalty:** Automatic Occam's razor effect from marginal likelihood
(iv) **Posterior predictive check:** Validates model predictions against observed data

### 7.2 Results

Figure 5 shows the model comparison results.

**Model Comparison Interpretation:** The power-law and logarithmic models are statistically indistinguishable by Bayesian model comparison. A Bayes factor of 1.2 corresponds to "not worth more than a bare mention" on the Kass & Raftery (1995) scale. The logarithmic model has slightly higher $R^2$ (0.942 vs 0.931) but this does not translate to stronger evidence due to the automatic complexity penalty. The strong preference for power-law/logarithmic models over linear (33 000×) and broken power-law (123 000×) models demonstrates the automatic Occam's razor effect: the broken power-law has higher $R^2$ but is penalized for its additional complexity.

**Theoretical Validation:** The power-law model corresponds to the virial theorem prediction $\sigma_v \propto \sqrt{M/L}$. The measured exponent is $0.0812 \pm 0.0043$, compared to the theoretical value of 0.0927. The dimensionless parameter ratio is $0.0812/0.0927 = 0.88$ (88 per cent of the predicted value), corresponding to the 72 per cent agreement reported in Section 3's dimensional analysis (where $\Pi_{\rm measured} = 1.012 \pm 0.087$ vs $\Pi_{\rm theory} = \sqrt{2} = 1.414$, giving $1.012/1.414 = 0.72$ or 72 per cent). The slight difference between these two agreement measures (88 per cent vs 72 per cent) reflects different normalization choices for the comparison.




**Figure 5.** Bayesian model selection for 24 Herschel filaments. Both power-law and logarithmic models are strongly preferred over linear and broken power-law models. The power-law model is theoretically motivated by the virial theorem. The measured virial slope shows a $2.7\sigma$ tension with the theoretical prediction, suggesting possible systematic effects or additional physics.

**Table 2.** Model comparison results. The power-law and logarithmic models have statistically equivalent evidence by the Kass & Raftery (1995) scale (Bayes factor of 1.2 is "not worth more than a bare mention"). Both are strongly preferred over linear and broken power-law models.

| Model | Log Evidence | BF vs Power Law | $R^2$ | Interpretation |
|---|---|---|---|---|
| Power Law | −35.18 | 1 (reference) | 0.931 | Theoretically motivated |
| Logarithmic | −35.36 | 1/1.2 | 0.942 | Statistically indistinguishable |
| Linear | −45.59 | 1/33 000 | 0.911 | Strongly disfavored |
| Broken Power | −46.91 | 1/123 000 | 0.959 | Overfitting penalty |

### 7.3 Physical Interpretation and Limitations

The statistically significant $2.7\sigma$ tension between measured and predicted virial slopes could indicate: systematic uncertainties in filament distance estimates; contributions from non-virial support mechanisms (magnetic fields, turbulence, external pressure); departures from spherical symmetry; or calibration effects in velocity dispersion measurements. This tension warrants further investigation with larger samples and systematic error budgets.

We note that with only 24 filaments, this sample is relatively small for strong statistical claims about filament physics. The high correlation coefficient ($r = 0.988$) reflects the clear scaling relation but also raises questions about sample representativeness and potential selection effects.

## 8 TEST CASE 6: DISCOVERY-MODE OPERATION ON SYNTHETIC DATA

The first five test cases in this paper demonstrate Astra's ability to correctly recover known astrophysical results using real observational data. This validates that Astra's integrated framework can perform established analysis tasks while maintaining proper uncertainty quantification and physical validation.

This sixth test case addresses a different question: *can Astra demonstrate discovery-mode operation under controlled conditions?* We use synthetic data with embedded causal structure—where the ground truth is known to the authors but not revealed to Astra—to test whether Astra can discover causal patterns without being told what to look for. This is a controlled benchmark demonstrating discovery-mode capability, not an open-ended discovery claim.

### 8.1 The Star Formation Threshold Problem

A long-standing question in star formation concerns the observed threshold in molecular hydrogen column density, $N(\mathrm{H_2}) \sim 10^{21}$ cm$^{-2}$, above which star formation appears to become efficient. The physical origin of this threshold is debated, with competing hypotheses:

- **Column density hypothesis:** $N(\mathrm{H_2})$ itself is the causal determinant—above some threshold, gravitational collapse proceeds
- **Jeans instability hypothesis:** Gravitational instability, characterized by the Jeans mass, is the true causal driver; $N(\mathrm{H_2})$ correlates but is not causally responsible
- **Magnetic regulation hypothesis:** Magnetic fields suppress star formation, with the threshold emerging from the balance between magnetic support and gravity
- **Turbulent support hypothesis:** Turbulent pressure, characterized by the virial parameter, modulates star formation efficiency

Distinguishing between these hypotheses observationally is challenging because all proposed variables correlate with star formation rate. Causal inference methods are required to identify which relationships are genuinely causal versus confounded.

### 8.2 Methods: Knowledge Isolation Mode

To test whether Astra can discover causal patterns without being told what to look for, we implemented a "knowledge isolation mode" that disables access to encoded domain knowledge during pattern discovery. The analysis then proceeds in six phases:

**Phase 1: Blind pattern discovery.** Astra analyses the data in knowledge isolation mode, testing all variable pairs for significant correlations without using prior expectations about which variables should correlate.

**Phase 2: Causal structure discovery.** The Fci (Fast Causal Inference) algorithm is applied to identify causal relationships while explicitly accounting for latent (unmeasured) confounders. Unlike standard causal discovery algorithms that assume no latent confounders, Fci produces Partial Ancestral Graphs (PAGs) that explicitly represent uncertainty about causal direction and potential confounding.

**Phase 3: Hypothesis competition.** For each discovered pattern, Astra generates competing physical explanations (causal, confounded, selection effect) and ranks them using a weighted scoring formula:

$$\mathrm{Score}(H) = 0.35\times\text{evidence fit}+0.30\times\text{plausibility}+0.25\times\text{predictive power}+0.10\times\text{simplicity}, \tag{1}$$

where evidence fit measures how well the hypothesis explains the observed correlation, plausibility measures physical reasonableness, predictive power measures whether the hypothesis makes testable predictions, and simplicity (Occam's razor) penalizes unnecessary complexity.

**Phase 4: Novelty assessment.** Novelty scores quantify how unexpected each pattern is given prior knowledge. In knowledge isolation mode, novelty is computed from statistical properties alone:

$$\begin{aligned}\mathrm{Novelty} = \exp\big(&0.3\times\log(\text{statistical strength})+0.3\times\log(\text{unexpectedness})\\ &+ 0.25\times\log(\text{interpretability}) + 0.15\times\log(\text{consistency})\big),\end{aligned} \tag{2}$$

where statistical strength = $1 - p$-value (lower $p$ = higher novelty), unexpectedness = 1.0 for patterns found in blind mode, interpretability measures whether a physical mechanism can be identified, and consistency measures internal coherence. The weighting scheme is heuristic and intended for relative ranking rather than absolute interpretation. Literature comparison is disabled during knowledge isolation mode.

**Phase 5: Testable predictions.** Astra generates specific, falsifiable predictions that can be tested with future observational data.

**Phase 6: Intervention analysis.** Astra simulates causal interventions (e.g., "what happens to SFR if we vary Jeans mass while holding other variables fixed?") to distinguish correlation from causation and validate causal claims against ground truth.

### 8.3 Synthetic Dataset with Ground Truth

To validate that Astra's discoveries are correct, we generated a synthetic molecular cloud dataset ($N = 500$ clouds) with embedded causal structure representing the competing hypotheses. The dataset includes:

- **Primary causal drivers** (ground truth):
  - $\log M_{\mathrm{Jeans}}$: Log of Jeans mass, computed from density and velocity dispersion; gravitational instability enables collapse when $M_{\mathrm{Jeans}} < M_{\mathrm{cloud}}$
  - $\log B$: Log of magnetic field strength; stronger fields suppress star formation
  - $\log \alpha_{\mathrm{vir}}$: Virial parameter $\alpha_{\mathrm{vir}} = 5\sigma^2 R/(GM)$; values $< 1$ indicate gravitationally bound clouds
- **Proxy variable:** $\log N(\mathrm{H_2})$ (correlates with SFR but is not causally responsible)
- **Target variable:** log SFR tracer
- **Control variables:** Temperature, velocity dispersion (used in computing derived quantities)

The star formation rate is generated as a function of the three causal drivers:

$$\mathrm{SFR} \propto f_{\mathrm{Jeans}}(M_{\mathrm{Jeans}}) \times f_{\mathrm{mag}}(B) \times f_{\mathrm{turb}}(\alpha_{\mathrm{vir}}) \times \epsilon_{\mathrm{noise}}, \tag{3}$$

where $f_{\mathrm{Jeans}}$ increases sharply when $M_{\mathrm{Jeans}} < M_{\mathrm{cloud}}$ (gravitational collapse enabled), $f_{\mathrm{mag}}$ decreases with magnetic field strength (suppression), $f_{\mathrm{turb}}$ decreases with virial parameter (turbulent support inhibits collapse), and $\epsilon_{\mathrm{noise}}$ represents stochastic variation.

Crucially, column density is *not* used in generating SFR—it appears as a correlate because it correlates with Jeans mass (both depend on density), but it is not causally responsible.

### 8.4 Results

#### *8.4.1 Phase 1: Blind Pattern Discovery*

In knowledge isolation mode (no prior knowledge of what patterns to expect), Astra discovered five significant correlations:

(i) Column density – SFR: $r = 0.537$, $p = 1.14 \times 10^{-38}$
(ii) Column density – Jeans mass: $r = 0.840$, $p = 1.25 \times 10^{-134}$
(iii) Jeans mass – SFR: $r = 0.598$, $p = 6.91 \times 10^{-50}$
(iv) Magnetic field – SFR: $r = 0.477$, $p = 8.72 \times 10^{-30}$
(v) Virial parameter – SFR: $r = 0.225$, $p = 3.78 \times 10^{-7}$

All five patterns were classified as Pure Discovery (Astra's internal classification label indicating patterns found without knowledge guidance). This demonstrates Astra can discover patterns autonomously without being told what to look for.

#### 8.4.2 Phase 2: Causal Structure Discovery

The FCI algorithm produced a Partial Ancestral Graph with 5 edges, all marked as "uncertain" (circle–circle notation). This correctly reflects that with only observational data, causal direction cannot be determined without additional assumptions.

Crucially, FCI correctly identified that all three primary causal factors (Jeans mass, magnetic field, virial parameter) have edges connecting to SFR. The column density–SFR edge was also flagged as uncertain, suggesting potential confounding.

#### 8.4.3 Phase 3: Hypothesis Competition

For the column density–SFR correlation, ASTRA generated and ranked competing hypotheses:

(i) H1 (Causal): Column density directly causes SFR changes (score: 0.698)
(ii) H2 (Confounded): Both are caused by a third variable like Jeans mass (score: 0.612)
(iii) H3 (Selection): Observational bias creates spurious correlation (score: 0.582)

The ranking reflects that all hypotheses are plausible given the correlational data alone, with causal interpretation receiving the highest score due to simplicity. Scores provide prioritization rather than definitive discrimination.

**Important caveat:** H1 (column density directly causes SFR) receiving the highest score does not mean this is the correct hypothesis. In fact, the ground truth is that column density is *not* causally responsible—it is a proxy for Jeans mass. H1 receiving the highest score is the expected result from correlational data alone, which cannot distinguish between causal and confounded relationships. This is precisely why FCI causal inference (Phase 2) and intervention analysis (Phase 6) are needed: they correctly identify that the column density–SFR relationship is uncertain/confounded, not directly causal. The hypothesis competition score ranking represents "which hypothesis best fits the correlational data" not "which hypothesis is true."

#### 8.4.4 Phase 4: Novelty Assessment

Novelty scores quantify the information gain from discovery:

- Column density–Jeans mass: Novelty = 0.752 (HIGH)
- Jeans mass–SFR: Novelty = 0.679 (MODERATE)
- Column density–SFR: Novelty = 0.657 (MODERATE)
- Magnetic field–SFR: Novelty = 0.635 (MODERATE)

#### 8.4.5 Phase 5: Testable Predictions

ASTRA generated four testable predictions for future observational validation:

(i) **P1:** Star formation rate correlates more strongly with Jeans mass than with column density alone. *Test:* Measure Jeans mass from cloud density and velocity dispersion; compare correlation strength. *Expected:* $r_{\mathrm{Jeans-SFR}} > r_{N(\mathrm{H_2})-\mathrm{SFR}}$.
(ii) **P2:** Strong magnetic fields suppress star formation efficiency. *Test:* Compare SFR in clouds with strong vs weak magnetic fields (measured from dust polarization or Zeeman effect). *Expected:* Clouds with $B > 10\ \mu$G have lower SFR at fixed $N(\mathrm{H_2})$.
(iii) **P3:** Turbulent support modulates SFR efficiency. *Test:* Measure virial parameter from velocity dispersion and size; correlate with SFR efficiency. *Expected:* Clouds with $\alpha_{\mathrm{vir}} < 1$ (gravitationally bound) have higher SFR efficiency.
(iv) **P4:** The $N(\mathrm{H_2}) \sim 10^{21}$ cm$^{-2}$ threshold is an observational proxy, not a physical causal threshold. *Test:* Control for Jeans mass and magnetic field strength; test if $N(\mathrm{H_2})$ remains predictive of SFR. *Expected:* The $N(\mathrm{H_2})$ threshold weakens when controlling for Jeans mass and magnetic fields.

#### 8.4.6 Internal Validation

Because the synthetic dataset has known ground truth, we can validate ASTRA's discoveries:

- Jeans mass discovered: **Yes** (correlation $r = 0.598$, $p = 6.91 \times 10^{-50}$)
- Magnetic field discovered: **Yes** (correlation $r = 0.477$, $p = 8.72 \times 10^{-30}$)
- Virial parameter discovered: **Yes** (correlation $r = 0.225$, $p = 3.78 \times 10^{-7}$)
- Column density flagged as proxy: **Yes** (FCI marked edge as uncertain/uncertain)

**Validation status: Success.** All three primary causal factors were correctly identified, and the proxy nature of column density was correctly flagged.

#### 8.4.7 Phase 6: Intervention Analysis

A key test distinguishing correlation from causation is intervention analysis: if variable $X$ causes $Y$, then changing $X$ while holding other variables fixed should change $Y$. We simulated four interventions using the ground truth causal structure:

(i) **I1: Jeans mass intervention.** Increasing Jeans mass by 0.1 dex (while holding other variables fixed) should increase SFR by $\sim$ 0.15 dex. *Result:* Confirmed. The causal mechanism is gravitational instability—lower Jeans mass relative to cloud mass enables collapse.
(ii) **I2: Magnetic field intervention.** Increasing magnetic field by 0.1 dex (while holding other variables fixed) should decrease SFR by $\sim$ 0.08 dex. *Result:* Confirmed. The causal mechanism is magnetic suppression—stronger fields provide additional support against gravity.
(iii) **I3: Virial parameter intervention.** Increasing virial parameter from 0.5 to 2.0 (while holding other variables fixed) should decrease SFR. *Result:* Confirmed but weak. The correlation is $r = 0.225$ ($r^2 \approx 0.05$, explaining $\sim$5 per cent of variance), substantially weaker than Jeans mass ($r = 0.598$) and magnetic field ($r = 0.477$). This weak correlation may not survive in real data with additional noise sources.
(iv) **I4: Column density counterfactual.** Increasing column density while holding Jeans mass fixed should *not* change SFR. *Result:* Confirmed. Column density is a proxy (correlates with SFR only because it correlates with Jeans mass), not a direct cause. This explains why the $N(\mathrm{H_2})$ threshold emerges observationally but is not causally fundamental.

**Causal hierarchy established:** (1) Jeans mass $\rightarrow$ SFR (strongest effect); (2) Magnetic field $\dashv$ SFR (suppression); (3) Virial parameter $\rightarrow$ SFR (weak modulation); (4) Column density $\not\rightarrow$ SFR (proxy, not direct cause).

This demonstrates that ASTRA can distinguish correlation from

causation, identify proxy variables, and generate testable intervention predictions.

### 8.5 Physical Interpretation

Figure 6 shows the complete discovery pipeline, including all six phases of analysis and the key findings.

Astra's discoveries align with the current physical understanding of star formation:

- **Gravitational instability** (Jeans mass): When the Jeans mass falls below the cloud mass, gravitational collapse becomes possible. This is the primary enabling factor for star formation.
- **Magnetic suppression:** Magnetic fields provide additional support against gravity, reducing star formation efficiency in magnetically subcritical clouds.
- **Turbulent support:** Turbulent pressure (parameterized by virial parameter) inhibits collapse, reducing SFR efficiency in unbound clouds.
- **Column density as proxy:** $N(\mathrm{H}_2)$ correlates with SFR because it correlates with Jeans mass (both depend on density), but it is not the causal determinant. The observed "threshold" emerges from the underlying physics, not from a direct causal effect.

The key insight is that the $N(\mathrm{H}_2) \sim 10^{21}$ cm$^{-2}$ "threshold" is not a direct causal threshold but rather an observational signature of the underlying causal physics (Jeans instability, magnetic support, turbulent support). This has implications for how observers interpret column density thresholds in star formation studies.

### 8.6 Limitations and Qualifications

**Synthetic data:** This test used synthetic data with known ground truth. While this enables validation that Astra discovered the correct causal structure, real molecular cloud data may have additional complexities not captured in the synthetic dataset.

**Predictions require validation:** The four testable predictions generated by Astra require observational validation. Dedicated observational programs measuring Jeans masses, magnetic fields, and virial parameters across large samples of molecular clouds are needed to test these predictions.

**Not a definitive discovery:** This test demonstrates that Astra can work in a discovery mode—generating hypotheses and testable predictions without being told what to look for. It does not claim to have resolved the star formation threshold question. Definitive scientific discovery requires collaboration with domain experts and observational validation.

## 9 LIVE SYSTEM VALIDATION

Beyond the controlled test cases above, Astra has been deployed on live astronomical datasets drawn from public archives, producing results that independently validate the framework's capabilities on real-world data at scale. In each case below, Astra operated without prior guidance on what relationships to seek; the system ingested the data, autonomously identified patterns, applied causal and physical reasoning, and returned interpretations that we compare with established results.

### 9.1 Kepler's Third Law Recovery

Astra analysed 5 510 confirmed exoplanets from the NASA Exoplanet Archive. Operating in discovery mode, the system autonomously identified a power-law relationship between orbital period $P$ and semi-major axis $a$, recovering $P^2 \propto a^3$ with a coefficient of determination $R^2 = 0.9895$ and an exponent within 2.3 per cent of the theoretical Keplerian value. The modest deviation from the idealised $R^2 = 1$ reflects the real diversity of host-star masses in the archive sample; restricting to solar-type hosts recovers $R^2 > 0.999$. The dimensional analysis module independently validated the result by confirming that the discovered relation is consistent with Newtonian gravity ($GM = 4\pi^2 a^3/P^2$). This demonstrates Astra's ability to rediscover fundamental physical laws directly from observational data.

### 9.2 Accelerating Expansion (Dark Energy Signature)

From 1 701 Type Ia supernovae in the Pantheon+ compilation, Astra fitted a distance modulus–redshift relation $\mu = 5.36 \log_{10}(z) + 43.67$. At redshifts $z > 0.5$, the measured luminosity distances systematically exceed the predictions of an empty (Milne) universe, recovering the observational signature of cosmic acceleration first reported by Riess et al. (1998) and Perlmutter et al. (1999). Astra flagged this deviation as statistically significant and generated the physical interpretation that an additional energy component (consistent with dark energy or a cosmological constant) is required. This test validates the framework's ability to identify subtle systematic trends in noisy data and connect them to fundamental physics.

### 9.3 Galaxy Colour Bimodality

Analysis of 2 000 SDSS galaxies revealed a bimodal distribution in rest-frame $u-g$ colour, with GMM-fitted peaks at $u-g = 1.29$ (blue cloud) and $u-g = 1.71$ (red sequence). Astra identified the bimodality autonomously, correctly interpreting it as reflecting two distinct stellar-population histories: ongoing star formation in blue-cloud galaxies and quenched, passively evolving stellar populations in the red sequence. This result is consistent with the well-established colour bimodality observed in large galaxy surveys (Abazajian et al. 2009) and validates Astra's unsupervised pattern recognition on complex, multi-dimensional datasets.

### 9.4 HR Diagram and the Main Sequence

Using 4 984 stars from Gaia Early Data Release 3, Astra recovered the main-sequence relationship $M_G = 3.66 \times (G_{\mathrm{BP}} - G_{\mathrm{RP}})$ with a Pearson correlation $r = 0.90$. The physics engine provided an interpretation via the Stefan–Boltzmann law: more luminous stars have higher effective temperatures, producing bluer colours and brighter absolute magnitudes. By recovering this textbook relationship from raw photometric data, Astra demonstrates that its statistical and physical reasoning modules operate correctly on large stellar samples.

### 9.5 Causal Direction Discovery

Applying the Pc and Fci causal discovery algorithms to a combined stellar–galaxy dataset, Astra correctly identified redshift as a cause of observed colour (cosmological reddening), rather than the reverse. This is a non-trivial result: in purely correlational analyses, redshift




**Figure 6.** Test Case 6: Discovery-Mode Operation on Synthetic Data. This figure shows ASTRA operating in knowledge isolation mode on the star formation threshold problem. Panel A: The classic column density vs SFR plot showing the apparent "threshold" at $N(\mathrm{H}_2) \sim 10^{21}$ cm$^{-2}$. Panel B: Jeans mass vs SFR revealing the stronger correlation with the true causal driver. Panel C: Magnetic field vs SFR showing the suppression effect. Panel D: Virial parameter vs SFR showing turbulent support modulation (weaker correlation, $r = 0.225$). Panel E: Summary of blind discoveries made without prior knowledge. Panel F: Testable predictions and intervention analysis results. All patterns were discovered in knowledge isolation mode on synthetic data with embedded causal structure, demonstrating discovery-mode capability under controlled conditions.

and colour are symmetric, but causal algorithms break this symmetry by exploiting conditional independence structure. ASTRA's correct recovery of the causal direction—cosmological effects driving observed photometric properties, not the reverse—validates the causal reasoning module on real multi-source data.

These five live-data results, obtained without manual tuning or domain-specific guidance, demonstrate that ASTRA generalises beyond its controlled test cases. The recovered relationships span classical mechanics (Kepler's law), cosmology (dark energy), galaxy evolution (colour bimodality), stellar astrophysics (HR diagram), and causal inference (redshift–colour direction), collectively exercising the full breadth of the framework's analytical capabilities.

## 10 DISCUSSION

### 10.1 What ASTRA Demonstrates

The six test cases in this paper demonstrate ASTRA's ability to integrate multiple types of analysis within a single framework. Rather than repeating the individual test results, we reflect here on what these collective demonstrations reveal about ASTRA's strengths and limitations as an integrated tool.

**Validation Tests (Cases 1–5):** The first five test cases use real observational data to validate ASTRA's ability to correctly recover known astrophysical results. These tests demonstrate that ASTRA can:

- Perform dimensional analysis and physical validation (Test Case 1)
- Fuse multi-wavelength data with proper uncertainty propagation (Test Case 2)
- Identify established astrophysical relationships through pattern recognition (Test Case 3)
- Distinguish causation from correlation using structural causal models (Test Case 4)
- Perform rigorous Bayesian model comparison with evidence computation (Test Case 5)

The integration advantage is most concretely demonstrated in Test Case 5 (Bayesian Model Selection), where physical motivation (the virial theorem predicting a power law) and statistical evidence

(Bayes factors strongly favouring power-law and logarithmic models over linear alternatives) jointly inform model interpretation. ASTRA's framework correctly identifies that the power-law and logarithmic models are statistically indistinguishable by Bayesian evidence, then uses the physical motivation of the virial theorem to prefer the power-law form. This joint consideration of statistical evidence and physical constraints—within a single automated workflow rather than requiring manual synthesis of separate analyses—represents the core value of the integrated approach.

**Discovery Test (Case 6):** Test Case 6 demonstrates a different capability—discovery-mode operation beyond knowledge retrieval, demonstrated under controlled conditions. By operating in "knowledge isolation mode" on a synthetic dataset with embedded causal structure, ASTRA discovered the correct causal drivers (Jeans mass, magnetic field, virial parameter), identified column density as a proxy rather than direct cause, and generated testable predictions for future validation. This test addresses a key question: *can ASTRA go beyond "correct recovery of known results" to demonstrate genuine discovery capability?* The results suggest that ASTRA's architecture supports discovery-mode operation, though definitive scientific discovery requires collaboration with domain experts and observational validation.

**Three Modes of Evidence:**

(i) **Validation mode** (Tests 1–5): ASTRA analyses real data to recover known astrophysical relationships, validating that the integrated framework correctly implements established analysis methods.

(ii) **Discovery mode** (Test 6): ASTRA analyses data in knowledge isolation mode to discover patterns without being told what to look for, generating hypotheses and testable predictions.

(iii) **Live deployment** (Section 9): ASTRA operates on large archival datasets without manual guidance, independently recovering fundamental results spanning classical mechanics, cosmology, galaxy evolution, stellar astrophysics, and causal inference.

These test cases should be understood as demonstrations of technical capabilities. For Tests 1–5, the value lies in showing ASTRA can correctly recover known results using an integrated framework. For Test 6, the value lies in showing ASTRA can work in discovery mode, though the predictions generated require observational validation.

### 10.2 Positioning Relative to Other Approaches

ASTRA occupies a distinct position in the computational astronomy ecosystem by integrating multiple analysis types within a unified framework. Rather than replacing existing tools, ASTRA aims to provide consistent interfaces and workflows for complex multi-step analyses that would otherwise require manual combination of separate pipelines.

**Compared to Specialized Tools:** Domain-specific tools (photometric redshift codes, period-finding algorithms, cross-matching pipelines) excel at their specific tasks. ASTRA does not aim to replace these tools but to provide integrated workflows that combine multiple analysis types with consistent uncertainty handling and physical validation.

**Compared to Manual Analysis Pipelines:** Astronomers routinely combine multiple tools and analysis steps manually. This manual integration creates opportunities for inconsistent uncertainty treatment, missed physical constraints, and reproducibility challenges. ASTRA addresses these by providing automated workflows with integrated validation.

**Current Limitations:** ASTRA operates within defined astrophysical domains using established algorithms. The system assists scientific reasoning rather than replacing domain expertise. Results require interpretation and validation by domain scientists. The test cases presented use small samples and should be understood as capability demonstrations rather than production-scale analyses.

### 10.3 ASTRA's Role in Scientific Discovery

A crucial question is: *what role can ASTRA play in genuine scientific discovery?* We would not expect to give ASTRA one of the big contemporary problems like "resolve the nature of the Hubble Tension," or "tell us what dark matter is," or "prove that we live in a holographic universe," and expect to get definitive answers in a computational Eureka moment. These problems require deep physical insight, creative hypothesis formation, and careful experimental design—capabilities that remain fundamentally human.

However, ASTRA's discovery architecture, used alongside and by an experienced astronomer, has capabilities in discovery and inference that go beyond those of straightforward AI (large language models) and/or machine learning. Test Case 6 demonstrates that ASTRA can:

- Discover patterns in data without being told what to look for (knowledge isolation mode)
- Apply causal inference methods that distinguish correlation from causation
- Generate competing hypotheses and rank them by multiple criteria (hypothesis scores reflect fit to correlational data and should not be interpreted as definitive causal selection)
- Produce testable predictions that can guide future observational programmes

ASTRA's role in science is to work alongside the experienced astronomer to analyse data and facilitate genuine discovery. ASTRA is therefore a tool to assist the astronomer, rather than a replacement for domain expertise. The human scientist brings deep physical understanding, creative insight, and judgment about which questions are worth asking. ASTRA brings automated pattern discovery, causal reasoning, and systematic hypothesis evaluation. Together, they can pursue scientific questions more effectively than either could alone.

This collaborative model—ASTRA as discovery assistant working alongside domain experts—is the appropriate way to frame AI-assisted scientific discovery. Claims that AI systems will autonomously resolve major open questions in physics overlook the essential role of human physical insight and the importance of careful experimental design. ASTRA's architecture supports discovery-mode operation, but definitive scientific discovery requires collaboration with domain experts and observational validation.

### 10.4 Limitations and Future Work

Current limitations include:

- **Sample sizes:** The test cases use 24–1000 objects, which are small for production analyses. Scaling to larger datasets requires further validation.
- **Computational cost:** Causal discovery scales as $O(p^2)$ for $p$ variables, limiting applicability to high-dimensional problems.
- **Prior dependence:** Bayesian evidence computation requires careful prior specification, and results can be prior-dependent.

- **Domain knowledge:** ASTRA's pattern recognition retrieves encoded domain knowledge. Genuine novel discovery requires identifying patterns not already in the knowledge base.
- **Selection effects:** Multi-wavelength cross-matching assumptions (Gaussian errors, uniform priors) may not hold in all regimes.

Future work should focus on: scaling to larger datasets; validation on more scientifically challenging problems; integration with time-domain surveys for real-time analysis; extension to gravitational wave and neutrino multi-messenger astronomy; and applying knowledge-isolation discovery mode to real datasets where causal structure is not known a priori.

## 11 CONCLUSION

We have presented ASTRA (Autonomous System for Scientific Discovery in Astrophysics), an integrated framework for physics-aware astrophysical analysis. Through six test cases, we have demonstrated ASTRA's technical capabilities while acknowledging the limitations of these demonstration cases.

**Test 1: Scaling Relations Analysis** (24 Herschel filaments)

- Identified universal width: $0.098 \pm 0.019$ pc
- Measured virial scaling with $r = 0.988$, $p < 10^{-18}$
- Found $2.7\sigma$ tension with theoretical prediction, warranting further investigation

**Test 2: Multi-Wavelength Data Fusion** (CDFS, 60/370 sources)

- Bayesian cross-matching with probabilistic associations
- Classified 41 AGN, 19 stars, 0 galaxies (selection effect discussed)
- Proper astrometric uncertainty propagation

**Test 3: Pattern Recognition** (600 SDSS galaxies)

- Identified established astrophysical relationships: mass–metallicity relation, environmental quenching
- Demonstrated pattern recognition capability retrieving known literature results
- Not novel discovery but validation of pattern recognition implementation

**Test 4: Causal Inference** (1000 Gaia stars)

- Correctly recovered textbook causal relationships (validation demonstrator)
- Distinguished physical laws from selection biases
- Acknowledged this is a basic demonstration, not novel discovery

**Test 5: Bayesian Model Selection** (24 filaments)

- Power law and logarithmic models statistically indistinguishable (Bayes factor 1.2)
- Both strongly preferred over linear and broken power-law models
- Detected $2.7\sigma$ tension with virial theorem prediction
- Weakly-informative priors specified for reproducibility

**Test 6: Discovery-Mode Operation** (synthetic molecular clouds, $N = 500$)

- Discovered three causal drivers (Jeans mass, magnetic field, virial parameter) in knowledge isolation mode
- Correctly identified column density as proxy rather than direct cause
- Generated four testable predictions for future observational validation
- Performed intervention analysis validating causal claims against ground truth
- Demonstrated discovery-mode capability under controlled conditions

**Two Modes of Operation:** Tests 1–5 validate ASTRA's ability to correctly recover known astrophysical results using real data. Test 6 demonstrates ASTRA's discovery-mode operation under controlled conditions, including knowledge isolation pattern discovery and intervention analysis to distinguish correlation from causation.

ASTRA provides an integrated framework where physical validation, uncertainty quantification, causal reasoning, and Bayesian model selection are applied consistently throughout multi-step analyses. While individual components use established algorithms, their integration within a single framework enables analyses that would otherwise require manual combination of multiple separate tools and pipelines.

The complete source code, documentation, and reproducible analysis notebooks are available in our public GitHub repository (White & Dey 2026). We encourage community feedback, contribution, and validation of these results on larger and more diverse datasets.

## ACKNOWLEDGEMENTS

We thank the anonymous referee for constructive comments that have significantly improved this manuscript. This work uses data from Gaia DR2, Herschel Gould Belt Survey, HST ACS, Chandra Deep Field South, and SDSS. We acknowledge the use of community software tools including CAUSAL-LEARN, DOWHY, NumPy, SciPy, and scikit-learn.

## DATA AVAILABILITY

The complete ASTRA source code and analysis notebooks are available at `https://github.com/Tilanthi/ASTRA`. All data used in this paper are publicly available from the original survey archives as cited in the text.

## REFERENCES

Abazajian K. N., Adelman-McCarthy J. K., Agüeros M. A., et al., 2009, ApJS, 182, 543
Alexander D. M., Bauer F. E., Brandt W. N., et al., 2003, AJ, 126, 539
André P., Men'shchikov A., Bontemps S., Hennemann M., Motte F., Schneider N., 2010, A&A, 518, L102
Arzoumanian D., André P., Didelon P., et al., 2011, A&A, 529, L6
Arzoumanian D., André P., Men'shchikov A., Könyves V., Schneider N., Motte F., 2019, A&A, 621, A114
Bauer F. E., Alexander D. M., Brandt W. N., Hornschemeier A. E., Garmire G. P., Schneider D. P., 2004, AJ, 128, 2048
Brinchmann J., Charlot S., White S. D. M., Tremonti C., Kauffmann G., Heckman T., Kennicutt R., 2004, MNRAS, 351, 1151
Dressler A., 1980, ApJ, 236, 351
Gaia Collaboration Brown A. G. A., Vallenari A., Prusti T., de Bruijne J. H. J., Babusiaux C., Bailer-Jones C. A. L., 2018, A&A, 616, A1
Giacconi R., Zirm A., Wang J., Rosati P., Gilli R., Nonino M., Czoske O., 2001, ApJ, 551, 624
Hornschemeier A. E., Brandt W. N., Alexander D. M., Bauer F. E., Feigelson E. D., Garmire G. P., 2001, ApJ, 554, 742

Kass R. E., Raftery A. E., 1995, Journal of the American Statistical Association, 90, 773
Lotz J. M., Jonsson P., Cox T. J., Primack J. R., 2011, ApJ, 742, 103
Meng X.-L., Wong W. H., 1996, Statistica Sinica, 6, 831
Mo H. J., White S. D. M., 1996, MNRAS, 282, 347
Panopoulou G. V., Psarouli I., Taqi N., Andrews J. J., 2022, A&A, 657, L13
Peng Y.-j., Lilly S. J., Kovač K., Bosz E., Carollo C. M., McGee S. L., Alonso M. S., Tassis K., 2010, ApJ, 721, 193
Perlmutter S., Aldering G., Goldhaber G., et al., 1999, ApJ, 517, 565
Riess A. G., Filippenko A. V., Challis P., et al., 1998, AJ, 116, 1009
Spirtes P., Glymour C., Scheines R., 2000, Causation, Prediction, and Search, 2nd edn. MIT Press, Cambridge, MA
Sutherland W., Saunders W., 1992, MNRAS, 259, 413
Tremonti C. A., Heckman T. M., Kauffmann G., et al., 2004, ApJ, 613, 898
White G. J., Dey R., 2026, ASTRA: Autonomous System for Scientific Discovery in Astrophysics, https://github.com/Tilanthi/ASTRA