

Numerical Solutions for Ideal MHD

On the essential ingredients of the method

Kyle Webster, Department of Mathematics
MATH 521 Project Presentation



The Ideal MHD Equations

Can be written as conservations of mass, momentum, and energy

$$\partial_t \rho + \nabla \cdot [\rho \mathbf{u}] = 0$$

$$\partial_t (\rho \mathbf{u}) + \nabla \cdot [\rho \mathbf{u} \mathbf{u}^T + p_{\text{tot}} \mathbb{I} - \mathbf{B} \mathbf{B}^T] = 0$$

$$\partial_t E + \nabla \cdot [(E + p_{\text{tot}}) \mathbf{u} - (\mathbf{u} \cdot \mathbf{B}) \mathbf{B}] = 0$$

and an evolution of the magnetic field

$$\partial_t \mathbf{B} + \nabla \cdot [\mathbf{u} \mathbf{B}^T - \mathbf{B} \mathbf{u}^T] = 0$$

with $p_{\text{tot}} = p + \frac{1}{2} |\mathbf{B}|^2$ and $E = \frac{1}{2} \rho |\mathbf{u}|^2 + \frac{1}{2} |\mathbf{B}|^2 + \frac{p}{\gamma - 1}$ where γ is the ratio of specific heats.



The Entire Interest

Consider writing a compact form of Ideal MHD with $\mathbf{U} = (\rho, \mathbf{u}, \mathbf{B}, E)$

$$\partial_t \mathbf{U} + \nabla \cdot \mathbf{F}(\mathbf{U}) = 0$$

and $\mathbf{F}(\mathbf{U})$ the appropriate flux functions.

What are the essentials for simulating these equations?



Discretization & Constraints

Discretization

- Space \longrightarrow Discontinuous Galerkin with Numerical Flux
- Time \longrightarrow Runge-Kutta / Linear Multistep

Constraint

- 1D \longrightarrow Constant B_x
- Multi-D $\longrightarrow \nabla \cdot \mathbf{B} = 0 \longrightarrow$ Projection Method



1D 1st Order Discontinuous Galerkin Method

Consider 1-dimensional Ω so that $\Omega = \cup_j I_j$

Find $U \in V_h^k$ such that

$$\frac{\partial}{\partial t} \int_{I_j} U v_h \, dx - \int_{I_j} f(U) \frac{\partial}{\partial x} v_h \, dx + f_{j+\frac{1}{2}} v_{h,j+\frac{1}{2}}^- - f_{j-\frac{1}{2}} v_{h,j-\frac{1}{2}}^+ = 0$$

for all $v_h \in V_h^k$ and for all j .


$$\frac{\partial}{\partial t} U_j^h = \frac{1}{\Delta x_j} \left(f_{j-\frac{1}{2}} - f_{j+\frac{1}{2}} \right)$$



HLL Numerical Flux

- Single Intermediate State Riemann Solver
- s_{\pm} from eigenvalues of flux Jacobian

$$A_{\rho} = \begin{bmatrix} u_1 & \rho & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & u_1 & 0 & 0 & \frac{-B_1}{\rho} & \frac{B_2}{\rho} & \frac{B_3}{\rho} & \frac{1}{\rho} \\ 0 & 0 & u_1 & 0 & \frac{-B_2}{\rho} & \frac{-B_1}{\rho} & 0 & 0 \\ 0 & 0 & 0 & u_1 & \frac{-B_3}{\rho} & 0 & \frac{-B_1}{\rho} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & B_2 & -B_1 & 0 & -u_2 & u_1 & 0 & 0 \\ 0 & B_3 & 0 & -B_1 & -u_3 & 0 & u_1 & 0 \\ 0 & \gamma p & 0 & 0 & \bar{\gamma} \mathbf{u} \cdot \mathbf{B} & 0 & 0 & u_1 \end{bmatrix}$$



$$\tilde{f}(U^-, U^+) = \frac{s_+ f_- - s_- f_+ + s_+ s_- (U^+ - U^-)}{s_+ - s_-}$$

2nd Order Adams-Bashforth

- Initialized with single FE step





$$U^{n+1} = U^n + \frac{\Delta t}{2} (F^n - F^{n-1})$$

$$U^{n+1} = U^n + \Delta t F^n$$



- CFL Condition

$$\Delta t \leq \frac{\Delta x}{s_{max}}, \quad s_{max} = \max_j (\max(|\lambda_-(U_j)|, |\lambda_+(U_j)|))$$



Questions?

