

## Viscoelastometric-guided early fibrinogen concentrate replacement during postpartum haemorrhage: OBS2, a double-blind randomized controlled trial

P. W. Collins<sup>1,\*</sup>, R. Cannings-John<sup>2</sup>, D. Bruynseels<sup>3</sup>, S. Mallaiah<sup>4</sup>, J. Dick<sup>5</sup>, C. Elton<sup>6</sup>, A. D. Weeks<sup>7</sup>, J. Sanders<sup>8</sup>, N. Aawar<sup>2</sup>, J. Townson<sup>2</sup>, K. Hood<sup>2</sup>, J. E. Hall<sup>9</sup> and R. E. Collis<sup>3</sup> on behalf the OBS2 study team<sup>†</sup>

<sup>1</sup>Institute of Infection and Immunity, School of Medicine Cardiff University, UK, <sup>2</sup>Centre for Trials Research, College of Biomedical and Life Sciences, Cardiff University, UK, <sup>3</sup>Department of Anaesthetics and Pain Control, Cardiff and Vale University Health Board, UK, <sup>4</sup>Tom Byson Department of Anaesthesia, Liverpool Women's Hospital, Liverpool, UK, <sup>5</sup>Department of Anaesthetics, University College Hospital London, UK, <sup>6</sup>Department of Anaesthetics, Leicester Royal Infirmary, Leicester, UK, <sup>7</sup>Department of Women's and Children's Health, Institute of Translational Medicine, University of Liverpool, Liverpool, UK, <sup>8</sup>School of Healthcare Sciences, Cardiff University, Cardiff, UK and <sup>9</sup>Department of Anaesthetics and Pain Control, School of Medicine Cardiff University, Heath Park, UK

\*Corresponding author. E-mail: peter.collins@wales.nhs.uk

<sup>†</sup>The OBS2 study team is listed in the Acknowledgements section.

# METHODS

- MC, DB, placebo controlled trial set in teaching hospital obstetric units
- women with PPH 1500-2000ml enrolled
- excluded placenta accreta, suspected AFE, if declined transfusion
- if Fibtem A5 was  $\leq 15\text{mm}$  and bleeding continued, subjects randomised to FC or placebo
- primary outcome: # PRC, FFP, cryo and platelets transfused