

AUTHORS' MANAGEMENT

- Cardiff
- MOH protocol —> FIBTEM, POCT Hb, FBC, lab coagulation screen
- Management based on FIBTEM results as in flowchart




Figure 1 Point-of-care FIBTEM algorithm used at The University Hospital of Wales in 2014. PPH, postpartum haemorrhage; FFP, fresh frozen plasma.

Viscoelastometric-guided early fibrinogen concentrate replacement during postpartum haemorrhage: OBS2, a double-blind randomized controlled trial

P. W. Collins^{1,*}, R. Cannings-John², D. Bruynseels³, S. Mallaiah⁴, J. Dick⁵, C. Elton⁶, A. D. Weeks⁷, J. Sanders⁸, N. Aawar², J. Townson², K. Hood², J. E. Hall⁹ and R. E. Collis³ on behalf the OBS2 study team[†]

¹Institute of Infection and Immunity, School of Medicine Cardiff University, UK, ²Centre for Trials Research, College of Biomedical and Life Sciences, Cardiff University, UK, ³Department of Anaesthetics and Pain Control, Cardiff and Vale University Health Board, UK, ⁴Tom Byson Department of Anaesthesia, Liverpool Women's Hospital, Liverpool, UK, ⁵Department of Anaesthetics, University College Hospital London, UK, ⁶Department of Anaesthetics, Leicester Royal Infirmary, Leicester, UK, ⁷Department of Women's and Children's Health, Institute of Translational Medicine, University of Liverpool, Liverpool, UK, ⁸School of Healthcare Sciences, Cardiff University, Cardiff, UK and ⁹Department of Anaesthetics and Pain Control, School of Medicine Cardiff University, Heath Park, UK

*Corresponding author. E-mail: peter.collins@wales.nhs.uk

[†]The OBS2 study team is listed in the Acknowledgements section.