

Facile exchange of arsenic between adducts and  
implications to drug discovery

By Yuxuan Gu

A thesis submitted to McGill University in partial fulfillment of the requirements  
for the degree of:

MASTER OF SCIENCE

Department of Chemistry, Faculty of Science

McGill University

Montréal, Québec, Canada

©Yuxuan Gu 2013



## Abstract

Arsenic is a well-known poison that lives a double life as a therapeutic agent. Recently, arsenic trioxide has been used to treat the cancer acute promyelocytic leukemia with high efficacy. Unfortunately, arsenic drug development has been road blocked by a lack of understanding of the mechanisms of interaction. This thesis attempts to elucidate these mechanisms by looking at the reaction kinetics and thermodynamics of various arsenic species, including potent new organic arsenicals such as S-dimethylarsino-glutathione (ZIO-101) and S-dimethylarsino-cysteine (DMAC).

Data shows that rapid thiolate exchange of dimethylarsonim,  $\text{Me}_2\text{As}^+$ , occurs when the two compounds are dissolved in aqueous solution. The equilibrium constants of this interthiol transfer were characterised by the integrals of the diastereotopic methylarsonium resonances of the species in  $^1\text{H}$  NMR. Dynamic NMR was used to characterise rapid intra molecular conformational dynamics which lead to the coalescence of diastereotopic methyl resonances. In addition, rapid thiolate exchange was also shown to occur in monomethylarsonium species as well.

The discovery and characterisation of these facile arsenium transfers allows us to think of the arsenicals in a different manner, instead of binding statically to vicinal thiols, arsenic could hop around various thiolates around the cell.

## Abstrait

L'Arsenic est un poison bien connu qui mène une double vie en tant qu'agent thérapeutique. Récemment, le trioxide d'arsenic a été utilisé pour le traitement du cancer de la leucémie aiguë promyélocytaire avec une grande efficacité. Malheureusement, le développement des médicaments basé sur l'arsenic connaît un ralentissement à cause du manque de compréhension de leurs mécanismes d'interaction. Cette thèse tente d'élucider ces mécanismes en regardant les réactions cinétiques et thermodynamiques de plusieurs espèces de molécules contenant de l'arsenic, incluant de nouveaux composés organique arsenicales comme le S-diméthylarsinoglutathione (ZIO-101) and S-diméthylarsino-cysteine (DMAC).

Les données démontrent que l'échange rapide du thiolate de diméthylarsonium,  $\text{Me}_2\text{As}^+$ , survient lorsque les deux composés sont dissous en solution aqueuse. Les constantes d'équilibres de cet échange interthiole ont été caractérisées par les intégraux des résonances diastéréotopiques du méthylarsonium des espèces en conformation dynamique intramoléculaire qui ont mené à la coalescence des méthyles diastéréotopiques en résonance des espèces en RMN de  $^1\text{H}$ . La RMN dynamique a été utilisée pour caractériser les conformations dynamiques intramoléculaires rapides qui ont menées à la coalescence des résonances des méthyles diastéréotopiques. De plus, il a été démontré que

l'échange rapide des thiolate survient aussi dans les espèces de monométhylarsoniums.

La découverte et caractérisation de ces échanges faciles d'espèces d'arseniums nous force à voir ces arsenicales d'un autre œil, au lieu de se lier statiquement aux thiols vicinaux, l'arsenic pourrait sauter entre différents thiolates autour de la cellule.

## Acknowledgements

I would like to thank my supervisor Prof D. Scott Bohle, for giving endless support, patience and guidance – even at times when I may be difficult to work with. Dr. Bohle has gone well beyond his role as a supervisor provided me with important life guidance and support. I would like to give special thanks and gratitude to Dr. Inna Perepichka for always being there. Special thanks also goes to ZhiJie Chua who has been my alter ego. My gratitude also goes to Tina Lam for the French translation of the abstract. Great thanks to past and present members of the Bohle research group: Erin Dodd, Kristopher Rosadiuk, Joël Poisson, Andrea Hill, Ivor Wharf, Mirna Veruca, Cheryl Bain, and Laura Brothers.

I would like to thank my parents, Maggie So and my friends for being supportive.

The following staff of the Department of Chemistry, McGill University: Dr Fred Morin, Dr Nadim Saade, Rick Rossi, Weihua Wang, Jean-Philippe Guay and Chantal Marotte own my gratitude for providing their expertise and prompt assistance during the course of my studies.

To end, thanks to McGill University and the supporting grants for funding my doctoral studies throughout.

### **Contributions of Authors**

This dissertation includes content which has been published: Bohle, D. S.; Gu, Y., Facile dimethylarsenic exchange and pyramidal inversion in its cysteine and glutathione adducts. *Organic & Biomolecular Chemistry* 2013, 11 (16), 2578-2581.

I hereby give copyright clearance for the inclusion of the following papers, of which I am a co-author, in the dissertation of Yuxuan Gu.

Bohle, D. S.; Gu, Y., Facile dimethylarsenic exchange and pyramidal inversion in its cysteine and glutathione adducts. *Organic & Biomolecular Chemistry* 2013, 11 (16), 2578-2581.

D. Scott Bohle

Professor and CRC Chair in Chemistry

Department of Chemistry

McGill University

801 Sherbrooke St West

Montreal

QC H3A 2K6

**Statement of originality and contribution to knowledge**

The author performed all work outlined in this thesis, including all the work presented in the paper as specified above, under the supervision of Professor D. Scott Bohle. All work presented in this thesis, with the exception of the introductory literature review, is declared by the author to be original scholarship and distinct contributions to knowledge as is mandatory for doctoral theses.



## Table of Contents

Table of Contents .....	ix
Table of Figures.....	xi
Chapter 1 .....	1
Thesis Introduction .....	1
1.1 The Bioinorganic Chemistry of Arsenic.....	1
1.1.1 Metabolism of arsenic.....	3
1.1.2 Arsenic as a poison.....	6
1.1.3 Treatment of arsenic poisoning .....	8
1.1.4 Medicinal applications of arsenic.....	9
1.2 Chemistry of Arsenic compounds .....	12
1.2.1 Redox of arsenic by GSH .....	12
1.2.2 Interaction of arsenic with thiols.....	15
1.2.3 Lability of the Arsenic Thiol bond.....	18
1.2.4 Co-ordination and geometry of arsenic compounds.....	19
1.2.5 Mechanism of arsenic bond lability.....	20
1.3 Conclusions.....	22
References .....	23
Chapter 2: .....	28
2.1 Kinetics and thermodynamics of rapid Me <sub>2</sub> As exchange by thiols .....	28
2.2 Supplementary material .....	33
2.3 References .....	36
Chapter 3: Rapid exchange in related arsenic derivatives.....	37
3.1 Mechanism elucidation of dimethyl arsenic exchange.....	37
3.2.1 Monomethylated derivatives .....	41
3.2.2 Examining the nature of Monomethyl arsinous acid.....	43

3.2.3 Interaction of MeAs(OH) <sub>2</sub> with cysteine .....	45
3.2.4 Temperature sensitivity of the methyl peak .....	47
3.3.1 Zinc Fingers .....	49
Experimental procedure for Zn 1b .....	50
3.3 Conclusions .....	59
3.4 References .....	60

## Table of Figures

Figure 1: Inorganic forms of arsenic .....	2
Figure 2: Challenger's proposed pathway of arsenic metabolism.....	5
Figure 3: Photo of arsenic induced arsenicosis, retrieved 2013 <sup>11</sup> .....	7
Figure 4: Structure of British Anti-Lewisite (BAL) .....	8
Figure 5: Structure of Darinparsin .....	10
Figure 6: Binding of zinc and arsenic to the PML-R-ZF1 and PML-R-ZF2 Purple: interaction of zinc Orange: Predicted interaction of arsenic (Diagram from Science <sup>13</sup> ).....	11
Figure 7: Reduction of dimethylarsinic acid and monomethylarsinic acid by thiols. ....	13
Figure 8: Structures of glutathione and alpha-glutathione. ....	15
Figure 9: Effect of pH on the speciation of arsenic (Figure from Rey et al paper <sup>21</sup> ) .....	16
Figure 10: Thermodynamic for the formation of As(III)-Thiolate complexes (from Wilcox et al. 2008 <sup>22</sup> ) .....	17
Figure 11: Interaction of H <sub>2</sub> S with arsenic V species .....	19
Figure 12: endo and exo binding forms of As(CH <sub>3</sub> S) <sub>3</sub> (Figure from Zampella et al <sup>29</sup> ) .....	21
Figure 14: Possible mechanisms for the observed coalescence of the dimethylarsonium peaks on <sup>1</sup> H NMR.....	38
Figure 15: Synthetic scheme for dimethylarseno-penicillamine .....	39
Figure 16: Synthetic scheme for the preparation of dimethylarseno-N-acetyl cysteine.....	40
Figure 17: Temperature variation on the sample of (MeAsO) <sub>x</sub> dissolved in CDCl <sub>3</sub> , 1) 273.15 K, 2) 283.15, 3)298.15 K, 4)313.15K, 5) 323.15K.....	44
Figure 18: Titration of Cysteine into a solution of (MeAsO) <sub>x</sub> in increments of 50ul. Top: No cysteine injected, Bottom 2.8 equivalents of cysteine injected. ....	46
Figure 19: NMR titration of Cysteine against MeAs(OH) <sub>2</sub> in D <sub>2</sub> O at pH 4.5.....	47

Figure 20: Temperature variation on a solution of $\text{MeAs}(\text{OH})_2$ and Cys. Top: 313K, bottom 293K .....	48
Figure 21: Structures of proposed zinc finger models.....	50
Figure 25: Titration of ligand against zinc salt.....	56
Figure 27: Excitation spectra of 8-aminoquinoline in ethanol (450 emission).....	58
Figure 28: Emission spectra with 365 nm excitation .....	59


## Chapter 1

### Thesis Introduction


#### **1.1 The Bioinorganic Chemistry of Arsenic**

Arsenic is a group 15 metalloid that is abundant in the earth's crust (ca, 1.8 ppm)<sup>1</sup> in the form of various minerals. In order to fully understand the bioinorganic chemistry of arsenic, the periodicity and chemical reactivity of arsenic must be appreciated. As a metalloid, arsenic exhibits a range of class defying properties and has the characteristics of both metal and non-metals. Elemental arsenic is semiconductor, non-ductile and has the capability of forming various allotropes with various crystal structures<sup>2</sup>. Common inorganic forms of


arsenic include arsenic trioxide ( $\text{As}_2\text{O}_3$ ) and various sulfur compounds such as realgar ( $\text{As}_4\text{S}_4$ ) and Orpiment ( $\text{As}_2\text{S}_3$ ).



Arsenic Trioxide



Realgar



Orpiment

**Figure 1: Inorganic forms of arsenic.**

In this way arsenic is similar to phosphorus which also forms a variety of sulfur bond compounds.

The chemistry of organic arsenic compounds is just as diverse and varied as its inorganic counterparts. As a pnictogen, arsenic can readily access V, III and -3 oxidation states. Arsenic is closely related to phosphorus in terms of bioinorganic chemistry, and as a result biologically arsenic can occasionally act as a phosphorus analogue. For example enzymes that use phosphate as a substrate recognise arsenate<sup>3</sup>. However there is one key difference between As and P which is that arsenic has filled inner  $3d$  orbitals and P does not. The  $3d$  orbital sits between  $4s$  and  $4p$  which makes the  $4s$  orbital less accessible<sup>4</sup>. The consequence when compared with P is that As has a more stable oxidation (III) state at the expense of a less stable (V) state. The increased accessibility of the

As (III) state gives arsenic access a wide range of different chemistry from phosphorus. This is one reason why that As is not as ubiquitous in biology whilst phosphorus is readily used in very inert components such as the phosphate backbone of DNA. In fact many arsenic compounds are toxic to biological species, the point where arsenic trioxide has gained the notoriety as the “*King of poisons*”. Despite its toxicity, arsenic compounds may be required by some biological species. This a contrast to the two heavier elements in group 15, antimony and bismuth, which have no known natural biological function<sup>2</sup>. In terms of toxicity, antimony has similar toxicity to arsenic, whilst bismuth has minimal toxicity. Bismuth has been used in over the counter products such as Pepto-Bismol to treat heartburn.

This section will focus on the interactions of arsenic within the human body and highlight recent interests and developments.

### **1.1.1 Metabolism of arsenic**


In order to fully understand arsenic's metabolism by the body, it is necessary to understand that many arsenic species are in rapidly established dynamic equilibria. It is therefore insufficient to just consider one single species, instead an easier way to rationalise the interaction is to consider arsenic as a general system that involves a rapid equilibrium all of the above states. The

dynamic nature of the arsenic species will be discussed in greater detail in section 1.2.3. It could be also argued that the chemistry of arsenic has mostly revolved around the stable forms of arsenic<sup>5</sup>, whilst the actual picture is much more distorted as arsenic goes through various unstable and hard to detect species.

Arsenic is ubiquitous in the environment and hence uptake of arsenic by humans is inevitable through food and water. The main form of arsenic that is being absorbed by the body in the inorganic form As (III)<sup>6</sup>. As arsenic is absorbed by the body, it goes through a complicated metabolism pathway that involves a range of oxidation states in addition to methylation on the arsenic. The concern of this section will be focused on the human metabolism of arsenic, however it is important to keep in mind that the metabolism of arsenic between different organisms follows similar patterns<sup>7</sup>. One of the earliest and most referenced studies into arsenic metabolism was done by Challenger in 1947 which introduced the formulation of the Challenger pathway. Despite its age, the Challenger pathway is the framework for which the metabolism of arsenic is rationalised.



Inorganic Arsenic (III)



**Figure 2: Challenger's proposed pathway of arsenic metabolism**

Challenger proposed that arsenic is absorbed as As (V), and is reduced to inorganic As (III) by glutathione (GSH). The arsenite is then methylated and oxidised by As-methyltransferase into mono-methylarsinic acid (V). This is followed by a subsequent reduction to monomethylarsenious acid by GSH. A second methylation then proceeds with the formation of dimethylarsenic acid.

Finally the dimethyl arsenious acid. Both methylated and non-methylated species are arsenite are capable of being removed from the body via the liver into urine. Recent developments<sup>8</sup> in the metabolism of arsenic has shown there are additional processes that allows arsenic to exchange between various states such as thiodimethylarsenopropanoic acid and thiodimethylarsenobutanic acid. There have be additional revisions to the pathway proposed by Naranmandura and Hayakawa<sup>9</sup>.

**Commented [a1]:** E. Schmeisser, A. Rumpler, M. Kollroser, G. Rechberger, W. Goessler, and K. A. Francesconi, "Arsenic fatty acids are human urinary metabolites of arsenolipids present in cod liver," *Angewandte Chemie International Edition*, vol. 45, no. 1, pp. 150–154, 2005.

It has been suggested that reason for the methylation is to reduce the toxicity of arsenic. However, recently this has be disputed as methylated forms of arsenic are shown to be just as toxic as its inorganic counterparts<sup>10</sup>.

### 1.1.2 Arsenic as a poison

Arsenic has an infamous history of being the "King of Poisons", and in famous deaths such as Napoleon and the horse Phar Lap. The LD<sub>50</sub> for arsenic trioxide is 35 mg per kg and a fatal dose of could be as low as 100 to 200 mg. The effects of arsenic poisoning appear from half an hour of ingestion. The immediate symptoms include abdominal pain, vomiting diarrhea and salivation and death. There are also long term effects to arsenic poison that include rashes in the form of "*arsenicosis*", cancer and damage to the cardiovascular system.



**Figure 3: Photo of arsenic induced arsenicosis, retrieved 2013<sup>11</sup>**


The exact mechanism of the arsenic interaction is not well established, but it is believed that different species of arsenic are responsible for a range of toxic responses. The initial symptoms have been proposed to be the result of arsenite (III) binding to vicinal thiols of active enzymes. The binding of arsenic may either cause conformational change in the enzyme or block active sites, hence inhibiting enzyme activity. Although has been widely regarded as the mechanism

of action, there has been little direct evidence to substantiate this<sup>12</sup>. Arsenate (V) have been proposed be toxic in another manner, they are transported into cells via phosphate transport pathways. They then to mimic phosphates and are then included into ATP and DNA<sup>7</sup>. As arsenic can do additional chemistry, this may cause mutations in the cell or cellular damage.

Arsenic's poisonous attributes is a large issue today as arsenic is found in ground water. This is a result of the leeching of arsenic from the soil into ground water. This is a major problem for in many countries, but most particularly in developing economies where there is little recourse but to these wells which brings this toxic ground water up and causes health problems in the local population.

### 1.1.3 Treatment of arsenic poisoning

Dimercaprol, also known as British anti-Lewisite (BAL), is used medically as a treatment against arsenic poisoning. BAL has a capability of binding very strongly to arsenic via a bi-dentate bond.




**Figure 4: Structure of British Anti-Lewisite (BAL)**

It has been proposed that BAL competes with metabolic enzymes for arsenic binding and hence reduces arsenic's toxicity. The chelation would result in subsequent removal of the bound compound from the body via urine.

#### **1.1.4 Medicinal applications of arsenic**

It has been proposed by Ehrlich that arsenic based drugs would get converted to "RAsO" and would interact with metallothionein. The exact mechanism of interaction is not known as little direct evidence was found to support this theory. The underlying pattern is that many enzymes are inhibited by arsenic, especially those containing thiol (SH) groups.


More recently arsenic trioxide ( $\text{As}_2\text{O}_3$ ) was discovered to be effective against the cancer Acute Promyelocytic Leukemia (APL). APL is a rare form of myeloid leukemia that affects an estimated 1500 new patients each year. The discovery of the drug initially came from traditional Chinese therapy followed by major research developments in the US, leading to FDA approval in 2000 under tradename of Trisenox. In the clinical trial for Trisenox, it was reported that 77% of the 582 patients treated were alive after 3 years, compared with just 50% for non-arsenic treatment. More recently, Ziopharm oncology has developed Darinaparsin, an oxidation state III organoarsenical bound to glutathione for the treatment of APL.



**Figure 5: Structure of Darinparsin**

Currently the drug is in Phase III trials and has been shown to be more effective than arsenic trioxide with reduced side effects. This is an interesting drug because it is a possible metabolite of arsenic trioxide (see section 1.1.1, the metabolism of arsenic) as  $\text{Me}_2\text{AsOH}$  forms an equilibrium with GSH to form Darinparsin in solution. In addition, there are two methyls bound to the arsenic already, preventing it from forming more than 1 bond to thiols in the As(III) ground state.

The mechanism for arsenic's activity in the treatment of APL is not fully understood. Luckily APL is a well-studied cancer and work has been done on the oncoprotein PML-RAR $\alpha$ . PML-RAR $\alpha$  is a fusion protein that consists of Promyelocytic Leukemia protein PML, and retinoic acid receptor RAR $\alpha$ . Zhang et al<sup>13</sup> proposed in 2010 that arsenic trioxide interacts directly with the two zinc fingers in PML by replacing the zinc in the zinc finger. They showed using EXAFS that arsenic (III) can interact directly with the sulfurs on the cysteine in the zinc finger.



**Figure 6: Binding of zinc and arsenic to the PML-R-ZF1 and PML-R-ZF2** Purple:  
interaction of zinc Orange: Predicted interaction of arsenic (Diagram from  
Science<sup>13</sup>)

They proposed that the coordination of the arsenic causes conformation changes in the protein. These changes would lead to aggregation of this protein and trigger SUMOylation and eventual cell death. The exact mechanism in which the arsenic displaces the zinc is relatively unknown and not much additional work has been done on the arsenic interaction in the zinc finger domain. In addition this proposal does not apply for dimethylated arsenic species such as Darinaparsin as the methyl groups take space in arsenic's co-ordination sphere.

Another proposal for the mechanism of arsenic<sup>14</sup> has been proposed by involves the ability of arsenic to bind to thiols on PML proteins. Once bound, arsenic would form intermolecular disulfide bridges that result in the

multimerization of PML into a mesh. It is unclear if arsenic has specificity for the thiols in PML.


## **1.2 Chemistry of Arsenic compounds**

The previous section focused on the biological interactions of arsenic with key one underlying theme: the binding of arsenic to thiols is the key interaction involved in its mechanism of action. This interaction has been profusely by biologists. Despite the frequent use of this concept to explain arsenic's poisonous attributes, the chemistry of this interaction is not well understood. This section will outline the approaches taken in chemistry to resolve the kinetics and thermodynamics of the arsenic thiol interaction.

### **1.2.1 Redox of arsenic by GSH**

Cullen et al.<sup>15</sup> first discovered the ability of thiols to reduce  $\text{Me}_2\text{AsOOH(V)}$  to form  $\text{Me}_2\text{AsGSH (III)}$ .





**Figure 7: Reduction of dimethylarsinic acid and monomethylarsinic acid by thiols.**

They used a range of thiols such as GSH, Cys and mercaptoethanol which had a range of pKas. They concluded that the thiols did not have to be deprotonated before this reduction took place as they could perform this reaction in a range of pH below that of the pKa of the thiol.

Delnomdedieu et al<sup>16</sup> built upon this work using modern NMR spectroscopy. They also noticed that as the pH increased, there is the possibility of the formation of free Me<sub>2</sub>As<sup>+</sup>. This is because even though there was no direct peak for that of free Me<sub>2</sub>As<sup>+</sup> they noticed the chemical shift of the methyl peaks would shift slightly as the pH was increased. They proposed change in the chemical shift was caused by the involvement of an increased amount of Me<sub>2</sub>As<sup>+</sup> as solution was basified. Based on this they proposed that the reduction occurred before binding to the thiol. They also noted changes in the spectrum at high pH;


the two diastereotopic methyl peaks would become a single peak. However no kinetic measurements were done on the system.

Carter et al<sup>17</sup> showed the reduction of arsenate to form As(GS)<sub>3</sub>





This reaction was found to happen readily at pH 7 and has been fully characterised by <sup>1</sup>H and <sup>13</sup>C NMR. They proposed that like for the methylated systems, the arsenate reduction occurred before the binding to GS as a when used a ratio of 1:2 or less, the formation of As(SG)<sub>3</sub> was not observed.

Arsenic (III) species are formed in the presence of a reducing environments. However, in non-reducing aqueous environments, arsenic (III) species are unstable. This has been noted by Cullen et al as they found the Me<sub>2</sub>AsCys and Me<sub>2</sub>AsGS were unstable in solution and rapidly decomposed to their parent oxidation state V acid along with the production of the disulfide.



As a result, they degassed all their solvents prior to use and noted this improved their yields when compared to that of Zingaro et al<sup>18</sup> who performed similar synthesis under aerobic conditions. However, the exact mechanism of this degradation was not fully understood. More recently Zhao et al<sup>19</sup> examined the

degradation of the similar system  $\text{As}(\text{GS})_3$  using HPLC-ESI-MS. They found that an isomer of  $\alpha$ -GSH that was the result of the cleavage of GSSG.



**Figure 8: Structures of glutathione and alpha-glutathione.**

The  $\alpha$ -GSH was identified comparing the fragmentation patterns of degradation products of  $\text{As}(\text{GS})_3$  compared to that of standard GSH. They noted that  $\alpha$ -GSH is not normally formed from the degradation of GSSG in the absence of arsenic. The mechanism and reason for the formation of this isomer is not known.

### 1.2.2 Interaction of arsenic with thiols

One of the well-studied systems in terms of thermodynamics is that of  $\text{As}(\text{OH})_3$  and its interaction with Glutathione to form  $\text{As}(\text{GSH})_3$ . Rey et al<sup>20</sup> used potentiometric and spectroscopic data to show that the formation constant,  $\log K$ , of  $\text{As}(\text{GS})_3$  to be 32. They noted that pH played important role in this interaction




Fig. 2. Species distribution curves for the systems: (a) As-Cys (1)  $\text{As}(\text{OH})_3$ ; (2)  $\text{As}(\text{OH})_2\text{O}^-$ ; (3)  $\text{As}(\text{OH})\text{O}_2^{2-}$ ; (4)  $[\text{As}(\text{HCys})_3]$ ; (5)  $\text{As}(\text{Cys})(\text{OH})_2^-$ ; and (b) As-GSH (1)  $\text{As}(\text{OH})_3$ ; (2)  $\text{As}(\text{OH})_2\text{O}^-$ ; (3)  $\text{As}(\text{OH})\text{O}_2^{2-}$ ; (6)  $[\text{As}(\text{HGS})_3]^-$ ; (7)  $[\text{As}(\text{GS})(\text{OH})_2]^-$ . In both cases the total ligand concentration is 15 mM and the total As(III) concentration is 5 mM.

#### Figure 9: Effect of pH on the speciation of arsenic (Figure from Rey et al paper<sup>21</sup>)

Further research has been done by Wilcox<sup>12</sup> et al. who used colormetric and calorimetric studies to quantify the formation constants of arsenothiolates.

**Table 3.** Thermodynamic Properties for the Formation of As(III)–Thiolate Complexes

	As(OH) <sub>3</sub>	MMA
GSH	1:3 $\beta_3 = 1.8 \times 10^6$ $\Delta G = -8.8 \text{ kcal/mol}$ $\Delta H = -38.7 \text{ kcal/mol}$ $\Delta S = -96 \text{ cal/(mol}\cdot\text{K)}$	1:2 $\beta_2 = 1.3 \times 10^7$ $\Delta G = -10.1 \text{ kcal/mol}$ $\Delta H = -17.8 \text{ kcal/mol}$ $\Delta S = -25 \text{ cal/(mol}\cdot\text{K)}$
DMSA (5) <sup>a</sup>	1:2 $\beta_2 = 8.3 \times 10^8$ $\Delta G = -12.7 \text{ kcal/mol}$ $\Delta H = -27.3 \text{ kcal/mol}$ $\Delta S = -47 \text{ cal/(mol}\cdot\text{K)}$	1:1 $K = 1.0 \times 10^7$ $\Delta G = -9.9 \text{ kcal/mol}$ $\Delta H = -13.2 \text{ kcal/mol}$ $\Delta S = -9 \text{ cal/(mol}\cdot\text{K)}$
DHLA (6) <sup>a</sup>	2:3 $\beta_{2:3} = 4 \times 10^{18} \text{ }^b$ $\Delta G = -25 \text{ kcal/mol}$ $\Delta H = -43 \text{ kcal/mol}^c$ $\Delta S = -59 \text{ cal/(mol}\cdot\text{K)}$	1:1 $K = 1.1 \times 10^7$ $\Delta G = -10.0 \text{ kcal/mol}$ $\Delta H = -17.0 \text{ kcal/mol}$ $\Delta S = -20 \text{ cal/(mol}\cdot\text{K)}$
DTT (7) <sup>a</sup>	1:1 $K = 9.5 \times 10^5$ $\Delta G = -8.5 \text{ kcal/mol}$ $\Delta H = -13.7 \text{ kcal/mol}$ $\Delta S = -17 \text{ cal/(mol}\cdot\text{K)}$	1:1 $K = 8.2 \times 10^5$ $\Delta G = -8.4 \text{ kcal/mol}$ $\Delta H = -15.9 \text{ kcal/mol}$ $\Delta S = -24 \text{ cal/(mol}\cdot\text{K)}$

**Figure 10: Thermodynamic for the formation of As(III)-Thiolate complexes (from Wilcox et al. 2008<sup>22</sup>)**

They found that for As(OH)<sub>3</sub> the binding of glutathione has a stability constant  $\beta_3 = 2 \times 10^6$ . In addition, they found a co-operative effect in the binding of thiols and the formation constant after each binding is increased. For example  $\Delta H_3$  was found to be -33.1 kcal/mol compared to  $\Delta H_1$  of -2.5 kcal/mol. They also looked at monomethylarsenite and its equivalent binding to GSH and found that the binding of GSH to a monomethylated species gave comparable numbers to that of the non-methylated species. Gailer<sup>23</sup> offered an alternative view and proposed after passing both methylated and non-methylated compounds through size exclusion chromatography, the methylated versions were found to be more stable.

Finally they worked out the enthalpy of thiolate displacement to be -2.8 kcal/mol. They also found that there is a large unfavourable  $\Delta S$  term involved in the binding as a result of the loss of conformational degrees of freedom. As a result they proposed that vicinal thiols that are conformationally constrained will have a higher affinity for arsenic III. This argument extends to conformationally unconstrained Cys residues found in zinc fingers would by this theory have a lower affinity for As (III).


Zingaro et al used the ability of thiols to displace the hydroxyl group in numerous synthesis of new dimethyl arsenious derivatives<sup>24</sup>.

### **1.2.3 Lability of the Arsenic Thiol bond**

It was found experimentally that as  $\text{As}(\text{SG})_3$  was passed through a size exclusion chromatography (SEC)<sup>23</sup>, an increase of temperature of the column would cause retention shifts of the arsenic peaks towards small-molecular-mass regions. This is indicative of labile arsenic sulfur bonds which could allow the GS to break away from the molecule.

Arsenic III compounds have the ability are known have labile bonds. The is shown by the ability of meso-2,3-dimercaptosuccinic acid to displace GSH from the complex from  $\text{As}(\text{GS})_3$ <sup>25</sup>

Arsenic systems have been studied by theoretical chemistry by Orthaber et al<sup>26</sup> using DFT calculations with B3LYP/6-31G basis set. They were interested in the interaction of H<sub>2</sub>S with Arsenic (V) species



**Figure 11: Interaction of H<sub>2</sub>S with arsenic V species**


They found that the formation constants is very favorable with  $\Delta G$  of up to 51 kJ/mol for dimethylarsinoylethanol. Interesting they found that the pentavalent state, the –OH and –SH could interconvert between axial and equatorial positions similar to that of the Berry pseudorotation. They found that the relative intermediate of the interconversion is 6.7 kJ/mol.

One analogous system for the lability of arsenic is that of Mercury. The mercury thiolate bond has been shown to be labile and capable of rapidly breaking and reforming<sup>27</sup>.

#### **1.2.4 Co-ordination and geometry of arsenic compounds**

The co-ordination chemistry of arsenic is both rich and diverse. As previously discussed in chapter 1.1, arsenic is a metalloid and can both accept and donate lone pairs. As it result it can accept co-ordination and act as a metal, in addition to donating its lone pair and acting as a ligand.

Edmonds et al<sup>28</sup> found that chiral arsenite species are capable rapidly racemizing. They studied this using the chiral arsenite, methylphenylarsinic acid and reacted it with (L)-glutathione to form two diastereomers. They then tried to separate the diastereomers using HPLC




They tried to resolve the diastereomers using NMR spectroscopy with the aid of lanthanide shift reagents and COSY 2D techniques. They were surprised to find that the species have racemized during the purification process (with heating was less than 40°C). They concluded that no pyramidal inversion of the arsenic was seen but could not give a definitive mechanism for this inversion.

### 1.2.5 Mechanism of arsenic bond lability

The mechanism under which arsenic can perform its bond lability is not fully understood. Zampella et al<sup>29</sup> tried to calculate the binding of structures of arsenite systems using DFT calculations in order to understand how arsenite



interacts with thiols of proteins. They modelled the arsenic binding systems with  $\text{As}(\text{CH}_3\text{S})_3$  and determined that there are two major forms of binding: the endo and exo isomers.



**Figure 12: endo and exo binding forms of  $\text{As}(\text{CH}_3\text{S})_3$  (Figure from Zampella et al<sup>29</sup>)**

They concluded that the calculated lowest energy forms were similar to that found in protein systems. This meant that the mode of binding was strongly influenced by the steric bulk of the substituents. They also determined that the direction of the lone pair was a crucial factor influencing the selectivity binding for particular metal ions.

### 1.3 Conclusions

The mechanism of arsenic based drugs is currently not well understood and poses a road-block to arsenic based drug development. It is hypothesised that whilst arsenic is often thought react in an analogous fashion phosphorus, the kinetics and thermodynamics of pnictogen bonds formed by these two species far from analogous. The Challenger pathway shows that arsenic can be prone to both oxidation and methylation, a sharp contrast to phosphorus bonds which are stable enough to be used in genetic material. In addition to having different reactivity than phosphorus, arsenic seems to have bond lability, where arsenic appears to move between different thiol containing compounds. The mechanism of this interaction has not been thoroughly characterised, the thermodynamics of arsenic thiol bonds remains unknown. This project aims at using modern kinetic techniques to study the kinetics and thermodynamics of arsenic compounds. This is a pivotal step in the greater picture of understanding of arsenic drug interaction and the transportation of arsenic within the body.

## References

1. Norman, N. C., *Chemistry of Arsenic, Antimony and Bismuth*. Springer: 1998.
2. Sun, H.; Editor, *Biological Chemistry Of Arsenic, Antimony And Bismuth*. John Wiley & Sons Ltd.: 2011; p 383 pp.
3. Tawfik, D. S.; Viola, R. E., Arsenate replacing phosphate: alternative life chemistries and ion promiscuity. *Biochemistry* **2011**, *50*, 1128-34.
4. Levason, W.; Reid, G., 3.6 - Arsenic, Antimony, and Bismuth. In *Comprehensive Coordination Chemistry II*, Editors-in-Chief: J. A. M.; Meyer, T. J., Eds. Pergamon: Oxford, 2003; pp 465-544.
5. Raab, A.; Hansen, H.; Feldmann, J., Labile Arsenic Compounds in Biological Matrices, or Possible Problems Finding the Metal Species Present in Cells. *Special Publication-Royal Society of Chemistry* **2005**, *301*, 72.
6. Jomova, K.; Jenisova, Z.; Feszterova, M.; Baros, S.; Liska, J.; Hudecova, D.; Rhodes, C. J.; Valko, M., Arsenic: toxicity, oxidative stress and human disease. *Journal of applied toxicology : JAT* **2011**, *31* (2), 95-107.
7. Cullen, W. R., *Is Arsenic an Aphrodisiac?* The Royal Society of Chemistry: 2008; p P001-P412.
8. Thomas, D. J.; Li, J.; Waters, S. B.; Xing, W.; Adair, B. M.; Drobna, Z.; Devesa, V.; Styblo, M., Arsenic (+3 oxidation state) methyltransferase and the

methylation of arsenicals. *Experimental biology and medicine* (Maywood, N.J.)

**2007**, *232*, 3-13.

9. Rehman, K.; Naranmandura, H., Arsenic metabolism and thioarsenicals.

*Metallomics : integrated biometal science* **2012**, *4* (9), 881-92.

10. Petrick, J. S.; Ayala-Fierro, F.; Cullen, W. R.; Carter, D. E.; Vasken

Aposhian, H., Monomethylarsonous acid (MMA(III)) is more toxic than arsenite in

Chang human hepatocytes. *Toxicol. Appl. Pharmacol.* **2000**, *163* (2), 203-207.

11. Arsenic Contamination of Groundwater in South and East Asian

Countries. <http://go.worldbank.org/WVH5RSX920> (accessed Jan. 2013).

12. Spuches, A. M.; Kruszyna, H. G.; Rich, A. M.; Wilcox, D. E.,

Thermodynamics of the As(III)-thiol interaction: arsenite and monomethylarsenite complexes with glutathione, dihydrolipoic acid, and other thiol ligands. *Inorg.*

*Chem.* **2005**, *44*, 2964-72.

13. Zhang, X. W.; Yan, X. J.; Zhou, Z. R.; Yang, F. F.; Wu, Z. Y.; Sun, H. B.;

Liang, W. X.; Song, A. X.; Lallemand-Breitenbach, V.; Jeanne, M.; others,

Arsenic trioxide controls the fate of the PML-RAR $\alpha$  oncoprotein by directly binding PML. *Science* **2010**, *328* (5975), 240-3.

14. Lallemand-Breitenbach, V.; Zhu, J.; Chen, Z.; de Thé, H., Curing APL

through PML/RARA degradation by As<sub>2</sub>O<sub>3</sub>. *Trends in molecular medicine* **2012**, *18*, 36-42.

15. Cullen, W.; McBride, B.; Reglinski, J., The reaction of methylarsenicals with thiols: some biological implications. *J. Inorg. Biochem.* **1984**, *21*, 179-193.
16. Delnomdedieu, M.; Basti, M. M.; Otvos, J. D.; Thomas, D. J., Reduction and binding of arsenate and dimethylarsinate by glutathione: a magnetic resonance study. *Chem. Biol. Interact.* **1994**, *90*, 139-55.
17. Scott, N.; Hatlelid, K. M.; MacKenzie, N. E.; Carter, D. E., Reactions of arsenic(III) and arsenic(V) species with glutathione. *Chem. Res. Toxicol.* **1993**, *6*, 102-6.
18. Chen, G. C.; Zingaro, R. A.; Thompson, C. R., 6-thio and-seleno-[beta]-glucose esters of dimethylarsinous acid. *Carbohydr. Res.* **1975**, *39*(1), 61-66.
19. Zhao, F.; Chen, Y.; Qiao, B.; Wang, J.; Na, P., Analysis of two new degradation products of arsenic triglutathione in aqueous solution. *Frontiers of Chemical Science and Engineering* **2012**, *6*, 292-300.
20. Rey, N.; Howarth, O., Equilibrium characterization of the As (III)-cysteine and the As (III)-glutathione systems in aqueous solution. *J. Inorg. Biochem.* **2004**, *98*, 1151-9.
21. Rey, N. A.; Howarth, O. W., Equilibrium characterization of the As (III)-cysteine and the As (III)-glutathione systems in aqueous solution. *J. Inorg. Biochem.* **2004**, *98*(6), 1151-9.

22. Wilcox, D. E., Isothermal titration calorimetry of metal ions binding to proteins: An overview of recent studies. *Inorg. Chim. Acta* **2008**, *361*, 857-867.
23. Percy, A. J.; Gailer, J., Methylated trivalent arsenic-glutathione complexes are more stable than their arsenite analog. *Bioinorganic chemistry and applications* **2008**, *2008*, 539082.
24. Banks, C. H.; Daniel, J. R.; Zingaro, R. A., Biomolecules Bearing the S- or SeAsMe<sub>2</sub> Function : Amino Acid and Steroid Derivatives. *J. Med. Chem.* **1979**, *22*, 572-575.
25. Delnomdedieu, M.; Basti, M. M.; Otvos, J. D.; Thomas, D. J., Transfer of arsenite from glutathione to dithiols: A model of interaction. *Chem. Res. Toxicol.* **1993**, *6*, 598-602.
26. Orthaber, A.; Sax, A. F.; Francesconi, K. a., Oxygen versus sulfur: Structure and reactivity of substituted arsine oxides and arsine sulfides. *J. Comput. Chem.* **2012**, *33*, 112-7.
27. Carson, G.; Dean, P., The Metal NMR Spectra of Thiolate. *Inorg. Chim. Acta* **1982**, *66*, 157-161.
28. Edmonds, J. S.; Nakayama, T.; Kondo, T.; Morita, M., Diastereoisomerism of thiol complexes of arsenic acids and pseudoasymmetry of arsenic: a <sup>1</sup>H and <sup>13</sup>C NMR study. *Magnetic resonance in chemistry : MRC* **2006**, *44*, 151-62.

29. Zampella, G.; Neupane, K. P.; De Gioia, L.; Pecoraro, V. L., The importance of stereochemically active lone pairs for influencing Pb(II) and As(III) protein binding. *Chemistry (Weinheim an der Bergstrasse, Germany)* **2012**, *18*, 2040-50.

## Chapter 2:

### **2.1 Kinetics and thermodynamics of rapid Me<sub>2</sub>As exchange by thiols.**

The research from this project is used in the following paper published on 05 March 2013 in the journal Organic & Biomolecular Chemistry.











## 2.2 Supplementary material

The following supplementary material details the methodology and calculations used in the chapter 2.1.

*Preparation of S-(dimethylarseno)cysteine (Me<sub>2</sub>AsCys).* The synthetic preparation was adapted from the Cullen method<sup>15</sup>, with the main change being nitrogen gas is used instead of carbon dioxide to produce an inert atmosphere. Cacodylic acid (0.445g) of and 0.929g of L-Cystine was mixed in distilled water under nitrogen for 16 hours. The precipitate was filtered dried under reduced pressure without heating. <sup>1</sup>H NMR (500 MHz, D<sub>2</sub>O) δ 3.99 – 3.91 (m, 1H), 3.24 – 3.13 (m, 1H), 1.37 (s, 1H), 1.36 (s, 1H).

*Preparation of Me<sub>2</sub>AsCys Solution.* 0.0115g of Me<sub>2</sub>AsCys was dissolved in 1 ml of Phosphate buffer at pH 4.2 (10% potassium phosphate buffer was prepared by the dissolution of potassium phosphate into water and subsequent adjustment of pH using NaOH. The solution was deuterated by drying the mixture and re-hydrating it with D<sub>2</sub>O). This solution was diluted to form a desired concentration of 6.40 mM by a 1/5 dilution using 200ul of the solution and 800 ul of buffer. To ensure the temperature is changing constantly and as expected, a plot of the


solvent chemical shift against temperature. This indicates minimal changes to the pH or random errors in the temperature readings, as the  $R^2$  is 0.999. This however does not rule out the possibility of systematic errors to the temperature readings. This could be remedied by using a temperature reference.

NMR data was acquired with a Varian 500 MHz instrument. Temperature calibration was done using standard samples of ethylene glycol and methanol<sup>30</sup>. All NMR acquisitions were made with 16 scans, transform size of 32k and shims were done with a gradient shimming on each measurement. Data was processed using Mnova 6.1.1 FID processing software.

To determine the rate constant of the reaction, the separation of the methyl peaks were used to calculate the rate constant using the following equation<sup>31</sup>:

$$k = \frac{\pi}{\sqrt{2}} \sqrt{\delta\nu^2 - \delta\nu_{\text{F}}^2}$$

This equation is applicable in this situation as  $\delta\nu$  and  $k$  are much larger than the bandwidth in absence of exchange (0.6 Hz). As it is not possible to lower the sample below 0 degrees due to the freezing point of water,  $\delta\nu$  was estimated using the lowest experimentally obtained separation of 7.8 Hz (solution of 1mM sample at 10 °C).



**Figure 13: Plot of rate constant against temperature of a 5mM solution of Me<sub>2</sub>AsCys**

We notice an almost linear build up of the rate constant.

The NMR dynamic exchange was modeled using DNMR3 with Spinworks 3 software<sup>32</sup>. The experimentally determined variables (low temperature limit) of the system at 5°C with a concentration of 1mM was used as the input parameter for the simulation (two spins at 1.3832 and 1.3698 with 0.6 line width). Other parameters include Permutation vector set as 2,1 (mutual exchange of the system), the relaxation time 1 sec (same as acquisition), populations were 0 as this is a case of mutual exchange and maximum iterations = 30. The rate constant RC (1,2) was gradually increased by 1 s<sup>-1</sup> until the simulated system exhibited coalescence. It was found that a rate constant of 16 s<sup>-1</sup> gave coalescence for the two peaks.

### 2.3 References

15. Cullen, W.; McBride, B.; Reglinski, J., The reaction of methylarsenicals with thiols: some biological implications. *J. Inorg. Biochem.* **1984**, *21*, 179-193.
30. Holz, M.; Heil, S. R.; Sacco, A., Temperature-dependent self-diffusion coefficients of water and six selected molecular liquids for calibration in accurate <sup>1</sup>H NMR PFG measurements. *PCCP* **2000**, *2*(20), 4740-4742.
31. Sandström, J., *Dynamic Nmr Spectroscopy*. Academic Press: 1982.
32. Spinworks, SpinWorks3 Manual. *Data Processing* **2010**, *7*, 1-117.




## Chapter 3: Rapid exchange in related arsenic derivatives.

### 3.1 Mechanism elucidation of dimethyl arsenic exchange

In Chapter 2 we showed that facile dimethylarsenic exchange occurred between dimethylarsenic cysteine and glutathione adducts. In addition to exchange between different dimethylarsenic adducts, we showed that the diastereotopic methyls of  $\text{Me}_2\text{AsCys}$  also underwent rapid exchange on the NMR timescale. The self-exchange was of particular interest because a good understanding of this was required to understand the intra-system exchange. In this second part of this project, I focused on elucidating the mechanism of the facile dimethylarsenic self-exchange.

There are several possible mechanisms for the observed effect. One possibility is a hydrolysis mechanism where  $\text{H}_2\text{O}$  would hydrolyse the arsenic species thus causing the formation of a  $\text{Me}_2\text{AsOH}$  which no longer contains diastereotopic methyls. Another possibility is an intermolecular attack from the nitrogen from another  $\text{Me}_2\text{AsCys}$  into the arsenic, thus forming pyramidal intermediate. This nucleophilic attack could also be intramolecular, causing the formation of a 5 membered ring.




**Figure 14: Possible mechanisms for the observed coalescence of the dimethylarsonium peaks on  $^1\text{H}$  NMR.**

In order to further validate this mechanism, we attempted to use chemical substitution to produce other dimethylarsenic derivatives and compare their kinetics. The first synthesis target is dimethylarseno-penicillamine which contains penicillamine, a cysteine derivative which contains two additional methyls between the thiol and the  $\beta$ -carbon. With additional methyls substituents different

rates of 5 membered ring formation are expected, and thus significantly different DNMR kinetics. Another synthetic target was dimethylarseno-N-acetyl cysteine where the nitrogen is tied up with an acetyl group, thus tying up its lone pairs and preventing it from nucleophilically attacking or chelating to the arsenic.

*Preparation of Dimethylarseniodide.*  $\text{Me}_2\text{AsI}$  was prepared using the Burrows method<sup>33</sup>. Potassium iodide, 15g, and 5g of  $\text{Me}_2\text{AsOOH}$  are dissolved in 45ml of distilled water. Concentrated HCl 5ml is added to make a clear colorless solution. Sulfur dioxide is bubbled for 15 minutes through the solution at which point the solution turned to light yellow. After around 5 minutes of bubbling the solution darkened to an opaque black, followed by the formation of a bottom layer which was clear yellow. The bottom layer was extracted and distilled under reduced pressure of 16 mm at 401K.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.01.

*Preparation of Dimethylarseno-penicillamine*




**Figure 15: Synthetic scheme for dimethylarseno-penicillamine**

Penicillamine, 0.5g, was suspended in dimethoxyethane. 1 ml of  $\text{Me}_2\text{AsI}$  was added by syringe causing the full dissolution was penicillamine. 1 ml of pyridine

was added and precipitation immediately occurred. The solution was refluxed for 15 minutes and left to stir for 2 hours. The solution was filtered and the filtrate was dried. Note this compound has an extremely unpleasant smell, Schlenk apparatus and proper fume hood containment methods are required. NMR revealed the filtrate to be pyridinium iodide.  $^1\text{H}$  NMR (400 MHz,  $\text{d}_2\text{O}$ )  $\delta$  3.86 (d,  $J$  = 2.3 Hz, 1H), 3.60 (s, 1H), 3.36 (s, 1H), 2.02 (d,  $J$  = 1.1 Hz, 1H), 1.60 (s, 4H), 1.44 (s, 4H), 1.38 (s, 3H), 1.35 – 1.28 (m, 6H). The NMR revealed additional unexpected peaks that could be attributed to cacodylic acid and the disulfide adduct of penicillamine. Attempts to further purify the product proved unsuccessful.

*Preparation of dimethylarseno-N-acetyl cysteine*



**Figure 16: Synthetic scheme for the preparation of dimethylarseno-N-acetyl cysteine**

0.5g of NAC was dissolved in dimethoxyethane and 1 ml of  $\text{Me}_2\text{AsI}$  was added by syringe. 1 ml of pyridine was added and precipitation immediately occurred. The solution was refluxed for 15 minutes and left to stir for 2 hours. The solution was filtered and the filtrate was dried. NMR revealed the filtrate to

be pyridinium iodide. Unfortunately this reaction mixture is contaminated by Me<sub>2</sub>AsOOH and N-acetylcysteine disulfide. This was difficult to purify as the product was a thick liquid and could not be recrystallized.

Unfortunately it proved difficult to synthesize the pure derivatives of Me<sub>2</sub>AsCys required for DNMR experimentation. Purity is especially important because contaminants of other thiol containing products are expected to significantly alter the kinetics.

### 3.2.1 Monomethylated derivatives

The kinetics and thermodynamics of monomethyl arsenic (III) derivatives is of immense interest and complements the work done on the dimethyl arsenic system. One particularly interesting property is the ability of monomethyl arsenic derivatives to bind to two thiols, thus allowing it to bind very strongly to vicinal dicysteine residues. The starting material for monomethyl derivatives is monomethyl arsenous acid sodium salt, (MeAsO)<sub>x</sub>, which was synthesised by literature methods<sup>34</sup>.

*Preparation of Methylarsenate(V) acid sodium salt.* 3g of Arsenic trioxide was dissolved in 10 ml of 10M NaOH. 15 ml of MeI was added, forming a bilayer

solution. The solution mixture was heated to reflux for 16 hours, which resulted in a white precipitate of methylarsenate(V) acid sodium salt in 72% yield.

*Preparation of Methyl arsonious acid sodium salt.*  $\text{MeAsO}(\text{ONa})_2$  was dissolved in 50 ml of  $\text{H}_2\text{O}$ . Dissolution of the initial salt was promoted by gradual heating of the solution. Once dissolved the solution is treated with sulfur dioxide which is bubbled through the solution. The solution quickly becomes clear (suggesting acid sensitivity) then light yellow after 2 minutes. After saturating with  $\text{SO}_2$ , the solution was quickly boiled for 2 minutes then cooled for 15 minutes. Neutralisation with sodium carbonate turned the solution from light yellow to clear. Solution was dried and  $(\text{MeAsO})_x$  was extracted with benzene. After removing the benzene in vacuo resulting in a white solid (70% yield).


$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.58 (d,  $J$  = 6.8 Hz, 5.9%), 1.52 – 1.49 (m, 27%), 1.48 (d,  $J$  = 2.1 Hz, 58%), 1.44 – 1.42 (m, 8%). This corresponds to the literature reference (Aposhian et al<sup>35</sup>) of ( $\text{CDCl}_3$ ):  $\delta$  1.58/1.59 (5.0%), 1.50/1.51 (26.8%), 1.48 (60.0%), 1.43 (8.1%).  $^1\text{H}$  NMR (400 MHz,  $\text{D}_2\text{O}$ )  $\delta$  1.17 (s, 1H).  $^1\text{H}$  NMR (500 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  1.21 (s, 1H).  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  1.25 (dd,  $J$  = 11.1, 7.1 Hz, 1H). ESI of the compound did not reveal any tetramer peaks as NMR later indicated that it possibly hydrolyses.

Monomethyl arsinous acid takes the form of cyclic and linear oligomers, hence resulting in the formula  $(\text{MeAsO})_x$  where the exact number of oligomers depends on the concentration of the solution. This is shown in the  $\text{CDCl}_3$  NMR which contains 4 sets of multiplets in  $\text{CDCl}_3$ .

### 3.2.2 Examining the nature of Monomethyl arsinous acid

Monomethyl arsinous acid exists in different oligomers and a more detailed study of this mixture is necessary. Marsmann and Wazer<sup>34</sup> proposed the possibility of the species oligomerizing at higher concentrations and temperatures, in particular the preference of the tetrameric form at 48% wt concentrations of arsenosomethane at  $120^\circ\text{C}$  (in diphenyl ether). It is interesting to note that the Wazer did not observe a hydrolysis with diphenyl ether.

To validate this possibility of oligomer, a temperature dependant NMR experiment was performed with the sample in  $\text{CDCl}_3$ . For this experiment 0.0975g of  $(\text{MeAsO})_x$  was dissolved in 1000  $\mu\text{l}$  of  $\text{CDCl}_3$ . The sample was initially cooled down to 273.15K and the temperature was slowly brought up in  $10^\circ$  increments.



**Figure 17: Temperature variation on the sample of (MeAsO)<sub>x</sub> dissolved in CDCl<sub>3</sub>, 1) 273.15 K, 2) 283.15, 3)298.15 K, 4)313.15K, 5) 323.15K.**

As the temperature increases, we notice a shoulder peak appearing at 1.48 ppm. We also notice an increase in the intensity of the peaks at 1.57 and 1.59. No coalescence of the peaks is observed suggesting that this process is slow on the NMR time scale. The integrals of the peaks stay constant during the heating and the mixture returns to its original composition upon cooling to room temperature. Additional experiments are necessary, in particular the use of different solvents (possibly diphenyl ether, the solvent used by Marsmann et al<sup>[34]</sup>). Concentration variation, especially that of high concentrations of (MeAsO)<sub>x</sub> may confirm the oligomerization. In addition it is difficult to heat the solvent to higher temperatures inside the NMR and this experiment could be extended by




heating the compounds separately other solvents. The oligomerization is might not be a reversible process in which cause this would not cause a coalescence of the peaks in the NMR.

### **3.2.3 Interaction of $\text{MeAs}(\text{OH})_2$ with cysteine**

Previously in the project we characterised the interaction of dimethyl arsenicals with cysteine. This provided us with a new and expected insight into the lability and kinetics of the As-S bond. With monomethyl derivatives the situation is more complicated as the arsenic can bind with two cysteines.


*Preparation of Monomethyl arsenious acid solution.* A solution of  $(\text{MeAsO})_x$  was prepared by dissolving 0.0245 g of the compound in 1.0 ml of  $\text{D}_2\text{O}$  (buffered with 10% deuterated sodium phosphate). A 231 mM solution of cysteine was prepared by dissolving 0.0277g of cysteine in 2000  $\mu\text{l}$  of the same buffered  $\text{D}_2\text{O}$ . All solutions were deoxygenated by bubbling nitrogen for 10 minutes. For the NMR titration, 500 $\mu\text{l}$  of the stock  $(\text{MeAsO})_x$  solution placed in a NMR tube. The Cysteine solution was titrated into the NMR tube at 100 $\mu\text{l}$  aliquots followed by 30 sec of mixing.



**Figure 18: Titration of Cysteine into a solution of (MeAsO)<sub>x</sub> in increments of 50ul.**


**Top: No cysteine injected, Bottom 2.8 equivalents of cysteine injected.**

The peaks corresponds to the following species, 1.76 ppm peak corresponds to MeAs(Cys)<sub>2</sub>, 1.64 ppm peak to MeAs(OH)(Cys) and 1.358 ppm peak to MeAs(OH)<sub>2</sub>. The system undergoes the following equilibrium:



From the integrals on the NMR spectra of the system under during the titrations it is possible to work out the concentration of each species after each addition.

As the concentration of the species change due to the dilution caused by the titration, it is easier to look the species in terms of molar ratios:



**Figure 19: NMR titration of Cysteine against MeAs(OH)<sub>2</sub> in D<sub>2</sub>O at pH 4.5**


We clearly see as more cysteine is titrated into the solution, cysteine would displace the hydroxyl group forming Me<sub>2</sub>As(Cys)<sub>2</sub> and Me<sub>2</sub>As(Cys)(OH). It is interesting to see that the concentration of Me<sub>2</sub>As(Cys)(OH) remains relatively constant whilst the preferred product Me<sub>2</sub>As(Cys)<sub>2</sub> increases over time until it becomes the dominant species.

### 3.2.4 Temperature sensitivity of the methyl peak

As arsenic has a lone pair, the arsenic species MeAs(OH)(Cys) is chiral and forms an overall diastereomer with the chiral α Carbon in the cysteine.

This should result in the presence of two peaks for this species as opposed to the singlet that we observe. This suggests there might be dynamic exchanges interactions occurring that is causing the signal to average out. In addition, if the cysteines are labile like in the Me<sub>2</sub>AsCys case, we might also observe the coalescence of all the methyl peaks.

A preliminary NMR experiment was done with a system with 65 mM of (MeAsO)<sub>x</sub> and 77mM of Cysteine at 25°C and 40°C.



**Figure 20: Temperature variation on a solution of MeAs(OH)<sub>2</sub> and Cys. Top: 313K, bottom 293K**

The results indicate that there is minimal change to the peaks of the arsenic bound methyls and the coalescence of these peaks was not observed.


Further experiments into lower temperatures are necessary observe if the peaks would split due chirality.

### 3.3.1 Zinc Fingers

Recent advances<sup>36,37</sup> into therapeutic arsenic trioxide chemistry has led to the identification of the zinc finger contained within the PML-RAR $\alpha$  as a possible site of interaction. PML-RAR $\alpha$  contains two zinc fingers, PML-R-ZF1 is a 4 cysteine zinc finger and **PML-R-ZF2** contains 3 cysteine<sup>38</sup> and 1 histidine. It is proposed that arsenic interacts directly with cysteine residues and it is this interaction which leads to the eventual death of the cell. Although the group showed there was interaction between the protein and arsenic, the mechanism and origin of the specificity of this interaction is unclear. Even more unclear is how arsenic could displace the zinc from a four co-ordinated site.

The key part of the zinc finger is a four co-ordinate zinc which holds the finger together by providing structural support<sup>39</sup>. The “finger” aspect is provided 16 generic residues in the form of a loop<sup>40</sup>. The zinc could be co-ordinated to either the sulfur on cysteine or the nitrogen on histidine molecules. As one of the zinc fingers in the PML-RAR $\alpha$  protein is a 4 cysteine zinc finger. After looking at existing zing finger analogues<sup>41,40</sup> and searching the Cambridge crystal database

for solved crystal structures of zinc fingers<sup>42,43</sup>, we propose the follow molecules as models for zinc fingers:




**Figure 21: Structures of proposed zinc finger models**

Compounds that contained tetrahedral zinc molecules bound to four sulfurs are preferred as they are better models for zinc fingers. Zinc 1,2 ethane dithiolate (**Zn1b**) is a charged species. We thought this would allow for reactions with charged arsenic species.

*O,O*-dialkyldithiophosphate zinc(II) (**Zn2a**) is a neutral species which contains a phosphorus linking the sulfurs together. Although phosphates are not present in zinc fingers, having such a group allows us the study the system using the phosphorus NMR. This is a major advantage as the phosphorus is very close to the sulphurs and will be a major indicator of interactions that occur.

*Experimental procedure for Zn1b*



**Figure 22: Preparation of Zn1b**


*Preparation of 1,2 ethanedithiol sodium salt<sup>44</sup>:* Dry tetrahydrofuran (200 ml, previously distilled from sodium and stored inside glove box) was added to finely divided sodium (5.0, 220 mmol). Ethanedithiol was added (10 ml, 95.5 mmols) and the solution was headed under reflux for 5 days until no pieces of sodium are seen. The white precipitate was filtered using Schlenk apparatus and dried under vacuum. The resulting white solid had a mass of 13.44g (92.8 mmols, 97% yield). The salt had very low solubility in all solvents and hence a NMR was not possible.

*Preparation of (Et<sub>4</sub>N)<sub>2</sub> [Zn(S<sub>2</sub>C<sub>2</sub>H<sub>4</sub>)<sub>2</sub>]:* The original preparation called for the use of tetraethyl ammonium chloride. This was used in the initial run of the experiment. The yield (c. 10%, purified but decomposed shortly after) of the first attempt was extremely low due to the water sensitive nature of the product. As a result the experiment was modified to use tetraethyl ammonium iodide which is

less hygroscopic than tetraethyl ammonium chloride. Zinc chloride (0.7699g, 5.66 mmols) and tetraethylammonium iodide (3.200g, 12.4 mmols) was stirred in a solution of dry acetonitrile (50ml), forming a clear solution. Ethanedithiol sodium salt (1.56g, 11.3 mmols). The white solid did not dissolve. The cloudy solution was left stirring for 1 day. The solution was filtered and the filtrate was concentrated to 10ml. Ether was added and the solution was cooled overnight. The 0.70 of solid crystals were collected with a 25% yield.  $^1\text{H}$  NMR (400 MHz, DMSO)  $\delta$  3.35 (s, 33H), 3.21 (d,  $J = 7.3$  Hz, 16H), 2.50 (s, 13H), 1.24 – 1.08 (m, 24H).

In the  $^1\text{H}$  NMR there is a peak at 3.21 ppm which can be assigned to be the  $-\text{CH}_2$  and the 1.16 peak is the  $-\text{CH}_3$  on the ethyl. The 3.35 peak could be assigned to be the  $\text{CH}_2$  peaks on the ethanedithiol. The integrals suggest there are more ( $\text{S}_2\text{C}_2\text{H}_4$ ) than the formula suggests. This is most likely due to remaining sodium ethanedithiol salt which is difficult to purify.

*Experimental procedure for **Zn2a**:*



**Figure 23: Reaction scheme for the preparation of Zn2a**

Potassium diethyl dithiophosphate 0.50g and zinc chloride, 0.20g, was dissolved in 20ml of acetonitrile. After 30 minutes, the colorless solution turned cloudy with a white precipitate. Stirring was continued for 5 hours. The solution was



subsequently filtered, and the filtrate was concentrated yielding the product as a colorless solid. Product was recrystallized into a colorless solid with 32% yield. The low yield is possibly attributed to an incomplete recrystallization.  $^1\text{H}$  NMR (200 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  4.14 (dq,  $J = 9.5, 7.1$  Hz, 2H), 1.32 (td,  $J = 7.1, 0.5$  Hz, 3H).  $^{31}\text{P}$  NMR (81 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  105.69 (s).


This reaction gave a much cleaner product than ethanedithiol and the purification was much easier. The yield is low, but this most likely due to product precipitating out of solution and getting filtered with the NaCl. This is verified by NMR the filtrand which was similar NMR peaks to the final product. Although this is not optimal in terms of being a model for zinc fingers, the zinc is still coordinated to 4 sulfurs and thus still serving as a chemical model.

Although this compound looks plausible, many five and six coordinate zinc complexes are known, the behaviour of the complex in solution is not. To begin with, in addition to being a strong chelating ligand, the dithiophosphate ligand can bridge between zinc centers and thus form either dimeric or polymeric structures<sup>45</sup>. If this process is fast compared to the NMR timescale, the spectroscopic result would be a single peak with an average chemical shift.

The initial attempt to co-ordinate arsenic with the zinc complex was by the use of triphenylarsine which is an excellent Lewis base. Triphenylarsine was added to a solution of **Zn2a** in  $\text{CDCl}_3$  and possible interactions were observed via

NMR.  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.32 – 7.30 (m, 2H), 4.33 – 4.14 (m, 2H), 1.50 – 1.17 (m, 3H).  $^{31}\text{P}$  NMR (81 MHz,  $\text{CDCl}_3$ )  $\delta$  97.68 (s). The  $^1\text{H}$  NMR spectrum indicates a multiplet at 7.3 ppm which corresponds to the hydrogens on triphenyl arsine. The ethyl peaks are present and in the correct integration. The resolution of the peaks seems reduced and hence we can no longer distinctly see the doublet of quartets and triplet of doublets seen in the pure sample of **Zn2a**. While this might be due to the shimming of the sample, both changes could also be due to signal averaging and rapid exchange. There is only one phosphorus signal suggesting there is only one averaged species present. Overall, the NMR does not suggest a strong interaction between the triphenylarsine and the complex. (However, given the interaction seen later on with  $\text{Me}_2\text{AsCys}$  and the fast exchange which results in a single peak for the phosphorus, this experiment should be looked at again).


The second attempt to co-ordinate arsenic to the zinc complex involved the use of  $\text{Me}_2\text{AsCys}$ .  $\text{Me}_2\text{AsCys}$  was added to a solution of **Zn2a** in Chloroform. The following stacked spectra shows the  $-\text{CH}_3$  group of **Zn2a** on top and the NMR of the mixture after the addition of  $\text{Me}_2\text{AsCys}$ .



**Figure 24: Stacked <sup>1</sup>H NMR of Zn2a (top) and the same solution following the addition of Me<sub>2</sub>AsCys**

The most important peak is the broad peak at 1.85 ppm which indicates an exchange process is happening. This peak is from the methyls on Me<sub>2</sub>AsCys. The phosphorus NMR, indicated a single peak at 98.0 ppm. This is quite surprising as interaction was seen in the <sup>1</sup>H NMR spectrum and one would expect multiple peaks in the phosphorus spectra if the diethyl dithiophosphate ligand was being displaced by Me<sub>2</sub>AsCys. It was decided free ligand and zinc complex interaction should be looked at in the form of a NMR titration.

NMR titration experimental: 0.011g of **Zn2a** was dissolved in 1ml of CD<sub>3</sub>OD. 0.4 ml of this solution was placed into the NMR tube. A solution of potassium diethylphosphoro dithiolate was made with 0.033g of the compound in 1.5 ml of CD<sub>3</sub>OD. 1ml of this solution was taken then in injected 0.05 ml aliquots to the NMR tube. The titration indicated that instead of having separate peaks for bound ligand and free ligand, exchange was occurring and hence only a single peak was observed. The following plot shows the chemical shift of the phosphorus signal after addition free ligand (<sup>31</sup>P NMR (81 MHz, CD<sub>3</sub>OD)  $\delta$  111.99 (s)).




**Figure 25: Titration of ligand against zinc salt**

The data could be fit to an exponential decay curve with the following formula:

$y = A1 \cdot \exp(-x/t1) + y0$  with  $y0 = 112.24509$ ,  $A1 = -6.24402$ ,  $t1 = 2.07497$  with  $R^2 = 0.9992$ .

The good fit to the exponential decay equation indicates that there is one dominating exchange reaction that is happening. In addition to this, the curve does not match that of a simple average of the shift of the free and bound ligand, indicating that there is indeed an interaction and a third species as opposed to simple averaging of the peaks.

Due to the slow timescale of NMR, a fluorescent zinc finger model composed of zinc complex to 8-mercaptoquinoline is proposed. This could be synthesized using the following scheme.



**Figure 26: Synthesis of 8-mercapto quinoline zinc salt.**


Starting material 8-aminoquinoline had a melting point of 82°C. <sup>1</sup>H NMR (400 MHz, dmsO) δ 8.72 (dd, J = 4.1, 1.6 Hz, 1H), 8.17 (dd, J = 8.3, 1.6 Hz, 1H), 7.68 – 6.54 (m, 4H), 5.90 (s, 2H).

Preparation of **Zn3a**. 8-aminoquinoline (Light green powder, 82°C melting point) (0.3460g) was placed in a round bottom flask. HBr (48%, from Gleason Group, light yellow in color) (1.7ml) was added. The solid turned to deep orange and dissolved upon addition of water (11 ml) forming a transparent orange solution which was cooled by the addition of ice. A solution of 0.21 g of sodium nitrite in 2 ml of water was added to the


cooled mixture. There is little sign of a visible reaction. The solution was then added to a warm solution of thiourea (0.19g in 5 ml of water). Gas evolution started upon warming to 40°C. The product was recrystallized using ethanol and ether yielding a deep orange crystals with a 89% yield.  $^1\text{H}$  NMR (400 MHz, DMSO- $\text{D}_6$ )  $\delta$  9.39 – 8.61 (m, 1H), 8.45 (d,  $J$  = 8.2 Hz, 1H), 8.03 – 7.21 (m, 4H).

#### Fluorescence data:

8-aminoquinoline sodium salt was dissolved in ethanol. The UV spectra indicated two peaks, one at 243 nm and 321 nm. Using this as a starting point, emission spectrum shows a maxima at 450 nm. The excitation spectrum showed 2 major excitation peaks at 270nm and 365nm. (see next page)



**Figure 27: Excitation spectra of 8-aminoquinoline in ethanol (450 emission)**



**Figure 28: Emission spectra with 365 nm excitation**

Given that the ligand is fluorescent, the next step is to synthesize the complex by reacting the 8-amino quinolone with  $\text{ZnCl}_2$ . Unfortunately due to time constraints, this part of could not be completed.

### 3.3 Conclusions

Throughout the course of this project it was shown that both methylarsenic and dimethylarsenic undergo facile could be exchange between different thiol adducts such as cysteine and glutathione. In the case of dimethylarsenic cysteine, the methyl exchange could be studied using DNMR – elucidating the entropy and enthalpy of the interaction. In addition to transfer between the thiols of cysteines, we also saw dimethylarsenic transfer between cysteine and glutathione groups. Whilst it was not possible to directly model this interaction, it was qualitatively shown that arsenic could move between various thiols. In

addition it was shown that monomethyl arsenic species also have bond lability by titrating cysteine to a solution of monomethylarsenious acid. I have identified the formation of both  $\text{MeAs}(\text{Cys})(\text{OH})$  and  $\text{MeAs}(\text{Cys})_2$ , and these concentrations would change depending on the amount of cysteine added. Unfortunately it was not possible to explore the kinetics of these species using the DNMR as line shapes did not change with temperature. Further kinetic studies of these systems could be done using stopped flow techniques.

The ability of arsenic to rapidly break and form new bonds has important biological implications. This rapid exchange mechanism gives rise to the possibility that arsenic is transported via a shuttle mechanism where it hops into to various species and is carried around the body. In addition to binding to small molecules such as cysteine and glutathione, arsenic could also bond to vicinal cysteines of large proteins, disrupting its function. Understanding this interaction is key to understanding the mechanism of arsenic based drugs such as ATO and Darinaparsin.

### 3.4 References

33. Burrows, G. J.; Turner, E. E., A new type of compound containing arsenic. *Journal of the Chemical Society, Transactions* **1920**, 117, 1373-1373.



34. Marsmann, H. C.; Wazer, J. R. V., Methylarsaoxanes. Structural chemistry of cacodyl oxide and arsenosomethane in the liquid state. *JACS* **1970**, *363* (2), 3969-3972.
35. Petrick, J. S.; Jagadish, B.; Mash, E. A.; Aposhian, H. V., Monomethylarsonous Acid (MMA III ) and Arsenite: LD 50 in Hamsters and In Vitro Inhibition of Pyruvate Dehydrogenase. *Chem. Res. Toxicol.* **2001**, *14* (6), 651-656.
36. Zhang, X.-W.; Yan, X.-J.; Zhou, Z.-R.; Yang, F.-F.; Wu, Z.-Y.; Sun, H.-B.; Liang, W.-X.; Song, A.-X.; Lallemand-Breitenbach, V.; Jeanne, M.; Zhang, Q.-Y.; Yang, H.-Y.; Huang, Q.-H.; Zhou, G.-B.; Tong, J.-H.; Zhang, Y.; Wu, J.-H.; Hu, H.-Y.; de The, H.; Chen, S.-J.; Chen, Z., Arsenic Trioxide Controls the Fate of the PML-RAR{alpha} Oncoprotein by Directly Binding PML. *Science* **2010**, *328* (5975), 240-243.
37. Borden, K. L.; Lally, J. M.; Martin, S. R.; O'Reilly, N. J.; Solomon, E.; Freemont, P. S., In vivo and in vitro characterization of the B1 and B2 zinc-binding domains from the acute promyelocytic leukemia protooncoprotein PML. *Proceedings of the National Academy of Sciences* **1996**, *93* (4), 1601-1601.
38. Borden, K. L. B.; Boddy, M. N.; Lally, J.; Oreilly, N. J.; Martin, S.; Howe, K.; Solomon, E.; Freemont, P. S., THE SOLUTION STRUCTURE OF THE RING

FINGER DOMAIN FROM THE ACUTE PROMYELOCYTIC LEUKEMIA PROTO-ONCOPROTEIN PML. *Embo J.* **1995**, *14* (7), 1532-1541.

39. Reddy, P. R.; Radhika, M., Synthesis and characterization of mixed ligand complexes of Zn (II) and Co (II) with amino acids: Relevance to zinc binding sites in zinc fingers. *Journal of Chemical Sciences* **2005**, *117* (3), 239-246.

40. Dudev, T.; Lim, C., Modeling Zn<sup>2+</sup>-cysteinate complexes in proteins. *The Journal of Physical Chemistry B* **2001**, *105* (43), 10709-10714.

41. Barnes, A. M.; Bartle, K. D.; Thibon, V. R. A., A review of zinc dialkyldithiophosphates (ZDDPS): characterisation and role in the lubricating oil. *Tribology International* **2001**, *34* (6), 389-395.

42. Segal, D. J.; Crotty, J. W.; Bhakta, M. S.; Barbas Iii, C. F.; Horton, N. C., Structure of Aart, a designed six-finger zinc finger peptide, bound to DNA. *J. Mol. Biol.* **2006**, *363* (2), 405-421.

43. Pavletich, N. P.; Pabo, C. O., Crystal structure of a five-finger GLI-DNA complex: new perspectives on zinc fingers. *Science* **1993**, *261* (5129), 1701-1701.

44. Rao, C. P.; Dorfman, J. R.; Holm, R. H., Synthesis and structural systematics of ethane-1,2-dithiolato complexes. *Inorganic Chemistry* **1986**, *25* (4), 428-439.

45. Ivanov, A. V.; Antzutkin, O. N.; Larsson, A.-C.; Kritikos, M.; Forsling, W., Polycrystalline and surface O,O'-dialkyldithiophosphate zinc(II) complexes: preparation,  $^{31}\text{P}$  CP/MAS NMR and single-crystal X-ray diffraction studies. *Inorganica Chimica Acta* **2001**, 315 (1), 26-35.