

Searches for Non-Resonant New Physics in the High Energy Di-Electron Spectrum with ATLAS at the LHC

Liam Duguid

Department of Physics,
Royal Holloway, University of London,
Egham, Surrey, UK, TW20 0EX.
`lduguid@cern.ch`

Supervisor: Dr. Tracey Berry

A thesis submitted to the University of London for the
Degree of Doctor of Philosophy

May 29, 2014

DECLARATION

I confirm that the work presented in this thesis is my own. Where information has been derived from other sources, I confirm that this has been indicated in the document.

Liam Duguid

Write thanks here

Abstract

Write abstract here

Preface

- **Chapter 1: Theory**

This chapter covers an overview of the Standard Model (SM) of particle physics and then continue on to Beyond the Standard model (BSM) phenomena. The main focus is on the idea of Non resonant excesses in the dilepton Drell-Yan (DY) spectrum of which two examples are discussed. The first example is Contact Interactions, a model which describes many BSM phenomena that can show as four fermion contact interaction that exhibit a divergence from the SM DY spectrum. The example shown is that of a quark-lepton composite model where at a certain energy level quarks and leptons can form composite particles. The second example given is the Arkani-Hamed, Dimopoulos, and Dvali (ADD) model. This is a graviton theory with the addition of large extra spacial dimensions to dilute gravity. These large extra spacial dimensions create Kaluza-Klein resonances of the graviton very close to each other and so exhibit signs of Non-resonance behaviour. A look at past results for similar searches is also discussed here.

- **Chapter 2: Experiment**

This chapter covers an overview of the ATLAS experiment and the LHC. A particular focus will be given to the inner tracking detector and energy Calorimeters of ATLAS as these systems are the parts used in the detection of di-electron events used in this analysis. Although parts of the detector will be discussed in some respect.

- **Chapter 3: Trigger**

This chapter focuses on the triggering system for selecting data events in the ATLAS detector. An overview of the whole system will be given but a focus made on the “egamma” part which selects electron and photon events. A slight detour will be made discussing the effect of increases in the luminosity of the LHC beam through the 2011-2012 data taking period and efforts taken to reduce high rates of data acquisition this entailed in the “egamma” chain.

- **Chapter 4: Reconstruction**

This chapter details the algorithms used in reconstructing electrons and photons from the detector

output. It also contains a discussion on ATLAS assignments of *tight*, *medium* and *loose* electrons.

- **Chapter 5: Event Selection**

This chapter covers the main event selection of di-electron events for the non-resonance analysis on the 20 fb^{-1} recorded in 2012. There will also be a discussion of and need for corrections applied to energy measurements.

- **Chapter 6: Background Estimate**

This chapter discusses the estimate made of the background processes to the non-resonant signal. It covers the Monte Carlo (MC) generated to estimate these backgrounds as well as corrections applied to match MC to the data collection conditions used and corrections to account for next to next to leading order calculation effects.

- **Chapter 7: Signal Search**

This chapter shows the search for new physics in the data collected in the 2012 data taking period. This includes a description of the MC used to predict the signals as well as comparison between the Data and the MC prediction of the background. Also looked at are the significance or p-value of any divergences from the SM background prediction.

- **Chapter 8: Statistical Analysis and Conclusion**

This final chapter discusses a statistical treatment of the results. First discussed is possible sources of systematic error in the analysis as well as levels of statistical error. Next a Bayesian approach is taken to setting lower limits on the scale of new physics predicted by this analysis. Last is a discussion of how this result can be interpreted and a comparison to past searches for similar phenomena and current ones. A_μ

Trigger work:

- optimisation of 2012 electron triggers at LV1 up to HLT for higher luminosity conditions and the required rate reductions.

Z Prime Analysis:

- Ran analysis of several MC sample for people. (including Black holes, pythia DY and samples for

Section 3.4.2 There was also a contribution to the maintenance of the e/γ trigger software run in the ATLAS detector, both these tasks forming the authorship qualification. The authorship task culminated

in presentation of a poster on behalf of the e/γ trigger group at the Computing and High Energy Physics (CHEP) conference held New York in May 2012.

Non-resonant Analysis:

- Full analysis coded and run by liam.
- production of reverse ID jets sample for Non-resonant analysis.
- optimisation of new isolation cut.
- Study of new opposite sign cut and effect on reverse ID jets sample.
- Study of $\cos\Theta^*$ variable data MC comparison in control region.
- optimisation of binning for statistical analysis.
- Limit setting via Bayesian analysis toolkit.

Contents

Preface	i
Contents	iv
1 Theory	1
1.1 Standard Model	1
1.2 Non-resonant new Physics	1
1.2.1 Contact Interaction Theory	2
1.2.2 ADD Theory	3
1.3 Past Searches	5
2 Experiment	6
2.1 The Large Hadron Collider	6
2.2 ATLAS - A Toroidal LHC Apparatus	8
2.2.1 Inner Detector	10
2.2.2 Calorimeters	12
2.2.3 Magnet System	14
2.2.4 Muon Spectrometer	15
3 The Trigger & Data Acquisition	17
3.1 Level-1 Trigger	17
3.2 Higher Level Trigger	19
3.3 Data Acquisition	20
3.4 Trigger Menu and Rates	20
3.4.1 The “ e/γ ” Trigger Menu	22
3.4.2 Trigger Rates in High Luminosity Regime	23

4	Reconstruction	25
4.1	Electron Reconstruction	26
5	Event Selection	28
5.1	Analysis Selection	28
5.1.1	Isolation Requirement	29
5.2	Opposite Sign requirement	31
5.3	Energy Resolution Correction	31
5.4	Selection Acceptance \times Efficiency	31
6	Background Estimate	34
6.1	Monte Carlo samples	34
6.1.1	MC Corrections	35
6.2	Fake Factor Multi-Jet Estimate	36
6.2.1	Real electron efficiency estimation	38
6.2.2	Fake electron rate estimation	38
6.2.3	Properties of Multi-Jet Background	40
6.2.4	Other methods and estimation of Error	40
7	Signal and Results	43
7.1	Signal Monte Carlo	43
7.2	Signal Parametrisation	44
7.3	Results	44
8	Statistical Analysis	49
8.1	Systematics	50
8.2	Signal Search & P-Values	51
8.3	Setting Limits	51
8.4	Combination with the Muon Search	52
9	Conclusion	56
9.1	Interpretation	56
9.2	Looking Forward	56
	Appendices	57

A Non-Resonance Analysis 2011	58
A.1 Data and Background Processes	58
A.1.1 Data	58
A.1.2 Background	58
A.1.3 Background Production	58
A.1.4 Signal	59
A.2 Electron Identification and Selection	59
A.2.1 Data and Background Comparison	61
A.2.2 New Physics Signal Expectation	61
A.3 Statistical Analysis	65
A.4 Results	66
Bibliography	67

Chapter 1

Theory

1.1 Standard Model

The Standard Model of particle physics has proven excellent at describing particle interactions up to the energy scale of modern colliders ($\sim \text{TeV}$) and with the discovery of a Standard Model like Higgs Boson at the LHC the theory will be able to claim completeness up to the energy scale of modern colliders. However the Standard Model is known to be incomplete, with observations such as neutrino mass, the lack of anti-matter in the observable universe and the lack of a quantum gravity description with the related hierarchy problem (drastic difference in force strength between gravity and the other fundamental forces seen in the standard model), the Standard Model is far from a theory of Everything. This then leaves the possibility of new physics appearing in the energy scope of the LHC.

1.2 Non-resonant new Physics

Beyond the Standard Model (BSM) or new physics models is a staple of the physics programs of the LHC detectors. Any theoretical models not contained within the Standard Model (SM) can fall in to this category and LHC experiments aim to search for as many of these models as are feasible within scope (Proton-Proton collisions and within the energy range of the LHC). Within the detection channel of two lepton decays (di-lepton), one evidence of new physics is non-resonant signals. This physics would show as a divergence from the SM prediction in the di-lepton mass spectrum unlike the resonant signals of particles such as the Z boson particle which shows as a peak in the di-lepton mass spectrum.

Non-resonant signals could be the results of many BSM theoretical models but two main theorys are presented here and their searches compose the rest of this thesis.

1.2.1 Contact Interaction Theory

The Standard Model (SM) assumes quarks and leptons to be fundamental particles in nature. This assumption is not without compelling argument but like the proton beforehand there is no reason quarks and leptons should not be composite structures or bound states of more fundamental particles, often referred to as Preons [1], at an energy scale Λ we have yet to reach.

One way quark-lepton compositeness would exhibit itself is in 4-fermion contact interactions between two quarks from the incoming protons producing two final state leptons ($q\bar{q} \rightarrow \ell^-\ell^+$). This is the compositeness signal searched for at the ATLAS detector and as can be seen the the Feynman diagrams in Fig. 1.1 compared to DY from which it is almost indistinguishable.




Figure 1.1: Feynman diagrams of a contact interaction (right) and the predominant background Drell-Yan production (left).

Without knowing the intermediate process one can write a Lagrangian describing the new interaction;

$$\mathcal{L} = \frac{g^2}{2\Lambda^2} [\eta_{LL}(\bar{\psi}_L \gamma_\mu \psi_L)(\bar{\psi}_L \gamma^\mu \psi_L) + \eta_{RR}(\bar{\psi}_R \gamma_\mu \psi_R)(\bar{\psi}_R \gamma^\mu \psi_R) + 2\eta_{LR}(\bar{\psi}_L \gamma_\mu \psi_L)(\bar{\psi}_R \gamma^\mu \psi_R)] \quad (1.1)$$

where g is the coupling constant, Λ is the energy scale of new physics and ψ_L and ψ_R are the left and right handed fermionic fields respectively. The sign of η defines whether the new interaction interferes constructively ($\eta = -1$) or destructively ($\eta = +1$) with DY and is always unity. For previous analyses [2, 3, 4] a benchmark model of just the Left-Left (LL) component has been used and defined by $\eta_{LL} = \pm 1$ and $\eta_{RR} = \eta_{LR} = 0$. Here however an investigation of each of the three parameters is done individually. Both the LL and RR cases are expected to behave similarly however the LR case exhibits a different angular dependence than either of the other formalisms or the DY background. This difference is the primary reason for the inclusion of the angular part of this analysis found later. The discriminating variables used are therefore dilepton invariant mass and cosine of the decay angle θ^* . The angle θ^* is defined in the Collins-Soper frame [5] which is defined with the x -axis perpendicular to the incoming parton momentum frame and the z -axis bisecting the angle between the two incoming parton momenta. Since the incoming parton information is understandably unavailable the z -axis is taken as the direction of the incoming quark (as opposed to anti-quark) obtained from the boost in to the dilepton frame. The angle θ^* is then defined as

the angle between this z -axis the momentum of the outgoing negatively charged lepton (or electron in this analysis).

Figure 1.2 shows the difference expected between the LR CI models and DY background from a truth Monte-Carlo study. The variables used are A_{FB} and dilepton invariant mass where A_{FB} is the forward-backwards asymmetry defined in relation to $\cos\theta^*$ as;

$$A_{FB} = \frac{N_F - N_B}{N_F + N_B} \quad (1.2)$$

where N_F and N_B are number of events found with $\cos\theta^*$ greater than 0 and less than 0 respectively.

Figure 1.2: MC truth level comparison between the forward backwards asymmetry of DY and of a CI LR signal.

A differential cross section for this interaction, $q\bar{q} \rightarrow \ell^-\ell^+$ ($qq\ell\ell$), is given by

$$\frac{d\sigma}{dm_{\ell\ell}} = \frac{d\sigma_{DY}}{dm_{\ell\ell}} - \eta \frac{F_I}{\Lambda^2} + \frac{F_C}{\Lambda^4}, \quad (1.3)$$

where $m_{\ell\ell}$ is the dilepton mass and Λ is the scale of the new physics. In the case of quark/lepton compositeness Λ refers to the point at which fermions stop being bound as SM quarks and leptons. F_I and F_C define the interference DY-CI term and the pure CI term respectively. The scale of the interference and pure term vary with both the dilepton invariant mass as well as the scale of new physics Λ .

Experimentally this interaction would be seen as a deviation from the Standard Model (SM) Drell-Yan (DY) ($q\bar{q} \rightarrow \gamma/Z \rightarrow \ell^+\ell^-$) dilepton mass spectrum as seen in Fig. 1.3).

Figure 1.3: MC truth level comparison between DY spectrum with and without CI signal.

1.2.2 ADD Theory

The large extra spacial dimensions theory described by Arkani-Hamed, Dimopoulos, and Dvali (or ADD theory) [6] predicts a Graviton producing a non-resonant excess in the dilepton mass spectrum. The ADD theory describes a graviton that propagates the extra dimensions and acquires closely backed Kaluza-Klein (KK) modes that show as a broad excess above the SM background. The ADD model was originally proposed to solve the hierarchy problem and bring the energy scale associated with gravity (the Planck scale $M_{Pl} \sim 10^{16}$ TeV) down to the level of electroweak energy scale ($M_{EM} \sim 10^{-1}$). This is achieved with the introduction of n additional compactified spacial dimensions with radius R . This then gives a new scale in the $4+n$ dimensional space, M_D , which is related to the Planck scale by $M_{Pl} = M_D^{n+2} R^n$. If both the radius

of the extra dimensions R and number n are large enough this solves the hierarchy problem by bringing M_D down to the level of M_{EM} . The Graviton is the only propagator in these extra n dimensions with each dimension resulting in a new KK mass splitting of the mass peak. The mass splitting occurs with an interval of $1/R$ and since R is required to be large by the theory this pushes the mass splitting together causing a continuous peak like structure analogous to a non-resonant excess. This sum over these virtual KK modes has to be regularised by an “ultra violet” cutoff (Λ_T) and it is convention to equate this cutoff to the onset of Quantum Gravity (M_S) only bellow which the theory is valid. This scale M_S is used as the scale of physics for the ADD theory which is a low energy effective theory bellow this scale. This scale can be related to the new n dimensional Planck scale (M_D) by;

$$M_S = 2 \sqrt{\pi} [\Gamma(n/2)]^{1/(n+2)} M_D \quad (1.4)$$

where Γ is the decay width. Bellow the scale M_S virtual Graviton exchange would lead to a broad excess over the SM Drell-Yan dilepton mass spectrum. The total differential cross-section for the dilepton SM DY and virtual Graviton exchange is then;

$$\frac{d\sigma}{dm_{\ell\ell}} = \frac{d\sigma_{DY}}{dm_{\ell\ell}} + \mathcal{F} \frac{F_I}{M_S^4} + \mathcal{F}^2 \frac{F_G}{M_S^8} \quad (1.5)$$

where σ_{DY} is the SM DY cross-section, F_I and F_G are the Graviton-DY interactions term and pure virtual Graviton exchange term respectively while \mathcal{F} is a formalism dependent parameter and also dimensionless. Three formalisms are commonly used to describe ADD theory, these are Giudice, Rattazzi, and Wells (GRW) [7], Han, Lykken, and Zhang (HLZ) [8] and Hewett [9]. Defining \mathcal{F} these formalisms alter the cross-section of virtual Graviton exchange with HLZ depending on the number of extra dimensions, n , introduced by the ADD theory. All three formalisms are detailed in Eq. 1.6

$$\begin{aligned} \mathcal{F} &= 1, & (\text{GRW}) \\ \mathcal{F} &= \begin{cases} \log\left(\frac{M_S^2}{m_{\ell\ell}^2}\right), & (n=2) \\ \frac{2}{n-2}, & (n>2) \end{cases}, & (\text{HLZ}) \\ \mathcal{F} &= \frac{2\lambda}{\pi} = \frac{\pm 2}{\pi}, & (\text{Hewett}) \end{aligned} \quad (1.6)$$

Of note the variable λ found in the Hewett formalism defines the constructive or destructive nature of the gravitational interaction with the SM DY processes. λ is always of order unity with +1 and -1 being constructive and destructive respectively. The GRW and HLZ with $n=2$ are the two formalisms explicitly searched for in this analysis with a conversion of limits done to asses the other formalisms in the Statistical Analysis chapter.

Experimentally this interaction would be seen as a deviation from the SM DY ($q\bar{q} \rightarrow \gamma/Z \rightarrow \ell^-\ell^+$) dilepton mass spectrum but with a cut-off where quantum gravity is assumed take effect. This can be seen in Fig. 1.4. It is important to note that no angular difference in the virtual graviton decay is expected from that of SM DY, therefore ADD is only searched for in the dilepton mass spectrum.

Figure 1.4: MC truth level comparison between DY spectrum with and without ADD signal.

1.3 Past Searches

Contact Interaction

Several previous CI analyses have been done at hadron colliders including the LHC [4, 10, 11, 12] and the Tevatron [2, 3, 13, 14, 15]. Searches were also performed at the electron-proton collider HERA [16, 17], previous lepton colliders [18, 19, 20, 21, 22] and neutrino scattering experiments []. Of the results comparable to this analysis searching for $qq\ell\ell$ Contact Interactions in the absence of signal the highest limits set on the scale of new physics Λ come from the previous ATLAS analysis [4] detailed in appendix A. This analysis set a limit of $\Lambda > 12.7$ TeV and $\Lambda > 9.63$ TeV for the dilepton LL CI model for constructive and destructive interference respectively. The limits obtained for the electron channel for comparison to this analysis were $\Lambda > 11.6$ TeV for constructive and $\Lambda > 8.76$ TeV for destructive interference.

ADD

The highest dilepton ADD limits set on the formalism normally used as a bench mark, GRW, are that of the previous ATLAS analysis [4] discussed in Appendix A which sets a limit of 3.0 TeV on the scale of new physics (M_5). Preliminary results from CMS on the LHC data set used in this analysis do however show higher limits but are not yet published. Other previous analyses have also been carried out searching for large extra dimensions with the ADD model. These analyses have come from the LHC [], from the Tevatron [], as well as from electron-proton collider HERA [] and electron-positron collider LEP [].

Chapter 2

Experiment

This chapter will explore the ATLAS experiment in order to explain how data specific to this analysis is obtained. First however is a discussion of the Large Hadron Collider which supplies the ATLAS experiment with proton collisions.


2.1 The Large Hadron Collider

The Large Hadron Collider (LHC) [24] is the largest and most powerful particle collider in the world with a circumference of 27 km and design centre of mass collision energy of 14 TeV. During the 2012 run the accelerator was run at a centre of mass energy of 8 TeV while providing an integrated luminosity of just above 20 fb^{-1} throughout the year to its two general purpose experiments, CMS and ATLAS, that latter of which provided data for this analysis.

The LHC itself is built in the same tunnel (see fig. 2.1) as was used by the Large Lepton-Positron (LEP) collider. Based at CERN (Centre of European Nuclear Research) the 27 km tunnel is between 50 to 175 m underground and like CERN itself crosses the French-Swiss border just outside Geneva. Construction of the LHC started in 2001 after the LEP collider was decommissioned and dismantled with excavation of the caverns for the LHC's four main experiments starting slightly before in 1998. The LHC is a synchrotron machine requiring 1,232 super-conducting Niobium-Titanium dipole magnets each providing an 8.33 T magnetic field to direct the proton beams around its loop and an additional 392 quadrupole magnets of the same type to focus the beams for the collision points. The super conducting magnets operate at 1.9 K with the whole accelerator requiring 96 tonnes of liquid helium to remain cooled.

For the 2012 run used in this analysis the LHC ran with xxxx proton bunches travelling in each direction which were accelerated around the LHC with an interval of 50 ns between bunches and with each bunch

LHC PROJECT



ST-CE/JLB-hlm
18/04/2003

Figure 2.1: Schematic of LHC tunnel with all its caverns showing those added in preparation for the LHC. This shows the position of the LHC's 4 main detectors and SPS ring [23].

composed of xxxx protons. These run conditions gave an instantaneous luminosity of xxxx at the start of a run which slowly degraded during a run as protons collided.

However the LHC can not run in isolation to provide beams for its 4 main experiments, instead it is the last and newest accelerator in a chain of accelerators which extract protons from a hydrogen canister with little to no momentum and inject them in to the LHC as a 450 GeV beam. The proton source is a device called a Duoplasmatron which injects hydrogen gas in to a strong electric field stripping electrons from their nuclei. The remaining protons are injected in to Linac 2, a linear accelerator which accelerates them to an energy of 50 MeV. The BOOSTER or Proton Synchrotron Booster (PBS) comes next in the chain and accelerates protons from 50 MeV to 1.4 GeV to be injected in to the main Proton Synchrotron (PS). The PS accelerates protons up to an energy of 25 GeV and again injects them in to another accelerator, the Super Proton Synchrotron (SPS). The SPS (seen in fig. 2.1) is the final stage before injection in to the LHC ring and pushes protons to an energy of 450 GeV. Protons from the SPS then get injected in to the LHC in both counter revolving directions and accelerated to their final collision energy. For the data used in the analysis

that follows (the 2012 data run) the final proton beam energy is 4 TeV giving a final centre of mass collision energy of 8 TeV.

Four collision points exist around the circumference of the LHC providing collisions to the four main experiments (see fig. 2.1); ATLAS (A Toroidal LHC Apparatus), CMS (Compact Muon Solenoid), ALICE (A Large Ion Collider Experiment) and LHCb (Large Hadron Collider beauty). ATLAS and CMS are both general purpose experiments designed to look for a variety of physics. ALICE is designed specifically to study quark-gluon plasma in heavy ion collisions scheduled for the end of each LHC run period while LHCb looks for beauty mesons in searches for CP-violation. There are also three additional LHC detectors in various stages of deployment without their own collision points; TOTEM (Total Elastic and diffractive cross section Measurement), LHCf (LHC forward) and MoEDAL (Monopole and Exotics Detector at the LHC) which measure separate beam properties. TOTEM shares CMS's collision point aiming to measure the proton cross-section very accurately while LHCf shares ATLAS's collision point measuring the very forward region of collision with the hope of investigating the source of ultra-high-energy cosmic rays. MoEDAL shares a cavern with LHCb and is targeted to search for magnetic monopoles and other highly ionising stable massive particles.

2.2 ATLAS - A Toroidal LHC Apparatus




Figure 2.2: Cut-away view of the ATLAS detector. The dimensions of the detector are 25 m in height and 44 m in length. The overall weight of the detector is approximately 7000 tonnes [25].

The ATLAS detector [25] sits 100 m underground just over the road from the main CERN site and at 45 m long, 25 m in diameter and weighing over 7,000 tons is one of largest and most complex particle physics experiments in the world. The Detector itself can be divided in to four main subsystems and from the interaction point out they are; the inner detector (ID) or tracking detector, the calorimeters both Electro-Magnetic (EM) and hadronic (HCAL), the magnet system and the muon spectrometer (MS). There is also a small set of forward detectors, not detailed here, for accurate measurement of the integrated luminosity provided to ATLAS by the LHC named ALFA, LUCID and ZDC [25].

As a whole the detector has several different sets of coordinate systems some of which are used in analysis and some used primarily in detector design and placement. The first is z or the z -axis. This runs along the beam line through the centre of the detector with 0 existing at the very centre of the detector. x and y -axes do exist but are rarely needed as radial coordinates serve the purpose better. Here R is then the radial distance out from the beam line and ϕ is the angle perpendicular to R and z measuring the angle around the barrel of the detector. The last coordinate is θ measuring the angle off of the z -axis. This angle however isn't often used and in stead the angle η or pseudorapidity is used. Defined in Eq. 2.1 this quantity has the benefit of being invariant under transformation.

$$\eta = -\ln\left[\tan\left(\frac{\theta}{2}\right)\right] \quad (2.1)$$

Broadly the detector is also divided in to the barrel region (cylinder surrounding the interaction point) and endcap regions (circles covering the ends of the barrel region) which use slightly different configurations and technology in order to cover a full range in η . Following is a description of each main subsystem while focusing particularly on both the Inner Detector and EM calorimeter as these are the important systems in identification of electrons used for this analysis. This section heavily uses the ATLAS technical design report [25] throughout.

Detector component	Required resolution	η coverage	
		Measurement	Trigger
Inner Detector	$\sigma_{p_T}/p_T = 0.05\% p_T \oplus 1\%$	$ \eta < 2.5$	N.A
EM calorimetry	$\sigma_E/E = 10\%/\sqrt{E} \oplus 0.7\%$	$ \eta < 3.2$	$ \eta < 2.5$
Hadronic calorimetry (jets) barrel and end-cap forward	$\sigma_E/E = 50\%/\sqrt{E} \oplus 3\%$	$ \eta < 3.2$	$ \eta < 3.2$
	$\sigma_E/E = 100\%/\sqrt{E} \oplus 10\%$	$3.1 < \eta < 4.9$	$3.1 < \eta < 4.9$
Muon spectrometer	$\sigma_{p_T}/p_T = 10\%$ at $p_T = 1$ TeV	$ \eta < 2.7$	$ \eta < 2.4$

Table 2.1: Table showing detector components resolution requirements and η ranges for triggering and readout [25].

2.2.1 Inner Detector




Figure 2.3: Cut-away view of the ATLAS inner detector [25].

The Inner Detector is ATLAS's main tracking detector which is fitted closest to the interaction point. A tracking detector is needed to trace charged particles from the interaction point out to the calorimetry system and give two bits of information; a charged particle's position to match with the calorimeters (or Muon Spectrometer in the case of muons) and when a magnetic field is present an estimate of a particle's momentum to compare with the calorimeter obtained from the radius of its curve. The ATLAS tracking system is composed of three different tracking technologies in order going out from the collision point; the Pixel Detector (PD), the Semiconductor Tracker (SCT) and the Transition Radiation Tracker (TRT). The Inner Detector was designed to precisely measure charged tracks in the energy range 0.5 GeV - 150 GeV while complimenting the energy measurements of the calorimetry system. Covering a range of $|\eta| < 2.5$ and full range in ϕ the Inner Detector with the help of the 2 T magnetic field imposed by the solenoid magnet (discussed bellow) boasts a momentum resolution of $\sigma_{p_T}/p_T = 0.05\% p_T \oplus 1\%$ for charged tracks. In its design it was also important for the Inner Detector to be able to distinguish between multiple primary vertices at the collision point, referred to as pile-up, as well as secondary vertices from sources such as the hadronisation of b quarks.




Figure 2.4: Plan view of a quarter-section of the ATLAS inner detector showing each of the major detector elements with its active dimensions and position in Z and R detector coordinates (envelopes) [25].

Pixel Detector

The Pixel Detector is the first layer and closest to the beam line consisting of three layers of silicon pixels. Because of its proximity to the beam line the pixel detector are designed to be heavily radiation hard and understood to the degree that its performance can be predicted over an extended period of radiation exposure. The Pixel Detector is made of a barrel and two endcaps composed of 1744 modules all together.

Semiconductor Tracker

The SCT consists of the same technology as the PD but is organised in to 4 layers in the barrel region and 9 layers in each endcap. Due to the packed nature of these electronics cooling is important in this layer and so the sensors in each module are glued to each side of a thermally conductive spine that gives the SCT both structure and allows transport of heat out via the mounting point of each module keeping them at their operating temperature of -7°C .

Transition Radiation Tracker

The TRT uses a completely different tracking technology to the rest of Inner Detector using straw detectors composed of 4 mm diameter polyimide tubes each with a 31 μm diameter gold plated Tungsten-Rhenium wire. Due to the small diameter of the straws the TRT can obtain the high read-out rate needed for experiments at the LHC. The barrel region consists of 50,000 of these straws with a readout at each end providing 100,000 readout channels. The endcaps contain another 320,000 straws only read out a one end giving the TRT a total of 420,000 channels. Each channel measures drift time giving a resolution of 170 μm in each straw. The straws are filled with a high Xenon concentration ($\text{Xe}(70\%)\text{CO}_2(27\%)\text{O}_2(3\%)$) of gas in order to detect electrons via radiated photons as they traverse the material between straws (-what is this material?). This is achieved by giving each straw two timing thresholds, the lower to discriminate tracking hits (direct hits) while the higher threshold discriminates transition radiation hits.

2.2.2 Calorimeters




Figure 2.5: Cut-away view of the ATLAS calorimeter system [25].

While the Inner Detector only measures charged particles, the calorimeters measure both neutral and charged particles and are split in to two sections for particles with differing properties. The inner Electromagnetic Calorimeter is designed primarily to measure electrons and photons as well as pions while the outer Hadronic Calorimeter looks for hadrons such as neutrons and protons. In analyses the Hadronic Calorimeter is primarily used to look for jet objects (a collection of particles issuing from the decay of one mother particle). The primary method of identifying charged particles is to look for an associated track within the Inner Detector although the shape of the energy deposit in the calorimeters also helps with identification.




Figure 2.6: Schematic of the transition region between the barrel and endcap cryostats [25].

Electromagnetic Calorimeter

The Electromagnetic Calorimeter (ECAL) is designed to fully stop all electromagnetic showers within its volume. Split in to a barrel section and two endcaps the ECAL uses Liquid Argon (LAr) as a detecting medium with lead as the absorber. The lead is arranged in an accordion fashion (seen in *) to ensure consistent performance throughout ϕ . In the barrel section a presampler of LAr type is found before the main calorimeter to correct later sub-detectors for dead material. The barrel contains three layers of LAr modules of decreasing size in towards the collision point in order to keep good position resolution. The endcap only contains two layers of modules with the the inner layer containing smaller modules for the same reason as the barrel region.

Hadronic Calorimeter

The Hadronic Calorimeter is designed to stop all hadronic showers within its volume and consists of two parts, the Tile Calorimeter (HCAL) in the barrel and the LAr Hadronic Endcap (HEC). The HCAL is a tile calorimeter consisting of alternating layers of scintillator and steel as the active medium and absorber respectively. The HEC on the other hand uses the same technology as the ECAL with copper plates filled with LAr as the detecting medium. As the Hadronic Calorimeter sits directly behind the ECAL it is used selecting good electron candidates using hadronic isolation or the amount of leakage in to the HCAL from a electron shower in the ECAL.

2.2.3 Magnet System




Figure 2.7: Geometry of magnet windings and tile calorimeter steel, end-cap toroid cold mass inserted into the cryostat and barrel toroid as installed in the underground cavern [25].

The ATLAS detector has two main magnet systems, the inner solenoid magnet found between the TRT and the ECAL and the outer toroid magnets found interleaved with the Muon Spectrometer.

The solenoid system is a super conducting magnet which is kept at 4.6 K to provide the 2 T magnetic field required by the inner detector to curve high energy particles found at the LHC. As the solenoid is found inside the calorimetry system it is important radiative thickness is minimised to reduce efficiency losses in energy measurements. In order to achieve this it was designed to minimise dead material and shares its cryostat vessel with the ECAL reducing the need for two and therefore contributing only 0.63 radiation lengths.

The outer toroid system provides a magnetic field for the muon spectrometer and consists of a barrel and two endcap systems each with eight coils assembled radially around the beam axis. The coils are all aluminium stabilised Niobium-Titanium (NbTi) superconductors with each coil in the barrel contained in its own cryostat while each of the coils in the endcap systems are contained in one single cryostat. The peak field provided by these toroids are 3.9 T and 4.1 T in the barrel and endcaps respectively.

2.2.4 Muon Spectrometer




Figure 2.8: Cut-away view of the ATLAS muon system [25].

Due to the penetrative nature of muons, all the layers of detector discussed above do not induce the showering of high energy muons. Therefore the outermost detector is another tracking detector specifically for muons. It uses the outer toroid magnet system to bend muon paths and measure muon momentum. The Muon Spectrometer is composed of 4 different technologies; Monitor Drift Tubes (MDT), Cathode Strip Chambers (CSC), Resistive Plate Chambers (RPC) and Thin Gap Chambers (TGC). Both the MDT and the CSC boast precision tracking but both have slow readout times. The RPCs and TGCs have the job of triggering muons and providing additional track measurements. The RPCs are found in the barrel region ($|\eta| < 1.05$) while the TGC trigger in the endcap region ($1.05 < |\eta| < 2.4$). The MDT covers a full range in η ($|\eta| < 2.7$) with complementary measurements from the CSC at $2.0 < |\eta| < 2.7$.

Chapter 3

The Trigger & Data Acquisition

The Trigger system within ATLAS is designed to manage the high rate of events produced by the LHC and bring them down to a total rate that can be written to permanent storage by selecting “interesting” events. The related Data Acquisition (DAQ) system controls the flow of data from detector hardware through the trigger system to permanent storage at CERN and the worldwide tier 1 grid sites.

The Trigger system is made up of three main decision levels; Level 1, Level 2 and Event Filter. Level 1 (L1) is mainly hardware based using limited detector information to locate regions of interest (RoI's) and pass them the Level 2. The Level 2 (L2) system checks the RoIs with full detector granularity and precision and the last stage the Event Filter (EF) uses analysis reconstruction techniques to further select “interesting” events down to the level of 400-500 Hz. Both the L2 and EF triggers compose what is called the High-Level-Trigger (HLT) together with the event building software needed by the EF. Figure 3.1 shows the over all trigger system and how data flows through it.

Figure 3.1: Diagram showing the different stages of the Trigger and how they interact.

Following is a description of each of the sections of the trigger while focusing in on the selection of electron objects that are relevant for this analysis. Following this is a discussion of how the trigger menu is formed so bandwidth can be shared between the differing physics goals as well as how ATLAS handles the continued high luminosity push of the LHC.

3.1 Level-1 Trigger

The Level 1 (L1) trigger searches for RoI's consisting of strong signatures, i.e. high p_T , muons, electron/photons or jets. The L1 trigger also searches for events with a large missing transverse energy (E_T^{miss}) or large total transverse energy (ΣE_T). Due to the decision speed required only some parts of the detector

can be used at L1 (and at a much coarser granularity than is possible). For muon acceptance only the RPC's and TGC's can be used while for electromagnetic clusters and jets as well as large E_T^{miss} and ΣE_T the full calorimetry system can be used. The Inner Detector can not be used in L1 decisions due to the time constraint. With a beam crossing interval of 50 ns latency is required to be less than $2.5\mu\text{s}$ with a target of $2.0\mu\text{s}$. However half of this quota, about $1.0\mu\text{s}$, is accounted for by the cable propagation of signals.

The L1Calo system uses trigger towers with a granularity reduced to roughly 0.1×0.1 in $\Delta\eta \times \Delta\phi$ in most of the detector range from both the electromagnetic and hadronic calorimeters. The ECAL produces almost 3500 of these trigger towers via summation of the analogue signals from a range of trigger cells. This trigger tower data is then sent to the Cluster Processor (CP) to identify electron/photon and tau candidates with E_T above a required threshold and passing isolation requirements, which are labelled as RoI's.




Figure 3.2: Electron/photon and tau trigger algorithms (left) and E_T local-maximum test for a cluster/RoI candidate (right). (The eta-axis runs from left to right, and the phi-axis from bottom to top. The symbol R refers to the candidate 2×2 region being tested.)

Figure 3.2 shows how the electron/photon trigger clustering algorithm works by identifying 2×2 clusters of trigger towers within which two adjacent towers sum to greater than the triggering threshold defined in the trigger menu (seen in section 3.4). Also shown is how three forms of isolation can be applied at this stage: the 12-tower surrounding ring, the 2×2 hadronic core behind the RoI and the 12-tower surrounding ring in the hadronic calorimeter. Only the hadronic core isolation has so far been used in electron/photon triggers within ATLAS. As all possible 2×2 clusters are observed in this way it is possible to have double

counting of RoIs and so the sum of each 2×2 RoI must be greater than each of its eight nearest overlapping neighbours. Figure 3.2 also shows how this local-maxima is tested to avoiding identical sums through use of ‘greater than’ and ‘greater than or equal to’ in differing η and ϕ directions. So if two adjacent 2×2 clusters have the same combined energy sum the one to the top or right is chosen so as not to delay the trigger process. The final L1 trigger decision is made by the Central Trigger Processor (CTP) which takes information from both the CP and jet algorithm as well as the L1 muon trigger. If an accept decision is made then RoI’s are sent to the RoI builder which seeds the L2 trigger system and all L1 sub-systems are read out via Readout Drives (ROD’s)(discussed in section 3.3) to the DAQ system for monitoring of the L1 trigger system.

3.2 Higher Level Trigger

Level-2 Trigger

The Level-2 (L2) trigger is seeded by and only makes decisions based on the RoI’s supplied by the L1 trigger. However it does this with full detector information and so the first stage of this trigger is a RoI builder. The RoI builder requests detector information for all relevant detectors for the observed RoI, including at this level the Inner Detector. In the case of electrons this includes the inner tracking detector, the electromagnetic calorimeter and the hadronic calorimeter. It is at L2 that a distinction between electrons and photons can be made due to existence of an associated track in the ID to the RoI in the ECAL. The RoI builder identifies calorimeter clusters and nearby tracks in order for the L2 trigger to make its decision based on algorithms reconstructing shower shapes, track-cluster matching and E_T thresholds with isolation. The list of these requirements are held within trigger chains each designed to accept specific physics signatures (see section 3.4.1). The general idea is simply to check if RoI’s still exist under closer inspection in order to reduce the rate of events before full event building takes place in the Event Filter.

Event Filter

The Event Filter (EF) doesn’t differ in method much to the L2 trigger it is purely a further test of the signals handed over from L2. At this level a full reconstruction of the event takes place and EF trigger requirements with slightly more stringent thresholds are applied to the event. This is the final decision for whether the event is going to be copied to permanent storage and so the EF reduces the final acceptance rate down to the 400 - 500 Hz required by CERN’s computing systems. The requirements at the EF level are also those used in ATLAS analysis so as to treat MC and data samples the same. These requirements are discussed in section 3.4.1.

3.3 Data Acquisition

The Data Acquisition (DAQ) is the set of systems that control the flow of data from detectors, through the trigger and in to permanent storage. The first stage of this process is the Readout System (ROS) a set of 145 PC's or nodes which manages the collection of all detector sub-system data and L1 trigger output from ATLAS. This system is helped by Readout Drives (ROD's) which interface directly with detector components and Readout Links (ROL's), direct point-to-point readout connecting the ROD's with the ROS's. Table 3.1 shows the number of readouts for each component of the detector and L1 system.

Each ROS PC contains Readout Buffer Module's (ROBIN's), custom PCI-X cards, each containing three Readout Buffers (ROB's), the other end of each ROL. The ROB's is where event data is stored while the L2 trigger makes its decision which comes from the set of 10 L2 Supervisor (L2SV) nodes. This decision is then made by the DataFlow Manager (DFM) on input from all the L2SV nodes and sends a command to the ROS's to either expunge data or forward it on to the event building nodes (or Sub farm Input, SFI). Once a event fully built it is sent forward to the HLT farm which makes the EF decision, then and only then is a message sent back down via the DFM for the ROS's to fully delete all data from the event. The HLT farm is the largest computing resource in the DAQ system with 1116 nodes each containing 8 CPU's. These nodes can either be configured to run as the EF or L2 Processing Units (L2PU's) for the L2SV and are reconfigured as need dictates. As the final step if an event is accepted by the EF all data is passed to the Sub Farm Output (SFO) where it is stored before transfer to CERN's central data-recording facility. In the case that this connection to CERN is offline for some reason ATLAS is able to store about 24 hours worth of data in the SFO's so no data is lost. Table 3.2 shows the number of each DAQ component used within ATLAS all of which are found in the USA15 service cavern next to the ATLAS cavern.

3.4 Trigger Menu and Rates

In its simplest form a single trigger is a energy threshold designed to select a high quantity of particles of of a selected type. ATLAS contains many of these thresholds to select many interesting physics objects which are roughly grouped in to similar signatures called streams. The trigger streams are Egamma triggers to select electrons and photons, JetTauEtMiss triggers to select hadronic decays, tau decays and large missing transverse energy, Muon triggers to select muons, MinBias trigger to check no bias's exist in other triggers and cosmics triggers to selected signals of cosmic radiation. Each stream has a given bandwidth allocated for readout from the trigger so all triggers need to be optimised so total acceptance rates are within requirements. Each trigger at the HLT level is designed to select a specific type of signal while those a L1 are more general and seed many HLT triggers. A full run through all three stages of the trigger is called a trigger chain. Each

Detector Partition			Number of RODs	Number of ROLs	Number of ROSs
Inner detector	Pixel	Layer 0	44	44	4
		Disks	24	24	2
		Layers 12	64	64	6
	SCT	End-cap A	24	24	2
		End-cap C	24	24	2
		Barrel A	22	22	2
		Barrel C	22	22	2
	TRT	End-cap A	64	64	6
		End-cap C	64	64	6
		Barrel A	32	32	3
		Barrel C	32	32	3
	Calorimetry	Tile	Barrel A	8	16
Barrel C			8	16	2
Extended barrel A			8	16	2
Extended barrel C			8	16	2
LAr		EM barrel A	56	224	20
		EM barrel C	56	224	20
		EM end-cap A	35	138	12
		EM end-cap C	35	138	12
		HEC	6	24	2
		FCal	4	14	2
Muon spectrometer	MDT	Barrel A	50	50	4
		Barrel C	50	50	4
		End-cap A	52	52	4
		End-cap C	52	52	4
	CSC	End-cap A	8	8	1
		End-cap C	8	8	1
L1	Calorimeter	CP	4	8	1
		JEP	2	8	1
		PP	8	32	3
	Muon RPC	Barrel A	16	16	2
		Barrel C	16	16	2
	Muon TGC	End-cap A	12	12	1
		End-cap C	12	12	1
	MUCTPI		1	1	1
CTP		1	1	1	
Total			932	1574	145

Table 3.1: Numbers of readout drivers (RODs), readout links (ROLs) and readout systems (ROs) per detector partition [25].

trigger in a trigger chain needs to not only be optimised to fulfil acceptance rate but also optimised to for high acceptance efficiency in the valid region. In terms of energy threshold this means an increasing threshold through the trigger chain so that each level is selecting within the range close to 100% efficient from the previous requirement when taking in to account the different accuracy of energy measurement provided by

Component	Number of nodes	Number of racks	Number of CPUs/node
ROS	145	16	1
DFM	12	1	2
L2SV	10	1	2
SFI	48	3	2
HLT	1116	36	8
SFO	6	2	2
Monitoring	32	4	4
Operations	20	4	2

Table 3.2: The main data-acquisition system components to be deployed for initial operation: the readout system (ROS), the event-building node (SFI), the data flow manager (DFM), the L2 supervisor (L2SV), the high-level trigger (HLT) and the event filter output nodes (SFO) [25].

each level. This section focuses on the Egamma trigger stream as all objects in this analysis were selected using it.

3.4.1 The “ e/γ ” Trigger Menu

The e/γ trigger menu that is used in this analysis refers to trigger chains designed to select electron and photon objects detailing requirements for all three stages of the trigger. ATLAS uses its own terminology to name these triggers with the name giving a description of the requirements used. At L1 electron and photon objects are selected with bearing the name EMXY where; ‘EM’ refers to EM calorimeter, X is the value of the energy threshold required of RoIs in GeV and Y refers to any other specification. Other specifications can be ‘V’ is a threshold varying in detector geometry (η) around the given value to optimise selection or ‘H’ indicating hadronic isolation applied behind the RoI, both of which are discussed in section 3.4.2. An example of a L1 trigger is then L1_EM18VH which is a trigger with an energy threshold of 18 GeV which is varied slightly through the detector and has an hadronic isolation requirement. L2 and EF use the same terminology but are prepended with either L2 or EF. They take the form such as e22vh_medium where ‘e’ represents an electron (g is used for photons), ‘vh’ represents the same as above and ‘medium’ refers to an associated set of shower shape and tracking requirements. As well as ‘medium’, ‘loose’ and ‘tight’ are also defined giving looser and tighter requirements respectively. These shower shape and tracking requirements are discussed in section 4.1. Section 3.4.2 discusses the development of the L1_EM16VH trigger which feeds in to L2_e20vh_medium and then in to EF_e22vh_medium.

The analysis discussed in the rest of this thesis uses a photon trigger even though searching for electrons. This is because photon and electron triggers are identical save for tracking requirements for electrons and it turned out for the 2012 run the lowest energy triggers without hadronic isolation that were applicable for high energy dielectron decays was a diphoton trigger chain. It is important that the trigger used did

not have hadronic isolation due to the very high energy nature of the electrons in this analysis which have a higher chance of leaking through in to the hadronic calorimeter. The trigger used is EF_g35_loose_g25_loose which selects two photon objects with thresholds of 35 GeV and 25 GeV while both requiring ‘loose’ shower shape requirements. This trigger is seeded by L2_g30_loose_g20_loose which itself is seeded by L1_2EM12_EM16V.

3.4.2 Trigger Rates in High Luminosity Regime

Due to the bandwidth limitations of the trigger each level is restricted to a certain output rate. During 2011 the L1 output rate was kept below 60 kHz, L2 below 5 kHz and the EF output rate at around 400 Hz averaged over the LHC fills. The bandwidth allocated to the e/γ triggers was approximately 30% of the total EF output rate however throughout 2011 the luminosity continued to increase putting pressure on the trigger’s ability to control the output rate. Several methods were employed to reduce the trigger rate and in the e/γ trigger a variable threshold and hadronic core isolation was investigated to reduce the rate of the Level-1 trigger. In order to keep within timing constraints only a low resolution of 0.4 η is available at L1. Threshold requirements were therefore investigated varying every 0.4 η . The effect of a hadronic core isolation was also investigated on the selection of electrons which defines a region in the hadronic calorimeter behind the e/γ candidate in which a minimum amount of energy is required to be deposited in order to distinguish between jets and e/γ objects.

A study in to introducing these new requirements [] at L1 was carried out using data from the trigger stream with a “tag & probe” study to calculate the efficiency of an array of new L1 requirements. These attempts where successful with a selection of a threshold that varied slightly with η and hadronic core isolation < 1 GeV. Rates were reduced and deemed low enough for the high luminosity environment for the rest of the year. Figure 3.3 shows the performance of the trigger after these changes had been made. It can be seen that a minimal impact of these new requirements is felt in efficiency in both eta and number of primary vertices and most importantly the “turn on” curve for efficiency as a function of E_T shows point at which 100% efficiency is reached is not that much higher than EM16. This then introduced the new lowest threshold electron trigger chain used in ATLAS. The final trigger being EF_e20vh_medium. For the 2012 run this trigger went through a revision raising the thresholds of each trigger in the chain to accommodate higher luminosity at 8 TeV, this chain can be seen discussed in section 3.4.1.




Figure 3.3: Performance of the first level of the ATLAS e/γ trigger before (EM16) and after (EM16VH) variable thresholds and hadronic core isolation are applied.

Chapter 4

Reconstruction

Reconstruction is the process and algorithms that attempt to reform information about collision events and their decay products from detector signals. This process is done at several points in the ATLAS analysis procedure. First partial reconstruction of RoI's is done at the Level-2 trigger while a mostly full detector reconstruction is done at the EF. After the data has been permanently stored full reconstruction of all possible signatures in each event as well as whole event variables can be completed if it failed to finish live during the trigger decision.

The other main source of reconstruction is done in a process called reprocessing. After data has been stored updates to sub-detector calibrations and optimisations can take place and so reconstruction of entire data sets takes place to update variables to more accurate measurements.

The Data format used in this analysis is an internal ATLAS format called a D3PD. This format is a type of ROOT [] ntuple, or sequence of ordered lists, which stores ATLAS event data. The data used has past through ATLAS software reconstruction while the Monte-Carlo (MC) background estimate samples have gone through GEANT [] detector simulation as well as ATLAS reconstruction. This means analysis of these D3PD's with root requires only minor corrections.

Some of the many variables reconstructed are event specific and one of the more important of this is the Number of interactions per bunch crossing. As more than one proton collision takes place with each bunch collision a lot of physics background to a search can appear in an event. This makes the search harder but it is also needs to be predicted properly by background estimates. The problem is referred to as pile-up and is discussed in section 6.

Bellow will mainly be a discussion of the reconstruction of electron (and related photon) objects as these are the decay products searched for in this analysis.

4.1 Electron Reconstruction

During reconstruction each EM calorimeter energy signature with an associated track in the inner detector get listed as an electron object. These objects get selected via a clustering algorithm in the EM calorimeter which defines the energy and is then matched to a track. These objects then have a list of associated variables derived from detector readouts on which an analysis selection for ‘good’ electrons can be made. These variables range from simple values of position and energy to more complex derived values such as isolation. A few variables will be listed bellow relevant to this analysis and how they are derived.

- η & ϕ , a particle trajectory or position in detector. These are the main variables for measuring the direction the particle went in the detector and for electrons can be measured in two ways. Either via cluster location in the EM calorimeter or measurement in the inner detector.
- p_T or transverse momentum. p_T is the main measure of energy used for particles where $p_T = |p| \cosh \eta$. Here η comes from either the inner detector or calorimeter cluster hit location and the choice is dependent on a how many hits the track made travelling though the inner detector and therefore how accurate the measurement is.
- $E_{T\text{cone}20}$. This is a variable measuring the sum of energy found around the region of interest minus the electron cluster for $R < 0.20$ where $R = \sqrt{\eta^2 + \phi^2}$. $E_{T\text{cone}20}$ is used to check cone isolation in order to eliminate jet like signatures from the analysis which often create large showers in the calorimeters.
- Electron Charge. This is simple matter of measuring the direction the electron curves in the inner detector. However as discussed in chapter 5 this can be hard for very high energy electrons.
- loose, medium, tight. This is a definition given to a set of selections defining how certain it is the object is an electron. The selections and variables associated with this are discussed bellow.

Loose, medium and tight define an increasing series of selections for identifying good electron signatures in the detector. The selections vary between shower shape variables to track quality and track cluster matching. The some of the important associated variables are discussed bellow in table 4.1 with the full selections found in appendix ?? as the full selections are two dimensional arrays of threshold for most selections. Also to note there are two different definitions of medium referred to as medium and medium++. The latter is a re-optimisation of the selection and slightly stricter than the original medium. medium++ is the definition used in this analysis and found in appendix ??.

Loose selection		
Type	Description	Name
Acceptance	$ \eta < 2.47$	η
Hadronic leakage	Ratio of E_T in the first layer of the hadronic calorimeter to E_T of the EM cluster (used over the range $ \eta < 0.8$ and $ \eta > 1.37$)	R_{had1}
	Ratio of E_T in the hadronic calorimeter to E_T of the EM cluster (used over the range $ \eta < 0.8$ and $ \eta > 1.37$)	R_{had}
Middle layer of EM calorimeter	Ratio of the energy in 3×7 cells over the energy in 7×7 cells centred at the electron cluster position	R_η
	Lateral shower width, $\sqrt{(\sum E_i \eta_i^2)/(\sum E_i) - ((\sum E_i \eta_i)/(\sum E_i))^2}$, where E_i is the energy and η_i is the pseudorapidity of cell i and the sum is calculated within a window of 3×5 cells	$\omega_{\eta 2}$
medium & medium++ selection (includes loose)		
Type	Description	Name
Strip layer of EM calorimeter	Shower width, $\sqrt{(\sum E_i (i - i_{max})^2)/(\sum E_i)}$, where i runs over all strips in a window of $\Delta\eta \times \Delta\phi \approx 0.0625 \times 0.2$, corresponding typically to 20 strips in η , and i_{max} is the index of the highest-energy strip	ω_{stot}
	Ratio of the energy difference between the largest and second largest energy deposits in the cluster over the sum of these energies	E_{ratio}
Track quality	Number of hits in the pixel detector (≥ 1)	n_{pixel}
	Number of total hits in the pixel and SCT detectors (≥ 7)	n_{Si}
	Transverse impact parameter ($ d_0 < 5$ mm)	d_0
Track-cluster matching	$\Delta\eta$ between the cluster position in the strip layer and the extrapolated track ($ \Delta\eta < 0.01$)	$\Delta\eta$
Tight selection (includes medium)		
Type	Description	Name
Track-cluster matching	$\Delta\phi$ between the cluster position in the middle layer and the extrapolated track ($ \Delta\phi < 0.02$)	$\Delta\phi$
	Ratio of the cluster energy to the track momentum	E/p
	Tighter $\Delta\eta$ requirement ($ \Delta\eta < 0.005$)	$\Delta\eta$
Track quality TRT	Tighter transverse impact parameter requirement ($ d_0 < 1$ mm)	d_0
	Total number of hits in the TRT	n_{TRT}
	Ratio of the number of high-threshold hits to the total number of hits in the TRT	f_{HT}
Conversions	Number of hits in the b-layer (≤ 1)	n_{BL}
	Veto electron candidates matched to reconstructed photon conversions	

Table 4.1: Table showing the variables associated with definitions of loose, medium and tight. Full thresholds found in appendix ??

Chapter 5

Event Selection

The main event selection for this analysis is based on a standard cut-flow selection used within ATLAS to select high energy di-electron events. Following will be the basic outline of each requirement an event must satisfy followed by a small discussion of optimisations done to some cuts for this analysis. Finally a discussion of corrections is included spanning minor variable corrections to data obtained by performance groups after reconstruction and more substantial corrections to MC samples to correctly estimate run conditions. The following analysis selection is applied equally the data and MC background samples unless were noted.

5.1 Analysis Selection

The following selection is made to all data and MC events for this analysis. Before selection several data flags are checked to insure full operation of the detector at time of data taking.

Event Selection

- Event is required have passed the chosen unscaled electron trigger (EF_g35_loose_g25_loose).
- Each event is required to contain at least one reconstructed primary vertex with at least 2 traceable charged tracks.

Electron Selection

- Electron $|\eta| < 2.47$ and not lie within the detector crack region $1.37 \leq |\eta| \leq 1.52$ due to a decreased energy resolution.
- Each electron is required to have a transverse momentum (p_T) greater than 30 GeV with the highest p_T electron lying above 40 GeV.

- Electrons are required to pass identification criteria on the transverse shower shape, the longitudinal leakage into the hadronic calorimeter, and the association to an inner detector track, defined together as a medium++ electron identification (see section 4.1).

Dielectron Selection

- Selection of two highest p_T electrons left in event.
- Lead Isolation (A cone around the candidate in the calorimeter is required to have $< 0.007 \times E_T + 5.0 \text{ GeV}$ deposited in it) of the highest p_T electron in the event is used to suppress jet's background.
- Sub-leading Isolation (A cone around the candidate in the calorimeter is required to have $< 0.0022 \times E_T + 6.0 \text{ GeV}$ deposited in it) of the highest p_T electron in the event is used to suppress more jet's background.
- Dielectron invariant mass (m_{ee}) is required to be greater than or equal 80 GeV.
- Opposite sign requirement. Require both electrons to have opposite charge.

5.1.1 Isolation Requirement

It was decided a re-investigation of the isolation requirement was needed, updating the cut from its previous iteration in the di-electron analysis on 2011 ATLAS data. The previous threshold was a flat, less than 7 GeV, cut on the p_T corrected (define pT corrected) electro-magnetic calorimeter cluster isolation (definition probably required) of the highest p_T electron in the selected pair. The first investigation was to see how this cut performed in the selection of MC signal at 8 TeV centre of mass energy. Due to the better statistics found in the $DY \rightarrow ee$ MC sample and this being an irreducible background and therefore indistinguishable from the signal this was used in the following investigation.

It can be seen in fig. # that this flat cut of 7 GeV causes an increasing efficiency loss at high energy and was deemed unsuitably high for this iteration of the analysis due to the higher reach in energy expected from the higher centre of mass energy. Two methods that could be used in conjunction were proposed to solve this issue as well as the possibility of introducing a requirement on the second highest energy electron in our selected pair.

The first alternative was a different algorithm of calculating the isolation variable, topo isolation (definition required). This was deemed unsuitable after a short investigation due the similarity of distribution of final result and problems with the implementation of this algorithm in ATLAS reconstruction code. (remember and find out the exact reason)

The second possibility was an isolation requirement varying with energy. Fig. 5.1 shows the distribution of DY MC events in E_T and cluster isolation. It can be seen that electrons become less isolated under this

definition of isolation as energy of the electron increases. This is to be expected as higher energy electrons produce larger showers in the EM calorimeter and so are less well restrained to only a few calorimeter cells in the electron shower.




Figure 5.1: Distribution of DY MC in E_T and cluster isolation for the highest energy electron. Colour density shows the fraction of electrons from that E_T column found in cell. The red line shows the 99% acceptance point of electrons in the E_T column. While the Black vertical and horizontal lines show the p_T and old isolation cut respectively.

In order to define a requirement varying in p_T the 99% acceptance point for each p_T column was looked at and a 1st order polynomial fit to these points by eye was done. The 99% acceptance points can be seen in fig. 5.2 as well as the estimated fit which would form the cut. The same thing was looked at for the second highest p_T electron and can be seen in fig. 5.3.

The two first order polynomials shown here correspond to isolation requirements of;

$$Lead\ Isolation < 0.007 \times E_T + 5.0\ GeV$$

$$Subleading\ Isolation < 0.022 \times E_T + 6.0\ GeV$$

for the highest and second highest energy electrons respectively.

An analysis of the efficiency of these cuts on signal can be seen in fig. 5.4 and fig. 5.5 where it can be seen they maintain a flat behaviour as E_T increases.

The main source of background the isolation cut is imposed to reduce is jets that fake an electron signal in the detector. It is therefore important to measure the effect of this new requirement. Jets background is




Figure 5.2: Similar plot to Fig. 5.1 but with a fit 99% as a possible isolation requirement.

estimated via a reverse ID method on data (see Section #) and so lacks statistics at high energy. For this reason it is impossible to optimise the isolation requirement against rejection of high energy fakes as seen in fig. #. Fig. # shows the efficiency of acceptance of jet fakes against pT and shows a sizeable increase in efficacy over the flat cut (fig. #) used previously.

5.2 Opposite Sign requirement

5.3 Energy Resolution Correction

5.4 Selection Acceptance \times Efficiency




Figure 5.3: Similar plot to Fig. 5.2 but for second highest energy electrons after the requirement proposed in Fig. 5.2 is applied.




Figure 5.4: Efficiency of new leading electron isolation cut on selection of signal MC. Red and orange lines indicate the 99% and 98% efficiency levels respectively.




Figure 5.5: Efficiency of new subleading electron isolation cut on selection of signal MC. Red and orange lines indicate the 99% and 98% efficiency levels respectively.

Chapter 6

Background Estimate

This chapter discusses methods used to estimate background processes the the signal. These background processes can be split up in to two categories; Reducible and irreducible. Irreducible backgrounds consist of those background almost indistinguishable from our signal process namely clean high energy dielectron decays such as decays of the Z boson and the Drell-Yan (DY) spectrum. DY ($q\bar{q} \rightarrow Z/\gamma^* \rightarrow \ell^-\ell^+$) is largest background process and also interferes with the signal processes. Another irreducible background also estimated is the Photon Induced processes ($\gamma\gamma \rightarrow \ell^-\ell^+$) coming from the collision of two photons. Reducible backgrounds are those that can be reduced through event selection and three are included in this analysis. These reducible backgrounds consist of Top precessed, collisions creating single top quarks and $t\bar{t}$ events which decay to include two electrons, Diboson events, the creation of WW, WZ and ZZ events that decay in to two electron events, and finally Multi-jet & W+jets events where one or more electron signature is faked by jet objects. All of these backgrounds are estimated via Monte Carlo generators except for the Multi-jet & W+jets background which is estimated via a data-driven fake factor method. Once completed all background samples are summed together to create the full background estimate. MC samples are then scaled to the integrated luminosity of data collected in 2012 which is 20.3 fb^{-1} . Within the Z boson peak region (80 - 120 GeV) were it is know no new physics is found MC samples are scaled to data minus multijet sample in order to rule out luminosity errors. This scale factor is found to be 1.048. Detailed bellow is the full derision of the background estimates ready for comparison to data.

6.1 Monte Carlo samples

Monte Carlo (MC) samples are produced centrally within ATLAS with each corresponding generator of choice. The generated events then undergo detector simulation using GEANT 4 [?]hich produces a data format identical to a readout from the ATLAS detector plus additional “truth” parameters from the original MC

generation. The samples then undergo the same reconstruction as data events within ATHENA producing a MC sample ready to be analysed the same as data.

Drell-Yan

The Drell-Yan (DY) background is produced using the POWHEG + PYTHIA generator which is a next to leading order (NLO) generation with POWHEG [] with event showering handled by PYTHIA 8 []. The parton density function (PDF) used is CT10 []. A K-factor is then used in order to weigh the cross-section from NLO up to next to next to leading order (NNLO). This NNLO K-factor is derived using FEWZ [] which uses the MSTW2008 NNLO PDF [] from which a QCD+EW mass-dependent K-factor is obtained. The DY sample is split in to 16 MC truth dilepton mass bins. The first bin from 60 - 120 GeV is a very high statistics sample providing a low statistical uncertainty for the region used for scaling MC in the Z boson peak.

Photon-Induced

The Photon-Induced (PI) fraction is estimated via PYTHIA 8 [] generator with the LO PDF MRST2004QED []. This sample is split in to 4 dilepton mass bins covering the region of this analysis.

Diboson

The Diboson MC sample is produced using HERWIG 6.510 [] with the LO PDF CTEQ6L1 []. The sample with split in to the three processed, WW, WZ and ZZ, with each process split in to 4 mass-binned samples. The sample is then scaled to NLO in a mass-independent way seen here [].

Top

The top sample is estimated using MC@NLO 3.41 [] with NLO PDF CT10 [] to generate matrix elements with JIMMY 4.31 [] describing parton interactions and HERWIG [] deriving the underlying event and parton showers. Both $t\bar{t}$ and single Top processes are generated in two inclusive samples. A NNLO QCD K-factor is also derived using Top++ 2.0 []. The top sample also undergoes a fit at high mass where statistics grow too low. A dijet function is used to fit between 200 - 700 GeV and then stitched above 500 GeV.

6.1.1 MC Corrections

Corrections are applied to MC sample due to many factors including unknown run conditions within the ATLAS detector due to MC samples being created earlier than data collection as well as known inefficiencies in the reconstruction of MC events. Bellow are listed all of the corrections which are applied on an event by

even basis during the analysis of MC samples.

Pile-up Correction

Pile-up (PU) or number of simultaneous proton-proton interactions within an event is hard to predict prior to running and it is therefore intentional that MC is produced with a broad range of PU values which then get weighted according to run conditions within the detector. PU conditions can also change throughout data taking and so the PU correction is specified for a particular set of ATLAS data.

Vertex Position Reweighting

Vertex position is again another variable hard to predict pre-run and can therefore be weighted later once run conditions are known. This correction is not widely used within ATLAS due to its minimal effect however it was found to add better data background agreement to the $\cos\theta^*$ distribution within the scaling and control region. Its effects on the analysis are minimal.

Energy Smearing Correction

The energy smearing correction is used to better estimate the energy of electron signatures. It comes about due to the detector simulation not fully estimating the smearing of energy values. This correction is derived from a Z peak calibration study [] done within the ATLAS electron photon performance group and matches MC to data. These corrections provide a η and E dependent smearing value applied to electron energy before electron selection.

Electron efficiency Scale Factor

The electron photon performance group also noticed some inefficiencies in electron reconstruction and identification. These form a set of scale factors applied in bins of E_T and η after event selection.

Isolation and Trigger Scale Factor

A study was done of the data/MC comparison for the isolation selections and the trigger requirements. The differences were found to be below 1% and are applied as a uniform scale factor after event selection.

6.2 Fake Factor Multi-Jet Estimate

One of the major sources of background to di-electron signals are di-jets or electron+jets (mainly W+jets) events where one or both selected leptons are jets faking electron signatures. The method for estimating this background, described here, is a “fake factor” or “matrix-method”. This is a data-driven method where

electrons are selected by a tight (N_{tight}) and loose (N_{loose}) selection. The tight selection is the standard electron selection used in this analysis while the loose selection has no isolation requirement and must only pass a loose++ egamma definition (see Chapter 4) with no track matching criteria. N_{tight} is therefore by design a subset of N_{loose} . Two more hidden values are also assigned *real* and *fake* referring to true source of each electron. This gives us two coefficients to determine from data.

$$f = \frac{N_{tight}^{fake}}{N_{loose}^{fake}} \quad r = \frac{N_{tight}^{real}}{N_{loose}^{real}} \quad (6.1)$$

The fake rate f denotes the probability that a *fake* electron which passes the loose requirement also passes tight while r refers to the probability that a *real* electron which passes the loose requirement also passes the tight. Reconstructed events are split in to two distinct groups, tight(T), and loose while failing tight(L), where *Tight* is now no longer a subset of *Loose*. This allows us to relate our reconstructed events to the underling truth events via a matrix of fake rates shown in Eq. 6.2.

$$\begin{pmatrix} N_{TT} \\ N_{TL} \\ N_{LT} \\ N_{LL} \end{pmatrix} = \begin{pmatrix} r_1 r_2 & r_1 f_2 & f_1 r_2 & f_1 f_2 \\ r_1(1-r_2) & r_1(1-f_2) & f_1(1-r_2) & f_1(1-f_2) \\ (1-r_1)r_2 & (1-r_1)f_2 & (1-f_1)r_2 & (1-f_1)f_2 \\ (1-r_1)(1-r_2) & (1-r_1)(1-f_2) & (1-f_1)(1-r_2) & (1-f_1)(1-f_2) \end{pmatrix} \begin{pmatrix} N_{RR} \\ N_{RF} \\ N_{FR} \\ N_{FF} \end{pmatrix} \quad (6.2)$$

The first index in Eq. 6.2 refers to the highest p_T electron while the second index refers to the second highest p_T electron. So N_{LT} indicates the reconstructed events with highest p_T electron only passing the *Loose* selection while the second highest p_T electron passes *Tight* selection. The indices 1 and 2 refer to fake rates (f) and efficiencies (r) on leading and sub-leading electrons respectively.

The interesting part for this study is the contribution to N_{TT} coming from sources other than N_{RR} , these can be seen in Eq. 6.3.

$$\begin{aligned} N_{TT}^{\ell+jets} &= r_1 f_2 N_{RF} + f_1 r_2 N_{FR} \\ N_{TT}^{di-jets} &= f_1 f_2 N_{FF} \\ N_{TT}^{\ell+jets \& di-jets} &= r_1 f_2 N_{RF} + f_1 r_2 N_{FR} + f_1 f_2 N_{FF} \end{aligned} \quad (6.3)$$

This function however contains hidden variables and so Eq. 6.2 is inverted to derive a better formalism.

$$\begin{pmatrix} N_{RR} \\ N_{RF} \\ N_{FR} \\ N_{FF} \end{pmatrix} = \alpha \begin{pmatrix} (f_1 - 1)(f_2 - 1) & (f_1 - 1)f_2 & f_1(f_2 - 1) & f_1f_2 \\ (f_1 - 1)(1 - r_2) & (1 - f_1)r_2 & f_1(1 - r_2) & -f_1r_2 \\ (r_1 - 1)(1 - f_2) & (1 - r_1)f_2 & r_1(1 - f_2) & -r_1f_2 \\ (1 - r_1)(1 - r_2) & (r_1 - 1)r_2 & r_1(r_2 - 1) & r_1r_2 \end{pmatrix} \begin{pmatrix} N_{TT} \\ N_{TL} \\ N_{LT} \\ N_{LL} \end{pmatrix} \quad (6.4)$$

where,

$$\alpha = \frac{1}{(r_1 - f_1)(r_2 - f_2)} \quad (6.5)$$

The fraction of selected events with at least one fake is then given by Eq. 6.2.

$$\begin{aligned} N_{TT}^{\ell+jets \& \ di-jets} = & \alpha r_1 f_2 [(f_1 - 1)(1 - r_2)N_{TT} + (1 - f_1)r_2 N_{TL} + f_1(1 - r_2)N_{LT} - f_1 r_2 N_{LL}] \\ & + \alpha f_1 r_2 [(r_1 - 1)(1 - f_2)N_{TT} + (1 - r_1)f_2 N_{TL} + r_1(1 - f_2)N_{LT} - r_1 f_2 N_{LL}] \\ & + \alpha f_1 f_2 [(1 - r_1)(1 - r_2)N_{TT} + (r_1 - 1)r_2 N_{TL} + r_1(r_2 - 1)N_{LT} + r_1 r_2 N_{LL}] \end{aligned} \quad (6.6)$$

$$\begin{aligned} = & \alpha [r_1 f_2 (f_1 - 1)(1 - r_2) + f_1 r_2 (r_1 - 1)(1 - f_2) + f_1 f_2 (1 - r_1)(1 - r_2)] N_{TT} \\ & + \alpha f_2 r_2 [r_1(1 - f_1) + f_1(1 - r_1) + f_1(r_1 - 1)] N_{TL} \\ & + \alpha f_1 r_1 [f_2(1 - r_2) + r_2(1 - f_2) + f_2(r_2 - 1)] N_{LT} \\ & - \alpha f_1 f_2 r_1 r_2 N_{LL} \end{aligned} \quad (6.7)$$

Equation 6.7 shows the derived formula relating the multi-jet background to fake rates, efficiencies and four independent samples selected from data. Detailed here is this method used on the full 20 fb^{-1} of integrated luminosity from ATLAS's 2012 run.

6.2.1 Real electron efficiency estimation

The real electron efficiency is defined as Eq. 6.1 $r = N_{tight}^{real}/N_{loose}^{real}$. This is determined from MC using a mass binned Drell-Yan sample. The efficiencies are found for both the leading and sub-leading electrons and binned in 8 p_T and three eta bins of $|\eta| < 1.37$ (barrel), $1.52 < |\eta| < 2.01$ and $2.01 < |\eta| < 2.47$ (endcap). The efficiency is distributed between 90 - 96% as can be seen in Fig. 6.1.

6.2.2 Fake electron rate estimation

The default method selected for analysing the fake rates is a single object method selection on the jet stream data. This gives the main advantage of more statistics and a higher energy reach compared to methods such




Figure 6.1: Real electron efficiencies obtained from Drell-Yan MC and binned in p_T and three coarse η bins covering the barrel and two endcap regions. Efficiencies for leading electrons are shown on the left while those for subleading electron are on the right.

as using tag and probe on the egamma stream data. An array of triggers are used for selecting suitable events based on the single jet trigger EF_jX_a4chad (where $X = 25, 35, 45, 55, 80, 110, 145, 180, 220, 280, 360$). Events are associated to groups with the lowest trigger threshold they pass as each trigger has a different prescale. Objects are selected with the AntiKt4TopoEMJets algorithm and then matched to objects in the egamma stream with a $\Delta R < 0.1$. Objects also have to pass the medium jet-cleaning criteria (define this). Two further steps are taken to suppress real electrons from W decays and real Drell-Yan events. A veto of $E_{Tmiss} > 25 \text{ GeV}$ is introduced to combat the former while events with two medium++ or loose++ electrons with $|m_{tag \& probe} - 91 \text{ GeV}| < 20 \text{ GeV}$ are vetoed to counter the real Drell-Yan.

The fake rate is then defined as Eq. 6.1 $f = N_{tight}^{fake} / N_{loose}^{fake}$ with distributions selected using the standard event selection on the matched egamma objects. Due to the different prescales of each trigger a separate set of fake rates are calculated for each trigger, these are then combined as a weighted average of all fake rates. Fig. 6.2 shows the distribution of fake rates for leading and subleading fakes which are distributed between 3 - 20%.




Figure 6.2: Fake rates obtained from data and binned in p_T and four coarse η bins covering the barrel and three endcap regions. Fake rates for leading electrons are shown on the left while those for subleading electron are on the right.

6.2.3 Properties of Multi-Jet Background

In order to compose the final sample events are organised by the distributions N_{TT} , N_{TL} , N_{LT} or N_{LL} and weights are applied according to each electrons p_T , η with respect to Eq. 6.7 and the corresponding efficiencies and fake rates. Fig. 6.3 shows these distributions before the efficiencies and fake rates are applied to weight to the final background prediction. In addition to these steps an extra fit is then applied at low invariant mass due to contamination due to the Z boson peak. This method is not suited to predicting the Multi-Jet background in the Z boson peak region and so a fit is obtained between 120 GeV and 400 GeV and stitched from 110 GeV and bellow. This gives a good estimate to the integral in this region for use in scaling MC's to luminosity but is not predicted to be good at predicting other variables in this region. At high-mass statistics of the sample decline and so an additional fit is made a high mass with the lower edge of the fit varied between 425 and 600 GeV and the upper edge from 700 to 1200 GeV, with the stitching point at 500 GeV.

6.2.4 Other methods and estimation of Error

Two other methods and variations upon them were used to test the validity of this method as well as estimate the systematic error of this background estimates procedure. These two methods are both tag and probe




Figure 6.3: Distribution of N_{TT} , N_{TL} , N_{LT} and N_{LL} from data with no weightings applied.

measurements on either the jet stream of data, or the egamma stream where the method is more an “inverse” tag and probe with the selection of a tag with high probability of being a jet. Variations are also made on the method by assuming r_1 and $r_2 = 1.0$ in all cases as well as changing the definition of loose but fail tight. These variations simplify the equations slightly but the method remains the same. Fig. 6.4 shows all of these variations compared to the default method used to obtain the estimation. This figure then gives us a good estimate to the systematic uncertainty of the multi-jet estimate which has been chosen to be 20%. (ref zprime support note)




Figure 6.4: Ratio of the final background estimate of all method variations to the default method. The ratio starts at 116 GeV and ends at 1500 GeV. Source: resonant support note, will not be able to use this.

Chapter 7

Signal and Results

This chapter discusses the generators used to estimate the signal models and how this is parametrised for the statistical analysis. Then a look at the results of the data background comparison looking for any sign of new physics signal is shown.

7.1 Signal Monte Carlo

All signal MC is produced in the same way as the background MC and then summed with the other background predictions to arrive at the full signal prediction. Since each sample already contains the DY prediction due to the interference nature of the new physics a method is used to remove this fraction and then use the SM DY background prediction instead is discussed for each signal. Each sample also gets scaled by the same factor as the background MC from the z peak scaling.

Contact Interaction

Contact Interaction samples are generated using PYTHIA 8 [1] with the leading order PDF MSTW2008LO [2]. The CI MC samples also have a K-factor applied that is derived in the same way as the DY K-factor but scaling from LO to NNLO instead (see section 6.1). A selection of Λ values was chosen to cover the reach in new physics for all the parametrisation of the model for the 2012 8 TeV data set. This constitutes signals for $\Lambda = 7, 10, 14, 20$ and 28 TeV. For each of these working points parametrisations of constructive and destructive interference and LL, LR and RR models are all generated. This makes 6 parametrisations with 5 Λ values produced for each one. Each MC sample is composed of three dilepton mass binned samples above 300 GeV in order to maintain statistics. Below 300 GeV negligible new physics is predicted and so the SM DY prediction is used below this point. Because this MC is LO a PYTHIA 8 DY sample is subtracted from this sample so that the background DY can be used instead.

ADD

The ADD samples used are produced using the SHERPA [] generator and NLO PDF CT10 []. Only two formalisms are produced as limits for other formalisms can be converted from the GRW one. The only formalism this is not possible for is HLZ ($n = 2$). For these 2 formalisms 4 values of M_s are generated of $M_s = 4.75, 4.0, 3.75$ and 3.5 TeV. Again 3 dilepton mass bins are used above 300 GeV with the SM DY replacing the distribution below this. Also like the CI samples a specific DY only SHERPA sample is produced which is then subtracted from the samples so the background DY can be used.

7.2 Signal Parametrisation

Each formalism of the CI and ADD model gets parametrised according to the form of their individual cross-sections (Eq's 1.3 and 1.5) and as a function of their parameter of interest (Λ, M_s). The parametrisations are a prediction of number of expected events N_{exp} where parameter of interest (Λ, M_s) set to inf equates to no signal and just the standard model background prediction, these can be seen in equations 7.1 and 7.2.

$$N_{\text{exp}}(\Lambda) = c_0 + \frac{c_1}{\Lambda^2} + \frac{c_2}{\Lambda^4} \quad (7.1)$$

$$N_{\text{exp}}(M_s) = c_0 + \frac{c_1}{M_s^4} + \frac{c_2}{M_s^8} \quad (7.2)$$

Here c_0 refers to the SM background prediction while c_1 and c_2 show the dependence of the scale of new physics on number of expected events. Each formalism gets parametrised in every search bin described at the start of chapter 8 and can be seen in figure 7.1 for CI and 7.2 for ADD.

Figure 7.1: Parametrisations of the CI signal for number of expected events as a function of Λ according to equation 7.1 and for each CI search bin.

Figure 7.2: Parametrisations of the ADD signal for number of expected events as a function of M_s according to equation 7.2 for the ADD search bin.

7.3 Results

Following are full results of the event selection for observed data, predicted background and some example signal models. Figures 7.3 and 7.5 show the distributions of the two main search variables dielectron invari-

ant mass and $\cos \theta^*$ with figure 7.4 showing a combination of these both in the forward backward asymmetry distribution as defined in equation 1.2. Following in figure 7.6 are some control plots namely electron p_T , η and ϕ . More results and control plots can be found in appendix ???. Lastly tables ??, ??, ?? and ?? show the full results of predicted and observed events within each of the search bins discussed in chapter 8.






Figure 7.3: Invariant mass comparison between data and MC with possible signal overlays of CI and ADD.

Figure 7.4: A_{FB} comparison between data and MC with possible signal overlay of CI.Figure 7.5: $\cos \theta^*$ comparison between data and MC in control and signal regions with possible signal overlay of CI.Figure 7.6: Control plots of p_T , η and ϕ distributions of selected electrons.

Process	m_{ee} [GeV]					
	400 - 550			550 - 800		
	All	Forward	Backward	All	Forward	Backward
Drell-Yan	72000 \pm 5000	41500 \pm 2600	31000 \pm 2200	13100 \pm 900	7900 \pm 500	5200 \pm 400
Top	6900 \pm 400	3480 \pm 210	3410 \pm 210	2840 \pm 170	1400 \pm 90	1440 \pm 90
Multijets & W+Jets	1650 \pm 330	900 \pm 180	780 \pm 160	670 \pm 130	330 \pm 70	340 \pm 70
Diboson	1330 \pm 70	710 \pm 40	619 \pm 33	583 \pm 31	331 \pm 19	252 \pm 15
Photon-Induced	1200 \pm 1200	600 \pm 600	600 \pm 600	400 \pm 400	230 \pm 230	220 \pm 220
Total SM	84000 \pm 5000	47200 \pm 2800	36400 \pm 2500	17600 \pm 1200	10200 \pm 600	7400 \pm 500
Data	83824	46910	36914	17525	10107	7418
Drell-Yan	910 \pm 70	580 \pm 40	333 \pm 32	302 \pm 25	193 \pm 13	109 \pm 12
Top	153 \pm 13	87 \pm 8	72 \pm 7	35.2 \pm 2.7	18.2 \pm 1.6	17.5 \pm 1.6
Multijets & W+Jets	88 \pm 18	43 \pm 9	45 \pm 9	27 \pm 6	13.0 \pm 3.0	13.0 \pm 3.1
Diboson	62.2 \pm 3.5	36.0 \pm 2.2	26.2 \pm 1.7	22.3 \pm 1.3	13.8 \pm 0.9	8.5 \pm 0.7
Photon-Induced	40 \pm 40	22 \pm 22	22 \pm 22	17 \pm 17	8 \pm 8	8 \pm 8
Total SM	1260 \pm 100	770 \pm 50	500 \pm 50	404 \pm 35	247 \pm 18	156 \pm 17
Data	1262	754	508	388	251	137
SM+CI(Λ_{LL}^{-14})	1310 \pm 110	810 \pm 60	510 \pm 50	440 \pm 40	276 \pm 22	167 \pm 18
SM+CI(Λ_{LL}^{-20})	1290 \pm 110	780 \pm 60	510 \pm 50	430 \pm 40	271 \pm 22	157 \pm 18
SM+CI(Λ_{LR}^{-14})	1340 \pm 110	790 \pm 60	550 \pm 50	460 \pm 40	266 \pm 22	195 \pm 19
SM+CI(Λ_{LR}^{-20})	1290 \pm 110	780 \pm 60	510 \pm 50	420 \pm 40	249 \pm 21	174 \pm 19
SM+CI(Λ_{RR}^{-14})	1310 \pm 110	810 \pm 60	510 \pm 50	440 \pm 40	276 \pm 22	167 \pm 18
SM+CI(Λ_{RR}^{-20})	1290 \pm 110	780 \pm 60	510 \pm 50	430 \pm 40	271 \pm 22	157 \pm 18
SM+CI(Λ_{LL}^{+14})	1230 \pm 110	730 \pm 60	510 \pm 50	380 \pm 40	227 \pm 21	155 \pm 18
SM+CI(Λ_{LL}^{+20})	1230 \pm 110	740 \pm 60	490 \pm 50	390 \pm 40	234 \pm 21	156 \pm 18
SM+CI(Λ_{LR}^{+14})	1200 \pm 110	740 \pm 60	470 \pm 50	400 \pm 40	247 \pm 21	154 \pm 18
SM+CI(Λ_{LR}^{+20})	1210 \pm 110	740 \pm 60	470 \pm 50	390 \pm 40	238 \pm 21	150 \pm 18
SM+CI(Λ_{RR}^{+14})	1230 \pm 110	730 \pm 60	510 \pm 50	380 \pm 40	227 \pm 21	155 \pm 18
SM+CI(Λ_{RR}^{+20})	1230 \pm 110	740 \pm 60	490 \pm 50	390 \pm 40	234 \pm 21	156 \pm 18

Table 7.1: Table comparing background prediction to data with prediction of several CI signal models. Bining used is the same as used for statistical analysis of CI sample.

Process	m_{ee} [GeV]					
	800 - 1200			1200 - 1800		
	All	Forward	Backward	All	Forward	Backward
Drell-Yan	63 ± 6	41.4 ± 3.4	22.1 ± 2.9	8.2 ± 1.2	5.3 ± 0.6	2.9 ± 0.6
Top	3.06 ± 0.18	1.58 ± 0.10	1.45 ± 0.09	0.140 ± 0.008	0.073 ± 0.004	0.065 ± 0.004
Multijets & W+Jets	5.8 ± 1.5	2.6 ± 0.9	2.5 ± 0.8	0.87 ± 0.32	0.35 ± 0.16	0.32 ± 0.24
Diboson	5.4 ± 0.4	3.41 ± 0.28	2.02 ± 0.17	0.83 ± 0.05	0.542 ± 0.035	0.287 ± 0.016
Photon-Induced	4 ± 4	2.2 ± 2.2	2.1 ± 2.1	0.7 ± 0.7	0.34 ± 0.34	0.4 ± 0.4
Total SM	82 ± 9	51 ± 5	30 ± 4	10.8 ± 1.6	6.6 ± 0.7	4.0 ± 0.8
Data	84	53	31	7	5	2
SM+CI(Λ_{LL}^{-14})	108 ± 10	68 ± 6	39 ± 5	20.9 ± 1.9	13.5 ± 1.0	7.2 ± 0.9
SM+CI(Λ_{LL}^{-20})	90 ± 10	58 ± 5	32 ± 4	14.4 ± 1.7	9.2 ± 0.9	5.0 ± 0.8
SM+CI(Λ_{LR}^{-14})	118 ± 10	62 ± 6	56 ± 5	26.3 ± 2.1	11.3 ± 1.0	14.8 ± 1.1
SM+CI(Λ_{LR}^{-20})	98 ± 10	57 ± 5	41 ± 5	15.7 ± 1.7	8.3 ± 0.9	7.2 ± 0.9
SM+CI(Λ_{RR}^{-14})	108 ± 10	68 ± 6	40 ± 5	20.8 ± 1.9	13.6 ± 1.0	6.9 ± 0.9
SM+CI(Λ_{RR}^{-20})	91 ± 10	58 ± 5	32 ± 4	14.3 ± 1.7	9.1 ± 0.9	5.0 ± 0.8
SM+CI(Λ_{LL}^{+14})	79 ± 9	47 ± 5	32 ± 4	12.2 ± 1.7	7.3 ± 0.8	4.7 ± 0.8
SM+CI(Λ_{LL}^{+20})	77 ± 9	48 ± 5	29 ± 4	10.0 ± 1.6	6.1 ± 0.8	3.7 ± 0.8
SM+CI(Λ_{LR}^{+14})	88 ± 10	55 ± 5	32 ± 4	18.9 ± 1.8	9.2 ± 0.9	9.5 ± 0.9
SM+CI(Λ_{LR}^{+20})	81 ± 9	52 ± 5	29 ± 4	11.5 ± 1.6	6.8 ± 0.8	4.5 ± 0.8
SM+CI(Λ_{RR}^{+14})	79 ± 9	47 ± 5	32 ± 4	12.1 ± 1.7	7.3 ± 0.8	4.6 ± 0.8
SM+CI(Λ_{RR}^{+20})	77 ± 9	48 ± 5	29 ± 4	10.2 ± 1.6	6.3 ± 0.8	3.8 ± 0.8

Table 7.2: Table comparing background prediction to data with prediction of several CI signal models. Bining used is the same as used for statistical analysis of CI sample.

Process	m_{ee} [GeV]					
	1800 - 3000			3000 - 4500		
	All	Forward	Backward	All	Forward	Backward
Drell-Yan	0.64 ± 0.17	0.41 ± 0.09	0.23 ± 0.08	0.006 ± 0.004	0.0039 ± 0.0021	0.0022 ± 0.0018
Top	< 0.004	< 0.002	< 0.002	< 0.001	< 0.001	< 0.001
Multijets & W+Jets	0.11 ± 0.04	0.040 ± 0.020	0.033 ± 0.027	0.0058 ± 0.0012	< 0.002	< 0.001
Diboson	0.075 ± 0.006	0.053 ± 0.004	0.0224 ± 0.0026	< 0.001	< 0.001	< 0.001
Photon-Induced	0.08 ± 0.08	0.04 ± 0.04	0.04 ± 0.04	0.0016 ± 0.0016	< 0.002	< 0.002
Total SM	0.91 ± 0.21	0.55 ± 0.10	0.33 ± 0.10	0.014 ± 0.005	0.0065 ± 0.0026	0.0042 ± 0.0022
Data	0	0	0	0	0	0
SM+CI(Λ_{LL}^{-14})	4.2 ± 0.4	2.75 ± 0.23	1.38 ± 0.15	0.141 ± 0.028	0.080 ± 0.020	0.058 ± 0.016
SM+CI(Λ_{LL}^{-20})	2.01 ± 0.25	1.26 ± 0.14	0.72 ± 0.12	0.045 ± 0.012	0.021 ± 0.007	0.022 ± 0.007
SM+CI(Λ_{LR}^{-14})	6.0 ± 0.5	2.31 ± 0.21	3.69 ± 0.30	0.28 ± 0.05	0.127 ± 0.030	0.146 ± 0.032
SM+CI(Λ_{LR}^{-20})	2.58 ± 0.28	1.01 ± 0.13	1.54 ± 0.16	0.078 ± 0.018	0.036 ± 0.011	0.039 ± 0.012
SM+CI(Λ_{RR}^{-14})	3.78 ± 0.34	2.51 ± 0.22	1.23 ± 0.15	0.23 ± 0.04	0.155 ± 0.031	0.069 ± 0.018
SM+CI(Λ_{RR}^{-20})	1.86 ± 0.24	1.11 ± 0.13	0.71 ± 0.12	0.072 ± 0.015	0.047 ± 0.011	0.022 ± 0.008
SM+CI(Λ_{LL}^{+14})	2.08 ± 0.25	1.30 ± 0.14	0.75 ± 0.12	0.075 ± 0.015	0.050 ± 0.012	0.023 ± 0.007
SM+CI(Λ_{LL}^{+20})	0.95 ± 0.22	0.55 ± 0.11	0.36 ± 0.11	0.029 ± 0.008	0.019 ± 0.006	0.0073 ± 0.0034
SM+CI(Λ_{LR}^{+14})	4.2 ± 0.4	1.60 ± 0.16	2.51 ± 0.22	0.191 ± 0.034	0.081 ± 0.020	0.107 ± 0.023
SM+CI(Λ_{LR}^{+20})	1.65 ± 0.24	0.82 ± 0.12	0.79 ± 0.12	0.058 ± 0.013	0.017 ± 0.006	0.039 ± 0.010
SM+CI(Λ_{RR}^{+14})	2.26 ± 0.26	1.44 ± 0.15	0.78 ± 0.12	0.098 ± 0.018	0.057 ± 0.012	0.038 ± 0.010
SM+CI(Λ_{RR}^{+20})	1.06 ± 0.22	0.65 ± 0.11	0.37 ± 0.11	0.036 ± 0.009	0.028 ± 0.008	0.0044 ± 0.0029

Table 7.3: Table comparing background prediction to data with prediction of several CI signal models. Bining used is the same as used for statistical analysis of CI sample.

Table 7.4: Table comparing background prediction to data with prediction of several ADD signal models. One bin used the same in the statistical analysis of ADD

Chapter 8

Statistical Analysis

The statistical analysis of results is done via a Bayesian approach where first a search for signs of new physics is done with a calculation of the significance of any excesses. Then in the absence of a signal exclusion limits on the scale of new physics (either Λ or M_s) are then set. A slightly different search approach is made between CI and ADD. In CI the shape of new physics is informative and therefore a series of invariant mass search bins are used with bin edges of 400, 550, 800, 1200, 1800, 3000 and 45000 GeV. With the addition of information from the $\cos\theta^*$ variable in this an analysis bins are also then split up in $\cos\theta^*$ as well as invariant mass. It was found (see Appendix ??) that most of the new information was obtained via using two bins in $\cos\theta^*$ making a total of 12 search bins distributed in invariant mass and $\cos\theta^*$. ADD on the other hand doesn't gain from the many search bin approach due to a sharper turn-on and undefined nature of the signal after the cut off point. Therefore only one search bin is used to search for ADD with a minimum invariant mass cut of 1900 GeV and upper cut of 4500 GeV. The ADD model gains no additional discriminating power from the $\cos\theta^*$ variable. For each search bin a parametrisation of new physics is produced (discussed in section 7.2) as well as the background predicted and data observed. Also included is the background and signal parametrisation varied by each of the appropriate sources of systematic error (discussed in section 8.1) for signal and background. This all composes the input of the event selection in the statistical analysis.

The statistical analysis is carried out using the ROOT package BAT or Bayesian Analysis Tool-kit [], this package allows for the integration over nuisance parameters (discussed below) using the Markov Chain Monte Carlo method. The statistical analysis starts off with the definition of the number of expected events μ found in the signal region as seen in Eq. 8.1.

$$\mu = n_s(\Theta, \overline{\Omega}) + n_b(\overline{\Omega}) \quad (8.1)$$

Here n_s is the number of signal events predicted by the model with a particular model parameter Θ and n_b is the total number of predicted background events. $\bar{\Omega}$ is then a set of Gaussian nuisance parameters or systematic uncertainties on the number of expected events for signal and background. A product of Poisson probabilities for each search bin k gives the Bayesian likelihood, seen in Eq. 8.2, of observing n events given the signal parameter Θ and nuisance parameters $\bar{\Omega}$.

$$\mathcal{L}(n | \Theta, \bar{\Omega}) = \prod_{k=1}^N \frac{\mu_k^{n_k} e^{-\mu_k}}{n_k!} \quad (8.2)$$

where $\mu_k^{n_k}$ and n_k are the total number of expected events and observed number of events in search bin k respectively.

$$\mathcal{P}(\Theta | n) = \frac{1}{\mathcal{Z}} \mathcal{L}_{\mathcal{M}}(n | \Theta) P(\Theta) \quad (8.3)$$

Equation 8.3 then shows the posterior probability using Bayes' theorem for the observation of Θ given n . Here \mathcal{Z} is the normalisation factor and $\mathcal{L}_{\mathcal{M}}$ is the marginalised likelihood after all nuisance parameters have been integrated out. It is assumed all nuisance parameters are correlated across all search bins. All full description of nuisance parameters can be found in section 8.1. Finally $P(\Theta)$ is the prior probability of Θ . For CI $P(\Theta)$ is chosen to be uniform and positive with respect to $1/\Lambda^2$ or $1/\Lambda^4$. This form of priors is chosen due to the form of differential cross-section (equation 1.3) and its dependence on Λ . It is not obvious which prior is more correct as these forms refer to the interference and pure CI terms respectively which change in dominance throughout the search bins and model parameters, therefore the search is done using both parameters for completeness. A similar effect is seen in the form of the ADD differential cross-section (equation 1.5) and so there a prior of $1/M_s^4$ or $1/M_s^8$ are used for the same reason. In order to check all signal formalisms 1000 background-like Pseudo-Experiments (PE) are run using BAT for each formalism. Each PE is then passed through the Bayesian statistical method above so they can be compared to data and signal predictions. Figures 8.1 and 8.2 show posterior probability density functions (pdf's), for the CI search and the ADD search respectively, of these 1000 PE's.

8.1 Systematics

The list of nuisance parameters used for this statistical analysis make up a list of all systematic errors thought of as relevant for this analysis. Table 8.1 lists all the systematic errors used for this analysis along with their size while figures 7.3, 7.4 and 7.5 in the previous chapter show total background systematic errors in their ratio's. Following is a brief description of each of the systematics including how it was derived.




Figure 8.1: Posterior pdf distributions for the CI model formalisms LL and LR with constructive interference and a uniform positive prior in $1/\Lambda^2$.

Figure 8.2: Posterior pdf distributions for the ADD model with a uniform positive prior in $1/M_s^8$.

8.2 Signal Search & P-Values

Consistency between data and background prediction is estimated by taking the likelihood of signal given n observed events (observed) and comparing this to the likelihood of signal given the outcome of a set of 1000 PE (expected given no signal) calculated above. A likelihood ratio is then calculated between the signal prediction and background only hypothesis where the signal predictions likelihood is maximised to the highest likelihood in Θ . This is done for both the observed likelihood and set of 1000 PE likelihoods for the expected result given no signal. These are converted to the distribution of negative Log Likelihood Ratio's (LLR) given in figures 8.3 and 8.4 comparing observed values to the expected values in the distribution of PE's. p -value's are also derived for each formalism quantifying the probability of observing a fluctuation in PE's at least as signal-like as is observed in data. A table of p -values for each formalism for CI and ADD can be found in Table 8.2 and 8.3 respectively.

8.3 Setting Limits

With no sign of new physics found limits are set on the lower limit for the scale of new physics for each CI and ADD formalism. Limits in Θ are extracted from each of the 1000 PE's for each formalism and the mean of this distribution is taken as the expected limit and converted in to a limit on Λ for CI and M_s for ADD.

Source	Signal	
	Forward	Backward
Normalization	4.0% (4.0%) [4.0%]	4.0% (4.0%) [4.0%]
PDF Variation	< 0.1% (0.2%) [0.5%]	< 0.1% (0.2%) [0.5%]
PDF Choice	NA	NA
α_s	NA	NA
EW Corrections	< 0.1% (< 0.1%) [0.1%]	< 0.1% (< 0.1%) [0.1%]
Photon-Induced	NA	NA
Efficiency	1.0% (2.0%) [3.0%]	1.0% (2.0%) [3.0%]
Scale/Resolution	1.2% (2.4%) [5.0%]	1.2% (2.4%) [5.0%]
Multi-jet/W+jets	NA	NA
Beam Energy	1.0% (3.0%) [5.0%]	1.0% (3.0%) [5.0%]
Charge MisID	1.2% (2.0%) [2.9%]	1.2% (2.0%) [2.9%]
Statistical	3.0% (3.0%) [3.0%]	3.0% (3.0%) [3.0%]
Total	5.5 (6.9%) [9.6%]	5.5% (6.9%) [9.6%]
	Background	
	Forward	Backward
Normalization	4.0% (4.0%) [4.0%]	4.0% (4.0%) [4.0%]
PDF Variation	6.0% (12.5%) [35.0%]	10.0% (28.0%) [62.5%]
PDF Choice	1.0% (7.0%) [22.0%]	1.0% (7.0%) [22.0%]
α_s	1.0% (3.0%) [5.0%]	1.0% (3.0%) [5.0%]
EW Corrections	1.0% (2.0%) [4.0%]	1.0% (2.0%) [4.0%]
Photon-Induced	6.0% (10.0%) [17.0%]	9.5% (16.5%) [29.0%]
Efficiency	1.0% (2.0%) [3.0%]	1.0% (2.0%) [3.0%]
Scale/Resolution	1.2% (2.4%) [5.0%]	1.2% (2.4%) [5.0%]
Multi-jet/W+jets	3.0% (5.0%) [21.0%]	3.0% (5.0%) [21.0%]
Beam Energy	1.0% (3.0%) [5.0%]	1.0% (3.0%) [5.0%]
Charge MisID	1.2% (2.0%) [2.9%]	1.2% (2.0%) [2.9%]
Statistical	0.5% (0.5%) [0.5%]	0.5% (0.5%) [0.5%]
Total	10.3% (19.6%) [50.6%]	14.9% (34.4%) [76.1%]

Table 8.1: Table listing all sources of systematic error and their approximate size for dielectron mass of 1 TeV (2 TeV) [3 TeV].

Figures 8.5 and 8.6 show the distribution of these PE's in Θ for both CI and ADD along with the mean value of the distribution taken as the expected limit. Tables 8.4 and 8.5 then show these expected limits converted into Λ and M_γ respectively. The final observed limits are also included in these figures and tables compared to the expected limits and extracted from the observed data.

8.4 Combination with the Muon Search

A similar analysis was carried out at the same time as this one looking at the dimuon decay channel instead. This analysis followed the same procedure and after failing to find a signal limits were set on the scale of new physics. Assuming lepton universality integrated luminosity can effectively be doubled by combining





Figure 8.3: Distribution of negative Log Likelihood Ratio's for the CI formalisms LL and LR with constructive interference given a uniform positive prior in $1/\Lambda^2$.

Figure 8.4: Distribution of negative Log Likelihood Ratio's for the ADD formalism GRW given a uniform positive prior in $1/M_s^8$.

the results from both channels. Therefore the posterior pdf's from each analysis were combined in BAT and new limits set on the scale of new physics. Care was taken to treat correctly sources of systematic uncertainty that are correlated between analyses. Combined limits of this form can be found in tables 8.6 and 8.7 for the CI and ADD models respectively. These limits mark the highest limits set for this analysis owing to the higher effective luminosity.

p -value [%]	$1/\Lambda^2$		$1/\Lambda^4$	
	Constructive	Destructive	Constructive	Destructive
LL: ee	58	60	> 76	> 58
LR: ee	> 35	36	> 85	> 62
RR: ee	> 35	68	> 75	> 62

Table 8.2: Table of p -values for all CI formalisms and prior's.Table 8.3: Table of p -values for ADD with each prior.Figure 8.5: Distribution of PE's with associated Limits for CI formalisms LL and LR with constructive interference given a uniform positive prior in $1/\Lambda^2$.Figure 8.6: Distribution of PE's with associated Limits for the ADD formalism GRW given a uniform positive prior in $1/M_s^8$.

Limits [TeV]	$1/\Lambda^2$		$1/\Lambda^4$	
	Constructive	Destructive	Constructive	Destructive
Expected LL: ee	19.11	14.02	17.44	13.02
Observed LL: ee	20.71	16.35	18.58	14.72
Expected LR: ee	22.01	17.37	20.09	16.26
Observed LR: ee	25.16	19.19	22.19	17.68
Expected RR: ee	18.97	14.23	17.23	13.14
Observed RR: ee	20.22	16.57	18.34	14.89

Table 8.4: Table of expected and observed Limits for all CI formalisms and prior's.

Table 8.5: Table of expected and observed Limits for ADD with each prior.

Limits [TeV]	$1/\Lambda^2$		$1/\Lambda^4$	
	Constructive	Destructive	Constructive	Destructive
Expected LL: $\ell\ell$	21.44	19.11	14.73	13.81
Observed LL: $\ell\ell$	21.55	19.61	17.15	15.35
Expected LR: $\ell\ell$	24.78	23.12	18.46	17.57
Observed LR: $\ell\ell$	26.25	23.77	18.95	17.79
Expected RR: $\ell\ell$	20.98	19.11	14.99	14.21
Observed RR: $\ell\ell$	21.11	19.31	17.50	15.58

Table 8.6: Combined expected and observed Limits for the 2D LL, LR, and RR Contact Interaction search using a uniform positive prior.

Table 8.7: Combined expected and observed Limits for the ADD search using a uniform positive prior.

Chapter 9

Conclusion

9.1 Interpretation

9.2 Looking Forward

Appendices

Appendix A

Non-Resonance Analysis 2011

A.1 Data and Background Processes

A.1.1 Data

All data used in the CI analysis is taken from the LHC 2011 $\sqrt{s} = 7\text{TeV}$ proton-proton collision data of which ATLAS recorded 4.9fb^{-1} of electron candidate data. Data was collected with stable LHC beams and a fully operational inner detector and calorimeter each being important in the identification of good electron candidates.

A.1.2 Background

The main background the CI signal in the electron channel is Drell-Yan (DY) $\rightarrow ee$ production mediated by a photon or the Z boson. However there are other small contributions from $t\bar{t}$, diboson, W + jets and QCD production. $t\bar{t}$ background consists of events with $t\bar{t}$ production decaying to amongst other things two electrons. Diboson background can involve either an event producing two W bosons, two Z boson or one W and one Z boson in which two electrons also exist in the final decay state. Both these productions can result in two electrons with a large combined invariant mass which could mimic CI production. W + jets production can result in an electron and a jet faking an electron left in the final state, while QCD refers to events where two jets fake electrons. These all combine to form the background sample.

A.1.3 Background Production

Contributions from SM processes were primarily simulated via leading-order (LO) *PYTHIA* [?] Monte Carlo (MC) event generation and using *Geant4* [?] to simulate the ATLAS detector. This method was used to generate a $Z \rightarrow ee$ sample for the low dielectron invariant mass region ($m_{ee} < 120\text{GeV}$) and a

$DY \rightarrow ee$ mass binned sample for the invariant high mass ($m_{ee} > 120\text{GeV}$) to keep high statistics at high invariant mass. Four other samples are included to produce the background estimate, these are $t\bar{t}$ (produced with *MC@NLO*), diboson(WW, WZ and ZZ decays produced with *HERWIG*), W + jets (produced with *ALPGEN*, *JIMMY* and *HERWIG*) and QCD (produced using a data driven method [?]).

A.1.4 Signal

Five benchmarks for the value of Λ where chosen for the CI signal samples for both constructive and destructive interference. Like the DY these were also produced with LO *PYTHIA* containing both the pure DY contribution as well as the interference and pure CI components. Samples where produced above dilepton invariant mass of 120 GeV to increase statistics above the Z^0 peak where new physics would appear.

m_{ee} [GeV]	120-300	300-600	≥ 600
$\Lambda^- = 3 \text{ TeV}$	9.8583	0.96765	0.77563
$\Lambda^- = 4 \text{ TeV}$	9.3400	0.50510	0.26647
$\Lambda^- = 5 \text{ TeV}$	9.0960	0.36935	0.12733
$\Lambda^- = 7 \text{ TeV}$	8.9686	0.28401	0.048406
$\Lambda^- = 12 \text{ TeV}$	8.9037	0.24774	0.021028
$\Lambda^+ = 3 \text{ TeV}$	8.9018	0.65247	0.66034
$\Lambda^+ = 4 \text{ TeV}$	8.8756	0.33164	0.20668
$\Lambda^+ = 5 \text{ TeV}$	8.7362	0.25646	0.085934
$\Lambda^+ = 7 \text{ TeV}$	8.8101	0.22656	0.028772
$\Lambda^+ = 12 \text{ TeV}$	8.9045	0.22774	0.014418

Table A.1: Table of CI sample cross sections [pb^{-1}].

Corrections are applied to the MC samples. A correction handling the number of proton-proton collisions seen in each event is added to all MC samples due to unknown ATLAS run conditions for the full data set. QCD and Electroweak K-factor corrections are applied as a function of invariant mass to the signal samples and SM DY samples.

A.2 Electron Identification and Selection

The selection of electron candidates for the CI analysis can be split in three main parts, selection of a good event, selection of a set of good electrons and selection of a good dielectron pair.

Event Selection

- Each event is required to contain at least one reconstructed primary vertex with more than 2 charged

tracks traceable to it.

- Event is required have passed the chosen unscaled electron trigger (EF_g20_loose).

Electron Selection

- Each electron is required to have a transverse momentum (p_T) greater than 25 GeV.
- Electron $|\eta| < 2.47$ and not lie within the detector crack region $1.37 \leq |\eta| \leq 1.52$ due to a decreased energy resolution.
- Electrons are required to pass identification criteria on the transverse shower shape, the longitudinal leakage into the hadronic calorimeter, and the association to an inner detector track, defined together as a medium electron identification.
- If expected electron is required to have signal in the inner most level of the tracking detector (B-layer).
Used to suppress background from photon conversions.

Dielectron Selection

- Selection of two highest p_T electrons left in event.
- Isolation (A cone around the candidate in the calorimeter is required to have $< 7\text{GeV}$ deposited in it) of the highest p_T electron in the event is required to suppress QCD jet background.
- Dielectron invariant mass (m_{ee}) is required to be greater than or equal 70 GeV.

These remaining candidates are then the results. The signal region is defined as $m_{ee} > 150\text{GeV}$ while the $70 \leq m_{ee} \leq 110\text{GeV}$ region is the control region.

m_{ee} [GeV]	70-110	110-200	200-400
DY	1231053.7 ± 1109.5	26756.7 ± 163.6	2964.0 ± 54.4
$t\bar{t}$	879.6 ± 29.7	1008.8 ± 31.8	315.8 ± 17.8
Dibosons	1827.1 ± 42.7	415.4 ± 20.4	146.6 ± 12.1
QCD + W+jets	2885.7 ± 53.7	1892.0 ± 43.5	510.5 ± 22.6
Total	1236646.0 ± 1112.0	30072.9 ± 173.4	3936.9 ± 62.7
Data	1236646	29816	4026

	400-800	800-1200	1200-3000
	266.0 ± 16.3	12.2 ± 3.5	1.5 ± 1.2
	20.5 ± 4.5	0.3 ± 0.6	0.0 ± 0.2
	16.5 ± 4.1	0.9 ± 0.9	0.1 ± 0.3
	49.5 ± 7.0	2.0 ± 1.4	0.3 ± 0.5
	352.4 ± 18.8	15.4 ± 3.9	1.9 ± 1.4
	358	17	3

Table A.2: Table of data yeild compared to background MC scaled to luminosity of data. Errors shown are statistical only.

A.2.1 Data and Background Comparison

Table A.2 shows the number of data events remaining after selection of dielectron candidates compared to all sources of MC background after the same candidate selection and scaled to the data luminosity. The simulated MC samples also undergo a scaling factor to scale within the Z boson peak in the control region. As can be seen the background prediction matches very closely to data within the statistical errors shown.

The QCD background shown is not predicted using MC simulation but instead by via a reverse identification criteria on data [?]. This reducible background is due to QCD multijet production where jets are mis-measured as electron candidates.

Control plots were produced to display that the distributions were behaving as predicted such as the p_T (Fig. A.1) and the η (Fig. A.2) distributions.

A.2.2 New Physics Signal Expectation

Tables A.3 and A.4 show the yeild from the CI and ADD MC signals used after scaling to data luminosity. The ADD yield is only shown in a single bin above 1300 GeV as the ADD statistical analysis uses only a one bin approach to set a limit of a general increase over SM background. Table A.5 shows the same one bin approach to the data MC comparison table.

Figures A.3 (A.5) show the dielectron invariant mass distribution comparing data to background MC




Figure A.1: p_T distribution of the leading (left) and subleading (right) electrons showing data, MC background and example CI signal samples compared to data.




Figure A.2: η distribution of the leading (left) and subleading (right) electrons showing data, MC background compared to data.

m_{ee} [GeV]	110-200	200-400	400-800	800-1200	1200-3000
$\Lambda^- = 3$ TeV	18790.8 ± 137.1	5022.4 ± 70.9	2766.3 ± 52.6	1089.2 ± 33.0	673.3 ± 25.9
$\Lambda^- = 4$ TeV	18212.5 ± 135.0	3707.1 ± 60.9	1102.5 ± 33.2	356.9 ± 18.9	214.3 ± 14.6
$\Lambda^- = 5$ TeV	17821.5 ± 133.5	3310.5 ± 57.5	653.1 ± 25.6	160.6 ± 12.7	97.7 ± 9.9
$\Lambda^- = 7$ TeV	17711.1 ± 133.1	3018.8 ± 54.9	385.0 ± 19.6	56.1 ± 7.5	26.5 ± 5.1
$\Lambda^- = 12$ TeV	17693.4 ± 133.0	2992.7 ± 54.7	296.5 ± 17.2	20.4 ± 4.5	5.6 ± 2.4
$\Lambda^+ = 3$ TeV	18106.6 ± 134.6	4063.8 ± 63.7	2103.3 ± 45.9	918.1 ± 30.3	621.4 ± 24.9
$\Lambda^+ = 4$ TeV	17958.1 ± 134.0	3178.6 ± 56.4	765.6 ± 27.7	288.0 ± 17.0	194.9 ± 14.0
$\Lambda^+ = 5$ TeV	18026.6 ± 134.3	2895.6 ± 53.8	432.1 ± 20.8	111.4 ± 10.6	78.8 ± 8.9
$\Lambda^+ = 7$ TeV	17926.4 ± 133.9	2857.5 ± 53.5	278.2 ± 16.7	34.3 ± 5.9	19.1 ± 4.4

Table A.3: Table of CI signal yields for 4.9 fb^{-1} .

while showing the effect CI (ADD) would have on this spectrum. Figures A.4 (A.6) then show the same spectrum but with an integrated invariant mass distribution instead which indicates better general increases in the dielectron spectrum.




Figure A.3: Dielectron invariant mass distribution for data and Monte Carlo simulation. Lines show expected distributions for the presence of Contact Interactions.




Figure A.4: Dielectron integrated invariant mass distribution for data and total background Monte Carlo simulation. Lines show expected distributions for the presence of Contact Interactions.




Figure A.5: Dielectron invariant mass distribution for data and Monte Carlo simulation. Lines show expected distributions for the presence of ADD.




Figure A.6: Dielectron integrated invariant mass distribution for data and total backtracking Monte Carlo simulation. Lines show expected distributions for the presence of ADD.

m_{ee} [GeV]	≥ 1300
$M_S = 1500$ GeV (GRW)	94.8 ± 9.7
$M_S = 2000$ GeV (GRW)	42.7 ± 6.5
$M_S = 2500$ GeV (GRW)	11.3 ± 3.4
$M_S = 3000$ GeV (GRW)	3.2 ± 1.8

Table A.4: Table of ADD analysis region yields for 4.9 fb^{-1} .

m_{ee} [GeV]	≥ 1300
DY	1.1 ± 1.1
$t\bar{t}$	0.0 ± 0.1
Dibosons	0.1 ± 0.3
QCD + W+jets	0.2 ± 0.4
Total	1.4 ± 1.2
Data	2.0

Table A.5: Table of data and MC yields for ADD analysis region.

A.3 Statistical Analysis

If no evidence for new physics is found then a Bayesian counting method can be used to set a limit on the scale (Λ) of new physics. A comparison between the observed events yields and expected yield for a range of different CI benchmarks is done using

$$\mu = n_{DY+CI}(\theta, \bar{\mathbf{v}}) + n_{non-DYbg}(\bar{\mathbf{v}}), \quad (\text{A.1})$$

where μ is the number of expected events in each mass bin and n_{DY+CI} and n_{non-DY} are the number of events predicted by a particular benchmark signal sample and the number of predicted non DY background events respectively. \mathbf{v} is a set of Gaussian nuisance parameters that account for systematic uncertainties in the analysis while θ corresponds to the energy scale Λ . The likelihood function for observing a set of \bar{n} events in N mass bins is therefore given by:

$$\mathcal{L}(\bar{n} | \theta, \bar{\mathbf{v}}) = \prod_{k=1}^N \frac{\mu_k^{n_k} e^{-\mu_k}}{n_k!} \quad (\text{A.2})$$

as a product of Poission probabilities for each mass bin k . Using Bayes' theorem this gives posterior probability

$$\mathcal{P}(\theta | \bar{n}) = \frac{1}{Z} \mathcal{L}_M(\bar{n} | \theta) P(\theta) \quad (\text{A.3})$$

where Z is a normalisation constant and $\mathcal{L}_{\mathcal{M}}$ is the marginalised likelihood after all nuisance parameters have been integrated out. A prior probability for $P(\theta)$ is chosen to be flat in $1/\Lambda^2$, motivated by the CI differential cross-section (Eq. 1.3).

A 95% confidence level (CL) limit is found by finding Λ_{lim} that satisfies $\int_0^{\theta_{lim}} P(\theta | \bar{n}) d\theta = 0.95$ with $\theta = 1/\Lambda^2$. For this analysis the Bayesian Analysis Toolkit (BAT) [?] was used to do this calculation.

A.4 Results

Results for the 2011 analysis are currently undergoing approval from ATLAS. Therefore the results presented here are provisional limits.

Channel	ee	$\mu\mu$	ee+ $\mu\mu$
Expected CI constructive	13.73 TeV	12.24 TeV	15.10 TeV
Expected CI destructive	10.41 TeV	10.23 TeV	11.42 TeV
Observed CI constructive	11.60 TeV	12.07 TeV	12.70 TeV
Observed CI destructive	8.76 TeV	9.17 TeV	9.63 TeV
Expected ADD	2.84 TeV	2.71 TeV	2.94 TeV
Observed ADD	2.71 TeV	2.72 TeV	2.94 TeV

Table A.6: Table of 95% confidence level limits found in the CI and ADD analyses

An expected limit was obtained for the data set by generating 1000 Standard Model like pseudoexperiments. The Bayesian limit setting method is then applied to each of these 1000 pseudoexperiments to get a distribution of 95% CL limits. The median of these distributions is then taken as the expected limit for each channel. These results can be seen in table A.6 in which the expected limits are generally found to be higher because they dictate a scenario where no deviation from the Standard model background is found. In reality slight statistical fluctuations are found in the data set and so limits seen are lower than expected.

These limits are found to be higher than the limits in the last iteration of this analysis seen in the paper provided and the first limits set by ATLAS on ADD.

Bibliography

- [1] E. Eichten, K. D. Lane, M. E. Peskin, New Tests for Quark and Lepton Substructure, *Phys.Rev.Lett.* 50 (1983) 811–814. doi:10.1103/PhysRevLett.50.811.
- [2] V. M. Abazov, et al., Measurement of dijet angular distributions at $s=1.96\text{TeV}$ and searches for quark compositeness and extra spatial dimensions, *Phys. Rev. Lett.* 103 (2009) 191803. doi:10.1103/PhysRevLett.103.191803.
URL <http://link.aps.org/doi/10.1103/PhysRevLett.103.191803>
- [3] A. Abulencia, et al., Search for $Z'\rightarrow e+e-$ using dielectron mass and angular distribution, *Phys. Rev. Lett.* 96 (2006) 211801. doi:10.1103/PhysRevLett.96.211801.
URL <http://link.aps.org/doi/10.1103/PhysRevLett.96.211801>
- [4] G. Aad, et al., Search for contact interactions and large extra dimensions in dilepton events from pp collisions at $s=7\text{ TeV}$ with the ATLAS detector, *Phys. Rev. D* 87 (2013) 015010. doi:10.1103/PhysRevD.87.015010.
URL <http://link.aps.org/doi/10.1103/PhysRevD.87.015010>
- [5] J. C. Collins, D. E. Soper, Angular distribution of dileptons in high-energy hadron collisions, *Phys. Rev. D* 16 (1977) 2219–2225. doi:10.1103/PhysRevD.16.2219.
URL <http://link.aps.org/doi/10.1103/PhysRevD.16.2219>
- [6] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, The Hierarchy problem and new dimensions at a millimeter, *Phys.Lett. B* 429 (1998) 263–272. arXiv:hep-ph/9803315, doi:10.1016/S0370-2693(98)00466-3.
- [7] G. F. Giudice, R. Rattazzi, J. D. Wells, Quantum gravity and extra dimensions at high-energy colliders, *Nucl.Phys. B* 544 (1999) 3–38. arXiv:hep-ph/9811291, doi:10.1016/S0550-3213(99)00044-9.
- [8] T. Han, J. D. Lykken, R.-J. Zhang, Kaluza-Klein states from large extra dimensions, *Phys. Rev. D* 59 (1999) 105006. doi:10.1103/PhysRevD.59.105006.
URL <http://link.aps.org/doi/10.1103/PhysRevD.59.105006>

- [9] J. L. Hewett, Indirect collider signals for extra dimensions, *Phys. Rev. Lett.* 82 (1999) 4765–4768. doi:10.1103/PhysRevLett.82.4765.
URL <http://link.aps.org/doi/10.1103/PhysRevLett.82.4765>
- [10] G. Aad, et al., ATLAS search for new phenomena in dijet mass and angular distributions using pp collisions at $\sqrt{s} = 7$ TeV, *JHEP* 1301 (2013) 029. arXiv:1210.1718, doi:10.1007/JHEP01(2013)029.
- [11] S. Chatrchyan, et al., Search for contact interactions in $\mu+\mu^-$ events in pp collisions at $s=7$ TeV, *Phys. Rev. D* 87 (2013) 032001. doi:10.1103/PhysRevD.87.032001.
URL <http://link.aps.org/doi/10.1103/PhysRevD.87.032001>
- [12] S. Chatrchyan, et al., Search for contact interactions using the inclusive jet p_T spectrum in pp collisions at $s=7$ TeV, *Phys. Rev. D* 87 (2013) 052017. doi:10.1103/PhysRevD.87.052017.
URL <http://link.aps.org/doi/10.1103/PhysRevD.87.052017>
- [13] T. Affolder, et al., Search for quark-lepton compositeness and a heavy W' boson using the $e\nu$ channel in $p\bar{p}$ collisions at $s=1.8$ TeV, *Phys. Rev. Lett.* 87 (2001) 231803. doi:10.1103/PhysRevLett.87.231803.
URL <http://link.aps.org/doi/10.1103/PhysRevLett.87.231803>
- [14] B. Abbott, et al., Measurement of the high-mass drell-yan cross section and limits on quark-electron compositeness scales, *Phys. Rev. Lett.* 82 (1999) 4769–4774. doi:10.1103/PhysRevLett.82.4769.
URL <http://link.aps.org/doi/10.1103/PhysRevLett.82.4769>
- [15] F. Abe, et al., Limits on quark-lepton compositeness scales from dileptons produced in 1.8 TeV $p\bar{p}$ collisions, *Phys. Rev. Lett.* 79 (1997) 2198–2203. doi:10.1103/PhysRevLett.79.2198.
URL <http://link.aps.org/doi/10.1103/PhysRevLett.79.2198>
- [16] S. Chekanov, et al., Search for contact interactions, large extra dimensions and finite quark radius in ep collisions at HERA, *Physics Letters B* 591 (12) (2004) 23 – 41. doi:http://dx.doi.org/10.1016/j.physletb.2004.03.081.
URL <http://www.sciencedirect.com/science/article/pii/S0370269304005660>
- [17] C. Adloff, et al., Search for new physics in $e\pm q$ contact interactions at HERA, *Physics Letters B* 568 (12) (2003) 35 – 47. doi:http://dx.doi.org/10.1016/j.physletb.2003.06.034.
URL <http://www.sciencedirect.com/science/article/pii/S0370269303009316>
- [18] J. Abdallah, et al., A study of $b\bar{b}$ production in $e+e^-$ collisions at $\sqrt{s} = 130 - 207$ GeV, *The European Physical Journal C* 60 (1) (2009) 1–15. doi:10.1140/epjc/s10052-009-0917-2.
URL <http://dx.doi.org/10.1140/epjc/s10052-009-0917-2>

- [19] S. Schael, et al., Fermion pair production in e^+e^- collisions at 189-209 GeV and constraints on physics beyond the standard model, *The European Physical Journal C* 49 (2) (2007) 411–437. doi:10.1140/epjc/s10052-006-0156-8.
URL <http://dx.doi.org/10.1140/epjc/s10052-006-0156-8>
- [20] J. Abdallah, et al., Measurement and interpretation of fermion-pair production at LEP energies above the Z resonance, *The European Physical Journal C* 45 (3) (2006) 589–632. doi:10.1140/epjc/s2005-02461-0.
URL <http://dx.doi.org/10.1140/epjc/s2005-02461-0>
- [21] G. Abbiendi, et al., Tests of the standard model and constraints on new physics from measurements of fermion-pair production at 189-209 GeV at LEP, *The European Physical Journal C* 33 (2) (2004) 173–212. doi:10.1140/epjc/s2004-01595-9.
URL <http://dx.doi.org/10.1140/epjc/s2004-01595-9>
- [22] M. Acciarri, et al., Search for manifestations of new physics in fermion-pair production at LEP, *Physics Letters B* 489 (12) (2000) 81 – 92. doi:[http://dx.doi.org/10.1016/S0370-2693\(00\)00887-X](http://dx.doi.org/10.1016/S0370-2693(00)00887-X).
URL <http://www.sciencedirect.com/science/article/pii/S037026930000887X>
- [23] L. Evans, The large hadron collider, *New Journal of Physics* 9 (9) (2007) 335.
URL <http://stacks.iop.org/1367-2630/9/i=9/a=335>
- [24] O. S. Brning, P. Collier, P. Lebrun, S. Myers, R. Ostojic, J. Poole, P. Proudlock, LHC Design Report, CERN, Geneva, 2004.
- [25] G. Aad, et al., The ATLAS Experiment at the CERN Large Hadron Collider, *Journal of Instrumentation* 3 (2008) S08003. 437 p, also published by CERN Geneva in 2010.