

Complex analysis and Fourier analysis.

Complex analysis.

Complex numbers appear already in Cardano's formula for the solution of third degree equations, where combinations of complex numbers could be used to get real solutions of an equation. They were also used in the middle 18:th century by de Moivre and Euler in their famous formulas

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta,$$

Complex analysis.

Complex numbers appear already in Cardano's formula for the solution of third degree equations, where combinations of complex numbers could be used to get real solutions of an equation. They were also used in the middle 18:th century by de Moivre and Euler in their famous formulas

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta,$$

and

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Complex analysis.

Complex numbers appear already in Cardano's formula for the solution of third degree equations, where combinations of complex numbers could be used to get real solutions of an equation. They were also used in the middle 18:th century by de Moivre and Euler in their famous formulas

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta,$$

and

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

It is unclear (to me) when complex numbers began to be considered as a well defined mathematical structure. Perhaps this did not happen until Gauss, who however had been reluctant to compute openly with complex numbers in his early work. Complex analysis, involving e g the Cauchy-Riemann equations goes back at least to d'Alembert, Euler and Riemann in the 18:th century, and became a fully developed theory with Cauchy's work.

The representation of complex numbers as points in \mathbb{R}^2 is usually ascribed to the Norwegian cartographer C Wessel who published his findings in 1797. His work went unnoticed for many years, but the idea was rediscovered by Argand in 1806 and later Gauss. Argand had a bookstore in Paris and was an amateur mathematician. Today he is also given credit for the first correct proof of the fundamental theorem of algebra in (1814). Previous proofs had been given by Euler, d'Alembert and Gauss, but none of them is considered complete.

The representation of complex numbers as points in \mathbb{R}^2 is usually ascribed to the Norwegian cartographer C Wessel who published his findings in 1797. His work went unnoticed for many years, but the idea was rediscovered by Argand in 1806 and later Gauss. Argand had a bookstore in Paris and was an amateur mathematician. Today he is also given credit for the first correct proof of the fundamental theorem of algebra in (1814). Previous proofs had been given by Euler, d'Alembert and Gauss, but none of them is considered complete. Gauss gave four different proofs of the theorem – probably a reflection of the fact that he was not content with his first attempts.

The representation of complex numbers as points in \mathbb{R}^2 is usually ascribed to the Norwegian cartographer C Wessel who published his findings in 1797. His work went unnoticed for many years, but the idea was rediscovered by Argand in 1806 and later Gauss. Argand had a bookstore in Paris and was an amateur mathematician. Today he is also given credit for the first correct proof of the fundamental theorem of algebra in (1814). Previous proofs had been given by Euler, d'Alembert and Gauss, but none of them is considered complete. Gauss gave four different proofs of the theorem – probably a reflection of the fact that he was not content with his first attempts. Today it is easy to see where the obstacle lay: The crucial point which makes the theorem hold is a property of the *real* number system, not the complex numbers. This property is the completeness, or in other words the supremum axiom, and these matters were not clarified until later.

Let us start by giving an idea of Gauss' first proof. In modern language he considered an algebraic equation

$$p(z) = z^n + a_1 z^{n-1} + \dots a_n = 0,$$

where a_j are *real*, and wanted to show that it has a complex solution.

Let us start by giving an idea of Gauss' first proof. In modern language he considered an algebraic equation

$$p(z) = z^n + a_1 z^{n-1} + \dots + a_n = 0,$$

where a_j are *real*, and wanted to show that it has a complex solution. Taking real and imaginary parts, and writing $z = r(\cos \theta + i \sin \theta)$ we get two equations

$$u(z) = r^n \cos n\theta + a_1 r^{n-1} \cos n\theta + \dots = 0$$

and

$$v(z) = r^n \sin n\theta + r^{n-1} \sin(n-1)\theta + \dots = 0.$$

Let us start by giving an idea of Gauss' first proof. In modern language he considered an algebraic equation

$$p(z) = z^n + a_1 z^{n-1} + \dots + a_n = 0,$$

where a_j are *real*, and wanted to show that it has a complex solution. Taking real and imaginary parts, and writing $z = r(\cos \theta + i \sin \theta)$ we get two equations

$$u(z) = r^n \cos n\theta + a_1 r^{n-1} \cos n\theta + \dots = 0$$

and

$$v(z) = r^n \sin n\theta + r^{n-1} \sin(n-1)\theta + \dots = 0.$$

He then argued that the solution set of the first equation is an algebraic curve C_1 , the solution set of the second equation is also an algebraic curve, C_2 , and we want to prove that these two curves intersect.

For this he took a large disk, $\{|z| < R\}$, and considered the intersection of the curves with the boundary of the disk.

For this he took a large disk, $\{|z| < R\}$, and considered the intersection of the curves with the boundary of the disk. He argued (correctly) that the first curve intersects the circle in $2n$ points, as does the second curve, and these intersection points are intertwined: First a C_1 -point, then a C_2 -point, then a C_1 -point etc. (This was based on comparison with the case $p(z) = z^n$ which approximates our p well when R is large.)

For this he took a large disk, $\{|z| < R\}$, and considered the intersection of the curves with the boundary of the disk. He argued (correctly) that the first curve intersects the circle in $2n$ points, as does the second curve, and these intersection points are intertwined: First a C_1 -point, then a C_2 -point, then a C_1 -point etc. (This was based on comparison with the case $p(z) = z^n$ which approximates our p well when R is large.) He then claimed:

“It seems to be well demonstrated that an algebraic curve neither ends abruptly (as it happens in the transcendental curve $y = 1/\log x$), nor loses itself after an infinite number of windings in a point (like a logarithmic spiral). As far as I know nobody has ever doubted this, but if anybody requires it, I take it on me to present, on another occasion, an indubitable proof.”

Accepting this, the argument is concluded by a combinatorical argument. The argument is clearly not complete by modern standards.

Accepting this, the argument is concluded by a combinatorical argument. The argument is clearly not complete by modern standards. Let us now see a modern proof of the theorem, which seems to be essentially what d'Alembert proposed:

Accepting this, the argument is concluded by a combinatorical argument. The argument is clearly not complete by modern standards. Let us now see a modern proof of the theorem, which seems to be essentially what d'Alembert proposed:

1. The function $|p(z)|$ tends to infinity at infinity, so it must have minimum some place.

Accepting this, the argument is concluded by a combinatorical argument. The argument is clearly not complete by modern standards. Let us now see a modern proof of the theorem, which seems to be essentially what d'Alembert proposed:

1. The function $|p(z)|$ tends to infinity at infinity, so it must have minimum some place.
2. If the minimum value is zero we are done, so assume it is not equal to zero.

Accepting this, the argument is concluded by a combinatorial argument. The argument is clearly not complete by modern standards. Let us now see a modern proof of the theorem, which seems to be essentially what d'Alembert proposed:

1. The function $|p(z)|$ tends to infinity at infinity, so it must have minimum some place.
2. If the minimum value is zero we are done, so assume it is not equal to zero.
3. We may assume that the minimum is attained at $z = 0$ (If the minimum is attained at a , consider $q(z) = p(a + z)$ which is also a polynomial. Then $|q|$ has a non zero minimum at $z = 0$.)

Accepting this, the argument is concluded by a combinatorial argument. The argument is clearly not complete by modern standards. Let us now see a modern proof of the theorem, which seems to be essentially what d'Alembert proposed:

1. The function $|p(z)|$ tends to infinity at infinity, so it must have minimum some place.
2. If the minimum value is zero we are done, so assume it is not equal to zero.
3. We may assume that the minimum is attained at $z = 0$ (If the minimum is attained at a , consider $q(z) = p(a + z)$ which is also a polynomial. Then $|q|$ has a non zero minimum at $z = 0$.)
4. Then $p(z) = a_n + z^k(b_0 + zb_1 + \dots)$. There are (small) values of z which makes this smaller than $|a_n|$.
5. This contradiction proves the theorem.

This was also the main scheme in Argand's proof. Both Argand and d'Alembert took 2. for granted: A continuous real valued function defined on a compact set has a minimum value. This was only proved rigorously much later!

Point 4. is known as d'Alembert's lemma, but it seems his argument was shaky.

This was also the main scheme in Argand's proof. Both Argand and d'Alembert took 2. for granted: A continuous real valued function defined on a compact set has a minimum value. This was only proved rigorously much later!

Point 4. is known as d'Alembert's lemma, but it seems his argument was shaky.

Exercise: Help him! In other words, prove that if $p(z) = a_1 + z^k(b_0 + b_1 z + \dots)$, $k = 1, 2, \dots$ and $a_1 \neq 0$, then $|p(0)| = |a_1|$ is *not* the minimum of $|p|$

This was also the main scheme in Argand's proof. Both Argand and d'Alembert took 2. for granted: A continuous real valued function defined on a compact set has a minimum value. This was only proved rigorously much later!

Point 4. is known as d'Alembert's lemma, but it seems his argument was shaky.

Exercise: Help him! In other words, prove that if $p(z) = a_1 + z^k(b_0 + b_1 z + \dots)$, $k = 1, 2, \dots$ and $a_1 \neq 0$, then $|p(0)| = |a_1|$ is *not* the minimum of $|p|$

It seems this was Argand's main contribution; to prove 4.

But, the main step forward was with the introduction of the *holomorphic functions*. The usual text book definition is :

But, the main step forward was with the introduction of the *holomorphic functions*. The usual text book definition is :A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic if the *complex derivative* of f ,

$$f'(z) = \lim_{w \rightarrow z} \frac{f(w) - f(z)}{w - z}$$

exists at every point (in its domain of definition).

But, the main step forward was with the introduction of the *holomorphic functions*. The usual text book definition is :A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic if the *complex derivative* of f ,

$$f'(z) = \lim_{w \rightarrow z} \frac{f(w) - f(z)}{w - z}$$

exists at every point (in its domain of definition).

This seemingly simple definition has many not so simple consequences:

1. f has a series expansion $f(z) = a_0 + a_1 z + a_2 z^2 + \dots$

But, the main step forward was with the introduction of the *holomorphic functions*. The usual text book definition is :A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic if the *complex derivative* of f ,

$$f'(z) = \lim_{w \rightarrow z} \frac{f(w) - f(z)}{w - z}$$

exists at every point (in its domain of definition).

This seemingly simple definition has many not so simple consequences:

1. f has a series expansion $f(z) = a_0 + a_1 z + a_2 z^2 + \dots$
2. Any function that is holomorphic in all of \mathbb{C} and bounded must be constant.

But, the main step forward was with the introduction of the *holomorphic functions*. The usual text book definition is :A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic if the *complex derivative* of f ,

$$f'(z) = \lim_{w \rightarrow z} \frac{f(w) - f(z)}{w - z}$$

exists at every point (in its domain of definition).

This seemingly simple definition has many not so simple consequences:

1. f has a series expansion $f(z) = a_0 + a_1 z + a_2 z^2 + \dots$
2. Any function that is holomorphic in all of \mathbb{C} and bounded must be constant.
3. If two holomorphic functions are equal on \mathbb{R} then they are equal everywhere.

But, the main step forward was with the introduction of the *holomorphic functions*. The usual text book definition is :A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is holomorphic if the *complex derivative* of f ,

$$f'(z) = \lim_{w \rightarrow z} \frac{f(w) - f(z)}{w - z}$$

exists at every point (in its domain of definition).

This seemingly simple definition has many not so simple consequences:

1. f has a series expansion $f(z) = a_0 + a_1 z + a_2 z^2 + \dots$
2. Any function that is holomorphic in all of \mathbb{C} and bounded must be constant.
3. If two holomorphic functions are equal on \mathbb{R} then they are equal everywhere. Example: $e^{x+y} = e^x e^y$ implies $e^{i\theta+i\phi} = e^{i\theta} e^{i\phi}$ implies

$$\sin(\theta + \phi) + i \cos(\theta + \phi) = (\sin \theta + i \cos \theta)(\sin \phi + i \cos \phi).$$

From a more modern view point the definition can be described as follows. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a smooth function. We can view it as a function from \mathbb{R}^2 to \mathbb{R}^2 . Then it has a differential at any point z

$$df|_z = f_x dx + f_y dy.$$

From a more modern view point the definition can be described as follows. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a smooth function. We can view it as a function from \mathbb{R}^2 to \mathbb{R}^2 . Then it has a differential at any point z

$$df|_z = f_x dx + f_y dy.$$

Fixing the point z the differential is a linear map from \mathbb{R}^2 to \mathbb{R}^2 :

$$df|_z(a + ib) = f_x a + f_y b.$$

From a more modern view point the definition can be described as follows. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a smooth function. We can view it as a function from \mathbb{R}^2 to \mathbb{R}^2 . Then it has a differential at any point z

$$df|_z = f_x dx + f_y dy.$$

Fixing the point z the differential is a linear map from \mathbb{R}^2 to \mathbb{R}^2 :

$$df|_z(a + ib) = f_x a + f_y b.$$

The function is holomorphic if the differential is **linear as a map from \mathbb{C} to \mathbb{C}** :

$$df(ib) = i df(b),$$

which means

$$f_y b = i f_x b,$$

i.e

$$f_x + i f_y = 0.$$

This is called the *Cauchy-Riemann equations* and usually written $f_{\bar{z}} = 0$.

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

Such linear maps preserve angles between lines.

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

Such linear maps preserve angles between lines. This is because A can be written $A = re^{i\phi}$, so the map $w \rightarrow Aw$ is the composition of two maps; $w \rightarrow rw$ and $w \rightarrow e^{i\phi}w$. The first map is just a dilation, so it preserves angles. The second map is a rotation, so it also preserves angles.

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

Such linear maps preserve angles between lines. This is because A can be written $A = re^{i\phi}$, so the map $w \rightarrow Aw$ is the composition of two maps; $w \rightarrow rw$ and $w \rightarrow e^{i\phi}w$. The first map is just a dilation, so it preserves angles. The second map is a rotation, so it also preserves angles.

Therefore holomorphic functions preserve angles between curves; they are *conformal*. (At least where $f'(z) \neq 0$.)

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

Such linear maps preserve angles between lines. This is because A can be written $A = re^{i\phi}$, so the map $w \rightarrow Aw$ is the composition of two maps; $w \rightarrow rw$ and $w \rightarrow e^{i\phi}w$. The first map is just a dilation, so it preserves angles. The second map is a rotation, so it also preserves angles.

Therefore holomorphic functions preserve angles between curves; they are *conformal*. (At least where $f'(z) \neq 0$.) This means that they preserve shapes, and for that reason they are often used when making maps.

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

Such linear maps preserve angles between lines. This is because A can be written $A = re^{i\phi}$, so the map $w \rightarrow Aw$ is the composition of two maps; $w \rightarrow rw$ and $w \rightarrow e^{i\phi}w$. The first map is just a dilation, so it preserves angles. The second map is a rotation, so it also preserves angles.

Therefore holomorphic functions preserve angles between curves; they are *conformal*. (At least where $f'(z) \neq 0$.) This means that they preserve shapes, and for that reason they are often used when making maps. All elementary functions are holomorphic, so there are many to make maps with.

If the Cauchy-Riemann equations hold, then

$$df|_z(a + ib) = f_x(a + ib) = A(a + ib), \quad df|_z(w) = Aw.$$

Such linear maps preserve angles between lines. This is because A can be written $A = re^{i\phi}$, so the map $w \rightarrow Aw$ is the composition of two maps; $w \rightarrow rw$ and $w \rightarrow e^{i\phi}w$. The first map is just a dilation, so it preserves angles. The second map is a rotation, so it also preserves angles.

Therefore holomorphic functions preserve angles between curves; they are *conformal*. (At least where $f'(z) \neq 0$.) This means that they preserve shapes, and for that reason they are often used when making maps. All elementary functions are holomorphic, so there are many to make maps with.

But, how do we map the surface of the earth (a sphere) to the plane?

This can be accomplished with the *stereographic projection*:

This can be accomplished with the *stereographic projection*:

Take the earth in form of a globe and put it on table (the complex plane), with the north pole facing up. For any other point p on the surface of the earth, draw a line through the north pole and p . That line (extended) intersects the table (complex plane) in one point, $s(p)$.

This can be accomplished with the *stereographic projection*:

Take the earth in form of a globe and put it on table (the complex plane), with the north pole facing up. For any other point p on the surface of the earth, draw a line through the north pole and p . That line (extended) intersects the table (complex plane) in one point, $s(p)$.

The map $p \rightarrow s(p)$ is the stereographic projection.

This can be accomplished with the *stereographic projection*:

Take the earth in form of a globe and put it on table (the complex plane), with the north pole facing up. For any other point p on the surface of the earth, draw a line through the north pole and p . That line (extended) intersects the table (complex plane) in one point, $s(p)$.

The map $p \rightarrow s(p)$ is the stereographic projection. This map is also conformal! The ‘Mercator projection’, often used in maps, is $\log s(p)$.

This can be accomplished with the *stereographic projection*:

Take the earth in form of a globe and put it on table (the complex plane), with the north pole facing up. For any other point p on the surface of the earth, draw a line through the north pole and p . That line (extended) intersects the table (complex plane) in one point, $s(p)$.

The map $p \rightarrow s(p)$ is the stereographic projection. This map is also conformal! The ‘Mercator projection’, often used in maps, is $\log s(p)$.

Note that the north pole corresponds to infinity!

This can be accomplished with the *stereographic projection*:

Take the earth in form of a globe and put it on table (the complex plane), with the north pole facing up. For any other point p on the surface of the earth, draw a line through the north pole and p . That line (extended) intersects the table (complex plane) in one point, $s(p)$.

The map $p \rightarrow s(p)$ is the stereographic projection. This map is also conformal! The ‘Mercator projection’, often used in maps, is $\log s(p)$.

Note that the north pole corresponds to infinity! The sphere looked upon this way, as \mathbb{C} together with infinity, is called the *Riemann sphere*. We can therefore talk about holomorphic functions on a sphere too.

This can be accomplished with the *stereographic projection*:

Take the earth in form of a globe and put it on table (the complex plane), with the north pole facing up. For any other point p on the surface of the earth, draw a line through the north pole and p . That line (extended) intersects the table (complex plane) in one point, $s(p)$.

The map $p \rightarrow s(p)$ is the stereographic projection. This map is also conformal! The 'Mercator projection', often used in maps, is $\log s(p)$.

Note that the north pole corresponds to infinity! The sphere looked upon this way, as \mathbb{C} together with infinity, is called the *Riemann sphere*. We can therefore talk about holomorphic functions on a sphere too.

Any other (orientable) closed surface without boundary (like the sphere) can also be looked at in a similar way, as a complex manifold. This leads to the concept of a Riemann surface.

Many things in real analysis can only be understood from a complex view point. Here is a very simple example :

Many things in real analysis can only be understood from a complex view point. Here is a very simple example :

Let

$$g(x) = \frac{1}{1+x^2}.$$

It is perfectly smooth on all of \mathbb{R} .

Many things in real analysis can only be understood from a complex view point. Here is a very simple example :

Let

$$g(x) = \frac{1}{1+x^2}.$$

It is perfectly smooth on all of \mathbb{R} . Its Taylor expansion is

$$g(x) = \sum_{k=0}^{\infty} (-1)^k x^{2k}.$$

It converges only for $|x| < 1$. Why?

Many things in real analysis can only be understood from a complex view point. Here is a very simple example :

Let

$$g(x) = \frac{1}{1+x^2}.$$

It is perfectly smooth on all of \mathbb{R} . Its Taylor expansion is

$$g(x) = \sum_{k=0}^{\infty} (-1)^k x^{2k}.$$

It converges only for $|x| < 1$. Why?

Because g has a singularity when $x = i$.

Fourier Analysis

Fourier Analysis

Fourier Analysis concerns the expression of arbitrary functions as a superposition – sum or integral – of trigonometric functions, or equivalently by Euler's formula of complex exponentials. The creation of the theory is certainly Fourier's achievement, even though special cases had previously been studied by others.

Fourier Analysis

Fourier Analysis concerns the expression of arbitrary functions as a superposition – sum or integral – of trigonometric functions, or equivalently by Euler's formula of complex exponentials. The creation of the theory is certainly Fourier's achievement, even though special cases had previously been studied by others.

It is also worth to point out that Ptolemaios' description of the motion of the planets as a sum of epicycles can be seen as an early version of the same idea.

Fourier Analysis

Fourier Analysis concerns the expression of arbitrary functions as a superposition – sum or integral – of trigonometric functions, or equivalently by Euler's formula of complex exponentials. The creation of the theory is certainly Fourier's achievement, even though special cases had previously been studied by others.

It is also worth to point out that Ptolemaios' description of the motion of the planets as a sum of epicycles can be seen as an early version of the same idea. This viewpoint is far from dead; it plays an important role in the modern mathematical theory of stability of the solar system.

Fourier Analysis

Fourier Analysis concerns the expression of arbitrary functions as a superposition – sum or integral – of trigonometric functions, or equivalently by Euler's formula of complex exponentials. The creation of the theory is certainly Fourier's achievement, even though special cases had previously been studied by others.

It is also worth to point out that Ptolemaios' description of the motion of the planets as a sum of epicycles can be seen as an early version of the same idea. This viewpoint is far from dead; it plays an important role in the modern mathematical theory of stability of the solar system.

J-B Fourier was born in 1768. He studied mathematics at a military academy, run by the Benedictine monks, and planned to become a monk himself, stopped only by the dissolution of the religious orders during the French revolution, two days before he were to make his oath. He was himself active in the revolution, and narrowly escaped to be executed during the reign of terror. He became a professor of mathematics at Ecole Polytechnique in 1797.

One year later he was sent by Napoleon with the expedition to Egypt, and became a prominent actor in the scientifique part of the expedition. His name is therefore also well known in Egyptology, and he wrote the preface to the treatise 'Description de l'Egypte', a general description of all the historical findings of the french in Egypt (many of which were exported to Paris).

One year later he was sent by Napoleon with the expedition to Egypt, and became a prominent actor in the scientifique part of the expedition. His name is therefore also well known in Egyptology, and he wrote the preface to the treatise 'Description de l'Egypte', a general description of all the historical findings of the french in Egypt (many of which were exported to Paris).

Coming back to France, Napoleon named him ('Citizen Fourier') 'prefect' of the department of Isere, in 1802.

One year later he was sent by Napoleon with the expedition to Egypt, and became a prominent actor in the scientific part of the expedition. His name is therefore also well known in Egyptology, and he wrote the preface to the treatise 'Description de l'Egypte', a general description of all the historical findings of the french in Egypt (many of which were exported to Paris).

Coming back to France, Napoleon named him ('Citizen Fourier') 'prefect' of the department of Isere, in 1802. His election to the French Academy of Sciences in 1816 was stopped by the king (Louis XVIII) – Fourier was not a nobleman – but he entered the academy one year later.

One year later he was sent by Napoleon with the expedition to Egypt, and became a prominent actor in the scientifique part of the expedition. His name is therefore also well known in Egyptology, and he wrote the preface to the treatise 'Description de l'Egypte', a general description of all the historical findings of the french in Egypt (many of which were exported to Paris).

Coming back to France, Napoleon named him ('Citizen Fourier') 'prefect' of the department of Isere, in 1802. His election to the French Academy of Sciences in 1816 was stopped by the king (Louis XVIII) – Fourier was not a nobleman – but he entered the academy one year later.

Then, in 1822, he published his 'Theorie analytique de la chaleur'. Here he introduced the *heat equation* and Fourier analysis as a tool to solve it.

So, what is Fourier analysis? We look at two cases:

Let f be a function on the interval $(-\pi, \pi)$. (If we want to we can extend it to a periodic function on all of \mathbb{R}).

So, what is Fourier analysis? We look at two cases:

Let f be a function on the interval $(-\pi, \pi)$. (If we want to we can extend it to a periodic function on all of \mathbb{R}). Let

$$c_k(f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt.$$

These numbers, for $k \in \mathbb{Z}$ are called the *Fourier coefficients* of the function f .

So, what is Fourier analysis? We look at two cases:

Let f be a function on the interval $(-\pi, \pi)$. (If we want to we can extend it to a periodic function on all of \mathbb{R}). Let

$$c_k(f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt.$$

These numbers, for $k \in \mathbb{Z}$ are called the *Fourier coefficients* of the function f . Then Fourier's main result is that

$$f = \sum_{-\infty}^{\infty} c_k(f) e^{ikt},$$

so we can write 'any' function as a sum of trigonometric functions.

So, what is Fourier analysis? We look at two cases:

Let f be a function on the interval $(-\pi, \pi)$. (If we want to we can extend it to a periodic function on all of \mathbb{R}). Let

$$c_k(f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt.$$

These numbers, for $k \in \mathbb{Z}$ are called the *Fourier coefficients* of the function f . Then Fourier's main result is that

$$f = \sum_{-\infty}^{\infty} c_k(f) e^{ikt},$$

so we can write 'any' function as a sum of trigonometric functions.

If we are a bit more precise and assume that f lies in

$$L^2(T) = \{f; \int |f|^2 dt < \infty\},$$

we can give this representation a meaning:

L^2 is a Hilbert space and $\{e_k = e^{ikt}\}_k$ is a basis for L^2 .

L^2 is a Hilbert space and $\{e_k = e^{ikt}\}_k$ is a basis for L^2 . Moreover, it is an orthonormal basis

$$\langle e_k, e_m \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ikt} e^{-imt} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(k-m)t} dt = \delta_{k,m}.$$

L^2 is a Hilbert space and $\{e_k = e^{ikt}\}_k$ is a basis for L^2 . Moreover, it is an orthonormal basis

$$\langle e_k, e_m \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ikt} e^{-imt} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(k-m)t} dt = \delta_{k,m}.$$

It is a special orthonormal basis. If A is the operator $Af = -i(d/dt)$, then $Ae_k = ke_k$, so it is an orthonormal basis of eigenvectors of A , with e_k corresponding to eigenvalue k .

L^2 is a Hilbert space and $\{e_k = e^{ikt}\}_k$ is a basis for L^2 . Moreover, it is an orthonormal basis

$$\langle e_k, e_m \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ikt} e^{-imt} dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(k-m)t} dt = \delta_{k,m}.$$

It is a special orthonormal basis. If A is the operator $Af = -i(d/dt)$, then $Ae_k = ke_k$, so it is an orthonormal basis of eigenvectors of A , with e_k corresponding to eigenvalue k .

Fourier's theorem then says that there is an orthonormal basis of eigenvectors for A , i.e. that A can be diagonalized.

The second version of Fourier analysis is for functions defined on \mathbb{R} , or why not \mathbb{R}^n . Let

$$L^2(\mathbb{R}^n) = \{f; \int_{\mathbb{R}^n} |f|^2 dt\} < \infty.$$

The second version of Fourier analysis is for functions defined on \mathbb{R} , or why not \mathbb{R}^n . Let

$$L^2(\mathbb{R}^n) = \{f; \int_{\mathbb{R}^n} |f|^2 dt\} < \infty.$$

Then the *Fourier transform* of f is

$$\hat{f}(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} f(t) e^{-ix \cdot t} dt,$$

where $x \cdot t = \sum x_j t_j$. It is a function on \mathbb{R}^n .

The second version of Fourier analysis is for functions defined on \mathbb{R} , or why not \mathbb{R}^n . Let

$$L^2(\mathbb{R}^n) = \{f; \int_{\mathbb{R}^n} |f|^2 dt\} < \infty.$$

Then the *Fourier transform* of f is

$$\hat{f}(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} f(t) e^{-ix \cdot t} dt,$$

where $x \cdot t = \sum x_j t_j$. It is a function on \mathbb{R}^n . Then

$$f(t) = \int_{\mathbb{R}^n} \hat{f}(x) e^{ix \cdot t} dx.$$

This is the Fourier inversion formula.

The second version of Fourier analysis is for functions defined on \mathbb{R} , or why not \mathbb{R}^n . Let

$$L^2(\mathbb{R}^n) = \{f; \int_{\mathbb{R}^n} |f|^2 dt\} < \infty.$$

Then the *Fourier transform* of f is

$$\hat{f}(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} f(t) e^{-ix \cdot t} dt,$$

where $x \cdot t = \sum x_j t_j$. It is a function on \mathbb{R}^n . Then

$$f(t) = \int_{\mathbb{R}^n} \hat{f}(x) e^{ix \cdot t} dx.$$

This is the Fourier inversion formula. It is less easy to interpret in the Hilbert space sense; we have now a ‘continuous basis’, and one needs von Neumann’s general operator theory to discuss such matters: The operator id/dx has *continuous spectrum* on \mathbb{R} , i.e. a continuous family of eigenvalues, namely \mathbb{R} .

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Say we have a function f in the plane and that for any line l in the plane we know the integral of f over l . Can we reconstruct f ?

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Say we have a function f in the plane and that for any line l in the plane we know the integral of f over l . Can we reconstruct f ? A special case, when f is the characteristic function of a set K in the plane: We know the length of the intersection $l \cap K$ for any line l . Can we reconstruct K ?

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Say we have a function f in the plane and that for any line l in the plane we know the integral of f over l . Can we reconstruct f ? A special case, when f is the characteristic function of a set K in the plane: We know the length of the intersection $l \cap K$ for any line l . Can we reconstruct K ? Yes, we can!

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Say we have a function f in the plane and that for any line l in the plane we know the integral of f over l . Can we reconstruct f ? A special case, when f is the characteristic function of a set K in the plane: We know the length of the intersection $l \cap K$ for any line l . Can we reconstruct K ? Yes, we can! It is enough to find the Fourier transform

$$\hat{f}(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(t) e^{-ix \cdot t} dt$$

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Say we have a function f in the plane and that for any line l in the plane we know the integral of f over l . Can we reconstruct f ? A special case, when f is the characteristic function of a set K in the plane: We know the length of the intersection $l \cap K$ for any line l . Can we reconstruct K ? Yes, we can! It is enough to find the Fourier transform

$$\hat{f}(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(t) e^{-ix \cdot t} dt$$

To illustrate, say $x = (x_1, 0)$. Then

$$\hat{f}(x_1, 0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ix_1 t_1} dt_1 \int_{-\infty}^{\infty} f(t_1, t_2) dt_2.$$

Fourier introduced his transforms to solve (partial) differential equations. But, it turned out to have many other magical applications. Here is one example, the *Radon transform*:

Say we have a function f in the plane and that for any line l in the plane we know the integral of f over l . Can we reconstruct f ? A special case, when f is the characteristic function of a set K in the plane: We know the length of the intersection $l \cap K$ for any line l . Can we reconstruct K ? Yes, we can! It is enough to find the Fourier transform

$$\hat{f}(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(t) e^{-ix \cdot t} dt$$

To illustrate, say $x = (x_1, 0)$. Then

$$\hat{f}(x_1, 0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ix_1 t_1} dt_1 \int_{-\infty}^{\infty} f(t_1, t_2) dt_2.$$

The integral with respect to t_2 is the integral over vertical lines, which we assume we know. So, we know \hat{f} , so we know f by Fourier inversion!

This method – tomography – is used in medical imaging. The set K is something inside the body (a tumor, organ or ...), the length of the intersection with a line is measured by how much of an X-ray in that direction is absorbed. It was awarded the Nobel prize in medicine in 1979.

This method – tomography – is used in medical imaging. The set K is something inside the body (a tumor, organ or ...), the length of the intersection with a line is measured by how much of an X-ray in that direction is absorbed. It was awarded the Nobel prize in medicine in 1979.

Our next example of applications of Fourier analysis is *the Shannon Sampling Theorem*:

This method – tomography – is used in medical imaging. The set K is something inside the body (a tumor, organ or ...), the length of the intersection with a line is measured by how much of an X-ray in that direction is absorbed. It was awarded the Nobel prize in medicine in 1979.

Our next example of applications of Fourier analysis is *the Shannon Sampling Theorem*:

Theorem

Let $f(t)$ be a function on \mathbb{R} which is such that its Fourier transform $\hat{f}(x)$ vanishes outside the interval $[-\pi, \pi]$. Then f is uniquely determined by its values at the integers, and more precisely

$$f(t) = \sum f(n) \frac{\sin \pi(t - n)}{\pi(t - n)}$$

We can also rescale the function and get equivalently

Theorem

Let $f(t)$ be a function on \mathbb{R} which is such that its Fourier transform $\hat{f}(x)$ vanishes outside the interval $[-B\pi, B\pi]$. Then f is uniquely determined by its values at the points n/B ,

$$f(t) = \sum f(n/B) \frac{\sin \pi(Bt - n)}{\pi(Bt - n)}$$

We can also rescale the function and get equivalently

Theorem

Let $f(t)$ be a function on \mathbb{R} which is such that its Fourier transform $\hat{f}(x)$ vanishes outside the interval $[-B\pi, B\pi]$. Then f is uniquely determined by its values at the points n/B ,

$$f(t) = \sum f(n/B) \frac{\sin \pi(Bt - n)}{\pi(Bt - n)}$$

Think of $f(t)$ as a signal varying with time, like the sound of music. ($f(t)$ is then the air pressure at a given fixed point at time t .) The theorem says that if the signal does not ‘contain frequencies’ higher than $B\pi$ then we can, from a sequence of discrete measurements with interval $1/B$, reconstruct the signal perfectly.

We can also rescale the function and get equivalently

Theorem

Let $f(t)$ be a function on \mathbb{R} which is such that its Fourier transform $\hat{f}(x)$ vanishes outside the interval $[-B\pi, B\pi]$. Then f is uniquely determined by its values at the points n/B ,

$$f(t) = \sum f(n/B) \frac{\sin \pi(Bt - n)}{\pi(Bt - n)}$$

Think of $f(t)$ as a signal varying with time, like the sound of music. ($f(t)$ is then the air pressure at a given fixed point at time t .) The theorem says that if the signal does not ‘contain frequencies’ higher than $B\pi$ then we can, from a sequence of discrete measurements with interval $1/B$, reconstruct the signal perfectly. This is what is done in digital recording of music.

We can also rescale the function and get equivalently

Theorem

Let $f(t)$ be a function on \mathbb{R} which is such that its Fourier transform $\hat{f}(x)$ vanishes outside the interval $[-B\pi, B\pi]$. Then f is uniquely determined by its values at the points n/B ,

$$f(t) = \sum f(n/B) \frac{\sin \pi(Bt - n)}{\pi(Bt - n)}$$

Think of $f(t)$ as a signal varying with time, like the sound of music. ($f(t)$ is then the air pressure at a given fixed point at time t .) The theorem says that if the signal does not ‘contain frequencies’ higher than $B\pi$ then we can, from a sequence of discrete measurements with interval $1/B$, reconstruct the signal perfectly. This is what is done in digital recording of music. It is also (I believe) what lies behind the term *broadband*.

We can also rescale the function and get equivalently

Theorem

Let $f(t)$ be a function on \mathbb{R} which is such that its Fourier transform $\hat{f}(x)$ vanishes outside the interval $[-B\pi, B\pi]$. Then f is uniquely determined by its values at the points n/B ,

$$f(t) = \sum f(n/B) \frac{\sin \pi(Bt - n)}{\pi(Bt - n)}$$

Think of $f(t)$ as a signal varying with time, like the sound of music. ($f(t)$ is then the air pressure at a given fixed point at time t .) The theorem says that if the signal does not 'contain frequencies' higher than $B\pi$ then we can, from a sequence of discrete measurements with interval $1/B$, reconstruct the signal perfectly. This is what is done in digital recording of music. It is also (I believe) what lies behind the term *broadband*. The broader the frequency band, i.e. the larger B is, the denser you need or can choose the points, which means that you can transmit more information per time unit.

Let us now sketch the proof of Shannon's theorem.

Let us now sketch the proof of Shannon's theorem. By the inversion formula for the Fourier *transform*

$$f(t) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{ixt} dx.$$

Let us now sketch the proof of Shannon's theorem. By the inversion formula for the Fourier *transform*

$$f(t) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{ixt} dx.$$

Hence

$$f(n) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{ixn} dx = c_{-n}(\hat{f}),$$

the *Fourier coefficients* of \hat{f} .

Let us now sketch the proof of Shannon's theorem. By the inversion formula for the Fourier *transform*

$$f(t) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{ixt} dx.$$

Hence

$$f(n) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{inx} dx = c_{-n}(\hat{f}),$$

the *Fourier coefficients* of \hat{f} . By the theorem on *Fourier series*,

$$\hat{f}(x) = \sum c_{-n} e^{-inx},$$

so

$$f(t) = (1/2\pi) \sum f(n) \int_{-\pi}^{\pi} e^{in(x-t)} dx = (1/\pi) \sum f(n) \sin \frac{n(x-t)\pi}{n(x-t)},$$

which is the claim.

Let us now sketch the proof of Shannon's theorem. By the inversion formula for the Fourier *transform*

$$f(t) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{ixt} dx.$$

Hence

$$f(n) = (1/2\pi) \int_{-\pi}^{\pi} \hat{f}(x) e^{inx} dx = c_{-n}(\hat{f}),$$

the *Fourier coefficients* of \hat{f} . By the theorem on *Fourier series*,

$$\hat{f}(x) = \sum c_{-n} e^{-inx},$$

so

$$f(t) = (1/2\pi) \sum f(n) \int_{-\pi}^{\pi} e^{in(x-t)} dx = (1/\pi) \sum f(n) \sin \frac{n(x-t)\pi}{n(x-t)},$$

which is the claim.