

The Perron-Frobenius theorem, Google and Google-like things.

Positive matrices

The Perron-Frobenius theorem is a theorem about eigenvalues and eigenvectors of *positive* matrices. We say that a matrix,

$$P = (p_{ij})_{N \times N},$$

is positive if all $p_{ij} > 0$ (and nonnegative if all $p_{ij} \geq 0$).

Positive matrices

The Perron-Frobenius theorem is a theorem about eigenvalues and eigenvectors of *positive* matrices. We say that a matrix,

$$P = (p_{ij})_{N \times N},$$

is positive if all $p_{ij} > 0$ (and nonnegative if all $p_{ij} \geq 0$).

We say that P is *stochastic* if it is non negative and

$$\sum_i P_{ij} = 1$$

for all j .

Stochastic matrices

Think of a stochastic process (X_k) that at any time k can be in one of the states $1, 2, \dots$ or N . Such a stochastic process is called a *Markov chain* if its probability distribution at a time $k + 1$ only depends on where it is at time k .

Stochastic matrices

Think of a stochastic process (X_k) that at any time k can be in one of the states $1, 2, \dots$ or N . Such a stochastic process is called a *Markov chain* if its probability distribution at a time $k + 1$ only depends on where it is at time k .

Let p_{ij} be the probability that the process at time $k + 1$ is in state i , given that it is in state j the previous time k . Then $\sum_i p_{ij} = 1$ for all j , and all $p_{ij} \geq 0$. So, it is a stochastic matrix.

Stochastic matrices

Think of a stochastic process (X_k) that at any time k can be in one of the states $1, 2, \dots$ or N . Such a stochastic process is called a *Markov chain* if its probability distribution at a time $k + 1$ only depends on where it is at time k .

Let p_{ij} be the probability that the process at time $k + 1$ is in state i , given that it is in state j the previous time k . Then $\sum_i p_{ij} = 1$ for all j , and all $p_{ij} \geq 0$. So, it is a stochastic matrix.

Say that the probability distribution at time k is given by $x = (x_1, \dots, x_N)$, so that the probability to be in the state i at time k is x_i .

Stochastic matrices

Think of a stochastic process (X_k) that at any time k can be in one of the states $1, 2, \dots$ or N . Such a stochastic process is called a *Markov chain* if its probability distribution at a time $k + 1$ only depends on where it is at time k .

Let p_{ij} be the probability that the process at time $k + 1$ is in state i , given that it is in state j the previous time k . Then $\sum_i p_{ij} = 1$ for all j , and all $p_{ij} \geq 0$. So, it is a stochastic matrix.

Say that the probability distribution at time k is given by $x = (x_1, \dots, x_N)$, so that the probability to be in the state i at time k is x_i .

Then the probability distribution at time $k + 1$ is given by the vector Px , i. e. the probability to be in state i is $\sum_j p_{ij}x_j$.

Stochastic matrices

Think of a stochastic process (X_k) that at any time k can be in one of the states $1, 2, \dots$ or N . Such a stochastic process is called a *Markov chain* if its probability distribution at a time $k + 1$ only depends on where it is at time k .

Let p_{ij} be the probability that the process at time $k + 1$ is in state i , given that it is in state j the previous time k . Then $\sum_i p_{ij} = 1$ for all j , and all $p_{ij} \geq 0$. So, it is a stochastic matrix.

Say that the probability distribution at time k is given by $x = (x_1, \dots, x_N)$, so that the probability to be in the state i at time k is x_i .

Then the probability distribution at time $k + 1$ is given by the vector Px , i. e. the probability to be in state i is $\sum_j p_{ij}x_j$.

The probability vector x is invariant if $Px = x$, i. e. x is an eigenvector of P with eigenvalue 1.

The Perron-Frobenius Theorem

We also say that a vector $x = (x_1, \dots, x_n)$ is positive if all $x_i > 0$ and nonnegative if all $x_i \geq 0$ (like probability distributions). We now state the first part of the Perron-Frobenius theorem.

Theorem

Let P be a positive matrix. Then P has exactly one positive eigenvector x^0 . If λ_0 is its eigenvalue, then $\lambda_0 > 0$, and if P is stochastic, $\lambda_0 = 1$.

The Perron-Frobenius Theorem

We also say that a vector $x = (x_1, \dots, x_n)$ is positive if all $x_i > 0$ and nonnegative if all $x_i \geq 0$ (like probability distributions). We now state the first part of the Perron-Frobenius theorem.

Theorem

Let P be a positive matrix. Then P has exactly one positive eigenvector x^0 . If λ_0 is its eigenvalue, then $\lambda_0 > 0$, and if P is stochastic, $\lambda_0 = 1$. If $\lambda > \lambda_0$, there is no nonnegative vector x with

$$Px \geq \lambda x.$$

Sketch of proof:

Let K_+ be the positive octant, i.e. the set of all non negative vectors.
That P is positive implies that

$$P(K_+) \subset K_+.$$

Sketch of proof:

Let K_+ be the positive octant, i.e. the set of all non negative vectors. That P is positive implies that

$$P(K_+) \subset K_+.$$

We say that a subset K of K_+ is a *cone* if $x \in K$ implies $sx \in K$ for all $s > 0$.

Sketch of proof:

Let K_+ be the positive octant, i.e. the set of all non negative vectors. That P is positive implies that

$$P(K_+) \subset K_+.$$

We say that a subset K of K_+ is a *cone* if $x \in K$ implies $sx \in K$ for all $s > 0$. We say that K is an invariant cone if $PK \subset K$.

Sketch of proof:

Let K_+ be the positive octant, i.e. the set of all non negative vectors. That P is positive implies that

$$P(K_+) \subset K_+.$$

We say that a subset K of K_+ is a *cone* if $x \in K$ implies $sx \in K$ for all $s > 0$. We say that K is an invariant cone if $PK \subset K$. So, K_+ is an invariant cone for P .

We are looking for an eigenvector of P in K_+ . If such an eigenvector x_0 exists, it would define a minimal invariant cone, $\{sx^0\}$. So, we look for small invariant cones.

We are looking for an eigenvector of P in K_+ . If such an eigenvector x_0 exists, it would define a minimal invariant cone, $\{sx^0\}$. So, we look for small invariant cones.

But, if P is positive $P(K_+)$ is strictly included in K_+ . In fact, it is a relatively closed subcone.

We are looking for an eigenvector of P in K_+ . If such an eigenvector x_0 exists, it would define a minimal invariant cone, $\{sx^0\}$. So, we look for small invariant cones.

But, if P is positive $P(K_+)$ is strictly included in K_+ . In fact, it is a relatively closed subcone. Note that

$$P^{k+1}(K_+) \subset P^k(K_+).$$

We are looking for an eigenvector of P in K_+ . If such an eigenvector x_0 exists, it would define a minimal invariant cone, $\{sx^0\}$. So, we look for small invariant cones.

But, if P is positive $P(K_+)$ is strictly included in K_+ . In fact, it is a relatively closed subcone. Note that

$$P^{k+1}(K_+) \subset P^k(K_+).$$

Now look at

$$\bigcap_k P^k(K_+).$$

It is an invariant cone.

One can then show that, since $P^k(K_+)$ is strictly decreasing, its limit is just a half line, $L = \{sx^0; s > 0\}$. Since L is an invariant cone, x^0 is an eigenvector.

One can then show that, since $P^k(K_+)$ is strictly decreasing, its limit is just a half line, $L = \{sx^0; s > 0\}$. Since L is an invariant cone, x^0 is an eigenvector. This means that $Px^0 = \lambda_0 x^0$ and clearly $\lambda_0 > 0$.

Clearly, any other non negative eigenvector lies in L as well (why?), so it must be equal to (a multiple of) x^0 . This proves the first part of the theorem.

One can then show that, since $P^k(K_+)$ is strictly decreasing, its limit is just a half line, $L = \{sx^0; s > 0\}$. Since L is an invariant cone, x^0 is an eigenvector. This means that $Px^0 = \lambda_0 x^0$ and clearly $\lambda_0 > 0$.

Clearly, any other non negative eigenvector lies in L as well (why?), so it must be equal to (a multiple of) x^0 . This proves the first part of the theorem. The second, more technical, part said that:

If $\lambda > \lambda_0$, there is no nonnegative vector x with

$$Px \geq \lambda x.$$

Proof of the technical part

Let K be any closed invariant subcone of K_+ and take $x \in K$. Let $x^k = P^k x$.

Proof of the technical part

Let K be any closed invariant subcone of K_+ and take $x \in K$. Let $x^k = P^k x$. Then $x^k/|x^k| \in K$ for all k .

Proof of the technical part

Let K be any closed invariant subcone of K_+ and take $x \in K$. Let $x^k = P^k x$. Then $x^k/|x^k| \in K$ for all k . Moreover

$$x^k/|x^k| \in P^k(K_+),$$

so it must converge to $x^0/|x^0|$. So $L \subset K$ i. e. L is a subset of any closed invariant subcone.

Proof of the technical part

Let K be any closed invariant subcone of K_+ and take $x \in K$. Let $x^k = P^k x$. Then $x^k / |x^k| \in K$ for all k . Moreover

$$x^k / |x^k| \in P^k(K_+),$$

so it must converge to $x^0 / |x^0|$. So $L \subset K$ i. e. L is a subset of any closed invariant subcone. Take $\lambda > 0$ and look at

$$\{x \geq 0; Px \geq \lambda x\}.$$

It is a closed invariant subcone of K_+ . Therefore, if it is not empty, it contains L and therefore x_0 . So

$$\lambda_0 x_0 = Px_0 \geq \lambda x_0.$$

Hence, $\lambda \leq \lambda_0$.

The next part of the Perron-Frobenius theorem is

Theorem

All (complex) eigenvalues of P satisfy $|\lambda| \leq \lambda_0$. Equality holds only if the eigenvector is positive.

Bevis.

If $z = (z_1, \dots, z_n)$ is a complex eigenvector, then

$$\sum_j p_{ij} z_j = \lambda z_i.$$

The triangle inequality gives,

$$|\lambda| |z_i| \leq \sum_j p_{ij} |z_j|.$$

By the previous part, $\lambda \leq \lambda_0$. Equality can hold only if all z_j have the same argument, so z can be taken positive. □

Finally

Theorem

If P is positive and stochastic, $\lambda_0 = 1$.

Finally

Theorem

If P is positive and stochastic, $\lambda_0 = 1$.

Bevis.

Let x be the eigenvector with $\sum x_i = 1$,

$$\lambda_0 x_i = \sum p_{ij} x_j.$$

Finally

Theorem

If P is positive and stochastic, $\lambda_0 = 1$.

Bevis.

Let x be the eigenvector with $\sum x_i = 1$,

$$\lambda_0 x_i = \sum p_{ij} x_j.$$

Sum over i :

$$\lambda_0 = \lambda_0 \sum x_i = \sum_{ij} p_{ij} x_j = 1.$$



If P is stochastic,

$$x_0 = \lim_{k \rightarrow \infty} P^k x',$$

for any $x' \neq 0$.

If P is stochastic,

$$x_0 = \lim_{k \rightarrow \infty} P^k x',$$

for any $x' \neq 0$. This is at least roughly clear since all the other eigenvalues have eigenvalues smaller than 1.

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem.

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage?

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage? It is a special case of a more general problem: How do you rank (the popularity of) anything?

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage? It is a special case of a more general problem: How do you rank (the popularity of) anything? Researchers, football teams, people?

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage? It is a special case of a more general problem: How do you rank (the popularity of) anything? Researchers, football teams, people? We start with the popularity of people.

A person is popular if (s)he has many friends.

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage? It is a special case of a more general problem: How do you rank (the popularity of) anything? Researchers, football teams, people? We start with the popularity of people.

A person is popular if (s)he has many friends. But, not all friends are equal; true popularity consists in having many *popular* friends.

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage? It is a special case of a more general problem: How do you rank (the popularity of) anything? Researchers, football teams, people? We start with the popularity of people.

A person is popular if (s)he has many friends. But, not all friends are equal; true popularity consists in having many *popular* friends. This sounds like a circular definition, but can be made precise. Label all the people from 1 to N , and say the popularity of individual j is x_j . Say j is friends with N_j other people (this may be a non symmetrical relation).

Application to Google

There is also a version of the theorem for non negative matrices, but I will not go into that. We will now discuss an application of the theorem. This is the *Page rank algorithm* of Google. It deals with the problem:

How do you rank the popularity of a webpage? It is a special case of a more general problem: How do you rank (the popularity of) anything? Researchers, football teams, people? We start with the popularity of people.

A person is popular if (s)he has many friends. But, not all friends are equal; true popularity consists in having many *popular* friends. This sounds like a circular definition, but can be made precise. Label all the people from 1 to N , and say the popularity of individual j is x_j . Say j is friends with N_j other people (this may be a non symmetrical relation). Then we say that each of them inherits a fraction of j 's popularity

$$(1/N_j)x_j.$$

The total popularity of i becomes

$$x_i = \sum_j p_{ij} x_j,$$

where $p_{ij} = (1/N_j)$ if j is friends with i and 0 if not. This is a stochastic matrix since

$$\sum_i p_{ij} = (1/N_j) \#\{i; j \text{ is friends with } i\} = 1.$$

The total popularity of i becomes

$$x_i = \sum_j p_{ij} x_j,$$

where $p_{ij} = (1/N_j)$ if j is friends with i and 0 if not. This is a stochastic matrix since

$$\sum_i p_{ij} = (1/N_j) \#\{i; j \text{ is friends with } i\} = 1.$$

(There is a problem that P is not strictly positive that we will come to later.)

The total popularity of i becomes

$$x_i = \sum_j p_{ij} x_j,$$

where $p_{ij} = (1/N_j)$ if j is friends with i and 0 if not. This is a stochastic matrix since

$$\sum_i p_{ij} = (1/N_j) \#\{i; j \text{ is friends with } i\} = 1.$$

(There is a problem that P is not strictly positive that we will come to later.)

By Perron-Frobenius, it has an essentially unique solution and this is the ranking.

The total popularity of i becomes

$$x_i = \sum_j p_{ij} x_j,$$

where $p_{ij} = (1/N_j)$ if j is friends with i and 0 if not. This is a stochastic matrix since

$$\sum_i p_{ij} = (1/N_j) \#\{i; j \text{ is friends with } i\} = 1.$$

(There is a problem that P is not strictly positive that we will come to later.)

By Perron-Frobenius, it has an essentially unique solution and this is the ranking.

The same argument applies to internet pages. Then we replace people by internet pages and the friendship relation is : *there is a link from j to i* . In this case, we can interpret the ranking differently.

Imagine a person that is surfing the internet. At any page, (s)he clicks on a random (uniformly distributed) link from that page to the next.

Imagine a person that is surfing the internet. At any page, (s)he clicks on a random (uniformly distributed) link from that page to the next.

Given that at one time he is in position j , the probability that in the next time he is in position i is $1/N_j$ or 0, i. e. p_{ij} . Then we get a Markov chain and

$$x^0 = \lim_{k \rightarrow \infty} P^k x',$$

where $x' \neq 0$ is an arbitrary starting point.

Imagine a person that is surfing the internet. At any page, (s)he clicks on a random (uniformly distributed) link from that page to the next.

Given that at one time he is in position j , the probability that in the next time he is in position i is $1/N_j$ or 0, i. e. p_{ij} . Then we get a Markov chain and

$$x^0 = \lim_{k \rightarrow \infty} P^k x',$$

where $x' \neq 0$ is an arbitrary starting point. So x_i^0 is the probability that our surfer is at page i after a long time, or the proportion of the time he is there.

In practice, one makes P strictly positive by replacing it by

$$P^\beta := (1 - \beta)P + \beta Q,$$

where Q is some arbitrary positive stochastic matrix (like $q_{ij} = 1/N$ for all i, j .)

In practice, one makes P strictly positive by replacing it by

$$P^\beta := (1 - \beta)P + \beta Q,$$

where Q is some arbitrary positive stochastic matrix (like $q_{ij} = 1/N$ for all i, j .)

Google updates its page rank monthly. This involves computing an eigenvector of an $N \times N$ where N is the number of web pages in the world.