Predicting Void ELF and Mapping Connectivity for Electride Screening

Electron localization function (ELF) prediction with an atom–void graph network, and exact geometric void connectivity

Andrew Weiland · [Advisor name] · [Department], Purdue University · [email]


LOGO

QR CODE

BACKGROUND


What is an electride?

In an electride, extra electrons sit in the empty space between atoms and act as anions. Whether those regions are isolated (0D) or connected (1D, 2D) shapes the material's properties.




Connectivity defines dimensionality in electrides

0D Isolated pockets




Mayenite-type $Ca_6Al_7O_{16}$: separate cages

1D Channels



Y_5Si_3: channels along one direction

2D Sheets



Ca_2N: layers between the atoms


GOAL screen crystals for electrides from structure alone, without DFT on every candidate. We built two void-centered tools and tested each one separately.

TOOL 1 · PREDICT ELF


Can a graph network predict ELF at each candidate void from the surrounding atoms?


Test-set performance	0.82	0.059	88,680
	R^2	RMSE (ELF)	test voids from 5,000 crystals

ELF at each void from nearby atoms




One crystal → many voids → one ELF estimate per void






TOOL 2 · MAP HOW VOIDS CONNECT

Can geometry alone recover the dimensionality of known electrides?




Predicted vs actual void connectivity



PROPOSED COMBINATION · FUTURE WORK

Geometry alone can't predict ELF connectivity

TOOL 2 · GEOMETRY	►	TOOL 1 · GNN	►	COMBINED
Find the channel and its bottleneck		Predict ELF at that bottleneck		Bottleneck ELF decides 1D vs. 0D



Connected electron-rich region

High ELF at the bottleneck
→ electrons connect (1D)

Separate electron-rich pockets

Low ELF at the bottleneck
→ isolated pockets (0D)

TAKEAWAY

- **Geometry shows where a path could exist**
- **ELF shows whether electrons occupy it**
- **Together they separate structural from electronic dimensionality**

NEXT STEPS

- Predict ELF at bottlenecks, not just voids
- Validate against DFT on known electrides
- Combine both into one electride screen

References [1] Becke & Edgecombe, J. Chem. Phys. 92, 5397 (1990), ELF. [2] Ellis, Ellis & Miao, J. Mater. Inf. 6, 45 (2026), doi:10.20517/jmi.2026.44, ELFNet. [3] Registry labels: Magnetic electrides, J. Am. Chem. Soc. (2023), doi:10.1021/jacs.3c00284; Zhu, Frolov & Choudhary, Matter 1, 1293 (2019); Burton et al., Chem. Mater. 30, 7521 (2018). [4] Methods: Shannon, Acta Cryst. A32, 751 (1976), ionic radii; Willems et al., Micropor. Mesopor. Mater. 149, 134 (2012), Zeo++; Gilmer et al., ICML (2017), message passing. **Acknowledgements:** [advisor, funding]; Purdue RCAC (Negishi).