

MAT 653: Statistical Simulation

Instructor: Dr. Wei Li

Scribe: Jiangyu Yu

Sep 14th, 2021

Singular value decomposition

A , a $m \times n$ matrix, $A^T A$ is a symmetric matrix. Let $\{v_1, \dots, v_n\}$ consists of all orthonormal eigenvectors of $A^T A$. Let $\lambda_1, \dots, \lambda_n$ the associated eigenvalues, $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$.

We have $\|Av_i\|^2 = \lambda_i$. Since

$$A^T Av_i = \lambda_i v_i \Rightarrow v_i^T A^T Av_i = \lambda_i v_i^T v_i \lambda_i$$

Denote $\sigma_i = \sqrt{\lambda_i}$, λ_i is the i th eigenvalue of $A^T A$.

Fact: rank of A is equal to the number of positive singular value of A .

SVD: Let A be a rank r matrix. There exists a matrix $\Sigma_{m \times n}$ with diagonal entries in D are the first r singular value. That is

$$\Sigma_{m \times n} = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix}$$
$$D = \begin{bmatrix} \sigma_1 & & \\ & \ddots & \\ & & \sigma_n \end{bmatrix}$$

The decomposition of A is:

$$A = U_{m \times n} \Sigma_{m \times n} V_{n \times n}^T$$

Where U and V are both orthogonal matrices. V consists of all the eigenvectors of $A^T A$. U consists the column u_i as

$$u_i = \frac{Av_i}{\|Av_i\|} = \frac{Av_i}{\sigma_i}, 1 \leq i \leq r$$

Those $\{u_i\}_1^r$ can extend to $\{u_i\}_1^m$ as the orthonormal basis.

Matrix U and V have the property:

$$Col(U) = Col(A), Col(V) = Row(A)$$

Reduced SVD

For those $A_{m \times n}$, $U_{m \times m}$, $V_{n \times n}$, $\Sigma_{m \times n}$ above, we have the partition:

$$U = [U_r, U_{m-r}], V = [V_r, V_{n-r}], \Sigma = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix}$$

Consequently, the SVD of A can be represented as:

$$A = \begin{bmatrix} U_r & U_{m-r} \end{bmatrix} \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} V_r^T \\ V_{n-r}^T \end{bmatrix} = U_r D V_r^T$$

Application

1. Linear Least Squares

We can use SVD to solve the Least Squares Problems as following:

$$\begin{aligned} Ax &= b, \\ A^T Ax &= A^T b, \\ (UDV^T)^T (UDV^T)x &= (UDV^T)^T b, \\ \Rightarrow VD^2Dx &= VDU^T b, \\ \Rightarrow D^2V^T x &= DU^T b, \\ \Rightarrow DV^T x &= U^T b. \end{aligned}$$

Denote $w = V^T x, y = U^T b$, we have the following algorithm:

Algorithm

1. Find the SVD of $A = UDV^T$;
2. Compute $y = U^T b$;
3. Solve $w^* = D^{-1}y$;
4. $V^T x = w^* \Rightarrow x = Vw^*$;

The solution of the equation is $x = V_r D^{-1} U_r^T b$, and we can notice that this SVD method allows A,b to be arbitrary. Notice that $V_r D^{-1} U_r^T$ is the inverse of A. Here we have the concept of generalized inverse.

Moore-Penrose inverse: $A^+ = V_r D^{-1} U_r^T$

2.LS-Problem

In LS Problem

$$\min_x \|Ax - b\|$$

,
where A and b are not restricted at all, we have:

$$A^T Ax = A^T b.$$

Let \mathcal{L} is the set of all the minimizers to the LS problem. We have following facts.

Fact 1 $x^* = A^+b \in \mathfrak{L}$.

Fact 2 $A\tilde{x}_1 = A\tilde{x}_2$, for any $\tilde{x}_1, \tilde{x}_2 \in \mathfrak{L}$.

Fact 3 For the optimization problem $\min_{x \in \mathfrak{L}} \|x\|$, there is a unique solution $x^* = A^+b$.

Fact 4 We already have the result that if we choose some λ , LS problem will has a unique solution to the “modified” normal equation:

$$(A^T A + \lambda I)x = A^T b$$

that is,

$$\hat{x} = (A^T A + \lambda I)^{-1} A^T b$$

In fact, let $\lambda \rightarrow 0$, we have

$$(A^T A + \lambda I)^{-1} A^T \rightarrow A^+$$

Fact 5 The projection of b on $\text{Col}(A)$ is given by $A(A^T A)^{-1} A^T b$ (assuming $A^T A$ is invertible), here is a more general result

$$\hat{b} = AA^+b = \lim_{\lambda \rightarrow 0} [A(A^T A + \lambda I)^{-1} A^T]b$$