

Undetermined Coefficients

Department of Mathematics

Salt Lake Community College

(Slides by Adam Wilson)

Remember

If L is a linear differential operator defined by

$$L(y) = a_n(t)y^{(n)} + a_{n-1}(t)y^{(n-1)} + \cdots + a_1(t)y' + a_0(t)y$$

(where all functions of t are assumed to be defined over some interval I) then we can look at superposition for the DE $L(y) = f(t)$.

Remember

If L is a linear differential operator defined by

$$L(y) = a_n(t)y^{(n)} + a_{n-1}(t)y^{(n-1)} + \cdots + a_1(t)y' + a_0(t)y$$

(where all functions of t are assumed to be defined over some interval I) then we can look at superposition for the DE $L(y) = f(t)$.

Superposition Principle for Nonhomogeneous Linear DEs

If $y_i(t)$ is a solution of $L(y) = f_i(t)$, for $i = 1, 2, \dots, n$, and constants $c_1, c_2, \dots, c_n \in \mathbb{R}$, then

$$y(t) = c_1y_1(t) + c_2y_2(t) + \cdots + c_ny_n(t)$$

is a solution of

$$L(y) = c_1f_1(t) + c_2f_2(t) + \cdots + c_nf_n(t)$$

Nonhomogeneous Principle for Linear DEs

The general solution of the nonhomogeneous linear DE $L(y) = f$ is

$$y = y_h + y_p$$

where

- y_h is the general solution of $L(y) = 0$
- y_p is a particular solution of $L(y) = f$

Nonhomogeneous Principle for Linear DEs

The general solution of the nonhomogeneous linear DE $L(y) = f$ is

$$y = y_h + y_p$$

where

- y_h is the general solution of $L(y) = 0$
- y_p is a particular solution of $L(y) = f$

Note

This is just applying the superposition principle for $f_1(t) = 0$ and $f_2(t) = f$.

Example 1

Consider the nonhomogeneous second-order DE

$$y'' - y' - 2y = 2t + 1 - 2e^t$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$y_1 = -t \quad \text{is a solution to} \quad L(y) = f_1$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$\begin{array}{ll} y_1 = -t & \text{is a solution to } L(y) = f_1 \\ y_2 = e^t & \text{is a solution to } L(y) = f_2. \end{array}$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$\begin{aligned} y_1 = -t & \text{ is a solution to } L(y) = f_1 \\ y_2 = e^t & \text{ is a solution to } L(y) = f_2. \end{aligned}$$

We can then use superposition to build a particular solution

$$y_p = y_1 + y_2 = -t + e^t$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$\begin{aligned} y_1 = -t & \text{ is a solution to } L(y) = f_1 \\ y_2 = e^t & \text{ is a solution to } L(y) = f_2. \end{aligned}$$

We can then use superposition to build a particular solution

$$y_p = y_1 + y_2 = -t + e^t$$

Finally, we use characteristic roots to solve $L(y) = 0$

$$r^2 - r - 2 = 0$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$\begin{aligned} y_1 = -t & \text{ is a solution to } L(y) = f_1 \\ y_2 = e^t & \text{ is a solution to } L(y) = f_2. \end{aligned}$$

We can then use superposition to build a particular solution

$$y_p = y_1 + y_2 = -t + e^t$$

Finally, we use characteristic roots to solve $L(y) = 0$

$$r^2 - r - 2 = 0 \rightarrow r_1 = 2, r_2 = -1$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$\begin{array}{ll} y_1 = -t & \text{is a solution to } L(y) = f_1 \\ y_2 = e^t & \text{is a solution to } L(y) = f_2. \end{array}$$

We can then use superposition to build a particular solution

$$y_p = y_1 + y_2 = -t + e^t$$

Finally, we use characteristic roots to solve $L(y) = 0$

$$r^2 - r - 2 = 0 \rightarrow r_1 = 2, r_2 = -1 \rightarrow y_h = c_1 e^{2t} + c_2 e^{-t}$$

Example 1

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{2t + 1}_{f_1} \underbrace{- 2e^t}_{f_2}$$

We can verify the following following:

$$\begin{array}{ll} y_1 = -t & \text{is a solution to } L(y) = f_1 \\ y_2 = e^t & \text{is a solution to } L(y) = f_2. \end{array}$$

We can then use superposition to build a particular solution

$$y_p = y_1 + y_2 = -t + e^t$$

Finally, we use characteristic roots to solve $L(y) = 0$

$$r^2 - r - 2 = 0 \rightarrow r_1 = 2, r_2 = -1 \rightarrow y_h = c_1 e^{2t} + c_2 e^{-t}$$

Thus, the general solution is

$$y = y_h + y_p = c_1 e^{2t} + c_2 e^{-t} - t + e^t$$

Example 2

Consider the nonhomogeneous second-order DE

$$y'' - y' - 2y = t + \frac{1}{2} + 8e^t$$

Example 2

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{t + \frac{1}{2}}_{\frac{1}{2}f_1} + \underbrace{8e^t}_{-4f_2}$$

Example 2

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{t + \frac{1}{2}}_{\frac{1}{2}f_1} + \underbrace{8e^t}_{-4f_2}$$

Using the solutions found in the last example, we can use superposition to build a particular solution to this DE.

$$y_p = \frac{1}{2}y_1 - 4y_2$$

Example 2

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{t + \frac{1}{2}}_{\frac{1}{2}f_1} + \underbrace{8e^t}_{-4f_2}$$

Using the solutions found in the last example, we can use superposition to build a particular solution to this DE.

$$y_p = \frac{1}{2}y_1 - 4y_2 = -\frac{1}{2}t - 4e^t$$

Example 2

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{t + \frac{1}{2}}_{\frac{1}{2}f_1} + \underbrace{8e^t}_{-4f_2}$$

Using the solutions found in the last example, we can use superposition to build a particular solution to this DE.

$$y_p = \frac{1}{2}y_1 - 4y_2 = -\frac{1}{2}t - 4e^t$$

Finally, we have already solved $L(y) = 0$. So, the general solution is

$$y = y_h + y_p = c_1e^{2t} + c_2e^{-t} - \frac{1}{2}t - 4e^t$$

Example 2

Consider the nonhomogeneous second-order DE

$$\underbrace{y'' - y' - 2y}_{L(y)} = \underbrace{t + \frac{1}{2}}_{\frac{1}{2}f_1} + \underbrace{8e^t}_{-4f_2}$$

Using the solutions found in the last example, we can use superposition to build a particular solution to this DE.

$$y_p = \frac{1}{2}y_1 - 4y_2 = -\frac{1}{2}t - 4e^t$$

Finally, we have already solved $L(y) = 0$. So, the general solution is

$$y = y_h + y_p = c_1e^{2t} + c_2e^{-t} - \frac{1}{2}t - 4e^t$$

Note

After accumulating some experience, a solution can be guessed by just “inspecting” the equation. By recognizing the patterns.

Example 3

Consider the second-order DE

$$ay'' + by' + cy = d$$

where all the coefficients and forcing term are constant.

Example 3

Consider the second-order DE

$$ay'' + by' + cy = d$$

where all the coefficients and forcing term are constant.

We can see that, when $c \neq 0$, $y_p = \frac{d}{c}$ is a particular solution.

Example 3

Consider the second-order DE

$$ay'' + by' + cy = d$$

where all the coefficients and forcing term are constant.

We can see that, when $c \neq 0$, $y_p = \frac{d}{c}$ is a particular solution.

Note

This idea works well for the n th-order equation

$$a_n(t)y^{(n)} + a_{n-1}(t)y^{(n-1)} + \cdots + a_1(t)y' + a_0(t)y = d$$

provided that $a_0 \neq 0$.

Example 4

Inspection of

$$y'' + y' = 1$$

Example 4

Inspection of

$$y'' + y' = 1$$

leads to the solution $y_p = t$.

Example 4

Inspection of

$$y'' + y' = 1$$

leads to the solution $y_p = t$.

Example 5

Inspection of

$$y'' - y = \sin(t)$$

Example 4

Inspection of

$$y'' + y' = 1$$

leads to the solution $y_p = t$.

Example 5

Inspection of

$$y'' - y = \sin(t)$$

leads to the solution $y_p = -\frac{1}{2} \sin(t)$

Example 4

Inspection of

$$y'' + y' = 1$$

leads to the solution $y_p = t$.

Example 5

Inspection of

$$y'' - y = \sin(t)$$

leads to the solution $y_p = -\frac{1}{2} \sin(t)$

Example 6

Inspection of

$$y'' + y' - 3y = 9e^{3t}$$

Example 4

Inspection of

$$y'' + y' = 1$$

leads to the solution $y_p = t$.

Example 5

Inspection of

$$y'' - y = \sin(t)$$

leads to the solution $y_p = -\frac{1}{2} \sin(t)$

Example 6

Inspection of

$$y'' + y' - 3y = 9e^{3t}$$

leads to the solution $y_p = e^{3t}$

Note

There are a few limitations of this method:

It only works for linear differential equations with specific forcing terms.

Note

There are a few limitations of this method:

It only works for linear differential equations with specific forcing terms.

Forcing Terms That Work With Undetermined Coefficients

Any finite products or sums of:

- Polynomials in t .
- Exponentials e^{at} .
- Sinusoidal functions of the form $\cos(kt)$ and $\sin(kt)$.

Note

There are a few limitations of this method:

It only works for linear differential equations with specific forcing terms.

Forcing Terms That Work With Undetermined Coefficients

Any finite products or sums of:

- Polynomials in t .
- Exponentials e^{at} .
- Sinusoidal functions of the form $\cos(kt)$ and $\sin(kt)$.

Note

Even with these limitations, undetermined coefficients is widely used, given that many functions are built from the above parts.

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Let us look for y_p in \mathbb{P}_2 . Which means y_p will be of the form

$$y_p = At^2 + Bt + C$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Let us look for y_p in \mathbb{P}_2 . Which means y_p will be of the form

$$y_p = At^2 + Bt + C$$

We can then calculate:

$$y'_p = 2At + B$$

$$y''_p = 2A$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Plugging these into the DE gives

$$2A - (2At + B) - 2(At^2 + Bt + C) = 3t^2 - 1$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Plugging these into the DE gives

$$2A - (2At + B) - 2(At^2 + Bt + C) = 3t^2 - 1$$

$$(-2A)t^2 + (-2A - 2B)t + (2A - B - 2C) = 3t^2 - 1$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Plugging these into the DE gives

$$\begin{aligned} 2A - (2At + B) - 2(At^2 + Bt + C) &= 3t^2 - 1 \\ (-2A)t^2 + (-2A - 2B)t + (2A - B - 2C) &= 3t^2 - 1 \end{aligned}$$

So, equating both sides gives the system

$$-2A = 3, \quad -2A - 2B = 0, \quad 2A - B - 2C = -1$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Plugging these into the DE gives

$$\begin{aligned} 2A - (2At + B) - 2(At^2 + Bt + C) &= 3t^2 - 1 \\ (-2A)t^2 + (-2A - 2B)t + (2A - B - 2C) &= 3t^2 - 1 \end{aligned}$$

So, equating both sides gives the system

$$-2A = 3, \quad -2A - 2B = 0, \quad 2A - B - 2C = -1$$

Which has solution $A = -\frac{3}{2}$, $B = \frac{3}{2}$, and $C = -\frac{7}{4}$.

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Thus, the particular solution is

$$y_p = -\frac{3}{2}t^2 + \frac{3}{2}t + \frac{7}{4}$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Thus, the particular solution is

$$y_p = -\frac{3}{2}t^2 + \frac{3}{2}t + \frac{7}{4}$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

Example 7

Consider

$$y'' - y' - 2y = 3t^2 - 1$$

Thus, the particular solution is

$$y_p = -\frac{3}{2}t^2 + \frac{3}{2}t + \frac{7}{4}$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

The general solution is

$$y = c_1 e^{2t} + c_2 e^{-t} - \frac{3}{2}t^2 + \frac{3}{2}t + \frac{7}{4}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Let us look for y_p of the form

$$y_p = Ae^{-3t}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Let us look for y_p of the form

$$y_p = Ae^{-3t}$$

We can then calculate:

$$y'_p = -3Ae^{-3t}$$

$$y''_p = 9Ae^{-3t}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Plugging these into the DE gives

$$9Ae^{-3t} + 3Ae^{-3t} - 2Ae^{-3t} = 2e^{-3t}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Plugging these into the DE gives

$$9Ae^{-3t} + 3Ae^{-3t} - 2Ae^{-3t} = 2e^{-3t}$$

$$10Ae^{-3t} = 2e^{-3t}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Plugging these into the DE gives

$$9Ae^{-3t} + 3Ae^{-3t} - 2Ae^{-3t} = 2e^{-3t}$$

$$10Ae^{-3t} = 2e^{-3t}$$

So, equating both sides gives

$$10A = 2 \quad \rightarrow \quad A = \frac{1}{5}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Thus, the particular solution is

$$y_p = \frac{1}{5}e^{-3t}$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Thus, the particular solution is

$$y_p = \frac{1}{5}e^{-3t}$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

Example 8

Consider

$$y'' - y' - 2y = 2e^{-3t}$$

Thus, the particular solution is

$$y_p = \frac{1}{5}e^{-3t}$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

The general solution is

$$y = c_1 e^{2t} + c_2 e^{-t} + \frac{1}{5} e^{-3t}$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Let us look for y_p of the form

$$y_p = A \cos(3t) + B \sin(3t)$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Let us look for y_p of the form

$$y_p = A \cos(3t) + B \sin(3t)$$

We can then calculate:

$$y_p' = -3A \sin(3t) + 3B \cos(3t)$$

$$y_p'' = -9A \cos(3t) - 9B \sin(3t)$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Plugging these into the DE gives

$$\begin{aligned} &(-9A \cos(3t) - 9 \sin(3t)) \\ &\quad - (-3A \sin(3t) + 3B \cos(3t)) \\ &\quad - 2(A \cos(3t) + B \sin(3t)) = 2 \cos(3t) \end{aligned}$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Plugging these into the DE gives

$$\begin{aligned} &(-9A \cos(3t) - 9 \sin(3t)) \\ &\quad - (-3A \sin(3t) + 3B \cos(3t)) \\ &\quad - 2(A \cos(3t) + B \sin(3t)) = 2 \cos(3t) \\ &(-11A - 3B) \cos(3t) + (3A - 11B) \sin(3t) = 2 \cos(3t) \end{aligned}$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Plugging these into the DE gives

$$\begin{aligned} &(-9A \cos(3t) - 9 \sin(3t)) \\ &\quad - (-3A \sin(3t) + 3B \cos(3t)) \\ &\quad - 2(A \cos(3t) + B \sin(3t)) = 2 \cos(3t) \\ &(-11A - 3B) \cos(3t) + (3A - 11B) \sin(3t) = 2 \cos(3t) \end{aligned}$$

So, equating both sides gives the system

$$-11A - 3B = 2, \quad 3A - 11B = 0$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Plugging these into the DE gives

$$\begin{aligned} &(-9A \cos(3t) - 9 \sin(3t)) \\ &\quad - (-3A \sin(3t) + 3B \cos(3t)) \\ &\quad - 2(A \cos(3t) + B \sin(3t)) = 2 \cos(3t) \\ &(-11A - 3B) \cos(3t) + (3A - 11B) \sin(3t) = 2 \cos(3t) \end{aligned}$$

So, equating both sides gives the system

$$-11A - 3B = 2, \quad 3A - 11B = 0$$

Which has solution $A = -\frac{11}{65}$ and $B = -\frac{3}{65}$.

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Thus, the particular solution is

$$y_p = -\frac{11}{65} \cos(3t) - \frac{3}{65} \sin(3t)$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Thus, the particular solution is

$$y_p = -\frac{11}{65} \cos(3t) - \frac{3}{65} \sin(3t)$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

Example 9

Consider

$$y'' - y' - 2y = 2 \cos(3t)$$

Thus, the particular solution is

$$y_p = -\frac{11}{65} \cos(3t) - \frac{3}{65} \sin(3t)$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

The general solution is

$$y = c_1 e^{2t} + c_2 e^{-t} - \frac{11}{65} \cos(3t) - \frac{3}{65} \sin(3t)$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Let us look for y_p of the form

$$y_p = (At^2 + Bt + C) e^t$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Let us look for y_p of the form

$$y_p = (At^2 + Bt + C) e^t$$

We can then calculate:

$$y'_p = (At^2 + (2A + B)t + (B + C)) e^t$$

$$y''_p = (At^2 + (4A + B)t + (2A + 2B + C)) e^t$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Plugging these into the DE gives

$$\begin{aligned} & (At^2 + (4A + B)t + (2A + 2B + C)) e^t \\ & - (At^2 + (2A + B)t + (B + C)) e^t \\ & + 2(At^2 + Bt + C) e^t = t^2 e^t \end{aligned}$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Plugging these into the DE gives

$$\begin{aligned} & (At^2 + (4A + B)t + (2A + 2B + C)) e^t \\ & - (At^2 + (2A + B)t + (B + C)) e^t \\ & + 2 (At^2 + Bt + C) e^t = t^2 e^t \\ & ((-2A)t^2 + (2A - 2B)t + (2A + B - 2C)) e^t = t^2 e^t \end{aligned}$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Plugging these into the DE gives

$$\begin{aligned} & (At^2 + (4A + B)t + (2A + 2B + C)) e^t \\ & - (At^2 + (2A + B)t + (B + C)) e^t \\ & + 2 (At^2 + Bt + C) e^t = t^2 e^t \\ & ((-2A)t^2 + (2A - 2B)t + (2A + B - 2C)) e^t = t^2 e^t \end{aligned}$$

So, equating both sides gives the system

$$-2A = 1, \quad 2A - 2B = 0, \quad 2A + B - 2C = 0$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Plugging these into the DE gives

$$\begin{aligned} & (At^2 + (4A + B)t + (2A + 2B + C)) e^t \\ & - (At^2 + (2A + B)t + (B + C)) e^t \\ & + 2(At^2 + Bt + C) e^t = t^2 e^t \\ & ((-2A)t^2 + (2A - 2B)t + (2A + B - 2C)) e^t = t^2 e^t \end{aligned}$$

So, equating both sides gives the system

$$-2A = 1, \quad 2A - 2B = 0, \quad 2A + B - 2C = 0$$

Which has solution $A = -\frac{1}{2}$, $B = -\frac{1}{2}$, and $C = -\frac{3}{4}$.

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Thus, the particular solution is

$$y_p = \left(-\frac{1}{2}t^2 - \frac{1}{2}t - \frac{3}{4} \right) e^t$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Thus, the particular solution is

$$y_p = \left(-\frac{1}{2}t^2 - \frac{1}{2}t - \frac{3}{4} \right) e^t$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

Example 10

Consider

$$y'' - y' - 2y = t^2 e^t$$

Thus, the particular solution is

$$y_p = \left(-\frac{1}{2}t^2 - \frac{1}{2}t - \frac{3}{4} \right) e^t$$

Since the homogeneous equation has characteristic equation

$$r^2 - r - 2 = (r - 2)(r + 1) = 0$$

The general solution is

$$y = c_1 e^{2t} + c_2 e^{-t} + \left(-\frac{1}{2}t^2 - \frac{1}{2}t - \frac{3}{4} \right) e^t$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ae^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ae^{2t}$$

We can then calculate:

$$y_p' = 2Ae^{2t}$$

$$y_p'' = 4Ae^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ae^{2t}$$

We can then calculate:

$$y_p' = 2Ae^{2t}$$

$$y_p'' = 4Ae^{2t}$$

Substituting into the DE gives

$$4Ae^{2t} - 2Ae^{2t} - 2Ae^{2t} = 5e^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ae^{2t}$$

We can then calculate:

$$y_p' = 2Ae^{2t}$$

$$y_p'' = 4Ae^{2t}$$

Substituting into the DE gives

$$4Ae^{2t} - 2Ae^{2t} - 2Ae^{2t} = 5e^{2t}$$

$$0 = 5e^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ae^{2t}$$

We can then calculate:

$$y_p' = 2Ae^{2t}$$

$$y_p'' = 4Ae^{2t}$$

Substituting into the DE gives

$$4Ae^{2t} - 2Ae^{2t} - 2Ae^{2t} = 5e^{2t}$$

$$0 = 5e^{2t}$$

That's not good. We'll have to try something else.

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ate^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Let us look for y_p of the form

$$y_p = Ate^{2t}$$

We can then calculate:

$$y_p' = (2At + A)e^{2t}$$

$$y_p'' = (4A + 4A)e^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Substituting into the DE gives

$$(4A + 4A)e^{2t} - 2Ae^{2t} - 2Ate^{2t} = 5e^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Substituting into the DE gives

$$(4A + 4A)e^{2t} - 2Ae^{2t} - 2Ate^{2t} = 5e^{2t}$$

$$3Ae^{2t} = 5e^{2t}$$

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Substituting into the DE gives

$$(4A + 4A)e^{2t} - 2Ae^{2t} - 2Ate^{2t} = 5e^{2t}$$

$$3Ae^{2t} = 5e^{2t}$$

When we equate both sides we get $3A = 5$ and so $A = \frac{5}{3}$.

Example 11

Consider

$$y'' - y' - 2y = 5e^{2t}$$

Substituting into the DE gives

$$\begin{aligned}(4A + 4A)e^{2t} - 2Ae^{2t} - 2Ate^{2t} &= 5e^{2t} \\ 3Ae^{2t} &= 5e^{2t}\end{aligned}$$

When we equate both sides we get $3A = 5$ and so $A = \frac{5}{3}$.

And so, the particular solution is

$$y_p = \frac{5}{3}te^{2t}$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ae^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ae^t$$

We can then calculate:

$$y_p' = Ae^t$$

$$y_p'' = Ae^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ae^t$$

We can then calculate:

$$y_p' = Ae^t$$

$$y_p'' = Ae^t$$

Substituting into the DE gives

$$Ae^t - 2Ae^t + Ae^{2t} = 3e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ae^t$$

We can then calculate:

$$y_p' = Ae^t$$

$$y_p'' = Ae^t$$

Substituting into the DE gives

$$Ae^t - 2Ae^t + Ae^{2t} = 3e^t$$

$$0 = 3e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ae^t$$

We can then calculate:

$$y_p' = Ae^t$$

$$y_p'' = Ae^t$$

Substituting into the DE gives

$$Ae^t - 2Ae^t + Ae^{2t} = 3e^t$$

$$0 = 3e^t$$

Thats not good. We'll have to try something else.

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ate^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ate^t$$

We can then calculate:

$$y_p' = Ae^t + Ate^t$$

$$y_p'' = 2Ae^t + Ate^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ate^t$$

We can then calculate:

$$y_p' = Ae^t + Ate^t$$

$$y_p'' = 2Ae^t + Ate^t$$

Substituting into the DE gives

$$2Ae^t + Ate^t - 2(Ae^t + Ate^t) + Ate^t = 3e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ate^t$$

We can then calculate:

$$y_p' = Ae^t + Ate^t$$

$$y_p'' = 2Ae^t + Ate^t$$

Substituting into the DE gives

$$2Ae^t + Ate^t - 2(Ae^t + Ate^t) + Ate^t = 3e^t$$

$$0 = 3e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = Ate^t$$

We can then calculate:

$$y_p' = Ae^t + Ate^t$$

$$y_p'' = 2Ae^t + Ate^t$$

Substituting into the DE gives

$$2Ae^t + Ate^t - 2(Ae^t + Ate^t) + Ate^t = 3e^t$$

$$0 = 3e^t$$

This too is a problem. We'll have to try something else.

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = At^2e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Let us look for y_p of the form

$$y_p = At^2e^t$$

We can then calculate:

$$y'_p = 2Ate^t + At^2e^t$$

$$y''_p = 2Ae^t + 4Ate^t + At^2e^t$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Substituting into the DE gives

$$2Ae^t + 4Ate^t + At^2e^t - 2(2Ate^t + At^2e^t) + At^2e^t = 5e^{2t}$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Substituting into the DE gives

$$2Ae^t + 4Ate^t + At^2e^t - 2(2Ate^t + At^2e^t) + At^2e^t = 5e^{2t}$$
$$2Ae^t = 5e^{2t}$$

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Substituting into the DE gives

$$2Ae^t + 4Ate^t + At^2e^t - 2(2Ate^t + At^2e^t) + At^2e^t = 5e^{2t}$$
$$2Ae^t = 5e^{2t}$$

When we equate both sides we get $2A = 5$ and so $A = \frac{5}{2}$.

Example 12

Consider

$$y'' - 2y' + y = 3e^t$$

Substituting into the DE gives

$$2Ae^t + 4Ate^t + At^2e^t - 2(2Ate^t + At^2e^t) + At^2e^t = 5e^{2t}$$
$$2Ae^t = 5e^{2t}$$

When we equate both sides we get $2A = 5$ and so $A = \frac{5}{2}$.

And so, the particular solution is

$$y_p = \frac{5}{2}te^{2t}$$