

# Separable Equations

Colby Community College

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\frac{dy}{dt} = 3t^2 \cdot (1 + y)$$

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\begin{aligned}\frac{dy}{dt} &= 3t^2 \cdot (1 + y) \\ \frac{dy}{1 + y} &= 3t^2 dt\end{aligned}$$

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\begin{aligned}\frac{dy}{dt} &= 3t^2 \cdot (1 + y) \\ \int \frac{dy}{1 + y} &= \int 3t^2 dt\end{aligned}$$

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\frac{dy}{dt} = 3t^2 \cdot (1 + y)$$

$$\int \frac{dy}{1 + y} = \int 3t^2 dt$$

$$\ln |1 + y| = t^3 + c$$

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\frac{dy}{dt} = 3t^2 \cdot (1 + y)$$

$$\int \frac{dy}{1 + y} = \int 3t^2 dt$$

$$\ln |1 + y| = t^3 + c$$

$$|1 + y| = e^c e^{t^3}$$

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\frac{dy}{dt} = 3t^2 \cdot (1 + y)$$

$$\int \frac{dy}{1 + y} = \int 3t^2 dt$$

$$\ln |1 + y| = t^3 + c$$

$$|1 + y| = e^c e^{t^3}$$

$$y = -1 \pm e^c e^{t^3}$$



## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\frac{dy}{dt} = 3t^2 \cdot (1 + y)$$

$$\int \frac{dy}{1 + y} = \int 3t^2 dt$$

$$\ln |1 + y| = t^3 + c$$

$$|1 + y| = e^c e^{t^3}$$

$$y = -1 \pm e^c e^{t^3}$$

$$y = -1 + ke^{t^3}, \quad k \in \mathbb{R}$$

## Separable Differential Equations

A **separable** differential equation is one that can be written  $y' = f(t)g(y)$ . Constant solutions  $y = c$  can be found by solving  $g(y) = 0$ .

### Example 1

Let us consider

$$\begin{aligned}\frac{dy}{dt} &= 3t^2 \cdot (1 + y) \\ \int \frac{dy}{1 + y} &= \int 3t^2 dt \\ \ln |1 + y| &= t^3 + c \\ |1 + y| &= e^c e^{t^3} \\ y &= -1 \pm e^c e^{t^3} \\ y &= -1 + ke^{t^3}, \quad k \in \mathbb{R}\end{aligned}$$

The method we just used is called **Separation of Variables**.

## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

Suppose also that  $y = y(t)$  is a function defined **implicitly** by

$$G(y) = F(t) + c$$

on some appropriate  $t$ -interval for some real constant  $c$ .

## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

Suppose also that  $y = y(t)$  is a function defined **implicitly** by

$$G(y) = F(t) + c \quad \Rightarrow \quad G(y(t)) = F(t) + c$$

on some appropriate  $t$ -interval for some real constant  $c$ .

## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

Suppose also that  $y = y(t)$  is a function defined **implicitly** by

$$G(y) = F(t) + c \quad \Rightarrow \quad G(y(t)) = F(t) + c$$

on some appropriate  $t$ -interval for some real constant  $c$ .

Then, if  $y$  is differentiable, we can differentiate both sides with respect to  $t$ .

$$G'(y(t))y'(t) = F'(t)$$

## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

Suppose also that  $y = y(t)$  is a function defined **implicitly** by

$$G(y) = F(t) + c \quad \Rightarrow \quad G(y(t)) = F(t) + c$$

on some appropriate  $t$ -interval for some real constant  $c$ .

Then, if  $y$  is differentiable, we can differentiate both sides with respect to  $t$ .

$$G'(y(t))y'(t) = F'(t)$$

$$\frac{y'(t)}{g(y(t))} = f(t)$$

## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

Suppose also that  $y = y(t)$  is a function defined **implicitly** by

$$G(y) = F(t) + c \quad \Rightarrow \quad G(y(t)) = F(t) + c$$

on some appropriate  $t$ -interval for some real constant  $c$ .

Then, if  $y$  is differentiable, we can differentiate both sides with respect to  $t$ .

$$G'(y(t))y'(t) = F'(t)$$

$$\frac{y'(t)}{g(y(t))} = f(t)$$

$$y'(t) = f(t)g(y(t))$$



## Why does it work?

Suppose  $G(y)$  and  $F(t)$  are antiderivatives of  $\frac{1}{g(y)}$  and  $f(t)$ , respectively.

Suppose also that  $y = y(t)$  is a function defined **implicitly** by

$$G(y) = F(t) + c \quad \Rightarrow \quad G(y(t)) = F(t) + c$$

on some appropriate  $t$ -interval for some real constant  $c$ .

Then, if  $y$  is differentiable, we can differentiate both sides with respect to  $t$ .

$$G'(y(t))y'(t) = F'(t)$$

$$\frac{y'(t)}{g(y(t))} = f(t)$$

$$y'(t) = f(t)g(y(t))$$

This has shown that  $y$  is a solution to  $y' = f(t)g(y)$  and explains why the previous example works.

## Method of Separation of Variables

**Step 1:** Set  $g(y) = 0$  and solve to find any equilibria.

**Step 2:** Now, assume that  $g(y) \neq 0$ . Rewrite the equation in separated form:

$$\frac{dy}{g(y)} = f(t)dt$$

**Step 3:** Integrate, if possible, each side:

$$\int \frac{dy}{g(y)} = \int f(t)dt$$

(obtaining the implicit one-parameter family of solutions.)

**Step 4:** If possible, solve for  $y$  in terms of  $t$ , getting the explicit solution  $y = y(t)$

**Step 5:** If you have an IVP, use the initial condition to evaluate  $c$ .

## Example 2

Are the following differentiable equations are separable?

①  $\frac{dy}{dt} = -\frac{t}{y}$

## Example 2

Are the following differentiable equations are separable?

$$\textcircled{1} \quad \frac{dy}{dt} = -\frac{t}{y} \quad \Rightarrow \quad y \, dy = -t \, dt$$

## Example 2

Are the following differentiable equations are separable?

①  $\frac{dy}{dt} = -\frac{t}{y} \Rightarrow y \, dy = -t \, dt$

②  $\frac{dy}{dt} = t^2 y$

## Example 2

Are the following differentiable equations are separable?

$$\textcircled{1} \quad \frac{dy}{dt} = -\frac{t}{y} \quad \Rightarrow \quad y \, dy = -t \, dt$$

$$\textcircled{2} \quad \frac{dy}{dt} = t^2 y \quad \Rightarrow \quad \frac{dy}{y} = t^2 dt$$

## Example 2

Are the following differentiable equations are separable?

$$\textcircled{1} \quad \frac{dy}{dt} = -\frac{t}{y} \quad \Rightarrow \quad y \, dy = -t \, dt$$

$$\textcircled{2} \quad \frac{dy}{dt} = t^2 y \quad \Rightarrow \quad \frac{dy}{y} = t^2 dt$$

$$\textcircled{3} \quad \frac{dy}{dt} = y + 1$$

## Example 2

Are the following differentiable equations are separable?

$$\textcircled{1} \quad \frac{dy}{dt} = -\frac{t}{y} \quad \Rightarrow \quad y \, dy = -t \, dt$$

$$\textcircled{2} \quad \frac{dy}{dt} = t^2 y \quad \Rightarrow \quad \frac{dy}{y} = t^2 dt$$

$$\textcircled{3} \quad \frac{dy}{dt} = y + 1 \quad \Rightarrow \quad \frac{dy}{y + 1} = dt$$



## Example 2

Are the following differentiable equations are separable?

$$\textcircled{1} \quad \frac{dy}{dt} = -\frac{t}{y} \quad \Rightarrow \quad y \, dy = -t \, dt$$

$$\textcircled{2} \quad \frac{dy}{dt} = t^2 y \quad \Rightarrow \quad \frac{dy}{y} = t^2 dt$$

$$\textcircled{3} \quad \frac{dy}{dt} = y + 1 \quad \Rightarrow \quad \frac{dy}{y + 1} = dt$$

$$\textcircled{4} \quad \frac{dy}{dt} = t + y$$

## Example 2

Are the following differentiable equations are separable?

$$\textcircled{1} \quad \frac{dy}{dt} = -\frac{t}{y} \quad \Rightarrow \quad y \, dy = -t \, dt$$

$$\textcircled{2} \quad \frac{dy}{dt} = t^2 y \quad \Rightarrow \quad \frac{dy}{y} = t^2 dt$$

$$\textcircled{3} \quad \frac{dy}{dt} = y + 1 \quad \Rightarrow \quad \frac{dy}{y + 1} = dt$$

$$\textcircled{4} \quad \frac{dy}{dt} = t + y \quad \Rightarrow \quad (\text{not separable})$$

### Example 3

What are the solutions for the following separable differential equation?

$$y' = \frac{t^2}{1 - y^2}, \quad y \neq \pm 1$$

### Example 3

What are the solutions for the following separable differential equation?

$$y' = \frac{t^2}{1 - y^2}, \quad y \neq \pm 1$$

First we need to rewrite the equation in separated form.

$$(1 - y^2) dy = t^2 dt$$

### Example 3

What are the solutions for the following separable differential equation?

$$y' = \frac{t^2}{1 - y^2}, \quad y \neq \pm 1$$

First we need to rewrite the equation in separated form.

$$(1 - y^2) dy = t^2 dt$$

Next, we can integrate both sides to get:

$$y - \frac{y^3}{3} = \frac{t^3}{3} + c$$

### Example 3

What are the solutions for the following separable differential equation?

$$y' = \frac{t^2}{1 - y^2}, \quad y \neq \pm 1$$

First we need to rewrite the equation in separated form.

$$(1 - y^2) dy = t^2 dt$$

Next, we can integrate both sides to get:

$$y - \frac{y^3}{3} = \frac{t^3}{3} + c$$
$$-t^3 + 3y - y^3 = k \quad \text{where } k = 3c$$

### Example 3

What are the solutions for the following separable differential equation?

$$y' = \frac{t^2}{1 - y^2}, \quad y \neq \pm 1$$

First we need to rewrite the equation in separated form.

$$(1 - y^2) dy = t^2 dt$$

Next, we can integrate both sides to get:


$$y - \frac{y^3}{3} = \frac{t^3}{3} + c$$
$$-t^3 + 3y - y^3 = k \quad \text{where } k = 3c$$

Note that because of the restrictions, each solution curve in the direction field will be a piecewise combination of several functions. A particular solution of an IVP for this DE would only be *one* of these.

### Example 3

What are the solutions for the following separable differential equation?

$$y' = \frac{t^2}{1 - y^2}, \quad y \neq \pm 1$$





### Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

### Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

This separable equation has no equilibrium solutions. Moreover,  $y \neq -\frac{1}{2}$ .

### Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

This separable equation has no equilibrium solutions. Moreover,  $y \neq -\frac{1}{2}$ .

To find any other solutions we need to separate the DE.

$$(1 + 2y)dy = (3t^2 + 1)dt$$

## Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

This separable equation has no equilibrium solutions. Moreover,  $y \neq -\frac{1}{2}$ .

To find any other solutions we need to separate the DE.

$$\begin{aligned}(1 + 2y)dy &= (3t^2 + 1)dt \\ \int (1 + 2y)dy &= \int (3t^2 + 1)dt\end{aligned}$$

## Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

This separable equation has no equilibrium solutions. Moreover,  $y \neq -\frac{1}{2}$ .

To find any other solutions we need to separate the DE.

$$\begin{aligned}(1 + 2y)dy &= (3t^2 + 1)dt \\ \int (1 + 2y)dy &= \int (3t^2 + 1)dt \\ y + y^2 &= t^3 + t + c\end{aligned}$$

### Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

To satisfy the initial conditions we must have  $y = 1$  when  $t = 0$ .

$$1 + 1^2 = 0^3 + 0 + c$$

### Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

To satisfy the initial conditions we must have  $y = 1$  when  $t = 0$ .

$$1 + 1^2 = 0^3 + 0 + c \quad \rightarrow \quad c = 2$$

### Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

To satisfy the initial conditions we must have  $y = 1$  when  $t = 0$ .

$$1 + 1^2 = 0^3 + 0 + c \quad \rightarrow \quad c = 2$$

Which give the solution:

$$y + y^2 = t^3 + t + 2$$



## Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$

To satisfy the initial conditions we must have  $y = 1$  when  $t = 0$ .

$$1 + 1^2 = 0^3 + 0 + c \quad \rightarrow \quad c = 2$$

Which give the solution:


$$y + y^2 = t^3 + t + 2$$
$$y = \frac{1}{2} \left( -1 \pm \sqrt{4t^3 + 4t + 9} \right)$$

Again, the solution curve is made up of multiple parts.

## Example 4

Solve the following IVP.

$$\frac{dy}{dt} = \frac{3t^2 + 1}{1 + 2y}, \quad y(0) = 1$$



## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

There are no equilibrium solutions and  $y \neq 0$ .

We can separate the DE.

$$y \, dy = -t \, dt$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

There are no equilibrium solutions and  $y \neq 0$ .

We can separate the DE.

$$\begin{aligned} y \, dy &= -t \, dt \\ \int y \, dy &= \int -t \, dt \end{aligned}$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

There are no equilibrium solutions and  $y \neq 0$ .

We can separate the DE.

$$\begin{aligned} y \, dy &= -t \, dt \\ \int y \, dy &= \int -t \, dt \\ \frac{y^2}{2} &= -\frac{t^2}{2} + c \end{aligned}$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

There are no equilibrium solutions and  $y \neq 0$ .

We can separate the DE.

$$\begin{aligned} y \, dy &= -t \, dt \\ \int y \, dy &= \int -t \, dt \\ \frac{y^2}{2} &= -\frac{t^2}{2} + c \end{aligned}$$

We can then use the initial condition to find  $c$ .

$$\frac{1}{2} = -\frac{0^2}{2} + c \quad \Rightarrow \quad c = \frac{1}{2}$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

We have two explicit solutions:

$$\frac{y^2}{2} = -\frac{t^2}{2} + \frac{1}{2}$$



## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

We have two explicit solutions:

$$\begin{aligned}\frac{y^2}{2} &= -\frac{t^2}{2} + \frac{1}{2} \\ y^2 &= -t^2 + 1\end{aligned}$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

We have two explicit solutions:

$$\frac{y^2}{2} = -\frac{t^2}{2} + \frac{1}{2}$$

$$y^2 = -t^2 + 1$$

$$y = \pm\sqrt{1 - t^2}$$

## Example 5

Solve the IVP.

$$y' = -\frac{t}{y}, \quad y(0) = 1$$

