

Direction Fields and Euler's Method

Colby Community College

What is a Differential Equation?

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

What is a Differential Equation?

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

In this chapter we will focus on DEs that can be written as:

$$\frac{dy}{dt} = f(t, y) \quad \text{or} \quad y' = f(t, y)$$

Where the dependent variable y is an unknown function, the **solution**.

What is a Differential Equation?

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

In this chapter we will focus on DEs that can be written as:

$$\frac{dy}{dt} = f(t, y) \quad \text{or} \quad y' = f(t, y)$$

Where the dependent variable y is an unknown function, the **solution**.

Note

There may be more than one solution for a given differential equation.

What is a Differential Equation?

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

In this chapter we will focus on DEs that can be written as:

$$\frac{dy}{dt} = f(t, y) \quad \text{or} \quad y' = f(t, y)$$

Where the dependent variable y is an unknown function, the **solution**.

Note

There may be more than one solution for a given differential equation.

Analytic Definition of a Solution

Analytically, $y(t)$ is a **solution** of a differential equation if substituting $y(t)$ for y reduced the equation to an identity:

$$y'(t) = f(t, y(t))$$

on an appropriate domain for t .

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$y'(t) = \frac{d}{dt}e^{2t}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \end{aligned}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \\ &= 2y(t) \end{aligned}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \\ &= 2y(t) \\ &= f(t, y(t)) \end{aligned}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \\ &= 2y(t) \\ &= f(t, y(t)) \end{aligned}$$

Similarly, we could show that


$$y(t) = 2e^{2t} \quad \text{and} \quad y(t) = \frac{-3}{2}e^{2t}$$

are also solutions. In fact, any constant multiple of e^{2t} is a solution.

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$



Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1 - t^2}$$

Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1-t^2}$$

Substituting into the DE gives:

$$y'(t) = \frac{d}{dt} \left(\sqrt{1-t^2} \right)$$

Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1-t^2}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} \left(\sqrt{1-t^2} \right) \\ &= \frac{1}{2} (1-t^2)^{-\frac{1}{2}} \cdot (-2t) \end{aligned}$$

Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1-t^2}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} \left(\sqrt{1-t^2} \right) \\ &= \frac{1}{2} (1-t^2)^{-\frac{1}{2}} \cdot (-2t) \\ &= \frac{-t}{\sqrt{1-t^2}} \end{aligned}$$

Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1-t^2}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} \left(\sqrt{1-t^2} \right) \\ &= \frac{1}{2} (1-t^2)^{-\frac{1}{2}} \cdot (-2t) \\ &= \frac{-t}{\sqrt{1-t^2}} \\ &= -\frac{t}{y} \end{aligned}$$

Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1-t^2}$$

Substituting into the DE gives:


$$\begin{aligned} y'(t) &= \frac{d}{dt} \left(\sqrt{1-t^2} \right) \\ &= \frac{1}{2} (1-t^2)^{-\frac{1}{2}} \cdot (-2t) \\ &= \frac{-t}{\sqrt{1-t^2}} \\ &= -\frac{t}{y} \end{aligned}$$

Other solutions are of the form $y(t) = \sqrt{k-t^2}$.

Example 2

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = -\frac{t}{y}, \quad y(t) = \sqrt{1 - t^2}$$



Note

It is no coincidence that the two previous examples had multiple solutions. Most differential equations have an infinite number of solutions.

Note

It is no coincidence that the two previous examples had multiple solutions. Most differential equations have an infinite number of solutions.

Example 3

Consider

$$\frac{dy}{dt} = f(t)$$

Note

It is no coincidence that the two previous examples had multiple solutions. Most differential equations have an infinite number of solutions.

Example 3

Consider

$$\frac{dy}{dt} = f(t)$$

We can integrate both sides to get the solution to get

$$y = \int f(t)dt + c$$

where c is an arbitrary constant.

Note

It is no coincidence that the two previous examples had multiple solutions. Most differential equations have an infinite number of solutions.

Example 3

Consider

$$\frac{dy}{dt} = f(t)$$

We can integrate both sides to get the solution to get

$$y = \int f(t)dt + c$$

where c is an arbitrary constant.

Family of Solutions



In general, all solutions of a first-order DE form a **family** of solutions expressed with a single parameter c . Such a family is called the **general solution**. A member of the family that results from a specific value of c is called a **particular solution**.

Example 4

The general solution of $y' = 3t^2$ is

$$y = t^3 + c$$

where c may be any real value.



Initial-Value Problem

The combination of a first-order differential equation and an **initial condition**

$$\frac{dy}{dt} = f(t, y), \quad y(t_0) = y_0$$

is called an **initial-value problem**. It's solution will pass through the point (t_0, y_0) .

Initial-Value Problem

The combination of a first-order differential equation and an **initial condition**

$$\frac{dy}{dt} = f(t, y), \quad y(t_0) = y_0$$

is called an **initial-value problem**. It's solution will pass through the point (t_0, y_0) .

Note

While a family of solutions for a DE contains multiple solutions, an IVP usually has only one solution. That is, the solution to an IVP is a particular solution to the DE.

Example 5

The function $y(t) = t^3 + 1$ is a solution to the IVP

$$y' = 3t^2, \quad y(0) = 1$$


Example 5

The function $y(t) = t^3 + 1$ is a solution to the IVP

$$y' = 3t^2, \quad y(0) = 1$$

Differentiating $y(t)$ confirms that

$$y'(t) = (t^3 + 1)' = 3t^2, \quad \text{and} \quad y(0) = 0^3 + 1 = 1$$



Graphical Definition of Solutions

A **solution** to a first-order differential equation is a function whose slope at each point is specified by the derivative.

Graphical Definition of Solutions


A **solution** to a first-order differential equation is a function whose slope at each point is specified by the derivative.

Direction Fields

We can see what solution curves look like by, on regular intervals, drawing short line segments with slope determined by the DE for that point. The collection of these segments are called **direction field** (or a **slope field**).


Example 6

$$y' = -2ty + t$$




Example 6

$$y' = -2ty + t$$




Example 7

$$y' = y + t^2$$




Example 7

$$y' = y + t^2$$




Example 8

$$y' = t - y$$




Example 8

$$y' = t - y$$



Example 8

$$y' = t - y \quad \text{and} \quad y'' = 1 - y' = 1 - t + y$$



Equilibria

For a differential equation, a solution that does not change over time is called an **equilibrium solution**.

Equilibria

For a differential equation, a solution that does not change over time is called an **equilibrium solution**.

For a first-order DE $y' = f(t, y)$, an equilibrium solution is always a horizontal line $y(t) = C$, which can be obtained by setting $y' = 0$.

Equilibria

For a differential equation, a solution that does not change over time is called an **equilibrium solution**.

For a first-order DE $y' = f(t, y)$, an equilibrium solution is always a horizontal line $y(t) = C$, which can be obtained by setting $y' = 0$.

Stability

For a differential equation $y' = f(t, y)$, an equilibrium solution $y(t) = C$ is called

- **stable** if solutions “near” it tend toward it as $t \rightarrow \infty$.

Equilibria

For a differential equation, a solution that does not change over time is called an **equilibrium solution**.

For a first-order DE $y' = f(t, y)$, an equilibrium solution is always a horizontal line $y(t) = C$, which can be obtained by setting $y' = 0$.

Stability

For a differential equation $y' = f(t, y)$, an equilibrium solution $y(t) = C$ is called

- **stable** if solutions “near” it tend toward it as $t \rightarrow \infty$.
- **unstable** if solutions “near” it tend away from it as $t \rightarrow \infty$.

Equilibria

For a differential equation, a solution that does not change over time is called an **equilibrium solution**.

For a first-order DE $y' = f(t, y)$, an equilibrium solution is always a horizontal line $y(t) = C$, which can be obtained by setting $y' = 0$.

Stability

For a differential equation $y' = f(t, y)$, an equilibrium solution $y(t) = C$ is called


- **stable** if solutions “near” it tend toward it as $t \rightarrow \infty$.
- **unstable** if solutions “near” it tend away from it as $t \rightarrow \infty$.

Note

A equilibrium solution is often called **semistable** if it is stable on one side and unstable on the other.


Example 9

The DE $y' = -2ty + t$ has the constant solution $y(t) = \frac{1}{2}$.




Example 10

$$y' = y^2 - 4$$



Example 10


$$y' = y^2 - 4$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP


$$y' = t - y, \quad y(0) = 1$$



Example 11

We can approximate the solution of the IVP

$$y' = t - y, \quad y(0) = 1$$



Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can calculate the t -values, for $1, 2, 3, \dots, k$, with $t_n = t_0 + n \cdot h$.

Where h , called the **step size**, is the common difference between successive points.

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can calculate the t -values, for $1, 2, 3, \dots, k$, with $t_n = t_0 + n \cdot h$.

Where h , called the **step size**, is the common difference between successive points.

So, starting at (t_0, y_0) we want to follow the tangent line determined by

$$y - y_0 = (t - t_0)f(t_0, y_0)$$

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can calculate the t -values, for $1, 2, 3, \dots, k$, with $t_n = t_0 + n \cdot h$.

Where h , called the **step size**, is the common difference between successive points.

So, starting at (t_0, y_0) we want to follow the tangent line determined by

$$y - y_0 = (t - t_0)f(t_0, y_0)$$

to find the approximate solution $(t_1, y(t_1))$:

$$y_1 = y_0 + h \cdot f(t_0, y_0)$$

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can extend this process to find all k points.

$$y_1 = y_0 + h \cdot f(t_0, y_0)$$

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can extend this process to find all k points.

$$y_1 = y_0 + h \cdot f(t_0, y_0)$$

$$y_2 = y_1 + h \cdot f(t_1, y_1)$$

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can extend this process to find all k points.

$$y_1 = y_0 + h \cdot f(t_0, y_0)$$

$$y_2 = y_1 + h \cdot f(t_1, y_1)$$

$$y_3 = y_2 + h \cdot f(t_2, y_2)$$

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can extend this process to find all k points.

$$y_1 = y_0 + h \cdot f(t_0, y_0)$$

$$y_2 = y_1 + h \cdot f(t_1, y_1)$$

$$y_3 = y_2 + h \cdot f(t_2, y_2)$$

$$\vdots$$

$$y_k = y_{k-1} + h \cdot f(t_{k-1}, y_{k-1})$$

Consider

Given the IVP

$$y' = f(t, y), \quad y(t_0) = y_0$$

We want to compute approximate values for $y(t_n)$ at the (finite) set of points $t_1, t_2, t_3, \dots, t_k$.

We can extend this process to find all k points.

$$y_1 = y_0 + h \cdot f(t_0, y_0)$$

$$y_2 = y_1 + h \cdot f(t_1, y_1)$$

$$y_3 = y_2 + h \cdot f(t_2, y_2)$$

$$\vdots$$

$$y_k = y_{k-1} + h \cdot f(t_{k-1}, y_{k-1})$$

The resulting piecewise-linear function (i.e. play connect-the-dots) is called the **Euler-approximate** solution.

Euler's Method

For the Initial-value problem

$$y' = f(t, y), \quad y(t_0) = y_0$$

use the formulas

$$t_{n+1} = t_n + h$$

$$y_{n+1} = y_n + h \cdot f(t_n, y_n)$$

to iteratively compute the points, using step size h ,

$$(t_1, y_1), (t_2, y_2), \dots, (t_k, y_k).$$

The piecewise-linear function connecting these points is the Euler approximation to the solution $y(t)$ of the IVP for $t_0 \leq t \leq t_k$.

Example 12

Let us obtain the Euler-approximate solution of the IVP

$$y' = -2ty + t, \quad y(0) = -1$$

with step size 0.1 on $[0, 0.4]$.

Example 12

Let us obtain the Euler-approximate solution of the IVP

$$y' = -2ty + t, \quad y(0) = -1$$

with step size 0.1 on $[0, 0.4]$.

In other words:

$$f(t, y) = -2ty + t = t(1 - 2y)$$

$$t_0 = 0$$

$$y_0 = -1$$

$$h = 0.1$$

$$k = 1, 2, 3, 4$$

Example 12

$$t_1 = t_0 + h = 0 + 0.1 = 0.1$$

$$y_1 = y_0 + h \cdot f(t_0, y_0) = -1 + (0.1)(0)(1 - 2(-1)) = -1$$

Example 12

$$t_1 = t_0 + h = 0 + 0.1 = 0.1$$

$$y_1 = y_0 + h \cdot f(t_0, y_0) = -1 + (0.1)(0)(1 - 2(-1)) = -1$$

$$t_2 = t_1 + h = 0.1 + 0.1 = 0.2$$

$$\begin{aligned} y_2 &= y_1 + h \cdot f(t_1, y_1) \\ &= -1 + (0.1)(0.1)(1 - 2(-1)) = -0.97 \end{aligned}$$

Example 12

$$t_1 = t_0 + h = 0 + 0.1 = 0.1$$

$$y_1 = y_0 + h \cdot f(t_0, y_0) = -1 + (0.1)(0)(1 - 2(-1)) = -1$$

$$t_2 = t_1 + h = 0.1 + 0.1 = 0.2$$

$$\begin{aligned} y_2 &= y_1 + h \cdot f(t_1, y_1) \\ &= -1 + (0.1)(0.1)(1 - 2(-1)) = -0.97 \end{aligned}$$

$$t_3 = t_2 + h = 0.2 + 0.1 = 0.3$$

$$\begin{aligned} y_3 &= y_2 + h \cdot f(t_2, y_2) \\ &= -0.97 + (0.1)(0.2)(1 - 2(-0.97)) = -0.9112 \end{aligned}$$

Example 12

$$t_1 = t_0 + h = 0 + 0.1 = 0.1$$

$$y_1 = y_0 + h \cdot f(t_0, y_0) = -1 + (0.1)(0)(1 - 2(-1)) = -1$$

$$t_2 = t_1 + h = 0.1 + 0.1 = 0.2$$

$$\begin{aligned} y_2 &= y_1 + h \cdot f(t_1, y_1) \\ &= -1 + (0.1)(0.1)(1 - 2(-1)) = -0.97 \end{aligned}$$

$$t_3 = t_2 + h = 0.2 + 0.1 = 0.3$$

$$\begin{aligned} y_3 &= y_2 + h \cdot f(t_2, y_2) \\ &= -0.97 + (0.1)(0.2)(1 - 2(-0.97)) = -0.9112 \end{aligned}$$

$$t_4 = t_3 + h = 0.3 + 0.1 = 0.4$$

$$\begin{aligned} y_4 &= y_3 + h \cdot f(t_3, y_3) \\ &= -0.9112 + (0.1)(0.3)(1 - 2(-0.9112)) = -0.82652 \end{aligned}$$

Example 12

How does this compare to the exact solution $y(t) = 0.5 - 1.5e^{-t^2}$?


Example 12

How does this compare to the exact solution $y(t) = 0.5 - 1.5e^{-t^2}$?

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|------------------|
| 0 | 0.0 | -1.000000 | -1.000000 | 0.000000 |
| 1 | 0.1 | -1.000000 | -0.985075 | -0.014925 |
| 2 | 0.2 | -0.970000 | -0.941184 | -0.028815 |
| 3 | 0.3 | -0.911200 | -0.870897 | -0.040303 |
| 4 | 0.4 | -0.826528 | -0.778216 | -0.048312 |

Notice how the error grows rapidly.

Example 12



Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-------|----------|-------|
|-----|-------|-------|----------|-------|

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|------------------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |
| 5 | 1.0 | 0.3993830 | 0.3678794 | -0.031504 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |
| 5 | 1.0 | 0.3993830 | 0.3678794 | -0.031504 |
| 6 | 1.2 | 0.2396298 | 0.2369277 | -0.002702 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |
| 5 | 1.0 | 0.3993830 | 0.3678794 | -0.031504 |
| 6 | 1.2 | 0.2396298 | 0.2369277 | -0.002702 |
| 7 | 1.4 | 0.1246075 | 0.1408584 | 0.016251 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |
| 5 | 1.0 | 0.3993830 | 0.3678794 | -0.031504 |
| 6 | 1.2 | 0.2396298 | 0.2369277 | -0.002702 |
| 7 | 1.4 | 0.1246075 | 0.1408584 | 0.016251 |
| 8 | 1.6 | 0.0548273 | 0.0773047 | 0.022477 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.

| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |
| 5 | 1.0 | 0.3993830 | 0.3678794 | -0.031504 |
| 6 | 1.2 | 0.2396298 | 0.2369277 | -0.002702 |
| 7 | 1.4 | 0.1246075 | 0.1408584 | 0.016251 |
| 8 | 1.6 | 0.0548273 | 0.0773047 | 0.022477 |
| 9 | 1.8 | 0.0197378 | 0.0391639 | 0.019426 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.


| n | t_n | y_n | $y(t_n)$ | Error |
|-----|-------|-----------|-----------|-----------|
| 0 | 0.0 | 1.0000000 | 1.0000000 | 0.000000 |
| 1 | 0.2 | 1.0000000 | 0.9607894 | -0.039211 |
| 2 | 0.4 | 0.9200000 | 0.8521437 | -0.067856 |
| 3 | 0.6 | 0.7728000 | 0.6976763 | -0.075124 |
| 4 | 0.8 | 0.5873280 | 0.5272925 | -0.060036 |
| 5 | 1.0 | 0.3993830 | 0.3678794 | -0.031504 |
| 6 | 1.2 | 0.2396298 | 0.2369277 | -0.002702 |
| 7 | 1.4 | 0.1246075 | 0.1408584 | 0.016251 |
| 8 | 1.6 | 0.0548273 | 0.0773047 | 0.022477 |
| 9 | 1.8 | 0.0197378 | 0.0391639 | 0.019426 |
| 10 | 2.0 | 0.0055265 | 0.0183156 | 0.012789 |

Example 13

Find the Euler-approximation of

$$y' = -2ty, \quad y(0) = 1$$

using a step size of 0.2 over the range of $[0, 2]$.



There are two types of error:

There are two types of error:

- **Roundoff error** is the discrepancy arising from rounding numbers. This tends to snowball pretty fast when you have a great many calculations.

There are two types of error:

- **Roundoff error** is the discrepancy arising from rounding numbers. This tends to snowball pretty fast when you have a great many calculations.
- **Discretization error** is the error that results from the approximation method itself. For Euler's method this is caused by using the linear tangent lines to approximate a nonlinear curve.

There are two types of error:

- **Roundoff error** is the discrepancy arising from rounding numbers. This tends to snowball pretty fast when you have a great many calculations.
- **Discretization error** is the error that results from the approximation method itself. For Euler's method this is caused by using the linear tangent lines to approximate a nonlinear curve.

It can be shown, using Taylor series expansions, that the error is proportional to the square of the step size. (We will talk about Taylor series in chapter 11.)

$$|y_i - y(t_i)| \leq C \cdot h^2$$

Where the constant C depends on the size of the second derivative of the exact solution.

There are two types of error:

- **Roundoff error** is the discrepancy arising from rounding numbers. This tends to snowball pretty fast when you have a great many calculations.
- **Discretization error** is the error that results from the approximation method itself. For Euler's method this is caused by using the linear tangent lines to approximate a nonlinear curve.

It can be shown, using Taylor series expansions, that the error is proportional to the square of the step size. (We will talk about Taylor series in chapter 11.)

$$|y_i - y(t_i)| \leq C \cdot h^2$$

Where the constant C depends on the size of the second derivative of the exact solution.

We call this error the **local discretization error** because it estimates the error for a single step only. After n steps, we have n times the error. Which we call the **global discretization error**.