

# Linear Systems of Differential Equations

Department of Mathematics

Salt Lake Community College

## Linear First-Order DE System

An  $n$ -dimensional **linear first-order DE system** on open interval  $I$  is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

## Linear First-Order DE System

An  $n$ -dimensional **linear first-order DE system** on open interval  $I$  is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$  is an  $n \times n$  matrix of continuous functions on  $I$ .

## Linear First-Order DE System

An  $n$ -dimensional **linear first-order DE system** on open interval  $I$  is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$  is an  $n \times n$  matrix of continuous functions on  $I$ .
- $\vec{f}(t)$  is an  $n \times 1$  vector of continuous functions on  $I$ .

## Linear First-Order DE System

An  $n$ -dimensional **linear first-order DE system** on open interval  $I$  is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$  is an  $n \times n$  matrix of continuous functions on  $I$ .
- $\vec{f}(t)$  is an  $n \times 1$  vector of continuous functions on  $I$ .
- $\vec{x}(t)$  is an  $n \times 1$  **solutions vector** of differentiable functions on  $I$  that satisfies the DE.

## Linear First-Order DE System

An  $n$ -dimensional **linear first-order DE system** on open interval  $I$  is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$  is an  $n \times n$  matrix of continuous functions on  $I$ .
- $\vec{f}(t)$  is an  $n \times 1$  vector of continuous functions on  $I$ .
- $\vec{x}(t)$  is an  $n \times 1$  **solutions vector** of differentiable functions on  $I$  that satisfies the DE.

If  $\vec{f}(t) = \vec{0}$ , the system is **homogeneous**

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t)$$

## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$



## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution  $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution  $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}$$

## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution  $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} - 2e^{2t} \\ 2e^{2t} \\ -2e^{2t} + e^{2t} + 3e^{2t} \end{bmatrix}$$

## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution  $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} - 2e^{2t} \\ 2e^{2t} \\ -2e^{2t} + e^{2t} + 3e^{2t} \end{bmatrix} = \begin{bmatrix} 4e^{2t} \\ 2e^{2t} \\ 2e^{2t} \end{bmatrix}$$

## Example 1

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution  $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} - 2e^{2t} \\ 2e^{2t} \\ -2e^{2t} + e^{2t} + 3e^{2t} \end{bmatrix} = \begin{bmatrix} 4e^{2t} \\ 2e^{2t} \\ 2e^{2t} \end{bmatrix} = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}'$$

## Initial-Value Problem for a Linear DE System

For a linear DE system, an **initial-value problem** is the combination of a linear DE system and an initial value vector.

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t), \quad \vec{x}(t_0) = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

## Initial-Value Problem for a Linear DE System

For a linear DE system, an **initial-value problem** is the combination of a linear DE system and an initial value vector.

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t), \quad \vec{x}(t_0) = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

## Existence and Uniqueness Theorem for Linear DE Systems

Given an  $n \times n$  matrix function  $\mathbf{A}(t)$  and a  $n \times 1$  vector function  $\vec{f}(t)$ , both continuous on an open interval  $I$  containing  $t_0$ , and a constant  $n$ -vector  $\vec{x}_0$ , there exists a unique vector function  $\vec{x}(t)$  such that

$$\vec{x}' = \mathbf{A}(t)\vec{x} + \vec{f}(t) \quad \text{and} \quad \vec{x}(t_0) = \vec{x}_0$$

## The Superposition Principle for Homogeneous Linear DE Systems

Let  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$  be solution vectors for the homogenous equation

$$\vec{x}' = \mathbf{A}(t)\vec{x} \quad \text{on } I$$

Then, any linear combination of these solution vectors is also a solution vector for the system.

That is,

$$\vec{x} = c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_n\vec{x}_n$$

is also a solution on  $I$  for any  $c_1, c_2, \dots, c_n \in \mathbb{R}$ .



## Solution Space Theorem for Homogeneous Linear DE Systems

If

$$\vec{x}' = \mathbf{A}(t)\vec{x}$$

where  $\mathbf{A}$  is an  $n \times n$  matrix, then the set of solutions  $\vec{x}(t)$  is a vector space of dimension  $n$ .

## Solution Space Theorem for Homogeneous Linear DE Systems

If

$$\vec{x}' = \mathbf{A}(t)\vec{x}$$

where  $\mathbf{A}$  is an  $n \times n$  matrix, then the set of solutions  $\vec{x}(t)$  is a vector space of dimension  $n$ .

## Solution Theorem for Homogenous Linear DE Systems

For  $n$  linearly independent solutions  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$  of

$$\vec{x}' = \mathbf{A}(t)\vec{x}$$

the general solution is

$$\vec{x} = c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_n\vec{x}_n \quad \text{where} \quad c_1, c_2, \dots, c_n \in \mathbb{R}$$

## Example 2

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

## Example 2

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that  $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$  are linearly independent on  $(-\infty, \infty)$  choose a point, say  $t_0 = 0$ .

## Example 2

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that  $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$  are linearly independent on  $(-\infty, \infty)$  choose a point, say  $t_0 = 0$ .

Calculate  $\vec{x}_1(t_0)$ ,  $\vec{x}_2(t_0)$ , and  $\vec{x}_3(t_0)$ . Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

## Example 2

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that  $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$  are linearly independent on  $(-\infty, \infty)$  choose a point, say  $t_0 = 0$ .

Calculate  $\vec{x}_1(t_0)$ ,  $\vec{x}_2(t_0)$ , and  $\vec{x}_3(t_0)$ . Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

If the determinant is nonzero, then the solutions are linearly independent.

## Example 2

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that  $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$  are linearly independent on  $(-\infty, \infty)$  choose a point, say  $t_0 = 0$ .

Calculate  $\vec{x}_1(t_0)$ ,  $\vec{x}_2(t_0)$ , and  $\vec{x}_3(t_0)$ . Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

If the determinant is nonzero, then the solutions are linearly independent.

$$|\mathbf{C}| = 0(1 \cdot 0 - 1 \cdot 1) - 2(0 \cdot 0 - 1 \cdot 1) + 1(0 \cdot 1 - 1 \cdot 1) = 1$$

## Example 2

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that  $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$  are linearly independent on  $(-\infty, \infty)$  choose a point, say  $t_0 = 0$ .

Calculate  $\vec{x}_1(t_0)$ ,  $\vec{x}_2(t_0)$ , and  $\vec{x}_3(t_0)$ . Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

If the determinant is nonzero, then the solutions are linearly independent.

$$|\mathbf{C}| = 0(1 \cdot 0 - 1 \cdot 1) - 2(0 \cdot 0 - 1 \cdot 1) + 1(0 \cdot 1 - 1 \cdot 1) = 1$$

So, the general solution is  $\vec{x} = c_1\vec{x}_1 + c_2\vec{x}_2 + c_3\vec{x}_3$ .



## Note

We have a few ways to express solutions:

$$\vec{x} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_3 \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

## Note

We have a few ways to express solutions:

$$\vec{x} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_3 \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix} = \begin{bmatrix} 0 & 2e^{2t} & e^t \\ 0 & e^{2t} & e^t \\ e^{3t} & e^{2t} & 0 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

## Note

We have a few ways to express solutions:

$$\vec{x} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_3 \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix} = \begin{bmatrix} 0 & 2e^{2t} & e^t \\ 0 & e^{2t} & e^t \\ e^{3t} & e^{2t} & 0 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

## Fundamental Matrix

For a basis of  $n$  linearly independent solutions of  $\vec{x}' = \mathbf{A}\vec{x}$ , the matrix  $\mathbf{X}(t)$  whose *columns* are the vector solutions  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$  is called the **fundamental matrix** for the system.

$$\vec{x} = \underbrace{\begin{bmatrix} | & | & \cdots & | \\ \vec{x}_1 & \vec{x}_2 & \cdots & \vec{x}_n \\ | & | & \cdots & | \end{bmatrix}}_{\mathbf{X}(t)} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \quad c_1, c_2, c_3 \in \mathbb{R}$$

## Graphical Views

- The  $t$ - $x$  and  $t$ - $y$  graphs showing the individual solution functions  $x(t)$  and  $y(t)$  are called **component graphs**, **solution graphs**, or **time series**.
- The  $x$ - $y$  graph is the **phase plane**. The **trajectories** in the phase plane are the parametric curves described by  $x(t)$  and  $y(t)$ .

Trajectories on a phase plane create a **phase portrait**.

### Example 3

The familiar equation

$$x'' + 0.1x = 0$$

### Example 3

The familiar equation

$$x'' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

### Example 3

The familiar equation

$$x'' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Any version of the these equations produces solutions of the form



$$x(t) = c_1 \cos(t\sqrt{0.1}) + c_2 \sin(t\sqrt{0.1})$$

$$y(t) = x'(t) = -c_1\sqrt{0.1} \sin(t\sqrt{0.1}) + c_2\sqrt{0.1} \cos(t\sqrt{0.1})$$

### Example 3

The familiar equation

$$x'' + 0.1x = 0$$





## Example 4

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$

## Example 4

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x - 0.05y \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & -0.05 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

## Example 4

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x - 0.05y \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & -0.05 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$




With solutions of the (approximate) form

$$\begin{aligned} x(t) &\approx e^{-0.025t} (c_1 \cos(0.32t) + c_2 \sin(0.32t)) \\ y(t) &\approx e^{-0.025t} (-0.32c_1 \sin(0.32t) + 0.32c_2 \cos(0.23t)) \\ &\quad - 0.025e^{-0.025t} (c_1 \cos(0.32t) + c_2 \sin(0.32t)) \end{aligned}$$

## Example 4

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$



## Example 5

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

## Example 5

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

This DE represents a periodically forced harmonic oscillator and has system form:

$$\begin{aligned} x' &= y \\ y' &= -0.1x + 0.5 \cos(t) \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 0 \\ 0.5 \cos(t) \end{bmatrix}$$

## Example 5

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

This DE represents a periodically forced harmonic oscillator and has system form:

$$\begin{aligned} x' &= y \\ y' &= -0.1x + 0.5 \cos(t) \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 0 \\ 0.5 \cos(t) \end{bmatrix}$$

It is not easy to find an analytic solution to this DE, but we can draw the solutions using numerical calculations.

We can use Euler's method, which we have seen before.

## Example 5

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

$$x_{n+1} = x_n + h \cdot x'(t_n) = x_n + h \cdot y_n$$

$$y_{n+1} = y_n + h \cdot y'(t_n) = y_n + h \cdot (-0.1x + 0.5 \cos(t_n))$$

With step size  $h = 0.1$ ,  $x(0) = 1$ , and  $y(0) = 0$ .



$t_n$	$x_n$	$y_n$	$x'$	$y'$
0.0	1.0000	0.0000	0.0000	0.4000
0.1	1.0000	0.0400	0.0400	0.3975
0.2	1.0040	0.0798	0.0798	0.3896
0.3	1.0120	0.1187	0.1187	0.3765
0.4	1.0238	0.1564	0.1564	0.3581
0.5	1.0395	0.1922	0.1922	0.3348
0.6	1.0587	0.2257	0.2257	0.3068
0.7	1.0813	0.2563	0.2563	0.2743
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$



## Example 5

Let us consider a nonautonomous version of



$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$



## Example 5

Let us consider a nonautonomous version of



$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$



## Example 5

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$



## Graphical Properties of Uniqueness

- For a linear system of differential equations in  $\mathbb{R}^n$ , solutions do not cross in  $t, x_1, x_2, \dots, x_n$ -space (i.e.  $\mathbb{R}^{n+1}$ ).

## Graphical Properties of Uniqueness

- For a linear system of differential equations in  $\mathbb{R}^n$ , solutions do not cross in  $t, x_1, x_2, \dots, x_n$ -space (i.e.  $\mathbb{R}^{n+1}$ ).
- For an *autonomous* linear system in  $\mathbb{R}^n$ , trajectories *also* do not cross in  $x_1, x_2, \dots, x_n$ -space (i.e.  $\mathbb{R}^n$ ).