

Linear Systems of Differential Equations

Adam Wilson

Salt Lake Community College

Linear First-Order DE System

An n -dimensional **linear first-order DE system** on open interval I is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

Linear First-Order DE System

An n -dimensional **linear first-order DE system** on open interval I is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$ is an $n \times n$ matrix of continuous functions on I .

Linear First-Order DE System

An n -dimensional **linear first-order DE system** on open interval I is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$ is an $n \times n$ matrix of continuous functions on I .
- $\vec{f}(t)$ is an $n \times 1$ vector of continuous functions on I .

Linear First-Order DE System

An n -dimensional **linear first-order DE system** on open interval I is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$ is an $n \times n$ matrix of continuous functions on I .
- $\vec{f}(t)$ is an $n \times 1$ vector of continuous functions on I .
- $\vec{x}(t)$ is an $n \times 1$ **solutions vector** of differentiable functions on I that satisfies the DE.

Linear First-Order DE System

An n -dimensional **linear first-order DE system** on open interval I is one that can be written as a matrix-vector equation:

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t)$$

- $\mathbf{A}(t)$ is an $n \times n$ matrix of continuous functions on I .
- $\vec{f}(t)$ is an $n \times 1$ vector of continuous functions on I .
- $\vec{x}(t)$ is an $n \times 1$ **solutions vector** of differentiable functions on I that satisfies the DE.

If $\vec{f}(t) = \vec{0}$, the system is **homogeneous**

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t)$$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}$$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} - 2e^{2t} \\ 2e^{2t} \\ -2e^{2t} + e^{2t} + 3e^{2t} \end{bmatrix}$$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} - 2e^{2t} \\ 2e^{2t} \\ -2e^{2t} + e^{2t} + 3e^{2t} \end{bmatrix} = \begin{bmatrix} 4e^{2t} \\ 2e^{2t} \\ 2e^{2t} \end{bmatrix}$$

Example

Consider the homogeneous linear first-order system

$$x' = 3x - 2y$$

$$y' = x$$

$$z' = -x + y + 3z$$

Which can be written as:

$$\vec{x}' = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \vec{x} \quad \text{where} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

and has solution $\vec{x} = \langle 2e^{2t}, e^{2t}, e^{2t} \rangle$

We can verify this:

$$\begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} - 2e^{2t} \\ 2e^{2t} \\ -2e^{2t} + e^{2t} + 3e^{2t} \end{bmatrix} = \begin{bmatrix} 4e^{2t} \\ 2e^{2t} \\ 2e^{2t} \end{bmatrix} = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}'$$

Initial-Value Problem for a Linear DE System

For a linear DE system, an **initial-value problem** is the combination of a linear DE system and an initial value vector.

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t), \quad \vec{x}(t_0) = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

Initial-Value Problem for a Linear DE System

For a linear DE system, an **initial-value problem** is the combination of a linear DE system and an initial value vector.

$$\vec{x}'(t) = \mathbf{A}(t)\vec{x}(t) + \vec{f}(t), \quad \vec{x}(t_0) = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

Existence and Uniqueness Theorem for Linear DE Systems

Given an $n \times n$ matrix function $\mathbf{A}(t)$ and a $n \times 1$ vector function $\vec{f}(t)$, both continuous on an open interval I containing t_0 , and a constant n -vector \vec{x}_0 , there exists a unique vector function $\vec{x}(t)$ such that

$$\vec{x}' = \mathbf{A}(t)\vec{x} + \vec{f}(t) \quad \text{and} \quad \vec{x}(t_0) = \vec{x}_0$$

The Superposition Principle for Homogeneous Linear DE Systems

Let $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ be solution vectors for the homogenous equation

$$\vec{x}' = \mathbf{A}(t)\vec{x} \quad \text{on } I$$

Then, any linear combination of these solution vectors is also a solution vector for the system.

That is,

$$\vec{x} = c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_n\vec{x}_n$$

is also a solution on I for any $c_1, c_2, \dots, c_n \in \mathbb{R}$.

Solution Space Theorem for Homogeneous Linear DE Systems

If

$$\vec{x}' = \mathbf{A}(t)\vec{x}$$

where \mathbf{A} is an $n \times n$ matrix, then the set of solutions $\vec{x}(t)$ is a vector space of dimension n .

Solution Space Theorem for Homogeneous Linear DE Systems

If

$$\vec{x}' = \mathbf{A}(t)\vec{x}$$

where \mathbf{A} is an $n \times n$ matrix, then the set of solutions $\vec{x}(t)$ is a vector space of dimension n .

Solution Theorem for Homogenous Linear DE Systems

For n linearly independent solutions $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ of

$$\vec{x}' = \mathbf{A}(t)\vec{x}$$

the general solution is

$$\vec{x} = c_1\vec{x}_1 + c_2\vec{x}_2 + \cdots + c_n\vec{x}_n \quad \text{where} \quad c_1, c_2, \dots, c_n \in \mathbb{R}$$

Example

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

Example

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ are linearly independent on $(-\infty, \infty)$ choose a point, say $t_0 = 0$.

Example

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ are linearly independent on $(-\infty, \infty)$ choose a point, say $t_0 = 0$.

Calculate $\vec{x}_1(t_0)$, $\vec{x}_2(t_0)$, and $\vec{x}_3(t_0)$. Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

Example

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ are linearly independent on $(-\infty, \infty)$ choose a point, say $t_0 = 0$.

Calculate $\vec{x}_1(t_0)$, $\vec{x}_2(t_0)$, and $\vec{x}_3(t_0)$. Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

If the determinant is nonzero, then the solutions are linearly independent.

Example

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ are linearly independent on $(-\infty, \infty)$ choose a point, say $t_0 = 0$.

Calculate $\vec{x}_1(t_0)$, $\vec{x}_2(t_0)$, and $\vec{x}_3(t_0)$. Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

If the determinant is nonzero, then the solutions are linearly independent.

$$|\mathbf{C}| = 0(1 \cdot 0 - 1 \cdot 1) - 2(0 \cdot 0 - 1 \cdot 1) + 1(0 \cdot 1 - 1 \cdot 1) = 1$$

Example

For the system in the last example we have three solutions

$$\vec{x}_1 = \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix}, \quad \vec{x}_2 = \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix}, \quad \vec{x}_3 = \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

To show that $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ are linearly independent on $(-\infty, \infty)$ choose a point, say $t_0 = 0$.

Calculate $\vec{x}_1(t_0)$, $\vec{x}_2(t_0)$, and $\vec{x}_3(t_0)$. Then construct the column space matrix:

$$\mathbf{C} = \begin{bmatrix} 0 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

If the determinant is nonzero, then the solutions are linearly independent.

$$|\mathbf{C}| = 0(1 \cdot 0 - 1 \cdot 1) - 2(0 \cdot 0 - 1 \cdot 1) + 1(0 \cdot 1 - 1 \cdot 1) = 1$$

So, the general solution is $\vec{x} = c_1\vec{x}_1 + c_2\vec{x}_2 + c_3\vec{x}_3$.

We have a few ways to express solutions:

$$\vec{x} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_3 \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix}$$

We have a few ways to express solutions:

$$\vec{x} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_3 \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix} = \begin{bmatrix} 0 & 2e^{2t} & e^t \\ 0 & e^{2t} & e^t \\ e^{3t} & e^{2t} & 0 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

We have a few ways to express solutions:

$$\vec{x} = c_1 \begin{bmatrix} 0 \\ 0 \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{2t} \\ e^{2t} \\ e^{2t} \end{bmatrix} + c_3 \begin{bmatrix} e^t \\ e^t \\ 0 \end{bmatrix} = \begin{bmatrix} 0 & 2e^{2t} & e^t \\ 0 & e^{2t} & e^t \\ e^{3t} & e^{2t} & 0 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

Fundamental Matrix

For a basis of n linearly independent solutions of $\vec{x}' = \mathbf{A}\vec{x}$, the matrix $\mathbf{X}(t)$ whose *columns* are the vector solutions $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ is called the **fundamental matrix** for the system.

$$\vec{x} = \underbrace{\begin{bmatrix} | & | & \cdots & | \\ \vec{x}_1 & \vec{x}_2 & \cdots & \vec{x}_n \\ | & | & \cdots & | \end{bmatrix}}_{\mathbf{X}(t)} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \quad c_1, c_2, c_3 \in \mathbb{R}$$

Graphical Views

- The t - x and t - y graphs showing the individual solution functions $x(t)$ and $y(t)$ are called **component graphs**, **solution graphs**, or **time series**.
- The x - y graph is the **phase plane**. The **trajectories** in the phase plane are the parametric curves described by $x(t)$ and $y(t)$.

Trajectories on a phase plane create a **phase portrait**.

Example

The familiar equation

$$x'' + 0.1x = 0$$

Example

The familiar equation

$$x'' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Example

The familiar equation

$$x'' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Any version of the these equations produces solutions of the form


$$x(t) = c_1 \cos(t\sqrt{0.1}) + c_2 \sin(t\sqrt{0.1})$$

$$y(t) = x'(t) = -c_1\sqrt{0.1} \sin(t\sqrt{0.1}) + c_2\sqrt{0.1} \cos(t\sqrt{0.1})$$

Example

The familiar equation

$$x'' + 0.1x = 0$$



Example

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$

Example

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x - 0.05y \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & -0.05 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Example

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$

can be written in the system form as

$$\begin{aligned} x' &= y \\ y' &= -0.1x - 0.05y \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & -0.05 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$



With solutions of the (approximate) form

$$\begin{aligned} x(t) &\approx e^{-0.025t} (c_1 \cos(0.32t) + c_2 \sin(0.32t)) \\ y(t) &\approx e^{-0.025t} (-0.32c_1 \sin(0.32t) + 0.32c_2 \cos(0.23t)) \\ &\quad - 0.025e^{-0.025t} (c_1 \cos(0.32t) + c_2 \sin(0.32t)) \end{aligned}$$

Example

The second-order DE

$$x'' + 0.05x' + 0.1x = 0$$



Example

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

Example

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

This DE represents a periodically forced harmonic oscillator and has system form:

$$\begin{aligned} x' &= y \\ y' &= -0.1x + 0.5 \cos(t) \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 0 \\ 0.5 \cos(t) \end{bmatrix}$$

Example

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

This DE represents a periodically forced harmonic oscillator and has system form:

$$\begin{aligned} x' &= y \\ y' &= -0.1x + 0.5 \cos(t) \end{aligned} \quad \text{or} \quad \begin{bmatrix} x \\ y \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ -0.1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 0 \\ 0.5 \cos(t) \end{bmatrix}$$

It is not easy to find an analytic solution to this DE, but we can draw the solutions using numerical calculations.

We can use Euler's method, which we have seen before.

Example

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$

$$x_{n+1} = x_n + h \cdot x'(t_n) = x_n + h \cdot y_n$$

$$y_{n+1} = y_n + h \cdot y'(t_n) = y_n + h \cdot (-0.1x + 0.5 \cos(t_n))$$



With step size $h = 0.1$, $x(0) = 1$, and $y(0) = 0$.

t_n	x_n	y_n	x'	y'
0.0	1.0000	0.0000	0.0000	0.4000
0.1	1.0000	0.0400	0.0400	0.3975
0.2	1.0040	0.0798	0.0798	0.3896
0.3	1.0120	0.1187	0.1187	0.3765
0.4	1.0238	0.1564	0.1564	0.3581
0.5	1.0395	0.1922	0.1922	0.3348
0.6	1.0587	0.2257	0.2257	0.3068
0.7	1.0813	0.2563	0.2563	0.2743
\vdots	\vdots	\vdots	\vdots	\vdots

Example

Let us consider a nonautonomous version of



$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$



Example

Let us consider a nonautonomous version of



$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$



Example

Let us consider a nonautonomous version of

$$x'' + 0.1x = 0.5 \cos(t) \quad x(0) = 1, \quad x'(0) = 0$$



Graphical Properties of Uniqueness

- For a linear system of differential equations in \mathbb{R}^n , solutions do not cross in t, x_1, x_2, \dots, x_n -space (i.e. \mathbb{R}^{n+1}).

Graphical Properties of Uniqueness

- For a linear system of differential equations in \mathbb{R}^n , solutions do not cross in t, x_1, x_2, \dots, x_n -space (i.e. \mathbb{R}^{n+1}).
- For an *autonomous* linear system in \mathbb{R}^n , trajectories *also* do not cross in x_1, x_2, \dots, x_n -space (i.e. \mathbb{R}^n).