

Linear Equations: The Nature of Their Solutions

Department of Mathematics

Salt Lake Community College

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15$$

$$4x - 2y + 3\sqrt{z} = 12$$

$$2x - 3y + 4z + 3 = w$$

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15 \quad \text{No}$$

$$4x - 2y + 3\sqrt{z} = 12$$

$$2x - 3y + 4z + 3 = w$$

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15 \quad \text{No}$$

$$4x - 2y + 3\sqrt{z} = 12 \quad \text{No}$$

$$2x - 3y + 4z + 3 = w$$

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15 \quad \text{No}$$

$$4x - 2y + 3\sqrt{z} = 12 \quad \text{No}$$

$$2x - 3y + 4z + 3 = w \quad \text{Yes}$$

Definition

A differential equation $F(y, y', y'', \dots, y^{(n)}) = f(t)$ is **linear** if it is of the form

$$a_n(t) \frac{d^n y}{dt^n} + a_{n-1}(t) \frac{d^{n-1} y}{dt^{n-1}} + \cdots + a_1(t) \frac{dy}{dt} + a_0(t)y = f(t)$$

where all functions of t are assumed to be defined over some common interval I .

If $f(t) = 0$ over the interval I , the differential equation is said to be **homogeneous**.

Definition

A differential equation $F(y, y', y'', \dots, y^{(n)}) = f(t)$ is **linear** if it is of the form

$$a_n(t) \frac{d^n y}{dt^n} + a_{n-1}(t) \frac{d^{n-1} y}{dt^{n-1}} + \cdots + a_1(t) \frac{dy}{dt} + a_0(t)y = f(t)$$

where all functions of t are assumed to be defined over some common interval I .

If $f(t) = 0$ over the interval I , the differential equation is said to be **homogeneous**.

First and Second Order Notation

It is common to write first-order differential equations as

$$y' + p(t)y = f(t)$$

and second-order differential equations as

$$y'' + p(t)y' + q(t)y = f(t)$$

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
-----------------------	-------	---------	--------------	--------------

$$y' + ty = 1$$

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes		

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
-----------------------	-------	---------	--------------	--------------

$$y' + ty = 1$$

1

Yes

No

Variable

$$y'' + yy' + y = t$$

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes		

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3	Yes		

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3	Yes	No	

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3	Yes	No	Variable

Notation

We will use a **vector** notation to represent a whole set of variables:

Linear Algebraic Equations:

$$\vec{x} = [x_1, x_2, \dots, x_n]$$

Linear Differential Equations:

$$\vec{y} = [y^{(n)}, y^{(n-1)}, \dots, y', y]$$

Notation

We will use a **vector** notation to represent a whole set of variables:

Linear Algebraic Equations:

$$\vec{x} = [x_1, x_2, \dots, x_n]$$

Linear Differential Equations:

$$\vec{y} = [y^{(n)}, y^{(n-1)}, \dots, y', y]$$

Definition

A **linear operator** L is an entire operation performed on a set of variables.

Linear Algebraic Equations:

$$L(\vec{x}) = a_1x_1 + a_2x_2 + \dots + a_nx_n$$

Linear Differential Equations:

$$L(\vec{y}) = a_n(t) \frac{d^n y}{dt^n} + a_{n-1}(t) \frac{d^{n-1} y}{dt^{n-1}} + \dots + a_1(t) \frac{dy}{dt} + a_0(t)y$$

Example 3

What is the linear operator for the following linear differential equations?

$$y' + ty = 1$$

Example 3

What is the linear operator for the following linear differential equations?

$$y' + ty = 1 \quad \rightarrow \quad L(\vec{y}) = y' + ty$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{l} y' + ty = 1 \\ y'' + 2y = 3y' + t \end{array} \quad \rightarrow \quad L(\vec{y}) = y' + ty$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{l} y' + ty = 1 \\ y'' - 3y' + 2y = t \end{array} \quad \rightarrow \quad L(\vec{y}) = y' + ty$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \end{array}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \\ y^{(4)} + 3y = \sin(t) & \end{array}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \\ y^{(4)} + 3y = \sin(t) & \rightarrow L(\vec{y}) = y^{(4)} + 3y \end{array}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{aligned}y' + ty = 1 &\quad \rightarrow \quad L(\vec{y}) = y' + ty \\y'' - 3y' + 2y = t &\quad \rightarrow \quad L(\vec{y}) = y'' - 3y' + 2y \\y^{(4)} + 3y = \sin(t) &\quad \rightarrow \quad L(\vec{y}) = y^{(4)} + 3y\end{aligned}$$

Linear Operator Properties

$$\begin{aligned}L(k\vec{u}) &= kL(\vec{u}), \quad k \in \mathbb{R} \\L(\vec{u} + \vec{w}) &= L(\vec{u}) + L(\vec{w})\end{aligned}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{aligned}y' + ty &= 1 & \rightarrow & L(\vec{y}) = y' + ty \\y'' - 3y' + 2y &= t & \rightarrow & L(\vec{y}) = y'' - 3y' + 2y \\y^{(4)} + 3y &= \sin(t) & \rightarrow & L(\vec{y}) = y^{(4)} + 3y\end{aligned}$$

Linear Operator Properties

$$\begin{aligned}L(k\vec{u}) &= kL(\vec{u}), \quad k \in \mathbb{R} \\L(\vec{u} + \vec{w}) &= L(\vec{u}) + L(\vec{w})\end{aligned}$$

Proof

The properties can be proved directly for algebraic operators.

For differential operators, the proof follows from the derivative properties:

- $(kf)' = kf'$
- $(f + g)' = f' + g'$

Superposition Principle for Linear Homogeneous Equations

Let \vec{u}_1 and \vec{u}_2 be any solutions of the *homogeneous linear* equation

$$L(\vec{u}) = 0$$

- The sum $\vec{u} = \vec{u}_1 + \vec{u}_2$ is also a solution.
- For any constant k , $\vec{u} = k\vec{u}_1$ is also a solution.

Superposition Principle for Linear Homogeneous Equations

Let \vec{u}_1 and \vec{u}_2 be any solutions of the *homogeneous linear* equation

$$L(\vec{u}) = 0$$

- The sum $\vec{u} = \vec{u}_1 + \vec{u}_2$ is also a solution.
- For any constant k , $\vec{u} = k\vec{u}_1$ is also a solution.

Proof

The proof of the Superposition Principle follows directly from the properties of linear operators from the previous slides.

$$L(\vec{u}) = L(\vec{u}_1 + \vec{u}_2) = L(\vec{u}_1) + L(\vec{u}_2) = 0 + 0 = 0$$

$$L(\vec{u}) = L(k\vec{u}_1) = kL(\vec{u}_1) = k \cdot 0 = 0$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

$$y'' - 4y = (8e^{2t} + 12e^{-2t}) - 4(2e^{2t} + 3e^{-2t})$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

$$\begin{aligned} y'' - 4y &= (8e^{2t} + 12e^{-2t}) - 4(2e^{2t} + 3e^{-2t}) \\ &= 8e^{2t} + 12e^{-2t} - 8e^{2t} - 12e^{-2t} \end{aligned}$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

$$\begin{aligned} y'' - 4y &= (8e^{2t} + 12e^{-2t}) - 4(2e^{2t} + 3e^{-2t}) \\ &= 8e^{2t} + 12e^{-2t} - 8e^{2t} - 12e^{-2t} = 0 \end{aligned}$$

Nonhomogeneous Principle

Let \vec{u}_p be any solution (called a particular solution) to *linear nonhomogeneous* equation

$$L(\vec{u}) = C \quad (\text{algebraic})$$

or

$$L(\vec{u}) = f(t) \quad (\text{differential})$$

Then,

$$\vec{u} = \vec{u}_h + \vec{u}_p$$

is also a solution, here \vec{u}_h is a solution to the **associated homogeneous** equation

$$L(\vec{u}) = 0$$

Furthermore, *every solution of the nonhomogeneous equation must be of the form $\vec{u} = \vec{u}_h + \vec{u}_p$.*

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$L(\vec{u}_q - \vec{u}_p) =$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$L(\vec{u}_q - \vec{u}_p) = L(\vec{u}_q) + L(-\vec{u}_p)$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$\begin{aligned} L(\vec{u}_q - \vec{u}_p) &= L(\vec{u}_q) + L(-\vec{u}_p) \\ &= L(\vec{u}_q) - L(\vec{u}_p) \end{aligned}$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$\begin{aligned} L(\vec{u}_q - \vec{u}_p) &= L(\vec{u}_q) + L(-\vec{u}_p) \\ &= L(\vec{u}_q) - L(\vec{u}_p) \\ &= f(t) - f(t) = 0 \end{aligned}$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$\begin{aligned} L(\vec{u}_q - \vec{u}_p) &= L(\vec{u}_q) + L(-\vec{u}_p) \\ &= L(\vec{u}_q) - L(\vec{u}_p) \\ &= f(t) - f(t) = 0 \end{aligned}$$

Process for Solving Nonhomogeneous Linear Equations

Step 1: Find all solutions \vec{u}_h of $L(\vec{u}) = 0$.

Step 2: Find any solution \vec{u}_p of $L(\vec{u}) = f$.

Step 3: Add $\vec{u}_h + \vec{u}_p = \vec{u}$ to find all solutions of $L(\vec{u}) = f$.

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1:

Step 2:

Step 3:

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1: Solve the associated homogeneous equation $y' - y = 0$, or $y' = y$. (Note: first-order homogeneous linear differential equations are always separable.)

$$y_h = ce^t, \quad \text{for any } c \in \mathbb{R}$$

Step 2:

Step 3:

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1: Solve the associated homogeneous equation $y' - y = 0$, or $y' = y$. (Note: first-order homogeneous linear differential equations are always separable.)

$$y_h = ce^t, \quad \text{for any } c \in \mathbb{R}$$

Step 2: We can verify by differentiation and substitution that $y_p = -t - 1$ is a particular solution.

Step 3:

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1: Solve the associated homogeneous equation $y' - y = 0$, or $y' = y$. (Note: first-order homogeneous linear differential equations are always separable.)

$$y_h = ce^t, \quad \text{for any } c \in \mathbb{R}$$

Step 2: We can verify by differentiation and substitution that $y_p = -t - 1$ is a particular solution.

Step 3: Superposition tells us that


$$y = y_h + y_p = ce^t - t - 1$$

is a solution for any $c \in \mathbb{R}$.

Example 6


Consider

$$y' - y = t$$




$\{y_h\}$

+



y_p

=



$\{y_h\} + y_p$

Note

Many of you may be wondering how exactly we go about finding particular solutions?

Note

Many of you may be wondering how exactly we go about finding particular solutions?

In the next section we will start talking about different methods to do just that.

Note

Many of you may be wondering how exactly we go about finding particular solutions?

In the next section we will start talking about different methods to do just that.

But, sometimes a particular solution may be staring us in the face.

Example 7

Let us solve

$$y' + ay = b$$

where a and b are constants.

Step 1:

Step 2:

Step 3:

Example 7

Let us solve

$$y' + ay = b$$

where a and b are constants.

Step 1: The associated homogeneous equation $y' + ay = 0$ will soon become an old friend.

It has the solution $y_h = ce^{-at}$, where $c \in \mathbb{R}$.

Step 2:

Step 3:

Example 7

Let us solve

$$y' + ay = b$$

where a and b are constants.

Step 1: The associated homogeneous equation $y' + ay = 0$ will soon become an old friend.

It has the solution $y_h = ce^{-at}$, where $c \in \mathbb{R}$.

Step 2: Looking at the DE we can see that $y_p = \frac{b}{a}$ will satisfy this equation. (Recall that the derivative of a constant is zero.)

Step 3:

Example 7

Let us solve

$$y' + ay = b$$

where a and b are constants.

Step 1: The associated homogeneous equation $y' + ay = 0$ will soon become an old friend.

It has the solution $y_h = ce^{-at}$, where $c \in \mathbb{R}$.

Step 2: Looking at the DE we can see that $y_p = \frac{b}{a}$ will satisfy this equation. (Recall that the derivative of a constant is zero.)

Step 3: Superposition tells us that

$$y = y_h + y_p = ce^{-at} + \frac{b}{a}$$

is a solution for any $c \in \mathbb{R}$.

Example 7

Let us solve

$$y' + ay = b$$

where a and b are constants.

Alternatively, we can look for a horizontal line in the direction field.

