

Solution and Direction Fields: Qualitative Analysis

Department of Mathematics

Salt Lake Community College

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

In this chapter we will focus on DEs that can be written as:

$$\frac{dy}{dt} = f(t, y) \quad \text{or} \quad y' = f(t, y)$$

Where the dependent variable y is an unknown function, the **solution**.

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

In this chapter we will focus on DEs that can be written as:

$$\frac{dy}{dt} = f(t, y) \quad \text{or} \quad y' = f(t, y)$$

Where the dependent variable y is an unknown function, the **solution**.

Note

There may be more than one solution for a given differential equation.

A **differential equation** (or DE) is an equation containing derivatives. The **order** of the equation refers to the highest-order derivative that occurs.

In this chapter we will focus on DEs that can be written as:

$$\frac{dy}{dt} = f(t, y) \quad \text{or} \quad y' = f(t, y)$$

Where the dependent variable y is an unknown function, the **solution**.

Note

There may be more than one solution for a given differential equation.

Analytic Definition of a Solution

Analytically, $y(t)$ is a **solution** of a differential equation if substituting $y(t)$ for y reduced the equation to an identity:

$$y'(t) = f(t, y(t))$$

on an appropriate domain for t .

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$y'(t) = \frac{d}{dt}e^{2t}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \end{aligned}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \\ &= 2y(t) \end{aligned}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \\ &= 2y(t) \\ &= f(t, y(t)) \end{aligned}$$

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

Substituting into the DE gives:

$$\begin{aligned} y'(t) &= \frac{d}{dt} e^{2t} \\ &= 2e^{2t} \\ &= 2y(t) \\ &= f(t, y(t)) \end{aligned}$$

Similarly, we could show that

$$y(t) = 2e^{2t} \quad \text{and} \quad y(t) = \frac{-3}{2}e^{2t}$$

are also solutions. In fact, any constant multiple of e^{2t} is a solution.

Example 1

Verify that $y(t)$ is a solution to the DE.

$$y'(t) = 2y, \quad y(t) = e^{2t}$$

