

Matrix Algebra

Department of Mathematics

Salt Lake Community College

Matrix

A **matrix** is a rectangular array of **elements** or **entries** (numbers or functions) arranged in **rows** (horizontal) and **columns** (vertical).

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

The **order** of \mathbf{A} is $m \times n$. If $m = n$, we call the matrix **square**.

Matrix

A **matrix** is a rectangular array of **elements** or **entries** (numbers or functions) arranged in **rows** (horizontal) and **columns** (vertical).

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

The **order** of \mathbf{A} is $m \times n$. If $m = n$, we call the matrix **square**.

Equal Matrices

Two matrices of the same order are **equal** if their corresponding entries are equal. If matrices $\mathbf{A} = [a_{ij}]$ and $\mathbf{B} = [b_{ij}]$ are both $m \times n$, then

$$\mathbf{A} = \mathbf{B} \Leftrightarrow a_{ij} = b_{ij}, \quad 1 \leq i \leq m, 1 \leq j \leq n$$

Special Matrices

- The $m \times n$ **zero matrix**, denoted $\mathbf{0}_{mn}$, has all its entries equal to zero.

Special Matrices

- The $m \times n$ **zero matrix**, denoted $\mathbf{0}_{mn}$, has all its entries equal to zero.
- A **diagonal matrix** is:

$$\mathbf{D} = \begin{bmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{mn} \end{bmatrix}$$

Special Matrices

- The $m \times n$ **zero matrix**, denoted $\mathbf{0}_{mn}$, has all its entries equal to zero.
- A **diagonal matrix** is:

$$\mathbf{D} = \begin{bmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix}$$

- The $n \times n$ **identity matrix**, denoted \mathbf{I}_n is:

$$\mathbf{I}_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Matrix Addition

Two matrices of the same order are added (or subtracted) by adding (or subtracting) corresponding entries and recording the results in a matrix of the same size. Using matrix notation, if $A = [a_{ij}]$ and $B = [b_{ij}]$ are both $m \times n$.

$$\mathbf{A} + \mathbf{B} = [a_{ij}] + [b_{ij}] = [a_{ij} + b_{ij}]$$

$$\mathbf{A} - \mathbf{B} = [a_{ij}] - [b_{ij}] = [a_{ij} - b_{ij}]$$

Matrix Addition

Two matrices of the same order are added (or subtracted) by adding (or subtracting) corresponding entries and recording the results in a matrix of the same size. Using matrix notation, if $A = [a_{ij}]$ and $B = [b_{ij}]$ are both $m \times n$.

$$\mathbf{A} + \mathbf{B} = [a_{ij}] + [b_{ij}] = [a_{ij} + b_{ij}]$$

$$\mathbf{A} - \mathbf{B} = [a_{ij}] - [b_{ij}] = [a_{ij} - b_{ij}]$$

Multiplication by a Scalar

To find the product of a matrix and a scalar (a complex number), multiply each entry of the matrix by that number. This is called **multiplication by a scalar**. Using matrix notation, if $\mathbf{A} = [a_{ij}]$, then

$$c \cdot \mathbf{A} = [c \cdot a_{ij}] = [a_{ij} \cdot c] = \mathbf{A} \cdot c$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix}$$

$$\mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix}$$

$$\mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix}$$

$$\mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix}$$

$$\mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3 + 4 & 1 + 1 & 5 + 0 \\ -2 + 8 & 0 + 1 & 6 + -3 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

What is $3\mathbf{A} - 2\mathbf{B}$?

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

What is $3\mathbf{A} - 2\mathbf{B}$?

$$3 \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

What is $3\mathbf{A} - 2\mathbf{B}$?

$$\begin{bmatrix} 3 \cdot 3 & 3 \cdot 1 & 3 \cdot 5 \\ 3 \cdot (-2) & 3 \cdot 0 & 3 \cdot 6 \end{bmatrix} - \begin{bmatrix} 2 \cdot 4 & 2 \cdot 1 & 2 \cdot 0 \\ 2 \cdot 8 & 2 \cdot 1 & 2 \cdot (-3) \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

What is $3\mathbf{A} - 2\mathbf{B}$?

$$\begin{bmatrix} 9 & 3 & 15 \\ -6 & 0 & 18 \end{bmatrix} - \begin{bmatrix} 8 & 2 & 0 \\ 16 & 2 & -6 \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

What is $3\mathbf{A} - 2\mathbf{B}$?

$$\begin{bmatrix} 9-8 & 3-2 & 15-0 \\ (-6)-16 & 0-2 & 18-(-6) \end{bmatrix}$$

Example 1

Suppose that

$$\mathbf{A} = \begin{bmatrix} 3 & 1 & 5 \\ -2 & 0 & 6 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 4 & 1 & 0 \\ 8 & 1 & -3 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 9 & 0 \\ -3 & 6 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{B}$?

$$\begin{bmatrix} 3+4 & 1+1 & 5+0 \\ -2+8 & 0+1 & 6+(-3) \end{bmatrix} = \begin{bmatrix} 7 & 2 & 5 \\ 6 & 1 & 3 \end{bmatrix}$$

What is $\mathbf{A} + \mathbf{C}$?

\mathbf{A} and \mathbf{C} have different dimensions, so the sum cannot be performed.

What is $3\mathbf{C}$?

$$\begin{bmatrix} 3 \cdot 9 & 3 \cdot 0 \\ 3 \cdot (-3) & 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ -9 & 18 \end{bmatrix}$$

What is $3\mathbf{A} - 2\mathbf{B}$?

$$\begin{bmatrix} 1 & 1 & 15 \\ -22 & -2 & 24 \end{bmatrix}$$

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)
- $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$ (Associativity)

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)
- $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$ (Associativity)
- $c(k\mathbf{A}) = (ck)\mathbf{A}$ (Associativity)

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)
- $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$ (Associativity)
- $c(k\mathbf{A}) = (ck)\mathbf{A}$ (Associativity)
- $\mathbf{A} + \mathbf{0} = \mathbf{A}$ (Zero Element)

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)
- $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$ (Associativity)
- $c(k\mathbf{A}) = (ck)\mathbf{A}$ (Associativity)
- $\mathbf{A} + \mathbf{0} = \mathbf{A}$ (Zero Element)
- $\mathbf{A} + (-\mathbf{A}) = \mathbf{0}$ (Inverse Element)

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)
- $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$ (Associativity)
- $c(k\mathbf{A}) = (ck)\mathbf{A}$ (Associativity)
- $\mathbf{A} + \mathbf{0} = \mathbf{A}$ (Zero Element)
- $\mathbf{A} + (-\mathbf{A}) = \mathbf{0}$ (Inverse Element)
- $c(\mathbf{A} + \mathbf{B}) = c\mathbf{A} + c\mathbf{B}$ (Distributivity)

Properties of Matrix Addition and Scalar Multiplication

Suppose \mathbf{A} , \mathbf{B} , and \mathbf{C} are $m \times n$ matrices and c and k are scalars. Then the following properties hold:

- $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (Commutativity)
- $\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$ (Associativity)
- $c(k\mathbf{A}) = (ck)\mathbf{A}$ (Associativity)
- $\mathbf{A} + \mathbf{0} = \mathbf{A}$ (Zero Element)
- $\mathbf{A} + (-\mathbf{A}) = \mathbf{0}$ (Inverse Element)
- $c(\mathbf{A} + \mathbf{B}) = c\mathbf{A} + c\mathbf{B}$ (Distributivity)
- $(c + k)\mathbf{A} = c\mathbf{A} + k\mathbf{A}$ (Distributivity)

Vectors

A vector $\vec{v} = \langle v_1, \dots, v_n \rangle$ can be represented by either by a $1 \times n$ row matrix, or a $n \times 1$ column matrix.

Vectors

A vector $\vec{v} = \langle v_1, \dots, v_n \rangle$ can be represented by either by a $1 \times n$ row matrix, or a $n \times 1$ column matrix.

Vector addition and Scalar Multiplication

Let

$$\vec{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \quad \text{and} \quad \vec{y} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$$

be vectors in \mathbb{R}^n and c be any scalar. Then, we have:

$$\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{bmatrix} \quad \text{and} \quad c \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c \cdot x_1 \\ \vdots \\ c \cdot x_n \end{bmatrix}$$

Properties of Vector Addition and Multiplication

For vectors \vec{u} , \vec{v} , and \vec{w} in \mathbb{R}^n and scalars c and k .

- $\vec{u} + \vec{v} = \vec{v} + \vec{u}$ (Commutativity)
- $\vec{u} + (\vec{v} + \vec{w}) = (\vec{u} + \vec{v}) + \vec{w}$ (Associativity)
- $c(k\vec{v}) = (ck)\vec{v}$ (Associativity)
- $\vec{u} + \vec{0} = \vec{u}$ (Zero Element)
- $\vec{u} + (-\vec{u}) = \vec{0}$ (Inverse Element)
- $c(\vec{u} + \vec{v}) = c\vec{u} + c\vec{v}$ (Distributivity)
- $(c + k)\vec{u} = c\vec{u} + k\vec{u}$ (Distributivity)

Dot Product

The **dot product** of a row vector \vec{x} and a column vector \vec{y} of equal length n is the result of adding the products of the corresponding entries as follows:

$$\vec{x} \cdot \vec{y} = [x_1 \quad \cdots \quad x_n] \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$$

$$= x_1 \cdot y_1 + x_2 \cdot y_2 + \cdots + x_n \cdot y_n$$

Dot Product

The **dot product** of a row vector \vec{x} and a column vector \vec{y} of equal length n is the result of adding the products of the corresponding entries as follows:

$$\begin{aligned}\vec{x} \cdot \vec{y} &= [x_1 \quad \cdots \quad x_n] \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \\ &= x_1 \cdot y_1 + x_2 \cdot y_2 + \cdots + x_n \cdot y_n\end{aligned}$$

Example 2

Consider

$$\vec{r} = [3 \quad -5 \quad 2] \quad \text{and} \quad \vec{c} = \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix}$$

What is $\vec{r} \cdot \vec{c}$?

Dot Product

The **dot product** of a row vector \vec{x} and a column vector \vec{y} of equal length n is the result of adding the products of the corresponding entries as follows:

$$\begin{aligned}\vec{x} \cdot \vec{y} &= [x_1 \quad \cdots \quad x_n] \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \\ &= x_1 \cdot y_1 + x_2 \cdot y_2 + \cdots + x_n \cdot y_n\end{aligned}$$

Example 2

Consider

$$\vec{r} = [3 \quad -5 \quad 2] \quad \text{and} \quad \vec{c} = \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix}$$

What is $\vec{r} \cdot \vec{c}$?

$$[3 \quad -5 \quad 2] \cdot \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix}$$

Dot Product

The **dot product** of a row vector \vec{x} and a column vector \vec{y} of equal length n is the result of adding the products of the corresponding entries as follows:

$$\begin{aligned}\vec{x} \cdot \vec{y} &= [x_1 \quad \cdots \quad x_n] \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \\ &= x_1 \cdot y_1 + x_2 \cdot y_2 + \cdots + x_n \cdot y_n\end{aligned}$$

Example 2

Consider

$$\vec{r} = [3 \quad -5 \quad 2] \quad \text{and} \quad \vec{c} = \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix}$$

What is $\vec{r} \cdot \vec{c}$?

$$[3 \quad -5 \quad 2] \cdot \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix} = 3 \cdot 3 + (-5) \cdot 4 + 2 \cdot (-5)$$

Dot Product

The **dot product** of a row vector \vec{x} and a column vector \vec{y} of equal length n is the result of adding the products of the corresponding entries as follows:

$$\begin{aligned}\vec{x} \cdot \vec{y} &= [x_1 \quad \cdots \quad x_n] \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \\ &= x_1 \cdot y_1 + x_2 \cdot y_2 + \cdots + x_n \cdot y_n\end{aligned}$$

Example 2

Consider

$$\vec{r} = [3 \quad -5 \quad 2] \quad \text{and} \quad \vec{c} = \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix}$$

What is $\vec{r} \cdot \vec{c}$?

$$[3 \quad -5 \quad 2] \cdot \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix} = 3 \cdot 3 + (-5) \cdot 4 + 2 \cdot (-5) = 9 - 20 - 10$$

Dot Product

The **dot product** of a row vector \vec{x} and a column vector \vec{y} of equal length n is the result of adding the products of the corresponding entries as follows:

$$\begin{aligned}\vec{x} \cdot \vec{y} &= [x_1 \quad \cdots \quad x_n] \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \\ &= x_1 \cdot y_1 + x_2 \cdot y_2 + \cdots + x_n \cdot y_n\end{aligned}$$

Example 2

Consider

$$\vec{r} = [3 \quad -5 \quad 2] \quad \text{and} \quad \vec{c} = \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix}$$

What is $\vec{r} \cdot \vec{c}$?

$$[3 \quad -5 \quad 2] \cdot \begin{bmatrix} 3 \\ 4 \\ -5 \end{bmatrix} = 3 \cdot 3 + (-5) \cdot 4 + 2 \cdot (-5) = 9 - 20 - 10 = -21$$

Matrix Product

The **matrix product** of a $m \times r$ matrix **A** and a $r \times n$ matrix **B** is denoted

$$\mathbf{C} = \mathbf{A} \cdot \mathbf{B} = \mathbf{AB}$$

where the ij th entry of **C** is the dot product of the i th row vector of **A** and the j th column vector of **B**:

$$c_{ij} = [a_{i1} \quad a_{i2} \quad \cdots \quad a_{ir}] \cdot \begin{bmatrix} b_{1j} \\ \vdots \\ b_{rj} \end{bmatrix}$$

The matrix **C** has order $m \times n$.

Example 3

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix}$$

Example 3

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix} \\ \hline \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} & \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix} \end{array}$$

Example 3

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix} \\ \hline \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} & \begin{bmatrix} -2 & \end{bmatrix} \end{array}$$

Example 3

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix} \\ \hline \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} & \begin{bmatrix} -2 & 5 \end{bmatrix} \end{array}$$

Example 3

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix} \\ \hline \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} & \begin{bmatrix} -2 & 5 \\ 6 & \end{bmatrix} \end{array}$$

Example 3

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 3 & 1 \\ 2 & -4 \\ -1 & 0 \end{bmatrix} \\ \hline \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & 2 \end{bmatrix} & \begin{bmatrix} -2 & 5 \\ 6 & -16 \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} & & & \\ & & & \\ & & & \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} 23 & & & \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} 23 & 41 & & \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} 23 & 41 & 4 & \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} 23 & 41 & 4 & 33 \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} 23 & 41 & 4 & 33 \\ 42 & & & \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

$$\begin{array}{c|c} & \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} & \begin{bmatrix} 23 & 41 & 4 & 33 \\ 42 & 89 & & \end{bmatrix} \end{array}$$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

	$\begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$
$\begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix}$	$\begin{bmatrix} 23 & 41 & 4 & 33 \\ 42 & 89 & 5 & \end{bmatrix}$

Example 4

Perform \mathbf{AB} where

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$$

	$\begin{bmatrix} 2 & 5 & 1 & 4 \\ 4 & 8 & 0 & 6 \\ -3 & 1 & -2 & -1 \end{bmatrix}$
$\begin{bmatrix} 2 & 4 & -1 \\ 5 & 8 & 0 \end{bmatrix}$	$\begin{bmatrix} 23 & 41 & 4 & 33 \\ 42 & 89 & 5 & 68 \end{bmatrix}$

Properties of Matrix Multiplication

- $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$ (Associativity)
- $\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$ (Distributivity)
- $(\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA}$ (Distributivity)

Properties of Matrix Multiplication

- $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$ (Associativity)
- $\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$ (Distributivity)
- $(\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA}$ (Distributivity)
- $\mathbf{AB} \neq \mathbf{BA}$ (Generally Noncommutative)

Properties of Matrix Multiplication

- $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$ (Associativity)
- $\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$ (Distributivity)
- $(\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA}$ (Distributivity)
- $\mathbf{AB} \neq \mathbf{BA}$ (Generally Noncommutative)

Properties of Identity Matrices

For a $m \times n$ matrix \mathbf{A} :

- $\mathbf{A} \cdot \mathbf{I}_n = \mathbf{A}$ and $\mathbf{I}_m \cdot \mathbf{A} = \mathbf{A}$
- $\mathbf{A} \cdot \mathbf{0}_n = \mathbf{0}_{mn}$ and $\mathbf{0}_m \cdot \mathbf{A} = \mathbf{0}_{mn}$