

Basic Concepts of Probability

Colby Community College

Definition

The result of an procedure is called an **outcome**.

Definition

The result of an procedure is called an **outcome**.

Example 1

Rolling a die is an procedure, where the outcome is the number rolled.

Definition

The result of an procedure is called an **outcome**.

Example 1

Rolling a die is an procedure, where the outcome is the number rolled.

Definition

An **event** is any particular outcome or group of outcomes.

A **simple event** is an event that cannot be broken down further.

Definition

The result of an procedure is called an **outcome**.

Example 1

Rolling a die is an procedure, where the outcome is the number rolled.

Definition

An **event** is any particular outcome or group of outcomes.

A **simple event** is an event that cannot be broken down further.

Example 2

Drawing a hand for poker is an event, where the outcome is the hand of cards you drew.

This is not a simple event, since we could break it down to five individual card draws. Where each card draw would be a simple event.

Definition

The **sample space** of an procedure is the set of all possible simple events.

Definition

The **sample space** of an procedure is the set of all possible simple events.

Example 3

If we roll a standard 6-sided die, what is the sample space?

Definition

The **sample space** of an procedure is the set of all possible simple events.

Example 3

If we roll a standard 6-sided die, what is the sample space?

The sample space is $\{1, 2, 3, 4, 5, 6\}$.

Definition

The **sample space** of an procedure is the set of all possible simple events.

Example 3

If we roll a standard 6-sided die, what is the sample space?

The sample space is $\{1, 2, 3, 4, 5, 6\}$.

Classical Definition

Given that all out outcomes are equally likely, we can compute the probability of an event E using the formula

$$P(E) = \frac{\text{Number of outcomes corresponding to the event } E}{\text{Total number of outcomes}}$$

Example 4

What is the probability we will roll a 1 on a 6-sided die?

Example 4

What is the probability we will roll a 1 on a 6-sided die?

There is only one outcome to “rolling a 1”, so

$$P(\text{rolling a 1}) = \frac{1}{6}$$

Example 4

What is the probability we will roll a 1 on a 6-sided die?

There is only one outcome to “rolling a 1”, so

$$P(\text{rolling a 1}) = \frac{1}{6}$$

Example 5

What is the probability we will roll better than a 4 on a 6-sided die?

Example 4

What is the probability we will roll a 1 on a 6-sided die?

There is only one outcome to “rolling a 1”, so

$$P(\text{rolling a 1}) = \frac{1}{6}$$

Example 5

What is the probability we will roll better than a 4 on a 6-sided die?

There are two outcomes bigger than 4, so

$$P(\text{roll better than a 4}) = \frac{2}{6} = \frac{1}{3}$$

Example 4

What is the probability we will roll a 1 on a 6-sided die?

There is only one outcome to “rolling a 1”, so

$$P(\text{rolling a 1}) = \frac{1}{6}$$

Example 5

What is the probability we will roll better than a 4 on a 6-sided die?

There are two outcomes bigger than 4, so

$$P(\text{roll better than a 4}) = \frac{2}{6} = \frac{1}{3}$$

Note

Probabilities are ratios, and so can be reduced like fractions.

Example 6

Suppose you have a bag with 20 cherries, 14 sweet and 6 sour. If you pick a cherry at random, what is the probability that it will sweet?

Example 6

Suppose you have a bag with 20 cherries, 14 sweet and 6 sour. If you pick a cherry at random, what is the probability that it will sweet?

The total number of outcomes is 20, one for each cherry in the bag. 14 of those outcomes are sweet cherries, so the probability is

$$P(\text{sweet cherry}) = \frac{14}{20} = \frac{7}{10}$$

Example 6

Suppose you have a bag with 20 cherries, 14 sweet and 6 sour. If you pick a cherry at random, what is the probability that it will sweet?

The total number of outcomes is 20, one for each cherry in the bag. 14 of those outcomes are sweet cherries, so the probability is

$$P(\text{sweet cherry}) = \frac{14}{20} = \frac{7}{10}$$

Note

It is assumed that you cannot differentiate the cherries by touch. If, say, the sweet cherries were smaller than the sour ones, you could always pick a sweet cherry.

Relative Frequency Definition

Conduct an procedure and count the number of times that event A occurs. $P(A)$ is then *approximated* as follows:

$$P(A) \approx \frac{\text{number of times } A \text{ occurs}}{\text{number of times the procedure was repeated}}$$

Relative Frequency Definition

Conduct an procedure and count the number of times that event A occurs. $P(A)$ is then *approximated* as follows:

$$P(A) \approx \frac{\text{number of times } A \text{ occurs}}{\text{number of times the procedure was repeated}}$$

Example 7

To determine the probability that an individual car crashes in a year, we must examine past results to determine the number of cars in use in a year and the number of them that crashed.

Relative Frequency Definition

Conduct an procedure and count the number of times that event A occurs. $P(A)$ is then *approximated* as follows:

$$P(A) \approx \frac{\text{number of times } A \text{ occurs}}{\text{number of times the procedure was repeated}}$$

Example 7

To determine the probability that an individual car crashes in a year, we must examine past results to determine the number of cars in use in a year and the number of them that crashed.

Note

Relative frequency probabilities are often used in place of classical probabilities. The instruction “find the probability” often means “estimate the probability.”

Example 8

In a recent year, there were about 3,000,000 skydiving jumps and 21 of them result in deaths.

Example 8

In a recent year, there were about 3,000,000 skydiving jumps and 21 of them result in deaths.

The relative frequency probability is

$$P(\text{skydiving death}) = \frac{\text{number of skydiving deaths}}{\text{total number of skydiving jumps}}$$

Example 8

In a recent year, there were about 3,000,000 skydiving jumps and 21 of them result in deaths.

The relative frequency probability is

$$\begin{aligned} P(\text{skydiving death}) &= \frac{\text{number of skydiving deaths}}{\text{total number of skydiving jumps}} \\ &= \frac{21}{3,000,000} \end{aligned}$$

Example 8

In a recent year, there were about 3,000,000 skydiving jumps and 21 of them result in deaths.

The relative frequency probability is

$$\begin{aligned} P(\text{skydiving death}) &= \frac{\text{number of skydiving deaths}}{\text{total number of skydiving jumps}} \\ &= \frac{21}{3,000,000} \\ &= 0.000007 = 0.007\% \end{aligned}$$

Example 8

In a recent year, there were about 3,000,000 skydiving jumps and 21 of them result in deaths.

The relative frequency probability is

$$\begin{aligned} P(\text{skydiving death}) &= \frac{\text{number of skydiving deaths}}{\text{total number of skydiving jumps}} \\ &= \frac{21}{3,000,000} \\ &= 0.000007 = 0.007\% \end{aligned}$$

Note

We cannot use the classical definition of probability because the outcomes “dying” and “not dying” are not equally likely.

Law of Large Numbers

As a procedure is repeated again and again, the relative frequency probability of an event tends to approach the actual probability.

Law of Large Numbers

As a procedure is repeated again and again, the relative frequency probability of an event tends to approach the actual probability.

Cautions

- The law of large numbers applies to behavior over a large number of trials, and it does not apply to any one individual outcome.

Law of Large Numbers

As a procedure is repeated again and again, the relative frequency probability of an event tends to approach the actual probability.

Cautions

- The law of large numbers applies to behavior over a large number of trials, and it does not apply to any one individual outcome.
 - Gamblers sometimes foolishly lose large sums of money by incorrectly thinking that a string of losses increases the chances of a win on the next bet, or that a string of wins is likely to continue.

Law of Large Numbers

As a procedure is repeated again and again, the relative frequency probability of an event tends to approach the actual probability.

Cautions

- The law of large numbers applies to behavior over a large number of trials, and it does not apply to any one individual outcome.
 - Gamblers sometimes foolishly lose large sums of money by incorrectly thinking that a string of losses increases the chances of a win on the next bet, or that a string of wins is likely to continue.
- If we know nothing about the likelihood of different possible outcomes, we should not assume that they are equally likely.

Law of Large Numbers

As a procedure is repeated again and again, the relative frequency probability of an event tends to approach the actual probability.

Cautions

- The law of large numbers applies to behavior over a large number of trials, and it does not apply to any one individual outcome.
 - Gamblers sometimes foolishly lose large sums of money by incorrectly thinking that a string of losses increases the chances of a win on the next bet, or that a string of wins is likely to continue.
- If we know nothing about the likelihood of different possible outcomes, we should not assume that they are equally likely.
 - You should not think that the probability of passing the next exam is $\frac{1}{2}$, or 0.5. The actual probability depends on factors such as the amount of preparation and the difficulty of the exam.

Definition

A standard deck of 52 playing cards consists of four **suits** in two colors: Hearts ♥, Spades ♠, Diamonds ♦, and Clubs ♣

Each suit contains 13 cards, each of a different **rank**: Ace, 2 through 10, Jack, Queen, and King.

Definition

A standard deck of 52 playing cards consists of four **suits** in two colors: Hearts ♥, Spades ♠, Diamonds ♦, and Clubs ♣

Each suit contains 13 cards, each of a different **rank**: Ace, 2 through 10, Jack, Queen, and King.

Example 9

Let us compute the probability of drawing a card from a deck and getting an Ace.

Definition

A standard deck of 52 playing cards consists of four **suits** in two colors: Hearts ♥, Spades ♠, Diamonds ♦, and Clubs ♣

Each suit contains 13 cards, each of a different **rank**: Ace, 2 through 10, Jack, Queen, and King.

Example 9

Let us compute the probability of drawing a card from a deck and getting an Ace.

There are four aces in a deck of 52 cards. Which gives the probability

$$P(\text{Ace}) = \frac{4}{52} = \frac{1}{13}$$

Definition

A standard deck of 52 playing cards consists of four **suits** in two colors: Hearts ♥, Spades ♠, Diamonds ♦, and Clubs ♣

Each suit contains 13 cards, each of a different **rank**: Ace, 2 through 10, Jack, Queen, and King.

Example 9

Let us compute the probability of drawing a card from a deck and getting an Ace.

There are four aces in a deck of 52 cards. Which gives the probability

$$P(\text{Ace}) = \frac{4}{52} = \frac{1}{13} = 0.0769 = 7.67\%$$

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

We need to calculate the total number of high school drivers

$$3785 \text{ texted while driving} + 4720 \text{ did not text while driving} = 8505$$

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

We need to calculate the total number of high school drivers

$$3785 \text{ texted while driving} + 4720 \text{ did not text while driving} = 8505$$

We can now calculate the probability that a student texted while driving.

$$P(\text{texting while driving}) = \frac{\text{number of drivers who texted while driving}}{\text{total number of drivers in the sample}}$$

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

We need to calculate the total number of high school drivers

$$3785 \text{ texted while driving} + 4720 \text{ did not text while driving} = 8505$$

We can now calculate the probability that a student texted while driving.

$$\begin{aligned} P(\text{texting while driving}) &= \frac{\text{number of drivers who texted while driving}}{\text{total number of drivers in the sample}} \\ &= \frac{3785}{8505} \end{aligned}$$

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

We need to calculate the total number of high school drivers

$$3785 \text{ texted while driving} + 4720 \text{ did not text while driving} = 8505$$

We can now calculate the probability that a student texted while driving.

$$\begin{aligned} P(\text{texting while driving}) &= \frac{\text{number of drivers who texted while driving}}{\text{total number of drivers in the sample}} \\ &= \frac{3785}{8505} = 0.445 = 44.5\% \end{aligned}$$

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

We need to calculate the total number of high school drivers

$$3785 \text{ texted while driving} + 4720 \text{ did not text while driving} = 8505$$

We can now calculate the probability that a student texted while driving.

$$\begin{aligned} P(\text{texting while driving}) &= \frac{\text{number of drivers who texted while driving}}{\text{total number of drivers in the sample}} \\ &= \frac{3785}{8505} = 0.445 = 44.5\% \end{aligned}$$

Note

A common mistake is to use $\frac{3785}{4720}$ as the probability.

Example 10

In a study of U.S. high school drivers, it was found that 3785 texted while driving during the previous 30 days, and 4720 did not text while driving. What is the probability that a individual student texted while driving?

We need to calculate the total number of high school drivers

$$3785 \text{ texted while driving} + 4720 \text{ did not text while driving} = 8505$$

We can now calculate the probability that a student texted while driving.

$$\begin{aligned} P(\text{texting while driving}) &= \frac{\text{number of drivers who texted while driving}}{\text{total number of drivers in the sample}} \\ &= \frac{3785}{8505} = 0.445 = 44.5\% \end{aligned}$$

Note

A common mistake is to use $\frac{3785}{4720}$ as the probability.

Don't make the common mistake finding a probability by mindlessly dividing a smaller number by a larger number. You need to think carefully about the numbers involved and what they represent.

Definition

An impossible event has probability 0.

A certain event has a probability of 1.

The probability of any event must be $0 \leq P(E) \leq 1$.

Definition

An impossible event has probability 0.

A certain event has a probability of 1.

The probability of any event must be $0 \leq P(E) \leq 1$.

Example 11

If a year is selected at random, find the probability that Thanksgiving Day in the United States will be on a Wednesday.

Definition

An impossible event has probability 0.

A certain event has a probability of 1.

The probability of any event must be $0 \leq P(E) \leq 1$.

Example 11

If a year is selected at random, find the probability that Thanksgiving Day in the United States will be on a Wednesday.

In the United States, Thanksgiving Day always falls on the fourth Thursday in November. This means it is impossible for Thanksgiving Day to fall on a Wednesday.

$$P(\text{Thanksgiving on Wednesday}) = 0$$

$$P(\text{Thanksgiving on Thursday}) = 1$$

Definition

The **complement** of an event E is the event “ E doesn't happen.”

The notation \bar{E} is used to denote the complement of E .

Definition

The **complement** of an event E is the event “ E doesn’t happen.”

The notation \bar{E} is used to denote the complement of E .

Example 12

If we roll a 6-sided die, and

$$P(\text{roll a six}) = \frac{1}{6}$$

The complement is

$$P(\text{roll not a six}) = P(\text{roll a one, two, three, four, or five}) = \frac{5}{6}$$

Note

Either an even occurs or it does not. This means the sum of $P(E)$ and $P(\bar{E})$ always has to equal 1.

In other words,

$$P(\bar{E}) = 1 - P(E) \quad \text{or} \quad P(E) = 1 - P(\bar{E})$$

Note

Either an even occurs or it does not. This means the sum of $P(E)$ and $P(\bar{E})$ always has to equal 1.

In other words,

$$P(\bar{E}) = 1 - P(E) \quad \text{or} \quad P(E) = 1 - P(\bar{E})$$

Example 13

Let us calculate the probability of drawing a card from a deck and not getting a heart.

Note

Either an even occurs or it does not. This means the sum of $P(E)$ and $P(\bar{E})$ always has to equal 1.

In other words,

$$P(\bar{E}) = 1 - P(E) \quad \text{or} \quad P(E) = 1 - P(\bar{E})$$

Example 13

Let us calculate the probability of drawing a card from a deck and not getting a heart.

There are 13 hearts in a deck, giving $P(\text{heart}) = \frac{13}{52} = \frac{1}{4}$.

Note

Either an even occurs or it does not. This means the sum of $P(E)$ and $P(\bar{E})$ always has to equal 1.

In other words,

$$P(\bar{E}) = 1 - P(E) \quad \text{or} \quad P(E) = 1 - P(\bar{E})$$

Example 13

Let us calculate the probability of drawing a card from a deck and not getting a heart.

There are 13 hearts in a deck, giving $P(\text{heart}) = \frac{13}{52} = \frac{1}{4}$.

This means the probability of not drawing a heart is

$$P(\text{not heart}) = 1 - P(\text{heart}) = 1 - \frac{1}{4} = \frac{3}{4}$$

The Rare Even Rule for Inferential Statistics

If, under a given assumption, the probability of a particular observed event is very small and the observed even occurs *significantly less than* or *significantly more than* what we typically expect with that assumption, we conclude that the assumption is probably not correct.

The Rare Even Rule for Inferential Statistics

If, under a given assumption, the probability of a particular observed event is very small and the observed even occurs *significantly less than* or *significantly more than* what we typically expect with that assumption, we conclude that the assumption is probably not correct.

Worded differently:

- x successes among n trials is a **significantly high** number of successes if the probability of x or more successes is unlikely with a probability of 0.05 or less, i.e if $P(x \text{ or more}) \leq 0.05$.
- x successes among n trials is a **significantly low** number of successes if the probability of x or fewer successes is unlikely with a probability of 0.05 or less, i.e if $P(x \text{ or fewer}) \leq 0.05$.

Definition

The **actual odds against** event A occurring are the ratio $\frac{P(\bar{A})}{P(A)}$.

Definition

The **actual odds against** event A occurring are the ratio $\frac{P(\bar{A})}{P(A)}$.

- Usually expressed in the form $a : b$ where a and b are integers. Reduce using the largest common factor; If $a = 16$ and $b = 4$, express the odds as $4 : 1$ instead of $16 : 4$.

Definition

The **actual odds against** event A occurring are the ratio $\frac{P(\bar{A})}{P(A)}$.

- Usually expressed in the form $a : b$ where a and b are integers. Reduce using the largest common factor; If $a = 16$ and $b = 4$, express the odds as $4 : 1$ instead of $16 : 4$.

The **actual odds in favor** of event A occurring are the ratio $\frac{P(A)}{P(\bar{A})}$, which is the reciprocal of the actual odds against.

Definition

The **actual odds against** event A occurring are the ratio $\frac{P(\bar{A})}{P(A)}$.

- Usually expressed in the form $a : b$ where a and b are integers. Reduce using the largest common factor; If $a = 16$ and $b = 4$, express the odds as $4 : 1$ instead of $16 : 4$.

The **actual odds in favor** of event A occurring are the ratio $\frac{P(A)}{P(\bar{A})}$, which is the reciprocal of the actual odds against.

- If the odds against an event are $a : b$, then the odds in favor are $b : a$.

Definition

The **actual odds against** event A occurring are the ratio $\frac{P(\bar{A})}{P(A)}$.

- Usually expressed in the form $a : b$ where a and b are integers. Reduce using the largest common factor; If $a = 16$ and $b = 4$, express the odds as $4 : 1$ instead of $16 : 4$.

The **actual odds in favor** of event A occurring are the ratio $\frac{P(A)}{P(\bar{A})}$, which is the reciprocal of the actual odds against.

- If the odds against an event are $a : b$, then the odds in favor are $b : a$.

The **payoff odds** against event A occurring are the ratio of net profit (if you win) to the amount bet:

$$\text{Payoff odds against event } A = (\text{net profit}) : (\text{amount bet})$$

Definition

The **actual odds against** event A occurring are the ratio $\frac{P(\bar{A})}{P(A)}$.

- Usually expressed in the form $a : b$ where a and b are integers. Reduce using the largest common factor; If $a = 16$ and $b = 4$, express the odds as $4 : 1$ instead of $16 : 4$.

The **actual odds in favor** of event A occurring are the ratio $\frac{P(A)}{P(\bar{A})}$, which is the reciprocal of the actual odds against.

- If the odds against an event are $a : b$, then the odds in favor are $b : a$.

The **payoff odds** against event A occurring are the ratio of net profit (if you win) to the amount bet:

$$\text{Payoff odds against event } A = (\text{net profit}) : (\text{amount bet})$$

Note

The actual odds reflect the actual likelihood of an event. The payoff odds describe the payout amount determined by the casino, lottery, or racetrack operators. (i.e. based on how greedy the operators are.)

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as 35 : 1.

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as 35 : 1.

- What are the actual odd against the outcome of 13?

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as 35 : 1.

- What are the actual odd against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)}$$

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as 35 : 1.

- What are the actual odd against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)} = \frac{37/38}{1/38}$$

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as 35 : 1.

- What are the actual odds against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)} = \frac{37/38}{1/38} = \frac{37}{1}$$

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as $35 : 1$.

- What are the actual odds against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)} = \frac{37/38}{1/38} = \frac{37}{1} \quad \text{or} \quad 37 : 1$$

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as $35 : 1$.

- What are the actual odds against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)} = \frac{37/38}{1/38} = \frac{37}{1} \quad \text{or} \quad 37 : 1$$

- How much profit would a bet of \$5 make, if 13 won?

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as 35 : 1.

- What are the actual odds against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)} = \frac{37/38}{1/38} = \frac{37}{1} \quad \text{or} \quad 37 : 1$$

- How much profit would a bet of \$5 make, if 13 won?

For every \$1 bet, we would win \$5. So, betting \$5 would win $\$5 \cdot 35 = \175 .

Example 14

If you bet \$5 on the number 13 in roulette, your probability of winning is $1/38$, but the payoff odds are given by the casino as $35 : 1$.

- What are the actual odds against the outcome of 13?

$$\text{Actual odds against 13} = \frac{P(\text{not 13})}{P(13)} = \frac{37/38}{1/38} = \frac{37}{1} \quad \text{or} \quad 37 : 1$$

- How much profit would a bet of \$5 make, if 13 won?

For every \$1 bet, we would win \$5. So, betting \$5 would win $\$5 \cdot 35 = \175 .

- If the casino was not run for profit, the payoff odds would be the same as the actual odds in favor, $37 : 1$. So, the profit would be \$185.