

Counting

Colby Community College

Definition

The number of combinations when choosing k objects out of a pool of n objects is:

$${}_nC_k = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Definition

The number of combinations when choosing k objects out of a pool of n objects is:

$${}_nC_k = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Example 1

How many five-card poker hands are there?

Definition

The number of combinations when choosing k objects out of a pool of n objects is:

$${}_nC_k = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Example 1

How many five-card poker hands are there?

$${}_{52}C_5 = \frac{52!}{5!(52-5)!} = 2,598,960$$

Definition

The number of combinations when choosing k objects out of a pool of n objects is:

$${}_nC_k = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Example 1

How many five-card poker hands are there?

$${}_{52}C_5 = \frac{52!}{5!(52-5)!} = 2,598,960$$

Example 2

What is the probability that a poker hand will contain exactly two jacks?

Definition

The number of combinations when choosing k objects out of a pool of n objects is:

$${}_nC_k = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Example 1

How many five-card poker hands are there?

$${}_{52}C_5 = \frac{52!}{5!(52-5)!} = 2,598,960$$

Example 2

What is the probability that a poker hand will contain exactly two jacks?

$$\frac{{}_4C_2 {}_{48}C_3}{{}_{52}C_5} = \frac{6 \cdot 17,296}{2,598,960} = \frac{103,776}{2,598,960} \approx 4\%$$

Example 3

What is the probability that a poker hand will contain a straight?

Example 3

What is the probability that a poker hand will contain a straight?

The lowest ranked straight is A,2,3,4,5 and the highest ranked straight is 10,J,Q,K,A. Thus, for the any of the ten ranks A through 10, we can build a straight. Each card can be from any of the four suits.

This means the probability is:

$$P(\text{straight}) = \frac{10 ({}_4C_1) ({}_4C_1) ({}_4C_1) ({}_4C_1) ({}_4C_1)}{{}_{52}C_5} = \frac{10,240}{2,598,960} \approx 3.94\%$$

Example 3

What is the probability that a poker hand will contain a straight?

The lowest ranked straight is A,2,3,4,5 and the highest ranked straight is 10,J,Q,K,A. Thus, for the any of the ten ranks A through 10, we can build a straight. Each card can be from any of the four suits.

This means the probability is:

$$P(\text{straight}) = \frac{10 ({}_4C_1) ({}_4C_1) ({}_4C_1) ({}_4C_1) ({}_4C_1)}{{}_{52}C_5} = \frac{10,240}{2,598,960} \approx 3.94\%$$

Example 4

What is the probability that a poker hand will contain a straight, but not a straight flush?

Example 3

What is the probability that a poker hand will contain a straight?

The lowest ranked straight is A,2,3,4,5 and the highest ranked straight is 10,J,Q,K,A. Thus, for any of the ten ranks A through 10, we can build a straight. Each card can be from any of the four suits.

This means the probability is:

$$P(\text{straight}) = \frac{10 ({}_4C_1) ({}_4C_1) ({}_4C_1) ({}_4C_1) ({}_4C_1)}{{}_{52}C_5} = \frac{10,240}{2,598,960} \approx 3.94\%$$

Example 4

What is the probability that a poker hand will contain a straight, but not a straight flush?

We calculated in Example 2 the number of straights possible. Now, just need to subtract out the number of straight flushes.

$$P(\text{straight}) = \frac{10,240 - 40}{{}_{52}C_5} = \frac{10,200}{2,598,960} \approx 3.92\%$$

Example 5

In the casino game Roulette, a wheel with 38 spaces (18 red, 18 black, and 2 green) is spun. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

On average, how much money should a player expect to win or lose if they play this game repeatedly?

Example 5

In the casino game Roulette, a wheel with 38 spaces (18 red, 18 black, and 2 green) is spun. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

On average, how much money should a player expect to win or lose if they play this game repeatedly?

Any number you bet on will have the following probabilities:

Outcome	Probability
\$35 (win)	$1/38$
-\$1 (lose)	$37/38$

Example 5

In the casino game Roulette, a wheel with 38 spaces (18 red, 18 black, and 2 green) is spun. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

On average, how much money should a player expect to win or lose if they play this game repeatedly?

Any number you bet on will have the following probabilities:

Outcome	Probability
\$35 (win)	1/38
-\$1 (lose)	37/38

So, **on average**, we will have a net change of

$$\$35 \cdot \frac{1}{38} + -\$1 \cdot \frac{37}{38} = \$0.9211 - \$0.9737 \approx -\$0.053$$

That is, **on average**, we will lose 5.3 cents per space we bet on.