

Properties of Linear Transformations

Department of Mathematics

Salt Lake Community College


(Slides by Adam Wilson)

Injectivity


A function $f : \mathbb{X} \rightarrow \mathbb{Y}$ is **one-to-one**, or **injective**, provided that $f(u) = f(v)$ implies $u = v$.

Injectivity

A function $f : \mathbb{X} \rightarrow \mathbb{Y}$ is **one-to-one**, or **injective**, provided that $f(u) = f(v)$ implies $u = v$.



(a) $f(x) = x^3$ is injective




(b) $g(x) = x^2$ is not injective

Surjectivity


The set of output values of a function $f : \mathbb{X} \rightarrow \mathbb{Y}$ is a subset of the codomain \mathbb{Y} and is called the **image** of the function. If the image is all of \mathbb{Y} , the function f is said to map **onto** \mathbb{Y} , or to be **surjective**.

Surjectivity

The set of output values of a function $f : \mathbb{X} \rightarrow \mathbb{Y}$ is a subset of the codomain \mathbb{Y} and is called the **image** of the function. If the image is all of \mathbb{Y} , the function f is said to map **onto** \mathbb{Y} , or to be **surjective**.



(a) $f(x) = -x$ is surjective



(b) $g(x) = |x|$ is not surjective

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix}$$

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of T ?

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of T ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of T ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \text{ are independent.}$$

So, **Col** \mathbf{A} , the column space of \mathbf{A} , is the image of T .

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of T ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \text{ are independent.}$$

So, **Col** \mathbf{A} , the column space of \mathbf{A} , is the image of T .

We can derive the equation of this plane by looking at the system

$$u_1 = v_1 + v_2, \quad u_2 = v_1 - v_2, \quad u_3 = 2v_1 + v_2$$

Example 1

The linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$ where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of T ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \text{ are independent.}$$

So, **Col A**, the column space of \mathbf{A} , is the image of T .

We can derive the equation of this plane by looking at the system

$$u_1 = v_1 + v_2, \quad u_2 = v_1 - v_2, \quad u_3 = 2v_1 + v_2$$

Eliminating v_1 and v_2 gives us

$$3u_1 + u_2 - 2u_3 = 0$$

Image Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with image $\mathbf{Im}(T)$. Then:

Image Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with image $\mathbf{Im}(T)$. Then:

- $\mathbf{Im}(T)$ is a subspace of \mathbb{W} .

Image Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with image $\mathbf{Im}(T)$. Then:

- $\mathbf{Im}(T)$ is a subspace of \mathbb{W} .
- T is surjective if and only if $\mathbf{Im}(T) = \mathbb{W}$.

Image Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with image $\mathbf{Im}(T)$. Then:

- $\mathbf{Im}(T)$ is a subspace of \mathbb{W} .
- T is surjective if and only if $\mathbf{Im}(T) = \mathbb{W}$.

Rank of a Linear Transformation

The dimension of the image of a linear transformation T is called its **rank**

$$\mathbf{rank}(T) = \mathbf{dim}(\mathbf{Im}(T))$$

Example 2

For $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

Example 2

For $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Example 2

For $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Which means

$$\text{Im}(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \end{bmatrix} \right\}$$

which is a subset of \mathbb{R}^2 .

Example 2

For $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Which means

$$\text{Im}(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \end{bmatrix} \right\}$$

which is a subset of \mathbb{R}^2 . To determine the dimension we need to determine how many pivot columns \mathbf{A} has:

$$\begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

Example 2

For $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Which means

$$\text{Im}(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \end{bmatrix} \right\}$$

which is a subset of \mathbb{R}^2 . To determine the dimension we need to determine how many pivot columns \mathbf{A} has:

$$\begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 0 & 3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Thus, $\text{rank}(T) = \text{dim}(\text{Im}(T)) = \text{dim}(\text{Col } \mathbf{A}) = 2$.

Rank of a Matrix Multiplication Operator

For any linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\vec{x}) = \mathbf{A}\vec{x}$, where $\mathbf{A} \in \mathbb{M}_{mn}$ and $\vec{x} \in \mathbb{V}$, the image of T is the column space of A . (That's is, $\mathbf{Im}(T) = \mathbf{Col} \mathbf{A}$.)

The pivot columns of A form a basis for $\mathbf{Im}(T)$.

Consequently,

$$\begin{aligned}\mathbf{rank}(T) &= \mathbf{dim}(\mathbf{Im}(T)) \\ &= \mathbf{dim}(\mathbf{Col} \mathbf{A}) \\ &= \text{The number of pivot columns in } \mathbf{A}.\end{aligned}$$

Rank of a Matrix Multiplication Operator

For any linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by $T(\vec{x}) = \mathbf{A}\vec{x}$, where $\mathbf{A} \in \mathbb{M}_{mn}$ and $\vec{x} \in \mathbb{V}$, the image of T is the column space of A . (That's is, $\mathbf{Im}(T) = \mathbf{Col} \mathbf{A}$.)

The pivot columns of A form a basis for $\mathbf{Im}(T)$.

Consequently,

$$\begin{aligned}\mathbf{rank}(T) &= \mathbf{dim}(\mathbf{Im}(T)) \\ &= \mathbf{dim}(\mathbf{Col} \mathbf{A}) \\ &= \text{The number of pivot columns in } \mathbf{A}.\end{aligned}$$

Reminder

The basis of $\mathbf{Col} \mathbf{A}$ must come from \mathbf{A} , *not* from the RREF of \mathbf{A} .

Recall

A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ must map the zero vector of \mathbb{V} to the zero vector of \mathbb{W} .

Recall

A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ must map the zero vector of \mathbb{V} to the zero vector of \mathbb{W} .

Kernel of a Linear Transformation

The **kernel** (or **nullspace**) of a linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$, denoted $\ker(T)$, is the set of vectors in \mathbb{V} that are mapped by T to the zero vector of \mathbb{W} .

$$\ker(T) = \left\{ \vec{v} \in \mathbb{V} \mid T(\vec{v}) = \vec{0} \right\}$$

Recall

A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ must map the zero vector of \mathbb{V} to the zero vector of \mathbb{W} .

Kernel of a Linear Transformation

The **kernel** (or **nullspace**) of a linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$, denoted $\ker(T)$, is the set of vectors in \mathbb{V} that are mapped by T to the zero vector of \mathbb{W} .

$$\ker(T) = \left\{ \vec{v} \in \mathbb{V} \mid T(\vec{v}) = \vec{0} \right\}$$

Note

The kernel always contains at least one element.

Example 3

Consider the projection $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by

$$T(x, y, z) = (x, y, 0)$$

What is the kernel of T ?

Example 3

Consider the projection $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by

$$T(x, y, z) = (x, y, 0)$$

What is the kernel of T ?

$$\mathbf{ker}(T) = \{(0, 0, z) \mid z \in \mathbb{R}\}$$

Example 4

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

Example 4

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to $\vec{0}$, we have to solve the system:

$$\begin{array}{rcrcrcrcrcrcl} v_1 & + & v_2 & + & 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + & 5v_3 & = & 0 \end{array}$$

Example 4

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to $\vec{0}$, we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right]$$

Example 4

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to $\vec{0}$, we have to solve the system:

$$\begin{array}{rclcl} v_1 & + & v_2 & + 2v_3 & = 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = 0 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

Example 4

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to $\vec{0}$, we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

So, we see that v_3 is a free variable and if $v_3 = s$ is a parameter, we have $v_1 = -s$, $v_2 = -s$, and $v_3 = s$.

Example 4

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to $\vec{0}$, we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

So, we see that v_3 is a free variable and if $v_3 = s$ is a parameter, we have $v_1 = -s$, $v_2 = -s$, and $v_3 = s$.

Thus, the kernel is any scalar multiple of $\langle -1, -1, 1 \rangle$:

$$\ker(T) = \text{span} \left\{ \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix} \right\}$$

Example 5

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of T ?

Example 5

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Example 5

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right]$$

Example 5

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right]$$

Example 5

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right]$$

Thus,

$$\ker(T) = \{\vec{0}\}$$

Example 6

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of T ?

Example 6

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Example 6

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right]$$

Example 6

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Example 6

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of T ?

To find the kernel we need to solve $A\vec{v} = \vec{0}$:

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{array} \right]$$

In this case we have $v_1 = -\frac{1}{2}v_2$ and so, if we let our parameter be $v_2 = s$ we have

$$\ker(T) = \left\{ s \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right\}$$

Example 7

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Example 7

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

It should be clear that this transformation maps all vectors $\vec{v} \in \mathbb{R}^2$ to $\vec{0}$.

Example 7

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

It should be clear that this transformation maps all vectors $\vec{v} \in \mathbb{R}^2$ to $\vec{0}$.

Thus,

$$\ker(T) = \mathbb{R}^2$$

Example 7

Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

It should be clear that this transformation maps all vectors $\vec{v} \in \mathbb{R}^2$ to $\vec{0}$.

Thus,

$$\ker(T) = \mathbb{R}^2$$

Note

These examples seem to suggest that the kernel of the linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is a subspace of \mathbb{W} .

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\mathbf{ker}(T)$.

Then,

- 1 $\mathbf{ker}(T)$ is a subspace of \mathbb{V} .
- 2 T is injective if and only if $\mathbf{ker}(T) = \{\vec{0}\}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- 1 $\ker(T)$ is a subspace of \mathbb{V} .
- 2 T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify 1, so let us prove 2.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- ① $\ker(T)$ is a subspace of \mathbb{V} .
- ② T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify ①, so let us prove ②.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- ① $\ker(T)$ is a subspace of \mathbb{V} .
- ② T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify ①, so let us prove ②.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Then, for $\vec{w} \in \ker(T)$, we have $T(\vec{w}) = \vec{0}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\mathbf{ker}(T)$.

Then,

- ① $\mathbf{ker}(T)$ is a subspace of \mathbb{V} .
- ② T is injective if and only if $\mathbf{ker}(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify ①, so let us prove ②.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Then, for $\vec{w} \in \mathbf{ker}(T)$, we have $T(\vec{w}) = \vec{0}$.

Which means $T(\vec{w}) = T(\vec{0})$ and hence $\vec{w} = \vec{0}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- ① $\ker(T)$ is a subspace of \mathbb{V} .
- ② T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify ①, so let us prove ②.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Then, for $\vec{w} \in \ker(T)$, we have $T(\vec{w}) = \vec{0}$.

Which means $T(\vec{w}) = T(\vec{0})$ and hence $\vec{w} = \vec{0}$. Thus, $\ker(T) = \{\vec{0}\}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- 1 $\ker(T)$ is a subspace of \mathbb{V} .
- 2 T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify 1, so let us prove 2.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Then, for $\vec{w} \in \ker(T)$, we have $T(\vec{w}) = \vec{0}$.

Which means $T(\vec{w}) = T(\vec{0})$ and hence $\vec{w} = \vec{0}$. Thus, $\ker(T) = \{\vec{0}\}$.

Now, suppose that we know $\ker(T) = \{\vec{0}\}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- 1 $\ker(T)$ is a subspace of \mathbb{V} .
- 2 T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify 1, so let us prove 2.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Then, for $\vec{w} \in \ker(T)$, we have $T(\vec{w}) = \vec{0}$.

Which means $T(\vec{w}) = T(\vec{0})$ and hence $\vec{w} = \vec{0}$. Thus, $\ker(T) = \{\vec{0}\}$.

Now, suppose that we know $\ker(T) = \{\vec{0}\}$.

So $T(\vec{u}) = T(\vec{v})$ implies $T(\vec{u}) - T(\vec{v}) = T(\vec{u} - \vec{v}) = T(\vec{0}) = \vec{0}$.

Kernel Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from vector space \mathbb{V} to vector space \mathbb{W} with kernel $\ker(T)$.

Then,

- 1 $\ker(T)$ is a subspace of \mathbb{V} .
- 2 T is injective if and only if $\ker(T) = \{\vec{0}\}$.

Proof

It is straightforward to verify 1, so let us prove 2.

Suppose T is injective, which means that $T(\vec{u}) = T(\vec{v})$ implies $\vec{u} = \vec{v}$.

Then, for $\vec{w} \in \ker(T)$, we have $T(\vec{w}) = \vec{0}$.

Which means $T(\vec{w}) = T(\vec{0})$ and hence $\vec{w} = \vec{0}$. Thus, $\ker(T) = \{\vec{0}\}$.

Now, suppose that we know $\ker(T) = \{\vec{0}\}$.

So $T(\vec{u}) = T(\vec{v})$ implies $T(\vec{u}) - T(\vec{v}) = T(\vec{u} - \vec{v}) = T(\vec{0}) = \vec{0}$.

Thus, $\vec{u} - \vec{v}$ is in the kernel, which means $\vec{u} - \vec{v} = \vec{0}$ and thus T is injective.

Example 8

Consider the transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

Example 8

Consider the transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[\begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right]$$

Example 8

Consider the transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[\begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

Example 8

Consider the transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[\begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

So, we see that $v_1 = 2v_2 - 3v_4$ and $v_3 = 0$. If we let $v_2 = r$ and $v_4 = s$,

$$\vec{v} = \begin{bmatrix} 2r - 3s \\ r + 0s \\ 0r + 0s \\ 0r + s \end{bmatrix} = r \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Example 8

Consider the transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[\begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

So, we see that $v_1 = 2v_2 - 3v_4$ and $v_3 = 0$. If we let $v_2 = r$ and $v_4 = s$,

$$\vec{v} = \begin{bmatrix} 2r - 3s \\ r + 0s \\ 0r + 0s \\ 0r + s \end{bmatrix} = r \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \rightarrow \ker(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

The dimension of the kernel of T is 2.

Example 8

Consider the transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ defined by $T(\vec{v}) = \mathbf{A}\vec{v}$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[\begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

So, we see that $v_1 = 2v_2 - 3v_4$ and $v_3 = 0$. If we let $v_2 = r$ and $v_4 = s$,

$$\vec{v} = \begin{bmatrix} 2r - 3s \\ r + 0s \\ 0r + 0s \\ 0r + s \end{bmatrix} = r \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \rightarrow \ker(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

The dimension of the kernel of T is 2.

(Remember that the dimension of the image of T was 2.)

Dimension Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from a finite vector space \mathbb{V} .

Then

$$\mathbf{dim}(\ker(T)) + \mathbf{dim}(\mathbf{Im}(T)) = \mathbf{dim}(\mathbb{V})$$

Dimension Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from a finite vector space \mathbb{V} .

Then

$$\mathbf{dim}(\ker(T)) + \mathbf{dim}(\mathbf{Im}(T)) = \mathbf{dim}(\mathbb{V})$$

Example 9

From earlier examples:

Dimension Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from a finite vector space \mathbb{V} .

Then

$$\mathbf{\dim(\ker(T)) + \dim(\operatorname{Im}(T)) = \dim(\mathbb{V})}$$

Example 9

From earlier examples:

- $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(x, y, z) = (x, y, 0)$ had

$$\mathbf{\dim(\ker(T)) + \dim(\operatorname{Im}(T)) = 1 + 2 = 3 = \dim(\mathbb{R}^3)}$$

Dimension Theorem

Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation from a finite vector space \mathbb{V} .

Then

$$\mathbf{\dim(\ker(T)) + \dim(\text{Im}(T)) = \dim(\mathbb{V})}$$

Example 9

From earlier examples:

- $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(x, y, z) = (x, y, 0)$ had

$$\mathbf{\dim(\ker(T)) + \dim(\text{Im}(T)) = 1 + 2 = 3 = \dim(\mathbb{R}^3)}$$

- $D_2 : \mathbb{P}_3 \rightarrow \mathbb{P}_1$ had

$$\mathbf{\ker(D_2) = \{cx + d\} \quad \text{and} \quad \text{Im}(D_2) = \{6ax + 2b\}}$$

$$\mathbf{\dim(\ker(D_2)) + \dim(\text{Im}(D_2)) = 2 + 2 = 4 = \dim(\mathbb{P}_3)}$$