


Vector Spaces

Department of Mathematics

Salt Lake Community College



Proofs by Randall Munroe is licensed under CC BY-NC 2.5

Definition

A **vector space** \mathbb{V} is a nonempty collection of objects called **vectors** for which the following operations

- Vector addition, denoted $\vec{x} + \vec{y}$
- Scalar multiplication, denoted $c\vec{x}$

satisfy the nine properties on the following slide. (For all $\vec{x}, \vec{y}, \vec{z} \in \mathbb{V}$ and all $c, d \in \mathbb{R}$)

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

- ② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$
- ③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$
- ④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

- ⑥ $1\vec{x} = \vec{x}$
- ⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$

③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$

④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$

⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

⑥ $1\vec{x} = \vec{x}$

⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$

⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$

⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$

③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$

④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$

⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

⑥ $1\vec{x} = \vec{x}$

⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$

⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$

⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$

③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$

④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$

⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

⑥ $1\vec{x} = \vec{x}$

⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$

⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$

⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

- ② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$
- ③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$
- ④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

- ⑥ $1\vec{x} = \vec{x}$
- ⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

- ② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$
- ③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$
- ④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

- ⑥ $1\vec{x} = \vec{x}$
- ⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

- ② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$
- ③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$
- ④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

- ⑥ $1\vec{x} = \vec{x}$
- ⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

- ② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$
- ③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$
- ④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

- ⑥ $1\vec{x} = \vec{x}$
- ⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨ $c(d\vec{x}) = (cd)\vec{x}$

Closure

① $c\vec{x} + d\vec{y} \in \mathbb{V}$

Addition

- ② There exists a **zero vector** $\vec{0} \in \mathbb{V}$ such that $\vec{x} + \vec{0} = \vec{x}$
- ③ For all $\vec{x} \in \mathbb{V}$ there exists $-\vec{x} \in \mathbb{V}$ such that $\vec{x} + (-\vec{x}) = \vec{0}$
- ④ $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤ $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

Scalar Multiplication

- ⑥ $1\vec{x} = \vec{x}$
- ⑦ $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧ $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨ $c(d\vec{x}) = (cd)\vec{x}$

Example 1

All vectors $\langle x_1, x_2, \dots, x_n \rangle$ in \mathbb{R}^n satisfy these properties.
(It doesn't matter if you think of them as row or column vectors.)

Example 1

All vectors $\langle x_1, x_2, \dots, x_n \rangle$ in \mathbb{R}^n satisfy these properties.
(It doesn't matter if you think of them as row or column vectors.)

Definition

Let \mathbb{M}_{mn} denote the collection of all $m \times n$ matrices.

Example 1

All vectors $\langle x_1, x_2, \dots, x_n \rangle$ in \mathbb{R}^n satisfy these properties.
(It doesn't matter if you think of them as row or column vectors.)

Definition

Let \mathbb{M}_{mn} denote the collection of all $m \times n$ matrices.

Example 2

Thinking back, we can see that the properties for addition and scalar multiplication of matrices we saw in section 3.1 satisfy all nine requirements to be a vector space.
Which means, for any $m, n \in \mathbb{R}$, \mathbb{M}_{mn} is a vector space.

Definition

A **function space** is a vector space where the “vectors” are functions defined on an interval I . The addition and scalar multiplication operations are defined in the usual way:

- $(f + g)(t) = f(t) + g(t)$, for all $t \in I$
- $(cf)(t) = cf(t)$, for all $t \in I$

Definition

A **function space** is a vector space where the “vectors” are functions defined on an interval I . The addition and scalar multiplication operations are defined in the usual way:

- $(f + g)(t) = f(t) + g(t)$, for all $t \in I$
- $(cf)(t) = cf(t)$, for all $t \in I$

Note

Solutions to *linear homogeneous* DEs form a vector space.

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$(au + bv)' + p(t)(au + bv)$$

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$(au + bv)' + p(t)(au + bv) = au' + bv' + au \cdot p(t) + bv \cdot p(t)$$

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$\begin{aligned}(au + bv)' + p(t)(au + bv) &= au' + bv' + au \cdot p(t) + bv \cdot p(t) \\ &= a(u' + p(t)u) + b(v' + p(t)v)\end{aligned}$$

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$\begin{aligned}(au + bv)' + p(t)(au + bv) &= au' + bv' + au \cdot p(t) + bv \cdot p(t) \\ &= a(u' + p(t)u) + b(v' + p(t)v) \\ &= a \cdot 0 + b \cdot 0\end{aligned}$$

Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where p and y are defined on some interval I) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$\begin{aligned}(au + bv)' + p(t)(au + bv) &= au' + bv' + au \cdot p(t) + bv \cdot p(t) \\ &= a(u' + p(t)u) + b(v' + p(t)v) \\ &= a \cdot 0 + b \cdot 0 \\ &= 0\end{aligned}$$

Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where q , p and y are defined on some interval I) is a vector space.

Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where q , p and y are defined on some interval I) is a vector space.

Again, all we really need to check is that closure holds.

Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where q , p and y are defined on some interval I) is a vector space.

Again, all we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv)$$

Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where q , p and y are defined on some interval I) is a vector space.

Again, all we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$\begin{aligned}(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv) \\ = a(u'' + q(t)u' + p(t)u) + b(v'' + q(t)v' + p(t)v)\end{aligned}$$

Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where q , p and y are defined on some interval I) is a vector space.

Again, all we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$\begin{aligned}(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv) \\&= a(u'' + q(t)u' + p(t)u) + b(v'' + q(t)v' + p(t)v) \\&= a \cdot 0 + b \cdot 0\end{aligned}$$

Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where q , p and y are defined on some interval I) is a vector space.

Again, all we really need to check is that closure holds.

For solutions $u(t)$ and $v(t)$, as well as scalars a and b , we need to verify that $a \cdot u(t) + b \cdot v(t)$ is a solution.

$$\begin{aligned}(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv) \\&= a(u'' + q(t)u' + p(t)u) + b(v'' + q(t)v' + p(t)v) \\&= a \cdot 0 + b \cdot 0 \\&= 0\end{aligned}$$

Example 5

The set of all solutions of the first-order linear (but not homogeneous) DE

$$y' + 2ty = 1$$

is **not** a vector space. Why?

Example 5

The set of all solutions of the first-order linear (but not homogeneous) DE

$$y' + 2ty = 1$$

is **not** a vector space. Why?

There is no zero vector. That is, there is no solution $z(t)$ such that, for all solutions $u(t)$, $u(t) + z(t) = u(t)$.

Example 5

The set of all solutions of the first-order linear (but not homogeneous) DE

$$y' + 2ty = 1$$

is **not** a vector space. Why?

There is no zero vector. That is, there is no solution $z(t)$ such that, for all solutions $u(t)$, $u(t) + z(t) = u(t)$.

Example 6

Consider the collection of all polynomials of degree ≤ 3 . A vector in this space is given by

$$P(t) = a_3x^3 + a_2x^2 + a_1x + a_0$$

where $a_3, a_2, a_1, a_0 \in \mathbb{R}$.

This collection is a vector space, verifiable using basic algebra.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Prominent Vector Spaces

- \mathbb{R}^2 , the space of all real ordered pairs.
- \mathbb{R}^3 , the space of all real ordered triples.
- \mathbb{R}^n , the space of all real ordered n -tuples.
- \mathbb{C}^n , the space of all complex n -tuples.
- \mathbb{P} , the space of all polynomials.
- \mathbb{P}_n , the space of all polynomials of degree $\leq n$
- \mathbb{M}_{mn} , the space of all $m \times n$ matrices.
- $\mathcal{C}(I)$, the space of all continuous functions defined on the interval I .
- $\mathcal{C}^n(I)$, the space of all functions, defined on the interval I , having n continuous derivatives.
- \mathcal{C}^n , the space of all functions, defined on the interval $(-\infty, +\infty)$, having n continuous derivatives.

Theorem 7

A nonempty subset, \mathbb{W} , of a vector space \mathbb{V} is a **subspace** of \mathbb{V} if

- $\vec{u} + \vec{v} \in \mathbb{W}$ for all $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$ for all $\vec{u} \in \mathbb{W}$ and $c \in \mathbb{R}$

Theorem 7

A nonempty subset, \mathbb{W} , of a vector space \mathbb{V} is a **subspace** of \mathbb{V} if

- $\vec{u} + \vec{v} \in \mathbb{W}$ for all $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$ for all $\vec{u} \in \mathbb{W}$ and $c \in \mathbb{R}$

Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given $\vec{u}, \vec{v} \in \mathbb{W}$, consider $\vec{u} + \vec{v}$.

Theorem 7

A nonempty subset, \mathbb{W} , of a vector space \mathbb{V} is a **subspace** of \mathbb{V} if

- $\vec{u} + \vec{v} \in \mathbb{W}$ for all $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$ for all $\vec{u} \in \mathbb{W}$ and $c \in \mathbb{R}$

Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given $\vec{u}, \vec{v} \in \mathbb{W}$, consider $\vec{u} + \vec{v}$.
Since $\mathbb{W} \subseteq \mathbb{V}$ we have $\vec{u} + \vec{v} = \vec{v} + \vec{u}$.

Theorem 7

A nonempty subset, \mathbb{W} , of a vector space \mathbb{V} is a **subspace** of \mathbb{V} if

- $\vec{u} + \vec{v} \in \mathbb{W}$ for all $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$ for all $\vec{u} \in \mathbb{W}$ and $c \in \mathbb{R}$

Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given $\vec{u}, \vec{v} \in \mathbb{W}$, consider $\vec{u} + \vec{v}$.

Since $\mathbb{W} \subseteq \mathbb{V}$ we have $\vec{u} + \vec{v} = \vec{v} + \vec{u}$.

But, since $\vec{u} + \vec{v} \in \mathbb{W}$ we must have $\vec{v} + \vec{u} \in \mathbb{W}$.

Theorem 7

A nonempty subset, \mathbb{W} , of a vector space \mathbb{V} is a **subspace** of \mathbb{V} if

- $\vec{u} + \vec{v} \in \mathbb{W}$ for all $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$ for all $\vec{u} \in \mathbb{W}$ and $c \in \mathbb{R}$

Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given $\vec{u}, \vec{v} \in \mathbb{W}$, consider $\vec{u} + \vec{v}$.

Since $\mathbb{W} \subseteq \mathbb{V}$ we have $\vec{u} + \vec{v} = \vec{v} + \vec{u}$.


But, since $\vec{u} + \vec{v} \in \mathbb{W}$ we must have $\vec{v} + \vec{u} \in \mathbb{W}$.

Note

A vector space is a subspace of itself.

Example 8


Is the upper half plane a subspace of \mathbb{R}^2 ?



Example 8

Is the upper half plane a subspace of \mathbb{R}^2 ?

No, points in the upper half plane are not closed under scalar multiplication.




Example 8

Is the upper half plane a subspace of \mathbb{R}^2 ?

No, points in the upper half plane are not closed under scalar multiplication.

Consider $(1, 1)$.




Example 8

Is the upper half plane a subspace of \mathbb{R}^2 ?

No, points in the upper half plane are not closed under scalar multiplication.


Consider $(1, 1)$.

Multiplying by the scalar -1 gives $(-1 \cdot 1, -1 \cdot 1) = (-1, -1)$, a point in Q3.



Example 9


Is the set containing Q1 and Q3 a subspace of \mathbb{R}^2 ?



Example 9

Is the set containing Q_1 and Q_3 a subspace of \mathbb{R}^2 ?

No, points in the set containing Q_1 and Q_3 are not closed under addition.




Example 9

Is the set containing $Q1$ and $Q3$ a subspace of \mathbb{R}^2 ?

No, points in the set containing $Q1$ and $Q3$ are not closed under addition.

Consider $(1, 2)$ and $(-2, -1)$.




Example 9

Is the set containing Q1 and Q3 a subspace of \mathbb{R}^2 ?

No, points in the set containing Q1 and Q3 are not closed under addition.


Consider $(1, 2)$ and $(-2, -1)$.

Adding these points gives
 $(1 + (-2), 2 + (-1)) = (-1, 1)$, a point in Q2.



Example 10

Is the line $y = x$ a subspace of \mathbb{R}^2 ?




Example 10

Is the line $y = x$ a subspace of \mathbb{R}^2 ?

Yes. Given (s, s) and (t, t) , two points on the line, then

$$a \cdot (s, s) + b \cdot (t, t) =$$




Example 10

Is the line $y = x$ a subspace of \mathbb{R}^2 ?

Yes. Given (s, s) and (t, t) , two points on the line, then

$$a \cdot (s, s) + b \cdot (t, t) = (as, as) + (bt, bt)$$




Example 10

Is the line $y = x$ a subspace of \mathbb{R}^2 ?

Yes. Given (s, s) and (t, t) , two points on the line, then

$$\begin{aligned} a \cdot (s, s) + b \cdot (t, t) &= (as, as) + (bt, bt) \\ &= (as + bt, as + bt) \end{aligned}$$




Example 10

Is the line $y = x$ a subspace of \mathbb{R}^2 ?

Yes. Given (s, s) and (t, t) , two points on the line, then


$$\begin{aligned} a \cdot (s, s) + b \cdot (t, t) &= (as, as) + (bt, bt) \\ &= (as + bt, as + bt) \end{aligned}$$

which is a point on the line.



Example 11


Is the line $y = x + 1$ a subspace of \mathbb{R}^2 ?



Example 11

Is the line $y = x + 1$ a subspace of \mathbb{R}^2 ?

No, the zero vector, $(0, 0)$ is not on the line.



Corollary 12

The only subspaces of \mathbb{R}^2 are

- *The zero subspace $(0, 0)$*
- *Any line passing through the origin*
- \mathbb{R}^2

Corollary 12

The only subspaces of \mathbb{R}^2 are

- *The zero subspace $(0, 0)$*
- *Any line passing through the origin*
- \mathbb{R}^2

We call a subspace of \mathbb{V} **trivial** if it is the subspace containing just the zero vector, or \mathbb{V} itself. All other subspaces are called **nontrivial**.

Corollary 12

The only subspaces of \mathbb{R}^2 are

- *The zero subspace $(0, 0)$*
- *Any line passing through the origin*
- *\mathbb{R}^2*

We call a subspace of \mathbb{V} **trivial** if it is the subspace containing just the zero vector, or \mathbb{V} itself. All other subspaces are called **nontrivial**.

Theorem 13

The set of solutions of the linear system $A\vec{x} = \vec{0}$ is a subspace of \mathbb{R}^m , where A is a $m \times n$ matrix and $\vec{x} \in \mathbb{R}^m$, is a subspace of \mathbb{R}^m .

Corollary 12

The only subspaces of \mathbb{R}^2 are

- *The zero subspace $(0, 0)$*
- *Any line passing through the origin*
- \mathbb{R}^2

We call a subspace of \mathbb{V} **trivial** if it is the subspace containing just the zero vector, or \mathbb{V} itself. All other subspaces are called **nontrivial**.

Theorem 13

The set of solutions of the linear system $A\vec{x} = \vec{0}$ is a subspace of \mathbb{R}^m , where A is a $m \times n$ matrix and $\vec{x} \in \mathbb{R}^m$, is a subspace of \mathbb{R}^m .

Proof

Closure is given by the Superposition Principle from section 2.1.

Corollary 12

The only subspaces of \mathbb{R}^2 are

- *The zero subspace $(0, 0)$*
- *Any line passing through the origin*
- \mathbb{R}^2

We call a subspace of \mathbb{V} **trivial** if it is the subspace containing just the zero vector, or \mathbb{V} itself. All other subspaces are called **nontrivial**.

Theorem 13

The set of solutions of the linear system $A\vec{x} = \vec{0}$ is a subspace of \mathbb{R}^m , where A is a $m \times n$ matrix and $\vec{x} \in \mathbb{R}^m$, is a subspace of \mathbb{R}^m .

Proof

Closure is given by the Superposition Principle from section 2.1.

Since solutions to $A\vec{x} = \vec{0}$ are vectors in \mathbb{R}^m , the remaining properties are inherited from \mathbb{R}^m