


# Vector Spaces

Department of Mathematics

Salt Lake Community College



Proofs by Randall Munroe is licensed under CC BY-NC 2.5

## Definition

A **vector space**  $\mathbb{V}$  is a nonempty collection of objects called **vectors** for which the following operations

- Vector addition, denoted  $\vec{x} + \vec{y}$
- Scalar multiplication, denoted  $c\vec{x}$

satisfy the nine properties on the following slide. (For all  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{V}$  and all  $c, d \in \mathbb{R}$ )

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

- ② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$
- ③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$
- ④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

- ⑥  $1\vec{x} = \vec{x}$
- ⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$

③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$

④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$

⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

⑥  $1\vec{x} = \vec{x}$

⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$

⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$

⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$

③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$

④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$

⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

⑥  $1\vec{x} = \vec{x}$

⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$

⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$

⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$

③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$

④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$

⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

⑥  $1\vec{x} = \vec{x}$

⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$

⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$

⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

- ② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$
- ③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$
- ④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

- ⑥  $1\vec{x} = \vec{x}$
- ⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

- ② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$
- ③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$
- ④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

- ⑥  $1\vec{x} = \vec{x}$
- ⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨  $c(d\vec{x}) = (cd)\vec{x}$



## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

- ② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$
- ③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$
- ④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

- ⑥  $1\vec{x} = \vec{x}$
- ⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

- ② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$
- ③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$
- ④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

- ⑥  $1\vec{x} = \vec{x}$
- ⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨  $c(d\vec{x}) = (cd)\vec{x}$

## Closure

①  $c\vec{x} + d\vec{y} \in \mathbb{V}$

## Addition

- ② There exists a **zero vector**  $\vec{0} \in \mathbb{V}$  such that  $\vec{x} + \vec{0} = \vec{x}$
- ③ For all  $\vec{x} \in \mathbb{V}$  there exists  $-\vec{x} \in \mathbb{V}$  such that  $\vec{x} + (-\vec{x}) = \vec{0}$
- ④  $(\vec{x} + \vec{y}) + \vec{z} = \vec{x} + (\vec{y} + \vec{z})$
- ⑤  $\vec{x} + \vec{y} = \vec{y} + \vec{x}$

## Scalar Multiplication

- ⑥  $1\vec{x} = \vec{x}$
- ⑦  $c(\vec{x} + \vec{y}) = c\vec{x} + c\vec{y}$
- ⑧  $(c + d)\vec{x} = c\vec{x} + d\vec{x}$
- ⑨  $c(d\vec{x}) = (cd)\vec{x}$

### Example 1

All vectors  $\langle x_1, x_2, \dots, x_n \rangle$  in  $\mathbb{R}^n$  satisfy these properties.  
(It doesn't matter if you think of them as row or column vectors.)

### Example 1

All vectors  $\langle x_1, x_2, \dots, x_n \rangle$  in  $\mathbb{R}^n$  satisfy these properties.  
(It doesn't matter if you think of them as row or column vectors.)

### Definition

Let  $\mathbb{M}_{mn}$  denote the collection of all  $m \times n$  matrices.

### Example 1

All vectors  $\langle x_1, x_2, \dots, x_n \rangle$  in  $\mathbb{R}^n$  satisfy these properties.  
(It doesn't matter if you think of them as row or column vectors.)

### Definition

Let  $\mathbb{M}_{mn}$  denote the collection of all  $m \times n$  matrices.

### Example 2

Thinking back, we can see that the properties for addition and scalar multiplication of matrices we saw in section 3.1 satisfy all nine requirements to be a vector space.  
Which means, for any  $m, n \in \mathbb{R}$ ,  $\mathbb{M}_{mn}$  is a vector space.

## Definition

A **function space** is a vector space where the “vectors” are functions defined on an interval  $I$ . The addition and scalar multiplication operations are defined in the usual way:

- $(f + g)(t) = f(t) + g(t)$ , for all  $t \in I$
- $(cf)(t) = cf(t)$ , for all  $t \in I$

## Definition

A **function space** is a vector space where the “vectors” are functions defined on an interval  $I$ . The addition and scalar multiplication operations are defined in the usual way:

- $(f + g)(t) = f(t) + g(t)$ , for all  $t \in I$
- $(cf)(t) = cf(t)$ , for all  $t \in I$

## Note

Solutions to *linear homogeneous* DEs form a vector space.



### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$(au + bv)' + p(t)(au + bv)$$

### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$(au + bv)' + p(t)(au + bv) = au' + bv' + au \cdot p(t) + bv \cdot p(t)$$

### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$\begin{aligned}(au + bv)' + p(t)(au + bv) &= au' + bv' + au \cdot p(t) + bv \cdot p(t) \\ &= a(u' + p(t)u) + b(v' + p(t)v)\end{aligned}$$

### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$\begin{aligned}(au + bv)' + p(t)(au + bv) &= au' + bv' + au \cdot p(t) + bv \cdot p(t) \\ &= a(u' + p(t)u) + b(v' + p(t)v) \\ &= a \cdot 0 + b \cdot 0\end{aligned}$$

### Example 3

The set of all solutions of the first order linear homogeneous DE

$$y' + p(t)y = 0$$

(where  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

The addition and scalar multiplication properties are well known for functions. All we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$\begin{aligned}(au + bv)' + p(t)(au + bv) &= au' + bv' + au \cdot p(t) + bv \cdot p(t) \\ &= a(u' + p(t)u) + b(v' + p(t)v) \\ &= a \cdot 0 + b \cdot 0 \\ &= 0\end{aligned}$$

### Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where  $q$ ,  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.



### Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where  $q$ ,  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

Again, all we really need to check is that closure holds.

### Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where  $q$ ,  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

Again, all we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv)$$

### Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where  $q$ ,  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

Again, all we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$\begin{aligned}(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv) \\ = a(u'' + q(t)u' + p(t)u) + b(v'' + q(t)v' + p(t)v)\end{aligned}$$

### Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where  $q$ ,  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

Again, all we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$\begin{aligned}(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv) \\&= a(u'' + q(t)u' + p(t)u) + b(v'' + q(t)v' + p(t)v) \\&= a \cdot 0 + b \cdot 0\end{aligned}$$

### Example 4

The set of all solutions of the second order linear homogeneous DE

$$y'' + q(t)y' + p(t)y = 0$$

(where  $q$ ,  $p$  and  $y$  are defined on some interval  $I$ ) is a vector space.

Again, all we really need to check is that closure holds.

For solutions  $u(t)$  and  $v(t)$ , as well as scalars  $a$  and  $b$ , we need to verify that  $a \cdot u(t) + b \cdot v(t)$  is a solution.

$$\begin{aligned}(au + bv)'' + q(t)(au + bv)' + p(t)(au + bv) \\&= a(u'' + q(t)u' + p(t)u) + b(v'' + q(t)v' + p(t)v) \\&= a \cdot 0 + b \cdot 0 \\&= 0\end{aligned}$$

### Example 5

The set of all solutions of the first-order linear (but not homogeneous) DE

$$y' + 2ty = 1$$

is **not** a vector space. Why?

### Example 5

The set of all solutions of the first-order linear (but not homogeneous) DE

$$y' + 2ty = 1$$

is **not** a vector space. Why?

There is no zero vector. That is, there is no solution  $z(t)$  such that, for all solutions  $u(t)$ ,  $u(t) + z(t) = u(t)$ .

### Example 5

The set of all solutions of the first-order linear (but not homogeneous) DE

$$y' + 2ty = 1$$

is **not** a vector space. Why?

There is no zero vector. That is, there is no solution  $z(t)$  such that, for all solutions  $u(t)$ ,  $u(t) + z(t) = u(t)$ .

### Example 6

Consider the collection of all polynomials of degree  $\leq 3$ . A vector in this space is given by

$$P(t) = a_3x^3 + a_2x^2 + a_1x + a_0$$

where  $a_3, a_2, a_1, a_0 \in \mathbb{R}$ .

This collection is a vector space, verifiable using basic algebra.



## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.



## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Prominent Vector Spaces

- $\mathbb{R}^2$ , the space of all real ordered pairs.
- $\mathbb{R}^3$ , the space of all real ordered triples.
- $\mathbb{R}^n$ , the space of all real ordered  $n$ -tuples.
- $\mathbb{C}^n$ , the space of all complex  $n$ -tuples.
- $\mathbb{P}$ , the space of all polynomials.
- $\mathbb{P}_n$ , the space of all polynomials of degree  $\leq n$
- $\mathbb{M}_{mn}$ , the space of all  $m \times n$  matrices.
- $\mathcal{C}(I)$ , the space of all continuous functions defined on the interval  $I$ .
- $\mathcal{C}^n(I)$ , the space of all functions, defined on the interval  $I$ , having  $n$  continuous derivatives.
- $\mathcal{C}^n$ , the space of all functions, defined on the interval  $(-\infty, +\infty)$ , having  $n$  continuous derivatives.

## Theorem 7

A nonempty subset,  $\mathbb{W}$ , of a vector space  $\mathbb{V}$  is a **subspace** of  $\mathbb{V}$  if

- $\vec{u} + \vec{v} \in \mathbb{W}$  for all  $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$  for all  $\vec{u} \in \mathbb{W}$  and  $c \in \mathbb{R}$

## Theorem 7

A nonempty subset,  $\mathbb{W}$ , of a vector space  $\mathbb{V}$  is a **subspace** of  $\mathbb{V}$  if

- $\vec{u} + \vec{v} \in \mathbb{W}$  for all  $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$  for all  $\vec{u} \in \mathbb{W}$  and  $c \in \mathbb{R}$

## Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given  $\vec{u}, \vec{v} \in \mathbb{W}$ , consider  $\vec{u} + \vec{v}$ .

## Theorem 7

A nonempty subset,  $\mathbb{W}$ , of a vector space  $\mathbb{V}$  is a **subspace** of  $\mathbb{V}$  if

- $\vec{u} + \vec{v} \in \mathbb{W}$  for all  $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$  for all  $\vec{u} \in \mathbb{W}$  and  $c \in \mathbb{R}$

## Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given  $\vec{u}, \vec{v} \in \mathbb{W}$ , consider  $\vec{u} + \vec{v}$ .  
Since  $\mathbb{W} \subseteq \mathbb{V}$  we have  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$ .

## Theorem 7

A nonempty subset,  $\mathbb{W}$ , of a vector space  $\mathbb{V}$  is a **subspace** of  $\mathbb{V}$  if

- $\vec{u} + \vec{v} \in \mathbb{W}$  for all  $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$  for all  $\vec{u} \in \mathbb{W}$  and  $c \in \mathbb{R}$

## Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given  $\vec{u}, \vec{v} \in \mathbb{W}$ , consider  $\vec{u} + \vec{v}$ .

Since  $\mathbb{W} \subseteq \mathbb{V}$  we have  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$ .

But, since  $\vec{u} + \vec{v} \in \mathbb{W}$  we must have  $\vec{v} + \vec{u} \in \mathbb{W}$ .

## Theorem 7

A nonempty subset,  $\mathbb{W}$ , of a vector space  $\mathbb{V}$  is a **subspace** of  $\mathbb{V}$  if

- $\vec{u} + \vec{v} \in \mathbb{W}$  for all  $\vec{u}, \vec{v} \in \mathbb{W}$
- $c\vec{u} \in \mathbb{W}$  for all  $\vec{u} \in \mathbb{W}$  and  $c \in \mathbb{R}$

## Proof

The definition of a subspace guarantees closure, everything else is inherited from the parent vector space.

For example, given  $\vec{u}, \vec{v} \in \mathbb{W}$ , consider  $\vec{u} + \vec{v}$ .

Since  $\mathbb{W} \subseteq \mathbb{V}$  we have  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$ .


But, since  $\vec{u} + \vec{v} \in \mathbb{W}$  we must have  $\vec{v} + \vec{u} \in \mathbb{W}$ .

## Note

A vector space is a subspace of itself.

## Example 8

Is the upper half plane a subspace of  $\mathbb{R}^2$ ?






### Example 8

Is the upper half plane a subspace of  $\mathbb{R}^2$ ?

No, points in the upper half plane are not closed under scalar multiplication.




## Example 8

Is the upper half plane a subspace of  $\mathbb{R}^2$ ?

No, points in the upper half plane are not closed under scalar multiplication.

Consider  $(1, 1)$ .




## Example 8

Is the upper half plane a subspace of  $\mathbb{R}^2$ ?

No, points in the upper half plane are not closed under scalar multiplication.


Consider  $(1, 1)$ .

Multiplying by the scalar  $-1$  gives  $(-1 \cdot 1, -1 \cdot 1) = (-1, -1)$ , a point in Q3.



### Example 9


Is the set containing Q1 and Q3 a subspace of  $\mathbb{R}^2$ ?



### Example 9

Is the set containing  $Q_1$  and  $Q_3$  a subspace of  $\mathbb{R}^2$ ?

No, points in the set containing  $Q_1$  and  $Q_3$  are not closed under addition.




## Example 9

Is the set containing  $Q1$  and  $Q3$  a subspace of  $\mathbb{R}^2$ ?

No, points in the set containing  $Q1$  and  $Q3$  are not closed under addition.

Consider  $(1, 2)$  and  $(-2, -1)$ .




## Example 9

Is the set containing Q1 and Q3 a subspace of  $\mathbb{R}^2$ ?

No, points in the set containing Q1 and Q3 are not closed under addition.


Consider  $(1, 2)$  and  $(-2, -1)$ .

Adding these points gives  $(1 + (-2), 2 + (-1)) = (-1, 1)$ , a point in Q2.



## Example 10

Is the line  $y = x$  a subspace of  $\mathbb{R}^2$ ?






## Example 10

Is the line  $y = x$  a subspace of  $\mathbb{R}^2$ ?

Yes. Given  $(s, s)$  and  $(t, t)$ , two points on the line, then

$$a \cdot (s, s) + b \cdot (t, t) =$$




## Example 10

Is the line  $y = x$  a subspace of  $\mathbb{R}^2$ ?

Yes. Given  $(s, s)$  and  $(t, t)$ , two points on the line, then

$$a \cdot (s, s) + b \cdot (t, t) = (as, as) + (bt, bt)$$




## Example 10

Is the line  $y = x$  a subspace of  $\mathbb{R}^2$ ?

Yes. Given  $(s, s)$  and  $(t, t)$ , two points on the line, then

$$\begin{aligned} a \cdot (s, s) + b \cdot (t, t) &= (as, as) + (bt, bt) \\ &= (as + bt, as + bt) \end{aligned}$$




## Example 10

Is the line  $y = x$  a subspace of  $\mathbb{R}^2$ ?

Yes. Given  $(s, s)$  and  $(t, t)$ , two points on the line, then


$$\begin{aligned} a \cdot (s, s) + b \cdot (t, t) &= (as, as) + (bt, bt) \\ &= (as + bt, as + bt) \end{aligned}$$

which is a point on the line.



## Example 11


Is the line  $y = x + 1$  a subspace of  $\mathbb{R}^2$ ?



## Example 11

Is the line  $y = x + 1$  a subspace of  $\mathbb{R}^2$ ?

No, the zero vector,  $(0, 0)$  is not on the line.



## Corollary 12

*The only subspaces of  $\mathbb{R}^2$  are*

- *The zero subspace  $(0, 0)$*
- *Any line passing through the origin*
- $\mathbb{R}^2$

## Corollary 12

*The only subspaces of  $\mathbb{R}^2$  are*

- *The zero subspace  $(0, 0)$*
- *Any line passing through the origin*
- $\mathbb{R}^2$

We call a subspace of  $\mathbb{V}$  **trivial** if it is the subspace containing just the zero vector, or  $\mathbb{V}$  itself. All other subspaces are called **nontrivial**.



## Corollary 12

*The only subspaces of  $\mathbb{R}^2$  are*

- *The zero subspace  $(0, 0)$*
- *Any line passing through the origin*
- *$\mathbb{R}^2$*

We call a subspace of  $\mathbb{V}$  **trivial** if it is the subspace containing just the zero vector, or  $\mathbb{V}$  itself. All other subspaces are called **nontrivial**.

## Theorem 13

*The set of solutions of the linear system  $A\vec{x} = \vec{0}$  is a subspace of  $\mathbb{R}^m$ , where  $A$  is a  $m \times n$  matrix and  $\vec{x} \in \mathbb{R}^m$ , is a subspace of  $\mathbb{R}^m$ .*

## Corollary 12

*The only subspaces of  $\mathbb{R}^2$  are*

- *The zero subspace  $(0, 0)$*
- *Any line passing through the origin*
- *$\mathbb{R}^2$*

We call a subspace of  $\mathbb{V}$  **trivial** if it is the subspace containing just the zero vector, or  $\mathbb{V}$  itself. All other subspaces are called **nontrivial**.

## Theorem 13

*The set of solutions of the linear system  $A\vec{x} = \vec{0}$  is a subspace of  $\mathbb{R}^m$ , where  $A$  is a  $m \times n$  matrix and  $\vec{x} \in \mathbb{R}^m$ , is a subspace of  $\mathbb{R}^m$ .*

## Proof

Closure is given by the Superposition Principle from section 2.1.

## Corollary 12

*The only subspaces of  $\mathbb{R}^2$  are*

- *The zero subspace  $(0, 0)$*
- *Any line passing through the origin*
- $\mathbb{R}^2$

We call a subspace of  $\mathbb{V}$  **trivial** if it is the subspace containing just the zero vector, or  $\mathbb{V}$  itself. All other subspaces are called **nontrivial**.

## Theorem 13

*The set of solutions of the linear system  $A\vec{x} = \vec{0}$  is a subspace of  $\mathbb{R}^m$ , where  $A$  is a  $m \times n$  matrix and  $\vec{x} \in \mathbb{R}^m$ , is a subspace of  $\mathbb{R}^m$ .*

## Proof

Closure is given by the Superposition Principle from section 2.1.

Since solutions to  $A\vec{x} = \vec{0}$  are vectors in  $\mathbb{R}^m$ , the remaining properties are inherited from  $\mathbb{R}^m$