

# Variation of Parameters

Department of Mathematics

Salt Lake Community College

(Slides by Adam Wilson)

## Variation of Parameters

Let us extend Variation of Parameters to solve

$$y'' + p(t)y' + q(t)y = f(t)$$

## Variation of Parameters

Let us extend Variation of Parameters to solve

$$y'' + p(t)y' + q(t)y = f(t)$$

We first need to find two linearly independent solutions to the associated homogeneous equation

$$y'' + p(t)y' + q(t)y = 0$$

## Variation of Parameters

Let us extend Variation of Parameters to solve

$$y'' + p(t)y' + q(t)y = f(t)$$

We first need to find two linearly independent solutions to the associated homogeneous equation

$$y'' + p(t)y' + q(t)y = 0$$

Which gives the general solution

$$y_h = c_1 y_1(t) + c_2 y_2(t)$$

where  $c_1$  and  $c_2$  are arbitrary constants.

## Variation of Parameters

Let us extend Variation of Parameters to solve

$$y'' + p(t)y' + q(t)y = f(t)$$

We first need to find two linearly independent solutions to the associated homogeneous equation

$$y'' + p(t)y' + q(t)y = 0$$

Which gives the general solution

$$y_h = c_1 y_1(t) + c_2 y_2(t)$$

where  $c_1$  and  $c_2$  are arbitrary constants.

Just like with single order equations, we want to perturb the homogeneous solution into a particular solution to the nonhomogeneous DE.

## Variation of Parameters

We do so by replacing the constants  $c_1$  and  $c_2$  with unknown functions.

$$y_p = v_1(t)y_1(t) + v_2(t)y_2(t)$$

## Variation of Parameters

We do so by replacing the constants  $c_1$  and  $c_2$  with unknown functions.

$$y_p = v_1(t)y_1(t) + v_2(t)y_2(t)$$

To find  $v_1(t)$  and  $v_2(t)$  we substitute  $y_p$  into the nonhomogeneous DE.

## Variation of Parameters

We do so by replacing the constants  $c_1$  and  $c_2$  with unknown functions.

$$y_p = v_1(t)y_1(t) + v_2(t)y_2(t)$$

To find  $v_1(t)$  and  $v_2(t)$  we substitute  $y_p$  into the nonhomogeneous DE.

But, we need two equations and we only have  $L(y) = f$ . Thus, we must choose an auxiliary condition.



## Variation of Parameters

We do so by replacing the constants  $c_1$  and  $c_2$  with unknown functions.

$$y_p = v_1(t)y_1(t) + v_2(t)y_2(t)$$

To find  $v_1(t)$  and  $v_2(t)$  we substitute  $y_p$  into the nonhomogeneous DE.

But, we need two equations and we only have  $L(y) = f$ . Thus, we must choose an auxiliary condition.

Let us calculate

$$y_p' = v_1 y_1' + v_2 y_2' + v_1' y_1 + v_2' y_2$$

## Variation of Parameters

We do so by replacing the constants  $c_1$  and  $c_2$  with unknown functions.

$$y_p = v_1(t)y_1(t) + v_2(t)y_2(t)$$

To find  $v_1(t)$  and  $v_2(t)$  we substitute  $y_p$  into the nonhomogeneous DE.

But, we need two equations and we only have  $L(y) = f$ . Thus, we must choose an auxiliary condition.

Let us calculate

$$y_p' = v_1 y_1' + v_2 y_2' + v_1' y_1 + v_2' y_2$$

So, we can choose  $v_1 y_1' + v_2 y_2' = 0$  as our auxiliary condition, which reduces  $y_p'$  to:

$$y_p' = v_1' y_1 + v_2' y_2$$

## Variation of Parameters

We can then obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'$$

## Variation of Parameters

We can then obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'$$

We then substitute  $y_p$ ,  $y_p'$ , and  $y_p''$  into  $L(y) = f$ .

$$(v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2') + p \cdot (v_1' y_1 + v_2' y_2) + q \cdot (v_1 y_1 + v_2 y_2) = f$$

## Variation of Parameters

We can then obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'$$

We then substitute  $y_p$ ,  $y_p'$ , and  $y_p''$  into  $L(y) = f$ .

$$\begin{aligned}(v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2') + p \cdot (v_1' y_1 + v_2' y_2) + q \cdot (v_1 y_1 + v_2 y_2) &= f \\ v_1(y_1'' + p y_1' + q y_1) + v_2(y_2'' + p y_2' + q y_2) + (v_1' y_1' + v_2' y_2') &= f\end{aligned}$$

## Variation of Parameters

We can then obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'$$

We then substitute  $y_p$ ,  $y_p'$ , and  $y_p''$  into  $L(y) = f$ .

$$(v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2') + p \cdot (v_1' y_1 + v_2' y_2) + q \cdot (v_1 y_1 + v_2 y_2) = f$$

$$v_1(y_1'' + p y_1' + q y_1) + v_2(y_2'' + p y_2' + q y_2) + (v_1' y_1' + v_2' y_2') = f$$

$$v_1 \cdot 0 + v_2 \cdot 0 + v_1' y_1' + v_2' y_2' = f$$

## Variation of Parameters

We can then obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'$$

We then substitute  $y_p$ ,  $y_p'$ , and  $y_p''$  into  $L(y) = f$ .

$$(v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2') + p \cdot (v_1' y_1 + v_2' y_2) + q \cdot (v_1 y_1 + v_2 y_2) = f$$

$$v_1(y_1'' + p y_1' + q y_1) + v_2(y_2'' + p y_2' + q y_2) + (v_1' y_1' + v_2' y_2') = f$$

$$v_1 \cdot 0 + v_2 \cdot 0 + v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1' + v_2' y_2' = f$$

## Variation of Parameters

We can then obtain

$$y_p'' = v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2'$$

We then substitute  $y_p$ ,  $y_p'$ , and  $y_p''$  into  $L(y) = f$ .

$$(v_1 y_1'' + v_2 y_2'' + v_1' y_1' + v_2' y_2') + p \cdot (v_1' y_1 + v_2' y_2) + q \cdot (v_1 y_1 + v_2 y_2) = f$$

$$v_1(y_1'' + p y_1' + q y_1) + v_2(y_2'' + p y_2' + q y_2) + (v_1' y_1' + v_2' y_2') = f$$

$$v_1 \cdot 0 + v_2 \cdot 0 + v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1' + v_2' y_2' = f$$

So, we have the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$



## Variation of Parameters

Using Cramer's Rule, the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

has solution

$$v_1' = \frac{\begin{vmatrix} 0 & y_2 \\ f & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} \quad \text{and} \quad v_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & f \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}}$$

## Variation of Parameters

Using Cramer's Rule, the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

has solution

$$v_1' = \frac{\begin{vmatrix} 0 & y_2 \\ f & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} \quad \text{and} \quad v_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & f \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}}$$

The denominator is just the Wronskian  $W(y_1, y_2) = y_1 y_2' - y_2 y_1' \neq 0$ , because  $y_1$  and  $y_2$  are linearly independent.

## Variation of Parameters

Using Cramer's Rule, the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

has solution

$$v_1' = \frac{\begin{vmatrix} 0 & y_2 \\ f & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} \quad \text{and} \quad v_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & f \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}}$$

The denominator is just the Wronskian  $W(y_1, y_2) = y_1 y_2' - y_2 y_1' \neq 0$ , because  $y_1$  and  $y_2$  are linearly independent.

Thus, we can integrate to find  $v_1$  and  $v_2$ .

## Variation of Parameters

Using Cramer's Rule, the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

has solution

$$v_1' = \frac{\begin{vmatrix} 0 & y_2 \\ f & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}} \quad \text{and} \quad v_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & f \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}}$$

The denominator is just the Wronskian  $W(y_1, y_2) = y_1 y_2' - y_2 y_1' \neq 0$ , because  $y_1$  and  $y_2$  are linearly independent.

Thus, we can integrate to find  $v_1$  and  $v_2$ .

$$v_1 = - \int \frac{y_2 f}{W(y_1, y_2)} \quad \text{and} \quad v_2 = \int \frac{y_1 f}{W(y_1, y_2)}$$

## Example 1

Consider

$$y'' + y = \sec(t) \quad |t| < \frac{\pi}{2}$$

## Example 1

Consider

$$y'' + y = \sec(t) \quad |t| < \frac{\pi}{2}$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

## Example 1

Consider

$$y'' + y = \sec(t) \quad |t| < \frac{\pi}{2}$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is

$$W(y_1, y_2) = \begin{vmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{vmatrix} = \cos^2(t) + \sin^2(t) = 1$$

## Example 1

Consider

$$y'' + y = \sec(t) \quad |t| < \frac{\pi}{2}$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is

$$W(y_1, y_2) = \begin{vmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{vmatrix} = \cos^2(t) + \sin^2(t) = 1$$

So,

$$v_1' = -\frac{y_2 f}{1} = -\sin(t) \sec(t) = -\frac{\sin(t)}{\cos(t)}$$

$$v_2' = \frac{y_1 f}{1} = \cos(t) \sec(t) = 1$$



## Example 1

Consider

$$y'' + y = \sec(t) \quad |t| < \frac{\pi}{2}$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is

$$W(y_1, y_2) = \begin{vmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{vmatrix} = \cos^2(t) + \sin^2(t) = 1$$

So,

$$v_1' = -\frac{y_2 f}{1} = -\sin(t) \sec(t) = -\frac{\sin(t)}{\cos(t)}$$

$$v_2' = \frac{y_1 f}{1} = \cos(t) \sec(t) = 1$$

Integrating gives  $v_1 = \ln(\cos(t))$  and  $v_2 = t$ .

## Example 1

Consider

$$y'' + y = \sec(t) \quad |t| < \frac{\pi}{2}$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is

$$W(y_1, y_2) = \begin{vmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{vmatrix} = \cos^2(t) + \sin^2(t) = 1$$

So,

$$v_1' = -\frac{y_2 f}{1} = -\sin(t) \sec(t) = -\frac{\sin(t)}{\cos(t)}$$

$$v_2' = \frac{y_1 f}{1} = \cos(t) \sec(t) = 1$$

Integrating gives  $v_1 = \ln(\cos(t))$  and  $v_2 = t$ .

Thus, the general solution is

$$y = c_1 \cos(t) + c_2 \sin(t) + \ln(\cos(t)) \cos(t) + t \sin(t)$$

## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is  $W(y_1, y_2) = 1$ .

## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is  $W(y_1, y_2) = 1$ .

So,

$$v_1' = -\frac{y_2 f}{1} = -2(1 - \cos(2t)) \quad \text{and} \quad v_2' = \frac{y_1 f}{1} = 2 \sin(2t)$$

## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is  $W(y_1, y_2) = 1$ .

So,

$$v_1' = -\frac{y_2 f}{1} = -2(1 - \cos(2t)) \quad \text{and} \quad v_2' = \frac{y_1 f}{1} = 2 \sin(2t)$$

Integrating gives  $v_1 = -2t + 2 \sin(t) \cos(t)$  and  $v_2 = 1 - 2 \cos^2(t)$ .

## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is  $W(y_1, y_2) = 1$ .

So,

$$v_1' = -\frac{y_2 f}{1} = -2(1 - \cos(2t)) \quad \text{and} \quad v_2' = \frac{y_1 f}{1} = 2 \sin(2t)$$

Integrating gives  $v_1 = -2t + 2 \sin(t) \cos(t)$  and  $v_2 = 1 - 2 \cos^2(t)$ .

Thus, the particular solution is

$$y_p = (-2t + 2 \sin(t) \cos(t))(\cos(t)) + (1 - 2 \cos^2(t))(\sin(t))$$



## Example 2

Consider

$$y'' + y = 4 \sin(t)$$

The characteristic equation is  $r^2 + 1 = 0$ , so the characteristic roots are  $r = \pm i$ . Which means  $y_1 = \cos(t)$  and  $y_2 = \sin(t)$ .

The Wronskian is  $W(y_1, y_2) = 1$ .

So,

$$v_1' = -\frac{y_2 f}{1} = -2(1 - \cos(2t)) \quad \text{and} \quad v_2' = \frac{y_1 f}{1} = 2 \sin(2t)$$

Integrating gives  $v_1 = -2t + 2 \sin(t) \cos(t)$  and  $v_2 = 1 - 2 \cos^2(t)$ .

Thus, the particular solution is

$$\begin{aligned} y_p &= (-2t + 2 \sin(t) \cos(t))(\cos(t)) + (1 - 2 \cos^2(t))(\sin(t)) \\ &= -2t \cos(t) + \sin(t) \end{aligned}$$

## Method for Determining a General Solution of $L(y) = f(t)$

- 1 Determine two linearly independent solutions,  $y_1$  and  $y_2$ , of  $L(y) = 0$ .

## Method for Determining a General Solution of $L(y) = f(t)$

- 1 Determine two linearly independent solutions,  $y_1$  and  $y_2$ , of  $L(y) = 0$ .
- 2 Solve, for  $v_1'$ , and  $v_2'$ , the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

(Cramer's Rule is recommended.)

## Method for Determining a General Solution of $L(y) = f(t)$

- 1 Determine two linearly independent solutions,  $y_1$  and  $y_2$ , of  $L(y) = 0$ .
- 2 Solve, for  $v_1'$ , and  $v_2'$ , the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

(Cramer's Rule is recommended.)

- 3 Integrate to determine  $v_1$  and  $v_2$ .

## Method for Determining a General Solution of $L(y) = f(t)$

- 1 Determine two linearly independent solutions,  $y_1$  and  $y_2$ , of  $L(y) = 0$ .
- 2 Solve, for  $v_1'$ , and  $v_2'$ , the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

(Cramer's Rule is recommended.)

- 3 Integrate to determine  $v_1$  and  $v_2$ .
- 4 Compute  $y = y_h + y_p = c_1 y_1 + c_2 y_2 + v_1 y_1 + v_2 y_2$ .

## Method for Determining a General Solution of $L(y) = f(t)$

- 1 Determine two linearly independent solutions,  $y_1$  and  $y_2$ , of  $L(y) = 0$ .
- 2 Solve, for  $v_1'$ , and  $v_2'$ , the system

$$v_1' y_1' + v_2' y_2' = f$$

$$v_1' y_1 + v_2' y_2 = 0$$

(Cramer's Rule is recommended.)

- 3 Integrate to determine  $v_1$  and  $v_2$ .
- 4 Compute  $y = y_h + y_p = c_1 y_1 + c_2 y_2 + v_1 y_1 + v_2 y_2$ .

## Note

This method can be extended to higher orders.

### Example 3

Consider

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

### Example 3

Consider

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

The associated homogeneous system has solutions  $y_1 = e^t$  and  $y_2 = te^t$ .



### Example 3

Consider

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

The associated homogeneous system has solutions  $y_1 = e^t$  and  $y_2 = te^t$ .

The Wronskian is  $W(y_1, y_2) = \begin{vmatrix} e^t & te^t \\ e^t & (t+1)e^t \end{vmatrix} = (t+1)e^{2t} - te^{2t} = e^{2t}$

### Example 3

Consider

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

The associated homogeneous system has solutions  $y_1 = e^t$  and  $y_2 = te^t$ .

The Wronskian is  $W(y_1, y_2) = \begin{vmatrix} e^t & te^t \\ e^t & (t+1)e^t \end{vmatrix} = (t+1)e^{2t} - te^{2t} = e^{2t}$

So, using the Cramer's Rule formulas from before

$$v_1' = -\frac{y_2 f}{W(y_1, y_2)} = -\frac{t}{t^2 + 1} \quad \text{and} \quad v_2' = \frac{y_1 f}{W(y_1, y_2)} = \frac{1}{t^2 + 1}$$

### Example 3

Consider

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

The associated homogeneous system has solutions  $y_1 = e^t$  and  $y_2 = te^t$ .

The Wronskian is  $W(y_1, y_2) = \begin{vmatrix} e^t & te^t \\ e^t & (t+1)e^t \end{vmatrix} = (t+1)e^{2t} - te^{2t} = e^{2t}$

So, using the Cramer's Rule formulas from before

$$v_1' = -\frac{y_2 f}{W(y_1, y_2)} = -\frac{t}{t^2 + 1} \quad \text{and} \quad v_2' = \frac{y_1 f}{W(y_1, y_2)} = \frac{1}{t^2 + 1}$$

Integrating gives

$$v_1 = -\frac{1}{2} \ln(t^2 + 1) \quad \text{and} \quad v_2 = \tan^{-1}(t)$$

### Example 3

Consider

$$y'' - 2y' + y = \frac{e^t}{t^2 + 1}$$

The associated homogeneous system has solutions  $y_1 = e^t$  and  $y_2 = te^t$ .

The Wronskian is  $W(y_1, y_2) = \begin{vmatrix} e^t & te^t \\ e^t & (t+1)e^t \end{vmatrix} = (t+1)e^{2t} - te^{2t} = e^{2t}$

So, using the Cramer's Rule formulas from before

$$v_1' = -\frac{y_2 f}{W(y_1, y_2)} = -\frac{t}{t^2 + 1} \quad \text{and} \quad v_2' = \frac{y_1 f}{W(y_1, y_2)} = \frac{1}{t^2 + 1}$$

Integrating gives

$$v_1 = -\frac{1}{2} \ln(t^2 + 1) \quad \text{and} \quad v_2 = \tan^{-1}(t)$$

Thus, the general solution is

$$y = c_1 e^t + c_2 te^t - \frac{1}{2} \ln(t^2 + 1) e^t + \tan^{-1}(t) te^t$$

## Example 4

Consider

$$t^2 y'' - 2ty' + 2y = t \ln(t), \quad t > 0$$

## Example 4

Consider

$$t^2 y'' - 2ty' + 2y = t \ln(t), \quad t > 0$$

$$y'' - \frac{2}{t}y' + \frac{2}{t^2}y = \frac{\ln(t)}{t}$$

## Example 4

Consider

$$t^2 y'' - 2ty' + 2y = t \ln(t), \quad t > 0$$

$$y'' - \frac{2}{t}y' + \frac{2}{t^2}y = \frac{\ln(t)}{t}$$

Two solutions of  $L(y) = 0$  are  $y_1 = t$  and  $y_2 = t^2$ .

## Example 4

Consider

$$t^2 y'' - 2ty' + 2y = t \ln(t), \quad t > 0$$

$$y'' - \frac{2}{t}y' + \frac{2}{t^2}y = \frac{\ln(t)}{t}$$

Two solutions of  $L(y) = 0$  are  $y_1 = t$  and  $y_2 = t^2$ .

The Wronskian is  $W(y_1, y_2) = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix} = 2t^2 - t^2 = t^2$



## Example 4

Consider

$$t^2 y'' - 2ty' + 2y = t \ln(t), \quad t > 0$$

$$y'' - \frac{2}{t}y' + \frac{2}{t^2}y = \frac{\ln(t)}{t}$$

Two solutions of  $L(y) = 0$  are  $y_1 = t$  and  $y_2 = t^2$ .

The Wronskian is  $W(y_1, y_2) = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix} = 2t^2 - t^2 = t^2$

So, using Cramer's Rule

$$\begin{aligned} v_1' &= -\frac{y_2 f}{W(y_1, y_2)} = -\frac{\ln(t)}{t} \rightarrow v_1 = -\frac{1}{2} \ln^2(t) \\ v_2' &= \frac{y_1 f}{W(y_1, y_2)} = \frac{\ln(t)}{t^2} \rightarrow v_2 = -\frac{\ln(t) + 1}{t} \end{aligned}$$

## Example 4

Consider

$$t^2 y'' - 2ty' + 2y = t \ln(t), \quad t > 0$$

$$y'' - \frac{2}{t}y' + \frac{2}{t^2}y = \frac{\ln(t)}{t}$$

Two solutions of  $L(y) = 0$  are  $y_1 = t$  and  $y_2 = t^2$ .

$$\text{The Wronskian is } W(y_1, y_2) = \begin{vmatrix} t & t^2 \\ 1 & 2t \end{vmatrix} = 2t^2 - t^2 = t^2$$

So, using Cramer's Rule

$$\begin{aligned} v_1' &= -\frac{y_2 f}{W(y_1, y_2)} = -\frac{\ln(t)}{t} \rightarrow v_1 = -\frac{1}{2} \ln^2(t) \\ v_2' &= \frac{y_1 f}{W(y_1, y_2)} = \frac{\ln(t)}{t^2} \rightarrow v_2 = -\frac{\ln(t) + 1}{t} \end{aligned}$$

The general solution is

$$y = c_1 t + c_2 t^2 - \frac{t}{2} \ln^2(t) - t \ln(t) - t$$