

Eigenvalues and Eigenvectors

Adam Wilson

Salt Lake Community College

Example

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$


Example

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

We can see how T maps a few vectors:

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \longrightarrow$$




Example

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

We can see how T maps a few vectors:

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ -2 \end{bmatrix} \quad \longrightarrow$$




Example

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

We can see how T maps a few vectors:

$$T\left(\begin{bmatrix} -1 \\ -1 \end{bmatrix}\right) = \begin{bmatrix} -3 \\ 0 \end{bmatrix} \quad \longrightarrow$$




Example

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

But something interesting happens for some special vectors.

$$T\left(\begin{bmatrix} 2 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 2 \end{bmatrix} = 2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \longrightarrow$$




Example

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

But something interesting happens for some special vectors.

$$T\left(\begin{bmatrix} 1 \\ -2 \end{bmatrix}\right) = \begin{bmatrix} -3 \\ 6 \end{bmatrix} = -3 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \rightarrow$$



Eigenvalues and Eigenvectors

Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation from vector space \mathbb{V} into itself. A scalar λ is a **eigenvalue** of T if there is a *nonzero* vector $\vec{v} \in \mathbb{V}$ such that

$$T(\vec{v}) = \lambda \vec{v}$$

Such a nonzero vector \vec{v} is called an **eigenvector** of T corresponding to λ .

Eigenvalues and Eigenvectors

Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation from vector space \mathbb{V} into itself. A scalar λ is a **eigenvalue** of T if there is a *nonzero* vector $\vec{v} \in \mathbb{V}$ such that

$$T(\vec{v}) = \lambda \vec{v}$$

Such a nonzero vector \vec{v} is called an **eigenvector** of T corresponding to λ .

Eigenvalues are also called **proper values** or **characteristic values**.

Eigenvalues and Eigenvectors

Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation from vector space \mathbb{V} into itself. A scalar λ is a **eigenvalue** of T if there is a *nonzero* vector $\vec{v} \in \mathbb{V}$ such that

$$T(\vec{v}) = \lambda \vec{v}$$

Such a nonzero vector \vec{v} is called an **eigenvector** of T corresponding to λ .

Eigenvalues are also called **proper values** or **characteristic values**.

If the linear transformation T is represented by an $n \times n$ matrix \mathbf{A} , where $\mathbb{V} = \mathbb{R}^n$ and $T(\vec{v}) = \mathbf{A}\vec{v}$, then λ and \vec{v} are characterized by the equation

$$\mathbf{A}\vec{v} = \lambda \vec{v}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

While this equation always has the trivial solution $\vec{v} = \vec{0}$, we are looking for any non-zero solutions. Therefore, we are looking for when

$$|\mathbf{A} - \lambda\mathbf{I}_n| = 0$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

While this equation always has the trivial solution $\vec{v} = \vec{0}$, we are looking for any non-zero solutions. Therefore, we are looking for when

$$|\mathbf{A} - \lambda\mathbf{I}_n| = 0$$

This is called the **characteristic equation** of matrix A .

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

While this equation always has the trivial solution $\vec{v} = \vec{0}$, we are looking for any non-zero solutions. Therefore, we are looking for when

$$|\mathbf{A} - \lambda\mathbf{I}_n| = 0$$

This is called the **characteristic equation** of matrix \mathbf{A} .

The polynomial in λ denoted by

$$p(\lambda) = |\mathbf{A} - \lambda\mathbf{I}_n|$$

is called the **characteristic polynomial** of \mathbf{A} .

Summary of Steps for Finding Eigenvalues and Eigenvectors

- 1 Write the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}_n| = 0$.
- 2 Solve the characteristic equation for λ .
- 3 For each eigenvalue λ_i , find the corresponding eigenvector \vec{v}_i by solving the system of equations

$$(\mathbf{A} - \lambda_i \mathbf{I}_n) \vec{v}_i = \vec{0}$$

Summary of Steps for Finding Eigenvalues and Eigenvectors

- 1 Write the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}_n| = 0$.
- 2 Solve the characteristic equation for λ .
- 3 For each eigenvalue λ_i , find the corresponding eigenvector \vec{v}_i by solving the system of equations

$$(\mathbf{A} - \lambda_i \mathbf{I}_n) \vec{v}_i = \vec{0}$$

Note

Eigenvectors are *not* unique. An eigenvector is just a direction, any nonzero multiple of \vec{v}_i works just as well.

Summary of Steps for Finding Eigenvalues and Eigenvectors

- 1 Write the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}_n| = 0$.
- 2 Solve the characteristic equation for λ .
- 3 For each eigenvalue λ_i , find the corresponding eigenvector \vec{v}_i by solving the system of equations

$$(\mathbf{A} - \lambda_i \mathbf{I}_n) \vec{v}_i = \vec{0}$$

Note

Eigenvectors are *not* unique. An eigenvector is just a direction, any nonzero multiple of \vec{v}_i works just as well.

For large matrices the steps become cumbersome, so computer algebra systems are often employed.

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 2 \\ 2 & -2 - \lambda \end{vmatrix} = 0$$
$$\rightarrow (1 - \lambda)(-2 - \lambda) - 4 = 0$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0$$
$$\rightarrow (1-\lambda)(-2-\lambda) - 4 = 0 \rightarrow \lambda^2 + \lambda - 6 = 0$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \end{aligned}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 \\ 2 & -4 \end{bmatrix} \begin{vmatrix} 0 \\ 0 \end{vmatrix}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \left[\begin{array}{cc|c} -1 & 2 & 0 \\ 2 & -4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & 0 \\ 2 & -4 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 4 & 2 & | & 0 \\ 2 & 1 & | & 0 \end{bmatrix}$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \left[\begin{array}{cc|c} -1 & 2 & 0 \\ 2 & -4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & -2 & 0 \\ 0 & 0 & 0 \end{array} \right] \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \left[\begin{array}{cc|c} 4 & 2 & 0 \\ 2 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Example

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 4 & 2 & | & 0 \\ 2 & 1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_2 = s \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \left[\begin{array}{cc|c} -2 & 1 & 0 \\ 4 & -2 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 2 & 1 & | & 0 \\ 4 & 2 & | & 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 2 & 1 & | & 0 \\ 4 & 2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 2 & 1 & | & 0 \\ 4 & 2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_2 = s \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Which simplifies to:

$$\lambda^3 - 2\lambda^2 - \lambda + 2 = 0$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Which simplifies to:

$$\begin{aligned} \lambda^3 - 2\lambda^2 - \lambda + 2 &= 0 \\ (\lambda - 2)(\lambda - 1)(\lambda + 1) &= 0 \end{aligned}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Which simplifies to:

$$\begin{aligned} \lambda^3 - 2\lambda^2 - \lambda + 2 &= 0 \\ (\lambda - 2)(\lambda - 1)(\lambda + 1) &= 0 \end{aligned}$$

So, the eigenvalues are $\lambda_1 = 2$, $\lambda_2 = 1$, and $\lambda_3 = -1$.

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\begin{bmatrix} 1 - (2) & 1 & -2 \\ -1 & 2 - (2) & 1 \\ 0 & 1 & -1 - (2) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\begin{bmatrix} -1 & 1 & -2 \\ -1 & 0 & 1 \\ 0 & 1 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\left[\begin{array}{ccc|c} -1 & 1 & -2 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 1 & -3 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = v_3$ and $v_2 = 3v_3$. Replacing v_3 with parameter s gives

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - (1) & 1 & -2 \\ -1 & 2 - (1) & 1 \\ 0 & 1 & -1 - (1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\begin{bmatrix} 0 & 1 & -2 \\ -1 & 1 & 1 \\ 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & -2 & 0 \\ -1 & 1 & 1 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = 3v_3$ and $v_2 = 2v_3$. Replacing v_3 with parameter s gives

$$\vec{v}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\begin{bmatrix} 1 - (-1) & 1 & -2 \\ -1 & 2 - (-1) & 1 \\ 0 & 1 & -1 - (-1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\begin{bmatrix} 2 & 1 & -2 \\ -1 & 3 & 1 \\ 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 2 & 1 & -2 & 0 \\ -1 & 3 & 1 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = v_3$ and $v_2 = 0$. Replacing v_3 with parameter s gives

$$\vec{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Special Cases

Triangular Matrices: The eigenvalues of an upper (or lower) triangular matrix appear on the main diagonal.

Special Cases

Triangular Matrices: The eigenvalues of an upper (or lower) triangular matrix appear on the main diagonal.

2×2 Matrices: For

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The eigenvalues are the solutions to

$$\lambda^2 - (\text{tr } \mathbf{A})\lambda + |\mathbf{A}| = 0$$

Special Cases

Triangular Matrices: The eigenvalues of an upper (or lower) triangular matrix appear on the main diagonal.

2×2 Matrices: For

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The eigenvalues are the solutions to

$$\lambda^2 - (\text{tr } \mathbf{A})\lambda + |\mathbf{A}| = 0$$

Trace

The **trace** of a matrix, **tr \mathbf{A}** , is the sum of all elements in the diagonal.

Eigenspace Theorem for Linear Transformations

For each eigenvalue λ of a linear transformations $T : \mathbb{V} \rightarrow \mathbb{V}$, the **eigenspace**, defined by

$$\mathbb{E}_\lambda = \{\vec{v} \in \mathbb{V} \mid T(\vec{v}) = \lambda\vec{v}\}$$

is a subspace of \mathbb{V} .


Eigenspace Theorem for Linear Transformations

For each eigenvalue λ of a linear transformations $T : \mathbb{V} \rightarrow \mathbb{V}$, the **eigenspace**, defined by

$$\mathbb{E}_\lambda = \{\vec{v} \in \mathbb{V} \mid T(\vec{v}) = \lambda \vec{v}\}$$

is a subspace of \mathbb{V} .

Example



Example

For the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

we had the following eigenvectors:

$$\begin{aligned} \lambda_1 = 2 \quad \vec{v}_1 &= \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \\ \lambda_2 = 1 \quad \vec{v}_2 &= \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \\ \lambda_3 = -1 \quad \vec{v}_3 &= \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

Example

For the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

we had the following eigenvectors:

$$\begin{array}{lll} \lambda_1 = 2 & \vec{v}_1 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} & \mathbb{E}_{\lambda_1} = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \right\} \\ \lambda_2 = 1 & \vec{v}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} & \mathbb{E}_{\lambda_2} = \text{span} \left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \right\} \\ \lambda_3 = -1 & \vec{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} & \mathbb{E}_{\lambda_3} = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\} \end{array}$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\vec{v}_2 = c\vec{v}_1 \quad \text{where } c \neq 0$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

But, we could also have multiplied by \mathbf{A}

$$\mathbf{A}\vec{v}_2 = c\mathbf{A}\vec{v}_1$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

But, we could also have multiplied by \mathbf{A}

$$\begin{aligned}\mathbf{A}\vec{v}_2 &= c\mathbf{A}\vec{v}_1 \\ \lambda_2 \vec{v}_2 &= c\lambda_1 \vec{v}_1\end{aligned}$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

But, we could also have multiplied by \mathbf{A}

$$\begin{aligned}\mathbf{A}\vec{v}_2 &= c\mathbf{A}\vec{v}_1 \\ \lambda_2 \vec{v}_2 &= c\lambda_1 \vec{v}_1\end{aligned}$$

Which would imply that $\lambda_1 = \lambda_2$,

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| &= 0 \\ \lambda(\lambda + 3)^2 &= 0 \end{aligned}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| &= 0 \\ \lambda(\lambda + 3)^2 &= 0 \end{aligned}$$

So, the eigenvalues are $\lambda_1 = 0$, $\lambda_2 = -3$. (Note that -3 is a repeated root.)

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\begin{bmatrix} -2 - \lambda & 1 & 1 \\ 1 & -2 - \lambda & 1 \\ 1 & 1 & -2 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\begin{bmatrix} -2 - (0) & 1 & 1 \\ 1 & -2 - (0) & 1 \\ 1 & 1 & -2 - (0) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\left[\begin{array}{ccc|c} -2 & 1 & 1 & 0 \\ 1 & -2 & 1 & 0 \\ 1 & 1 & -2 & 0 \end{array} \right]$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = v_3$ and $v_2 = v_3$. Replacing v_3 with parameter s gives

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\begin{bmatrix} -2 - \lambda & 1 & 1 \\ 1 & -2 - \lambda & 1 \\ 1 & 1 & -2 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\begin{bmatrix} -2 - (-3) & 1 & 1 \\ 1 & -2 - (-3) & 1 \\ 1 & 1 & -2 - (-3) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \end{array} \right]$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = -v_2 - v_3$. This means we need two parameters, $v_2 = r$ and $v_3 = s$. Which means we have two linearly independent eigenvectors.

$$\vec{v}_2 = \begin{bmatrix} -r - s \\ 1 \\ 1 \end{bmatrix} = r \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + s \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

This means the eigenspace is

$$\mathbb{E}_{\lambda_2} = \mathbf{span} \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

which is a two-dimensional subspace of \mathbb{R}^3 .

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

This means the eigenspace is

$$\mathbb{E}_{\lambda_2} = \mathbf{span} \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

which is a two-dimensional subspace of \mathbb{R}^3 .

Any linear combination of these two vectors is also an eigenvector, which means that the eigenspace is a plane.

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Since this is an upper diagonal matrix, we know that the eigenvalue is $\lambda = 1$, with multiplicity of 3.

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & 1 \\ 0 & 1 - \lambda & 1 \\ 0 & 0 & 1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - (1) & 1 & 1 \\ 0 & 1 - (1) & 1 \\ 0 & 0 & 1 - (1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_2 + v_3 = 0$ and $v_3 = 0$. Replacing v_1 with parameter s gives

$$\vec{v} = \begin{bmatrix} s \\ 0 \\ 0 \end{bmatrix} = s \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Example

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_2 + v_3 = 0$ and $v_3 = 0$. Replacing v_1 with parameter s gives

$$\vec{v} = \begin{bmatrix} s \\ 0 \\ 0 \end{bmatrix} = s \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Which means the eigenspace has dimension 1.

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0$$

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0$$

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

The only solutions to this equation is $\lambda = \pm i$.

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

The only solutions to this equation is $\lambda = \pm i$.

The situation here is like that with quadratic equations: there are some things which just cannot be done in a real-valued world.

Example

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

The only solutions to this equation is $\lambda = \pm i$.

The situation here is like that with quadratic equations: there are some things which just cannot be done in a real-valued world.

We can continue in the same way and find that the eigenvectors are

$$\vec{\mathbf{v}}_1 = \begin{bmatrix} -i \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{\mathbf{v}}_2 = \begin{bmatrix} -1 \\ -i \end{bmatrix}$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \end{aligned}$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \end{aligned}$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \\ &\rightarrow \cos(\theta) - \lambda = \pm i \sin(\theta) \end{aligned}$$

Example

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \\ &\rightarrow \cos(\theta) - \lambda = \pm i \sin(\theta) \\ &\rightarrow \lambda = \cos(\theta) \pm i \sin(\theta) \end{aligned}$$

Example

Consider the rotation transformation


$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \\ &\rightarrow \cos(\theta) - \lambda = \pm i \sin(\theta) \\ &\rightarrow \lambda = \cos(\theta) \pm i \sin(\theta) \end{aligned}$$

Which means these eigenvalues rotate a vector, instead of scaling it.

Example



Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.
- \mathbf{A} has a zero eigenvalue if and only if $|\mathbf{A}| = 0$

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.
- \mathbf{A} has a zero eigenvalue if and only if $|\mathbf{A}| = 0$
- \mathbf{A} and \mathbf{A}^T have the same characteristic polynomials and the same eigenvalues.

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.
- \mathbf{A} has a zero eigenvalue if and only if $|\mathbf{A}| = 0$
- \mathbf{A} and \mathbf{A}^T have the same characteristic polynomials and the same eigenvalues.
- If λ is an eigenvalue of an invertible matrix \mathbf{A} , then $\frac{1}{\lambda}$ is an eigenvalue of \mathbf{A}^{-1} .

Characteristic roots of a linear homogeneous DE are eigenvalues.

Characteristic roots of a linear homogeneous DE are eigenvalues.

Properties of Linear Homogeneous Differential Equations with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

Characteristic roots of a linear homogeneous DE are eigenvalues.

Properties of Linear Homogeneous Differential Equations with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.

Characteristic roots of a linear homogeneous DE are eigenvalues.

Properties of Linear Homogeneous Differential Equations with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.
- The solution set is also a vector space of vector functions.

Characteristic roots of a linear homogeneous DE are eigenvalues.

Properties of Linear Homogeneous Differential Equations with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.
- The solution set is also a vector space of vector functions.
- The eigenspace for each eigenvalue is a one-dimensional line in the direction of a vector in \mathbb{R}^n .

Characteristic roots of a linear homogeneous DE are eigenvalues.

Properties of Linear Homogeneous Differential Equations with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.
- The solution set is also a vector space of vector functions.
- The eigenspace for each eigenvalue is a one-dimensional line in the direction of a vector in \mathbb{R}^n .

We will explore the connection between eigenvalues and solutions to differential equations in chapter 6