

# Inverses of Matrices and Matrix Equations

Department of Mathematics

Salt Lake Community College

## Inverse Matrix

If there exists, for an  $n \times n$  matrix  $\mathbf{A}$ , another matrix  $\mathbf{A}^{-1}$  of the same order such that

$$\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}_n$$

then  $\mathbf{A}^{-1}$  is called the **inverse** of matrix  $\mathbf{A}$ , and  $\mathbf{A}$  is called **invertible**.

## Inverse Matrix

If there exists, for an  $n \times n$  matrix  $\mathbf{A}$ , another matrix  $\mathbf{A}^{-1}$  of the same order such that

$$\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}_n$$

then  $\mathbf{A}^{-1}$  is called the **inverse** of matrix  $\mathbf{A}$ , and  $\mathbf{A}$  is called **invertible**.

## Vocabulary

- A square matrix that is not invertible is called **singular**.
- A square matrix that is invertible is called **nonsingular**.

## Inverse Matrix

If there exists, for an  $n \times n$  matrix  $\mathbf{A}$ , another matrix  $\mathbf{A}^{-1}$  of the same order such that

$$\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}_n$$

then  $\mathbf{A}^{-1}$  is called the **inverse** of matrix  $\mathbf{A}$ , and  $\mathbf{A}$  is called **invertible**.

## Vocabulary

- A square matrix that is not invertible is called **singular**.
- A square matrix that is invertible is called **nonsingular**.

## Invertible Matrix Properties

- If  $\mathbf{A}$  is invertible, then so is  $\mathbf{A}^{-1}$  and  $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$

## Inverse Matrix

If there exists, for an  $n \times n$  matrix  $\mathbf{A}$ , another matrix  $\mathbf{A}^{-1}$  of the same order such that

$$\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}_n$$

then  $\mathbf{A}^{-1}$  is called the **inverse** of matrix  $\mathbf{A}$ , and  $\mathbf{A}$  is called **invertible**.

## Vocabulary

- A square matrix that is not invertible is called **singular**.
- A square matrix that is invertible is called **nonsingular**.

## Invertible Matrix Properties

- If  $\mathbf{A}$  is invertible, then so is  $\mathbf{A}^{-1}$  and  $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$
- If  $\mathbf{A}$  and  $\mathbf{B}$  are invertible matrices of the same order, then their product  $\mathbf{AB}$  is invertible. In fact,  $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$

## Inverse of a $2 \times 2$ Matrix

If

$$\mathbf{A} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

then,

$$\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{vmatrix} d & -b \\ -c & a \end{vmatrix}$$

If  $ad - bc = 0$ , then  $\mathbf{A}$  is not invertible.

## Inverse of a $2 \times 2$ Matrix

If

$$\mathbf{A} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

then,

$$\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{vmatrix} d & -b \\ -c & a \end{vmatrix}$$

If  $ad - bc = 0$ , then  $\mathbf{A}$  is not invertible.

## Inverses by Reduced Row Echelon Form

For an  $n \times n$  matrix  $\mathbf{A}$ , the following process will calculate  $\mathbf{A}^{-1}$ , or show that  $\mathbf{A}$  is not invertible.

## Inverse of a $2 \times 2$ Matrix

If

$$\mathbf{A} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

then,

$$\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{vmatrix} d & -b \\ -c & a \end{vmatrix}$$

If  $ad - bc = 0$ , then  $\mathbf{A}$  is not invertable.

## Inverses by Reduced Row Echelon Form

For an  $n \times n$  matrix  $\mathbf{A}$ , the following process will calculate  $\mathbf{A}^{-1}$ , or show that  $\mathbf{A}$  is not invertible.

Step 1: Form the  $n \times 2n$  augmented matrix  $\mathbf{M} = [\mathbf{A} | \mathbf{I}_n]$ .



## Inverse of a $2 \times 2$ Matrix

If

$$\mathbf{A} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

then,

$$\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{vmatrix} d & -b \\ -c & a \end{vmatrix}$$

If  $ad - bc = 0$ , then  $\mathbf{A}$  is not invertable.

## Inverses by Reduced Row Echelon Form

For an  $n \times n$  matrix  $\mathbf{A}$ , the following process will calculate  $\mathbf{A}^{-1}$ , or show that  $\mathbf{A}$  is not invertible.

**Step 1:** Form the  $n \times 2n$  augmented matrix  $\mathbf{M} = [\mathbf{A} | \mathbf{I}_n]$ .

**Step 2:** Transform  $\mathbf{M}$  into Reduced Row Echelon Form.

## Inverse of a $2 \times 2$ Matrix

If

$$\mathbf{A} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

then,

$$\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{vmatrix} d & -b \\ -c & a \end{vmatrix}$$

If  $ad - bc = 0$ , then  $\mathbf{A}$  is not invertible.

## Inverses by Reduced Row Echelon Form

For an  $n \times n$  matrix  $\mathbf{A}$ , the following process will calculate  $\mathbf{A}^{-1}$ , or show that  $\mathbf{A}$  is not invertible.

**Step 1:** Form the  $n \times 2n$  augmented matrix  $\mathbf{M} = [\mathbf{A} | \mathbf{I}_n]$ .

**Step 2:** Transform  $\mathbf{M}$  into Reduced Row Echelon Form.

**Step 3:**

- If the left hand side of  $\mathbf{M}$  is the identity matrix, then the right hand side is  $\mathbf{A}^{-1}$ .
- Otherwise,  $\mathbf{A}$  is a non-invertible matrix.

## Example 1

Consider the matrix:

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

Find  $\mathbf{A}^{-1}$

## Example 1

Consider the matrix:

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

Find  $\mathbf{A}^{-1}$

Start by building the augmented matrix

$$\mathbf{M}_A = \left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

Then transform  $\mathbf{M}_A$  into Reduced Row Echelon Form.

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right] R_3 = r_3 - r_1$$

### Example 1

$$\begin{bmatrix} 1 & 1 & 1 & | & 1 & 0 & 0 \\ 0 & 2 & 1 & | & 0 & 1 & 0 \\ 1 & 0 & 1 & | & 0 & 0 & 1 \end{bmatrix} R_3 = r_3 - r_1$$
$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 & | & 1 & 0 & 0 \\ 0 & 2 & 1 & | & 0 & 1 & 0 \\ 0 & -1 & 0 & | & -1 & 0 & 1 \end{bmatrix}$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \end{array} \right]$$



### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \end{array} \right] \begin{array}{l} \\ R_2 = -r_3 \\ R_3 = r_2 \end{array}$$

### Example 1

$$\begin{aligned} & \left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 = -r_3 \\ R_3 = r_2 \end{array} \\ \Rightarrow & \left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 2 & 1 & 0 & 1 & 0 \end{array} \right] \end{aligned}$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 2 & 1 & 0 & 1 & 0 \end{array} \right]$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 2 & 1 & 0 & 1 & 0 \end{array} \right] R_3 = r_3 - 2r_2$$

### Example 1

$$\begin{aligned} & \left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 2 & 1 & 0 & 1 & 0 \end{array} \right] R_3 = r_3 - 2r_2 \\ \Rightarrow & \left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 & 1 & 2 \end{array} \right] \end{aligned}$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 & 1 & 2 \end{array} \right]$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 & 1 & 2 \end{array} \right] R_1 = r_1 - r_3$$

### Example 1

$$\begin{bmatrix} 1 & 1 & 1 & | & 1 & 0 & 0 \\ 0 & 1 & 0 & | & 1 & 0 & -1 \\ 0 & 0 & 1 & | & -2 & 1 & 2 \end{bmatrix} R_1 = r_1 - r_3$$
$$\Rightarrow \begin{bmatrix} 1 & 1 & 0 & | & 3 & -1 & -2 \\ 0 & 1 & 0 & | & 1 & 0 & -1 \\ 0 & 0 & 1 & | & -2 & 1 & 2 \end{bmatrix}$$



### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 0 & 3 & -1 & -2 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 & 1 & 2 \end{array} \right]$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 0 & 3 & -1 & -2 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 & 1 & 2 \end{array} \right] R_1 = r_1 - r_2$$

### Example 1

$$\begin{bmatrix} 1 & 1 & 0 & | & 3 & -1 & -2 \\ 0 & 1 & 0 & | & 1 & 0 & -1 \\ 0 & 0 & 1 & | & -2 & 1 & 2 \end{bmatrix} R_1 = r_1 - r_2$$
$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 & | & 2 & -1 & -1 \\ 0 & 1 & 0 & | & 1 & 0 & -1 \\ 0 & 0 & 1 & | & -2 & 1 & 2 \end{bmatrix}$$

### Example 1

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 2 & -1 & -1 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -2 & 1 & 2 \end{array} \right]$$

Since the left hand side is  $I_3$ , we know the right hand side is the inverse:

$$\mathbf{A}^{-1} = \begin{bmatrix} 2 & -1 & -1 \\ 1 & 0 & -1 \\ -2 & 1 & 2 \end{bmatrix}$$

## Example 2

Consider the matrix:

$$\mathbf{B} = \begin{bmatrix} 3 & 0 & 3 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Find  $\mathbf{B}^{-1}$

## Example 2

Consider the matrix:

$$\mathbf{B} = \begin{bmatrix} 3 & 0 & 3 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Find  $\mathbf{B}^{-1}$

Start by building the augmented matrix

$$\mathbf{M}_B = \left[ \begin{array}{ccc|ccc} 3 & 0 & 3 & 1 & 0 & 0 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

Then transform  $\mathbf{M}_B$  into Reduced Row Echelon Form.

## Example 2

$$\left[ \begin{array}{ccc|ccc} 3 & 0 & 3 & 1 & 0 & 0 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

## Example 2

$$\left[ \begin{array}{ccc|ccc} 3 & 0 & 3 & 1 & 0 & 0 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_1 = r_3 \\ \\ R_3 = r_1 \end{array}$$



## Example 2

$$\begin{aligned} & \left[ \begin{array}{ccc|ccc} 3 & 0 & 3 & 1 & 0 & 0 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 1 & 1 & 2 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_1 = r_3 \\ \\ R_3 = r_1 \end{array} \\ \Rightarrow & \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 3 & 0 & 3 & 1 & 0 & 0 \end{array} \right] \end{aligned}$$

## Example 2

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 3 & 0 & 3 & 1 & 0 & 0 \end{array} \right]$$

## Example 2

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 3 & 0 & 3 & 1 & 0 & 0 \end{array} \right] \begin{array}{l} \\ R_2 = r_2 + r_1 \\ R_3 = r_2 - 3r_1 \end{array}$$

## Example 2

$$\begin{aligned} & \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ -1 & 2 & 1 & 0 & 1 & 0 \\ 3 & 0 & 3 & 1 & 0 & 0 \end{array} \right] \begin{array}{l} R_2 = r_2 + r_1 \\ R_3 = r_2 - 3r_1 \end{array} \\ \Rightarrow & \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 3 & 3 & 0 & 1 & 1 \\ 0 & -3 & -3 & 1 & 0 & -1 \end{array} \right] \end{aligned}$$

## Example 2

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 3 & 3 & 0 & 1 & 1 \\ 0 & -3 & -3 & 1 & 0 & -1 \end{array} \right]$$

## Example 2

$$\left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 3 & 3 & 0 & 1 & 1 \\ 0 & -3 & -3 & 1 & 0 & -1 \end{array} \right] \begin{array}{l} \\ R_2 = \frac{1}{3}r_2 \\ R_3 = r_3 + r_2 \end{array}$$

## Example 2

$$\begin{aligned} & \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 3 & 3 & 0 & 1 & 1 \\ 0 & -3 & -3 & 1 & 0 & -1 \end{array} \right] \begin{array}{l} R_2 = \frac{1}{3}r_2 \\ R_3 = r_3 + r_2 \end{array} \\ \Rightarrow & \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & \frac{1}{3} & \frac{1}{3} \\ 0 & 0 & 0 & 1 & 1 & 0 \end{array} \right] \end{aligned}$$

## Example 2

$$\Rightarrow \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 3 & 3 & 0 & 1 & 1 \\ 0 & -3 & -3 & 1 & 0 & -1 \end{array} \right] \begin{array}{l} R_2 = \frac{1}{3}r_2 \\ R_3 = r_3 + r_2 \end{array}$$
$$\Rightarrow \left[ \begin{array}{ccc|ccc} 1 & 1 & 2 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & \frac{1}{3} & \frac{1}{3} \\ 0 & 0 & 0 & 1 & 1 & 0 \end{array} \right]$$

This means that  $\mathbf{B}$  is a non-invertible matrix.



## Invertibility and Solutions

Consider the matrix equation  $\mathbf{A}\vec{x} = \vec{b}$ .

Where  $\mathbf{A}$  is an  $n \times n$  matrix, and  $\vec{x}$  and  $\vec{b}$  are of length  $n$ .

- A unique solution exists if and only if  $\mathbf{A}$  is invertible.

## Invertibility and Solutions

Consider the matrix equation  $\mathbf{A}\vec{x} = \vec{b}$ .

Where  $\mathbf{A}$  is an  $n \times n$  matrix, and  $\vec{x}$  and  $\vec{b}$  are of length  $n$ .

- A unique solution exists if and only if  $\mathbf{A}$  is invertible.
- Otherwise there are either:
  - No solutions.
  - Infinitely many solutions.

(Another method must be used to determine which.)

### Example 3

Consider the system

$$\begin{array}{rcccccc} x & + & y & + & z & = & 2 \\ & & 2y & + & z & = & -1 \\ x & & & + & z & = & 3 \end{array}$$

### Example 3

Consider the system

$$\begin{array}{rcrcrcrcrcrl} x & + & y & + & z & = & 2 \\ & & 2y & + & z & = & -1 \\ x & & & & + & z & = & 3 \end{array}$$

We can write this as the matrix equation:

$$\underbrace{\begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 1 \end{bmatrix}}_{\mathbf{A}} \underbrace{\begin{bmatrix} x \\ y \\ z \end{bmatrix}}_{\vec{x}} = \underbrace{\begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}}_{\vec{b}}$$

### Example 3

So, if  $\mathbf{A}$  is invertible, then we can solve the matrix equation for  $\vec{x}$

$$\mathbf{A}\vec{x} = \vec{b}$$

### Example 3

So, if  $\mathbf{A}$  is invertible, then we can solve the matrix equation for  $\vec{x}$

$$\mathbf{A}\vec{x} = \vec{b}$$

$$\mathbf{A}^{-1}\mathbf{A}\vec{x} = \mathbf{A}^{-1}\vec{b}$$

### Example 3

So, if  $\mathbf{A}$  is invertible, then we can solve the matrix equation for  $\vec{x}$

$$\mathbf{A}\vec{x} = \vec{b}$$

$$\mathbf{A}^{-1}\mathbf{A}\vec{x} = \mathbf{A}^{-1}\vec{b}$$

$$\mathbf{I}_3\vec{x} = \mathbf{A}^{-1}\vec{b}$$

### Example 3

So, if  $\mathbf{A}$  is invertible, then we can solve the matrix equation for  $\vec{x}$

$$\mathbf{A}\vec{x} = \vec{b}$$

$$\mathbf{A}^{-1}\mathbf{A}\vec{x} = \mathbf{A}^{-1}\vec{b}$$

$$\mathbf{I}_3\vec{x} = \mathbf{A}^{-1}\vec{b}$$

$$\vec{x} = \mathbf{A}^{-1}\vec{b}$$



### Example 3

So, if  $\mathbf{A}$  is invertible, then we can solve the matrix equation for  $\vec{x}$

$$\mathbf{A}\vec{x} = \vec{b}$$

$$\mathbf{A}^{-1}\mathbf{A}\vec{x} = \mathbf{A}^{-1}\vec{b}$$

$$\mathbf{I}_3\vec{x} = \mathbf{A}^{-1}\vec{b}$$

$$\vec{x} = \mathbf{A}^{-1}\vec{b}$$

So, if we can compute  $\mathbf{A}^{-1}\vec{b}$  we will have solved the system.

### Example 3

$$\left[ \begin{array}{ccc|c} & & & 2 \\ & & & -1 \\ & & & 0 \\ \hline 2 & -1 & -1 & \\ 1 & 0 & -1 & \\ -2 & 1 & 2 & 5 \end{array} \right]$$

### Example 3

$$\left[ \begin{array}{ccc|c} & & & 2 \\ & & & -1 \\ & & & 0 \\ \hline 2 & -1 & -1 & \\ 1 & 0 & -1 & 5 \\ -2 & 1 & 2 & 2 \end{array} \right]$$

### Example 3

$$\begin{array}{ccc|c} & & & \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & -1 & -1 \\ 1 & 0 & -1 \\ -2 & 1 & 2 \end{bmatrix} & & \begin{bmatrix} 5 \\ 2 \\ -5 \end{bmatrix} \end{array}$$

### Example 3

$$\begin{array}{c|c} & \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} \\ \hline \begin{bmatrix} 2 & -1 & -1 \\ 1 & 0 & -1 \\ -2 & 1 & 2 \end{bmatrix} & \begin{bmatrix} 5 \\ 2 \\ -5 \end{bmatrix} \end{array}$$

So, we have

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \\ -5 \end{bmatrix}$$

## Invertible Matrix Characterization

Let  $\mathbf{A}$  be a  $n \times n$  matrix. The following are equivalent:

- $\mathbf{A}$  is an invertible matrix.

## Invertible Matrix Characterization

Let  $\mathbf{A}$  be a  $n \times n$  matrix. The following are equivalent:

- $\mathbf{A}$  is an invertible matrix.
- $\mathbf{A}$  is row equivalent to  $\mathbf{I}_n$ .

(This means when you put  $\mathbf{A}$  in RREF, you get  $\mathbf{I}_n$ )

## Invertible Matrix Characterization

Let  $\mathbf{A}$  be a  $n \times n$  matrix. The following are equivalent:

- $\mathbf{A}$  is an invertible matrix.
- $\mathbf{A}$  is row equivalent to  $\mathbf{I}_n$ .  
(This means when you put  $\mathbf{A}$  in RREF, you get  $\mathbf{I}_n$ )
- The equation  $\mathbf{A}\vec{x} = \vec{0}$  has only the trivial solution  $\vec{x} = \vec{0}$ .



## Invertible Matrix Characterization

Let  $\mathbf{A}$  be a  $n \times n$  matrix. The following are equivalent:

- $\mathbf{A}$  is an invertible matrix.
- $\mathbf{A}$  is row equivalent to  $\mathbf{I}_n$ .  
(This means when you put  $\mathbf{A}$  in RREF, you get  $\mathbf{I}_n$ )
- The equation  $\mathbf{A}\vec{x} = \vec{0}$  has only the trivial solution  $\vec{x} = \vec{0}$ .
- The equation  $\mathbf{A}\vec{x} = \vec{b}$  has a unique solution for every  $\vec{b} \in \mathbb{R}^n$ .