

# Binomial Distribution

Colby Community College

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ),    Brittany ( $B$ ),    Carlton ( $C$ ),    Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no})$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 \end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no})$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 \end{aligned}$$



## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no})$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no}) \\ = 0.15 \cdot 0.15 \cdot 0.85 \cdot 0.15 \end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no}) \\ = 0.15 \cdot 0.15 \cdot 0.85 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no}) \\ = 0.15 \cdot 0.15 \cdot 0.85 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$P(A = \text{no and } B = \text{no and } C = \text{no and } D = \text{yes})$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned}P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\= 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869\end{aligned}$$

$$\begin{aligned}P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\= 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869\end{aligned}$$

$$\begin{aligned}P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no}) \\= 0.15 \cdot 0.15 \cdot 0.85 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869\end{aligned}$$

$$\begin{aligned}P(A = \text{no and } B = \text{no and } C = \text{no and } D = \text{yes}) \\= 0.15 \cdot 0.15 \cdot 0.15 \cdot 0.85\end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no}) \\ = 0.15 \cdot 0.15 \cdot 0.85 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{no and } D = \text{yes}) \\ = 0.15 \cdot 0.15 \cdot 0.15 \cdot 0.85 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

## Example 1

Let us assume there is a 0.85 probability that a randomly chosen adult has heard of Twitter.

Four people are chosen at random:

Ariana ( $A$ ), Brittany ( $B$ ), Carlton ( $C$ ), Damian ( $D$ )

We want the probability exactly one of them will have heard of Twitter.

There are four combinations possible:

$$\begin{aligned} P(A = \text{yes and } B = \text{no and } C = \text{no and } D = \text{no}) \\ = 0.85 \cdot 0.15 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{yes and } C = \text{no and } D = \text{no}) \\ = 0.15 \cdot 0.85 \cdot 0.15 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{yes and } D = \text{no}) \\ = 0.15 \cdot 0.15 \cdot 0.85 \cdot 0.15 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

$$\begin{aligned} P(A = \text{no and } B = \text{no and } C = \text{no and } D = \text{yes}) \\ = 0.15 \cdot 0.15 \cdot 0.15 \cdot 0.85 = (0.85)^1 (0.15)^3 = 0.002869 \end{aligned}$$

So, the probability exactly one has heard of Twitter is

$$0.002869 + 0.002869 + 0.002869 + 0.002869 = 0.11475 = 11.475\%$$



## Definition

The **binomial distribution** is used to describe the number of successes in a fixed number of trials.

## Definition

The **binomial distribution** is used to describe the number of successes in a fixed number of trials.

## Notation

$p$	The probability of a success.
$q = 1 - p$	The probability of a failure.
$n$	The fixed number of trials.
$k$	The number of successes.

## Definition

The **binomial distribution** is used to describe the number of successes in a fixed number or trials.

## Notation

$p$	The probability of a success.
$q = 1 - p$	The probability of a failure.
$n$	The fixed number of trials.
$k$	The number of successes.

## Note

Example 1 is how to find a binomial distribution the hard way.

## Definition

The **binomial distribution** is used to describe the number of successes in a fixed number or trials.

## Notation

$p$	The probability of a success.
$q = 1 - p$	The probability of a failure.
$n$	The fixed number of trials.
$k$	The number of successes.

## Note

Example 1 is how to find a binomial distribution the hard way.

## Note

If all the scenarios are independent of each other, then we can calculate the final probability as:

$$[\text{\# of scenarios}] \cdot P(\text{single scenario})$$

## Definition

The **factorial**, for any positive integer  $n$ , is

$$0! = 1$$

$$1! = 1$$

$$2! = 2 \cdot 1 = 2$$

$$3! = 3 \cdot 2 \cdot 1 = 6$$

$$4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$$

$$\vdots$$

$$n! = n \cdot (n - 1) \cdot (n - 2) \cdot \dots \cdot 4 \cdot 3 \cdot 2 \cdot 1$$

## Definition

The **factorial**, for any positive integer  $n$ , is

$$0! = 1$$

$$1! = 1$$

$$2! = 2 \cdot 1 = 2$$

$$3! = 3 \cdot 2 \cdot 1 = 6$$

$$4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$$

$$\vdots$$

$$n! = n \cdot (n - 1) \cdot (n - 2) \cdot \dots \cdot 4 \cdot 3 \cdot 2 \cdot 1$$

## Note

Factorials can be calculated iteratively. i.e.

$$(n + 1)! = n! \cdot (n + 1)$$

## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Example 2

The number of ways to choose  $k = 3$  successes in  $n = 4$  trials:

$$\binom{4}{3}$$



## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Example 2

The number of ways to choose  $k = 3$  successes in  $n = 4$  trials:

$$\binom{4}{3} = \frac{4!}{3!(4-3)!}$$

## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Example 2

The number of ways to choose  $k = 3$  successes in  $n = 4$  trials:

$$\binom{4}{3} = \frac{4!}{3!(4-3)!} = \frac{4!}{3!1!}$$

## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Example 2

The number of ways to choose  $k = 3$  successes in  $n = 4$  trials:

$$\binom{4}{3} = \frac{4!}{3!(4-3)!} = \frac{4!}{3!1!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1 \cdot 1}$$

## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Example 2

The number of ways to choose  $k = 3$  successes in  $n = 4$  trials:

$$\binom{4}{3} = \frac{4!}{3!(4-3)!} = \frac{4!}{3!1!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1 \cdot 1} = \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{\cancel{3} \cdot \cancel{2} \cdot 1 \cdot 1}$$

## Definition

The **binomial coefficients** gives the number of ways to choose  $k$  successes in  $n$  trials.:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Read “ $n$  choose  $k$ .”

## Example 2

The number of ways to choose  $k = 3$  successes in  $n = 4$  trials:

$$\binom{4}{3} = \frac{4!}{3!(4-3)!} = \frac{4!}{3!1!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1 \cdot 1} = \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{\cancel{3} \cdot \cancel{2} \cdot 1 \cdot 1} = 4$$

## Binomial Distribution

Suppose the probability of a single trial being a success is  $p$ . Then the probability of observing exactly  $k$  successes in  $n$  independent trials is given by

$$\binom{n}{k} p^k (1 - p)^{n-k} = \frac{n!}{k!(n-k)!} p^k (1 - p)^{n-k}$$

The mean, variance, and standard deviation of the number of observed successes are

$$\mu = np, \quad \sigma^2 = np(1 - p), \quad \sigma = \sqrt{np(1 - p)}$$

## Binomial Distribution

Suppose the probability of a single trial being a success is  $p$ . Then the probability of observing exactly  $k$  successes in  $n$  independent trials is given by

$$\binom{n}{k} p^k (1 - p)^{n-k} = \frac{n!}{k!(n-k)!} p^k (1 - p)^{n-k}$$

The mean, variance, and standard deviation of the number of observed successes are

$$\mu = np, \quad \sigma^2 = np(1 - p), \quad \sigma = \sqrt{np(1 - p)}$$

## Is It Binomial?

Every binomial distribution has to satisfy the following:

- The trials are independent.

## Binomial Distribution

Suppose the probability of a single trial being a success is  $p$ . Then the probability of observing exactly  $k$  successes in  $n$  independent trials is given by

$$\binom{n}{k} p^k (1-p)^{n-k} = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}$$

The mean, variance, and standard deviation of the number of observed successes are

$$\mu = np, \quad \sigma^2 = np(1-p), \quad \sigma = \sqrt{np(1-p)}$$

## Is It Binomial?

Every binomial distribution has to satisfy the following:

- The trials are independent.
- The number of trials,  $n$ , is fixed.



## Binomial Distribution

Suppose the probability of a single trial being a success is  $p$ . Then the probability of observing exactly  $k$  successes in  $n$  independent trials is given by

$$\binom{n}{k} p^k (1 - p)^{n-k} = \frac{n!}{k!(n-k)!} p^k (1 - p)^{n-k}$$

The mean, variance, and standard deviation of the number of observed successes are

$$\mu = np, \quad \sigma^2 = np(1 - p), \quad \sigma = \sqrt{np(1 - p)}$$

## Is It Binomial?

Every binomial distribution has to satisfy the following:

- The trials are independent.
- The number of trials,  $n$ , is fixed.
- Each trial outcome can be classified as either a *success* or *failure*.

## Binomial Distribution

Suppose the probability of a single trial being a success is  $p$ . Then the probability of observing exactly  $k$  successes in  $n$  independent trials is given by

$$\binom{n}{k} p^k (1 - p)^{n-k} = \frac{n!}{k!(n-k)!} p^k (1 - p)^{n-k}$$

The mean, variance, and standard deviation of the number of observed successes are

$$\mu = np, \quad \sigma^2 = np(1 - p), \quad \sigma = \sqrt{np(1 - p)}$$

## Is It Binomial?

Every binomial distribution has to satisfy the following:

- The trials are independent.
- The number of trials,  $n$ , is fixed.
- Each trial outcome can be classified as either a *success* or *failure*.
- The probability of a success,  $p$ , is the same for each trial.

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$P(\text{exactly one has heard of twitter}) = \binom{n}{k} p^k (1 - p)^{n-k}$$

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$\begin{aligned} P(\text{exactly one has heard of twitter}) &= \binom{n}{k} p^k (1 - p)^{n-k} \\ &= \binom{4}{1} (0.85)^1 (1 - 0.85)^{4-1} \end{aligned}$$

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$\begin{aligned}P(\text{exactly one has heard of twitter}) &= \binom{n}{k} p^k (1 - p)^{n-k} \\&= \binom{4}{1} (0.85)^1 (1 - 0.85)^{4-1} \\&= \frac{4!}{1!(4-1)!} (0.85)^1 (0.15)^3\end{aligned}$$

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$\begin{aligned}P(\text{exactly one has heard of twitter}) &= \binom{n}{k} p^k (1 - p)^{n-k} \\&= \binom{4}{1} (0.85)^1 (1 - 0.85)^{4-1} \\&= \frac{4!}{1!(4-1)!} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1} (0.85)^1 (0.15)^3\end{aligned}$$

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$\begin{aligned}P(\text{exactly one has heard of twitter}) &= \binom{n}{k} p^k (1 - p)^{n-k} \\&= \binom{4}{1} (0.85)^1 (1 - 0.85)^{4-1} \\&= \frac{4!}{1!(4-1)!} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.85)^1 (0.15)^3\end{aligned}$$



### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$\begin{aligned}P(\text{exactly one has heard of twitter}) &= \binom{n}{k} p^k (1 - p)^{n-k} \\&= \binom{4}{1} (0.85)^1 (1 - 0.85)^{4-1} \\&= \frac{4!}{1!(4-1)!} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.85)^1 (0.15)^3 \\&= 4 \cdot (0.85)^1 (0.15)^3\end{aligned}$$

### Example 3

From Example 1 we have  $p = 0.85$ ,  $n = 4$ , and  $k = 1$ .

$$\begin{aligned}P(\text{exactly one has heard of twitter}) &= \binom{n}{k} p^k (1 - p)^{n-k} \\&= \binom{4}{1} (0.85)^1 (1 - 0.85)^{4-1} \\&= \frac{4!}{1!(4-1)!} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1} (0.85)^1 (0.15)^3 \\&= \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.85)^1 (0.15)^3 \\&= 4 \cdot (0.85)^1 (0.15)^3 \\&= 0.11475\end{aligned}$$

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\binom{n}{k} p^k (1 - p)^{n-k} = \binom{8}{5} (0.7)^5 (0.3)^{8-5}$$

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\begin{aligned} \binom{n}{k} p^k (1 - p)^{n-k} &= \binom{8}{5} (0.7)^5 (0.3)^{8-5} \\ &= \frac{8!}{5!(8-5)!} (0.7)^5 (0.3)^3 \end{aligned}$$

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\begin{aligned} \binom{n}{k} p^k (1 - p)^{n-k} &= \binom{8}{5} (0.7)^5 (0.3)^{8-5} \\ &= \frac{8!}{5!(8-5)!} (0.7)^5 (0.3)^3 = \frac{8!}{5!(3)!} (0.7)^5 (0.3)^3 \end{aligned}$$

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\begin{aligned} \binom{n}{k} p^k (1 - p)^{n-k} &= \binom{8}{5} (0.7)^5 (0.3)^{8-5} \\ &= \frac{8!}{5!(8-5)!} (0.7)^5 (0.3)^3 = \frac{8!}{5!(3)!} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \end{aligned}$$



### Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\begin{aligned} \binom{n}{k} p^k (1 - p)^{n-k} &= \binom{8}{5} (0.7)^5 (0.3)^{8-5} \\ &= \frac{8!}{5!(8-5)!} (0.7)^5 (0.3)^3 = \frac{8!}{5!(3)!} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot \cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{\cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \end{aligned}$$

### Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\begin{aligned} \binom{n}{k} p^k (1 - p)^{n-k} &= \binom{8}{5} (0.7)^5 (0.3)^{8-5} \\ &= \frac{8!}{5!(8-5)!} (0.7)^5 (0.3)^3 = \frac{8!}{5!(3)!} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot \cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{\cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \\ &= 56 \cdot (0.7)^5 (0.3)^3 \end{aligned}$$

## Example 4

Assume that 70% of customers won't exceed their car insurance deductible. Let's the probability that 5 of 8 randomly selected customers won't exceed their premium.

Start by identifying

$$p = 0.7, \quad q = 1 - p = 0.3, \quad n = 8, \quad k = 5$$

$$\begin{aligned} \binom{n}{k} p^k (1-p)^{n-k} &= \binom{8}{5} (0.7)^5 (0.3)^{8-5} \\ &= \frac{8!}{5!(8-5)!} (0.7)^5 (0.3)^3 = \frac{8!}{5!(3)!} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \\ &= \frac{8 \cdot 7 \cdot 6 \cdot \cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{\cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1 \cdot 3 \cdot 2 \cdot 1} (0.7)^5 (0.3)^3 \\ &= 56 \cdot (0.7)^5 (0.3)^3 \\ &= 0.254122 \end{aligned}$$

### Example 5

Assume the probability that a smoker will develop a severe lung condition in their life time is 0.3.

*If you have four friends who smoke, are the conditions for the binomial model satisfied?*

## Example 5

Assume the probability that a smoker will develop a severe lung condition in their life time is 0.3.

*If you have four friends who smoke, are the conditions for the binomial model satisfied?*

It is likely that independence is not satisfied, since they probably all know each other.

### Example 5

Assume the probability that a smoker will develop a severe lung condition in their life time is 0.3.

*If you have four friends who smoke, are the conditions for the binomial model satisfied?*

It is likely that independence is not satisfied, since they probably all know each other.

### Example 6

Suppose instead four people are randomly selected.

*Is the binomial model appropriate to find the probability that none of them will develop a severe lung condition?*

We are assuming that the four are randomly selected, yes.

$$\begin{aligned}\binom{n}{k} p^k (1-p)^{n-k} &= \binom{4}{0} (0.3)^0 (1-0.3)^{4-0} = \frac{4!}{0!(4-0)!} (0.3)^0 (0.7)^4 \\ &= 1 \cdot 1 \cdot (0.7)^4 = 0.2401\end{aligned}$$

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$P(\text{none}) + P(\text{exactly one})$$



## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$\begin{aligned} P(\text{none}) + P(\text{exactly one}) \\ = \binom{4}{0}(0.3)^0(1 - 0.3)^{4-0} + \binom{4}{1}(0.3)^1(1 - 0.3)^{4-1} \end{aligned}$$

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$\begin{aligned} &P(\text{none}) + P(\text{exactly one}) \\ &= \binom{4}{0}(0.3)^0(1 - 0.3)^{4-0} + \binom{4}{1}(0.3)^1(1 - 0.3)^{4-1} \\ &= \frac{4!}{0!(4 - 0)!}(0.3)^0(0.7)^4 + \frac{4!}{1!(4 - 1)!}(0.3)^1(0.7)^3 \end{aligned}$$

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$\begin{aligned} &P(\text{none}) + P(\text{exactly one}) \\ &= \binom{4}{0}(0.3)^0(1 - 0.3)^{4-0} + \binom{4}{1}(0.3)^1(1 - 0.3)^{4-1} \\ &= \frac{4!}{0!(4 - 0)!}(0.3)^0(0.7)^4 + \frac{4!}{1!(4 - 1)!}(0.3)^1(0.7)^3 \\ &= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 4 \cdot 3 \cdot 2 \cdot 1}(0.3)^0(0.7)^4 + \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1}(0.3)^1(0.7)^3 \end{aligned}$$

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$\begin{aligned} &P(\text{none}) + P(\text{exactly one}) \\ &= \binom{4}{0}(0.3)^0(1 - 0.3)^{4-0} + \binom{4}{1}(0.3)^1(1 - 0.3)^{4-1} \\ &= \frac{4!}{0!(4 - 0)!}(0.3)^0(0.7)^4 + \frac{4!}{1!(4 - 1)!}(0.3)^1(0.7)^3 \\ &= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 4 \cdot 3 \cdot 2 \cdot 1}(0.3)^0(0.7)^4 + \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1}(0.3)^1(0.7)^3 \\ &= \frac{\cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}(0.3)^0(0.7)^4 + \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}(0.3)^1(0.7)^3 \end{aligned}$$

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$P(\text{none}) + P(\text{exactly one})$$

$$= \binom{4}{0} (0.3)^0 (1 - 0.3)^{4-0} + \binom{4}{1} (0.3)^1 (1 - 0.3)^{4-1}$$

$$= \frac{4!}{0!(4-0)!} (0.3)^0 (0.7)^4 + \frac{4!}{1!(4-1)!} (0.3)^1 (0.7)^3$$

$$= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 4 \cdot 3 \cdot 2 \cdot 1} (0.3)^0 (0.7)^4 + \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1} (0.3)^1 (0.7)^3$$

$$= \frac{\cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.3)^0 (0.7)^4 + \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.3)^1 (0.7)^3$$

$$= 0.2401 + 0.4116$$

## Example 7

Let consider finding the probability than no more than one of the four people will develop a severe lung condition.

The events that “none of them develops a severe lung condition” and “exactly one develops a severe lung condition” are mutually exclusive.

$$P(\text{none}) + P(\text{exactly one})$$

$$= \binom{4}{0} (0.3)^0 (1 - 0.3)^{4-0} + \binom{4}{1} (0.3)^1 (1 - 0.3)^{4-1}$$

$$= \frac{4!}{0!(4-0)!} (0.3)^0 (0.7)^4 + \frac{4!}{1!(4-1)!} (0.3)^1 (0.7)^3$$

$$= \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 4 \cdot 3 \cdot 2 \cdot 1} (0.3)^0 (0.7)^4 + \frac{4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 3 \cdot 2 \cdot 1} (0.3)^1 (0.7)^3$$

$$= \frac{\cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.3)^0 (0.7)^4 + \frac{4 \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{1 \cdot \cancel{3} \cdot \cancel{2} \cdot 1} (0.3)^1 (0.7)^3$$

$$= 0.2401 + 0.4116$$

$$= 0.6517 = 65.17\%$$

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”



## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$P(\text{at least two}) = 1 - P(\text{no more than one})$$

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$\begin{aligned} P(\text{at least two}) &= 1 - P(\text{no more than one}) \\ &= 1 - 0.6517 \end{aligned}$$

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$\begin{aligned} P(\text{at least two}) &= 1 - P(\text{no more than one}) \\ &= 1 - 0.6517 = 0.3483 = 34.83\% \end{aligned}$$

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$\begin{aligned} P(\text{at least two}) &= 1 - P(\text{no more than one}) \\ &= 1 - 0.6517 = 0.3483 = 34.83\% \end{aligned}$$

## Example 9

*Out of seven randomly selected smokers, how many would we expect to develop a severe lung condition?*

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$\begin{aligned} P(\text{at least two}) &= 1 - P(\text{no more than one}) \\ &= 1 - 0.6517 = 0.3483 = 34.83\% \end{aligned}$$

## Example 9

*Out of seven randomly selected smokers, how many would we expect to develop a severe lung condition?*

The mean of the binomial model is

$$\mu = np$$

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$\begin{aligned} P(\text{at least two}) &= 1 - P(\text{no more than one}) \\ &= 1 - 0.6517 = 0.3483 = 34.83\% \end{aligned}$$

## Example 9

*Out of seven randomly selected smokers, how many would we expect to develop a severe lung condition?*

The mean of the binomial model is

$$\mu = np = 7 \cdot 0.3$$

## Example 8

Lets consider finding the probability that at least two of the four people will develop a severe lung condition.

The complement of “at least two will develop a severe lung condition” is “no more than one will develop a severe lung condition.”

We know from Example 7 that  $P(\text{no more than one}) = 0.6517$ .

So,

$$\begin{aligned} P(\text{at least two}) &= 1 - P(\text{no more than one}) \\ &= 1 - 0.6517 = 0.3483 = 34.83\% \end{aligned}$$

## Example 9

*Out of seven randomly selected smokers, how many would we expect to develop a severe lung condition?*

The mean of the binomial model is

$$\mu = np = 7 \cdot 0.3 = 2.1$$

On average, we would expect 2.1 of 7 randomly chosen smokers to develop a severe lung condition.