

Normal Distribution

Colby Community College

Example 1

Consider making popcorn.



Example 1

Consider making popcorn.

You put some oil and corn kernels in a pan and start heating.



Example 1



Consider making popcorn.

You put some oil and corn kernels in a pan and start heating.

For the first few minutes nothing happens, then a few kernels start to pop.

Example 1



Consider making popcorn.

You put some oil and corn kernels in a pan and start heating.

For the first few minutes nothing happens, then a few kernels start to pop.

A little while later more and more start to pop.

Example 1



Consider making popcorn.

You put some oil and corn kernels in a pan and start heating.

For the first few minutes nothing happens, then a few kernels start to pop.

A little while later more and more start to pop.

This goes on for a minute or so, and the popping gradually tapers off.

Example 1



Consider making popcorn.

You put some oil and corn kernels in a pan and start heating.

For the first few minutes nothing happens, then a few kernels start to pop.

A little while later more and more start to pop.

This goes on for a minute or so, and the popping gradually tapers off.

Most of the popping happens in that brief, noisy moment.

Example 1



Consider making popcorn.

You put some oil and corn kernels in a pan and start heating.

For the first few minutes nothing happens, then a few kernels start to pop.

A little while later more and more start to pop.


This goes on for a minute or so, and the popping gradually tapers off.

Most of the popping happens in that brief, noisy moment.

This demonstrates a typical pattern that is part of many phenomena.

Definition

A **normal distribution** is a perfectly symmetric, bell-shaped distribution. It is also referred to as a **normal curve** or a **bell curve**.



Note


The normal distribution with mean μ and standard deviation σ is denoted $N(\mu, \sigma)$, where μ and σ are called the parameters.

Note


The normal distribution with mean μ and standard deviation σ is denoted $N(\mu, \sigma)$, where μ and σ are called the parameters.

Example 2

Both are normal distributions, but with different center and spread.



(a) $N(\mu = 0, \sigma = 1)$



(b) $N(\mu = 19, \sigma = 4)$

Definition

The graph of any continuous probability distribution is called a **density curve** if the total area under the curve is exactly 1.

Definition

The graph of any continuous probability distribution is called a **density curve** if the total area under the curve is exactly 1.

Note

This means there is a correspondence between the area under a density curve and probabilities.

Definition


The graph of any continuous probability distribution is called a **density curve** if the total area under the curve is exactly 1.

Note

This means there is a correspondence between the area under a density curve and probabilities.

Definition


The special case $N(\mu = 0, \sigma = 1)$ is called the **standard normal distribution**. The total area under the curve is exactly equal to 1.



Definition

The **empirical rule**, or **68-95-99.7 rule**, states approximately how much of the area is contained when stepping one, two, or three standard deviations from the mean.


One standard deviation from the mean.



Definition

The **empirical rule**, or **68-95-99.7 rule**, states approximately how much of the area is contained when stepping one, two, or three standard deviations from the mean.


Two standard deviations from the mean.



Definition

The **empirical rule**, or **68-95-99.7 rule**, states approximately how much of the area is contained when stepping one, two, or three standard deviations from the mean.

Three standard deviations from the mean.



Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma}$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8}$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8} \approx 1.875$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8} \approx 1.875$$

Example 4

On the same exam, George has a z-score of -1.3 . What was his score?

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8} \approx 1.875$$

Example 4

On the same exam, George has a z-score of -1.3 . What was his score?

$$z = \frac{x - \mu}{\sigma}$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8} \approx 1.875$$

Example 4

On the same exam, George has a z-score of -1.3 . What was his score?

$$z = \frac{x - \mu}{\sigma} \Rightarrow z\sigma = x - \mu$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8} \approx 1.875$$

Example 4

On the same exam, George has a z-score of -1.3 . What was his score?

$$z = \frac{x - \mu}{\sigma} \Rightarrow z\sigma = x - \mu \Rightarrow x = z\sigma + \mu$$

Definition

A **z-score** is a measure of the number of standard deviations a particular data point is away from the mean.

$$z = \frac{(\text{data point}) - (\text{mean})}{\text{standard deviation}} = \frac{x - \mu}{\sigma}$$

Example 3

On a college entrance exam, the mean was 70, and the standard deviation was 8. Rose scored a 85, what is her z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{85 - 70}{8} \approx 1.875$$

Example 4

On the same exam, George has a z-score of -1.3 . What was his score?

$$z = \frac{x - \mu}{\sigma} \Rightarrow z\sigma = x - \mu \Rightarrow x = z\sigma + \mu = (-1.3)(8) + 70 = 59.6$$

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An "A" on the exam is a 93, what is the z-score?

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An "A" on the exam is a 93, what is the z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{93 - 82}{7} \approx 1.57$$

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An "A" on the exam is a 93, what is the z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{93 - 82}{7} \approx 1.57$$

Note

We know from the empirical rule that roughly 68% of the scores fall within one standard deviation of the mean.

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An "A" on the exam is a 93, what is the z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{93 - 82}{7} \approx 1.57$$

Note

We know from the empirical rule that roughly 68% of the scores fall within one standard deviation of the mean.

This means that 68% of the students scored between 75 and 89.

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An “A” on the exam is a 93, what is the z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{93 - 82}{7} \approx 1.57$$

Note

We know from the empirical rule that roughly 68% of the scores fall within one standard deviation of the mean.

This means that 68% of the students scored between 75 and 89.

Moreover, we know that roughly 95% of the scores fall within two standard deviations of the mean.

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An "A" on the exam is a 93, what is the z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{93 - 82}{7} \approx 1.57$$

Note

We know from the empirical rule that roughly 68% of the scores fall within one standard deviation of the mean.

This means that 68% of the students scored between 75 and 89.

Moreover, we know that roughly 95% of the scores fall within two standard deviations of the mean.

Which means that $95\% - 68\% = 27\%$ of the scores are more than one standard deviation from the mean, but less than two.

Example 5

The mean on a exam was 82, with a standard deviation of 7 points. An “A” on the exam is a 93, what is the z-score?

$$z = \frac{x - \mu}{\sigma} = \frac{93 - 82}{7} \approx 1.57$$

Note

We know from the empirical rule that roughly 68% of the scores fall within one standard deviation of the mean.

This means that 68% of the students scored between 75 and 89.

Moreover, we know that roughly 95% of the scores fall within two standard deviations of the mean.


Which means that $95\% - 68\% = 27\%$ of the scores are more than one standard deviation from the mean, but less than two.

Since the curve is symmetric, we know that 13.5% of the students scored between 89 and 96, as well as 13.5% between 68 and 75

Definition

The **empirical rule**, or **68-95-99.7 rule**, states approximately how much of the area is contained when stepping one, two, or three standard deviations from the mean.

For each standard deviation.



Definition


An **inflection point** is where a curve changes from being concave up to concave down, or vice versa

Definition


An **inflection point** is where a curve changes from being concave up to concave down, or vice versa

Note

A normal density curve always has two inflection points, each one standard deviation from the mean.




Example 6



The area of the shaded region is the probability that a z score is less than or equal to x , $P(z \leq x)$.

Example 6




The area of the shaded region is the probability that a z score is less than or equal to x , $P(z \leq x)$.

Note


Most statistical software, programming languages, spreadsheets programs, and calculators are able to calculate the area for you.

Example 7



The area of the shaded region is the probability that a z score is greater than or equal to x , $P(z \geq x)$.

Example 7




The area of the shaded region is the probability that a z score is greater than or equal to x , $P(z \geq x)$.

Note

The diagram shows three normal distribution curves. The first curve has the area to the right of x shaded blue, with the label $P(z \geq x)$ below it. This is followed by an equals sign. The second curve has the entire area under the curve shaded blue, with the label 1 below it. This is followed by a minus sign. The third curve has the area to the left of x shaded blue, with the label $P(z \leq x)$ below it.


$$P(z \geq x) = 1 - P(z \leq x)$$

Example 8



The area of the shaded region is the probability that a z score lies between x_1 and x_2 , $P(x_1 \leq z \leq x_2)$.

Example 8




The area of the shaded region is the probability that a z score lies between x_1 and x_2 , $P(x_1 \leq z \leq x_2)$.

Note

The diagram shows three normal distribution curves. The first curve has the area between x_1 and x_2 shaded blue. The second curve has the area to the left of x_2 shaded blue. The third curve has the area to the left of x_1 shaded blue. An equals sign and a minus sign are placed between the curves to show the relationship: the area between x_1 and x_2 is equal to the area to the left of x_2 minus the area to the left of x_1 .

$$P(x_1 \leq z \leq x_2) = P(z \leq x_2) - P(z \leq x_1)$$

Example 8



The area of the shaded region is the probability that a z score lies between x_1 and x_2 , $P(x_1 \leq z \leq x_2)$.

Note

The diagram shows three normal distribution curves. The first curve has the area between x_1 and x_2 shaded blue. The second curve has the area to the left of x_2 shaded blue. The third curve has the area to the left of x_1 shaded blue. The equation below these curves is:

$$P(x_1 \leq z \leq x_2) = P(z \leq x_2) - P(z \leq x_1)$$

Note

The diagram shows four normal distribution curves. The first curve has the area between x_1 and x_2 shaded blue. The second curve has the entire area under the curve shaded blue. The third curve has the area to the left of x_1 shaded blue. The fourth curve has the area to the right of x_2 shaded blue. The equation below these curves is:


$$P(x_1 \leq z \leq x_2) = 1 - P(z \leq x_1) - P(z \geq x_2)$$

Procedure for Finding Areas with a Nonstandard Normal Distribution

- 1 Sketch a normal curve, label the mean and any specific x values, and then shade the region representing the desired probability.


2

3




Procedure for Finding Areas with a Nonstandard Normal Distribution

- 1 Sketch a normal curve, label the mean and any specific x values, and then shade the region representing the desired probability.
- 2 For each relevant x value that is a boundary for the shaded region, convert that value to the equivalent z score.
- 3



Procedure for Finding Areas with a Nonstandard Normal Distribution

- 1 Sketch a normal curve, label the mean and any specific x values, and then shade the region representing the desired probability.
- 2 For each relevant x value that is a boundary for the shaded region, convert that value to the equivalent z score.
- 3 Use technology to find the area of the shaded region.



Example 9

The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Example 9

The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

Example 9

The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

We start by finding the z-value of the shower head.

$$z = \frac{x - \mu}{\sigma}$$

Example 9

The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

We start by finding the z-value of the shower head.

$$z = \frac{x - \mu}{\sigma} = \frac{72 - 68.6}{2.8}$$

Example 9

The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

We start by finding the z-value of the shower head.

$$z = \frac{x - \mu}{\sigma} = \frac{72 - 68.6}{2.8} = 1.21$$

Example 9


The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

We start by finding the z-value of the shower head.

$$z = \frac{x - \mu}{\sigma} = \frac{72 - 68.6}{2.8} = 1.21$$

Next, we sketch a picture and shade the area we wish to find:



Example 9


The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

We start by finding the z-value of the shower head.

$$z = \frac{x - \mu}{\sigma} = \frac{72 - 68.6}{2.8} = 1.21$$

Next, we sketch a picture and shade the area we wish to find:



We can then use technology to compute:

$$P(z \geq 1.21) \approx 0.1123 \quad (\text{rounded})$$

Example 9


The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men who are taller than a shower head installed at a height of 72 in.

We start by finding the z-value of the shower head.

$$z = \frac{x - \mu}{\sigma} = \frac{72 - 68.6}{2.8} = 1.21$$

Next, we sketch a picture and shade the area we wish to find:



We can then use technology to compute:

$$P(z \geq 1.21) \approx 0.1123 \quad (\text{rounded})$$

So, about 11.23% of men are taller than the shower head.

Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{X - \mu}{\sigma}$$

Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{x - \mu}{\sigma} = \frac{1030 - 1100}{200}$$

Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{x - \mu}{\sigma} = \frac{1030 - 1100}{200} = -0.35$$

Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{x - \mu}{\sigma} = \frac{1030 - 1100}{200} = -0.35$$

Recall that the percentile of a data value is the percentage of data less than the data value.


Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{x - \mu}{\sigma} = \frac{1030 - 1100}{200} = -0.35$$

Recall that the percentile of a data value is the percentage of data less than the data value.



This means $P(z \leq -0.35)$ is the percentile of Edwards SAT score.


Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{x - \mu}{\sigma} = \frac{1030 - 1100}{200} = -0.35$$

Recall that the percentile of a data value is the percentage of data less than the data value.



This means $P(z \leq -0.35)$ is the percentile of Edwards SAT score.

We can then use technology to compute:

$$P(z \leq -0.35) \approx 0.3632 \quad (\text{rounded})$$


Example 10

Cumulative SAT scores are approximately normal with a mean of 1100 and standard distribution of 200. Edward earned a 1030 on his SAT.

The z-value of his score is:

$$z = \frac{x - \mu}{\sigma} = \frac{1030 - 1100}{200} = -0.35$$

Recall that the percentile of a data value is the percentage of data less than the data value.



This means $P(z \leq -0.35)$ is the percentile of Edwards SAT score.

We can then use technology to compute:

$$P(z \leq -0.35) \approx 0.3632 \quad (\text{rounded})$$

So, Edward is in the 36th percentile.

Example 11

The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men meet that requirement.

Example 11

The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 68.6}{2.8} = -1.64$$

Example 11

The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 68.6}{2.8} = -1.64 \text{ and } z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 68.6}{2.8} = 3.00$$

Example 11


The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 68.6}{2.8} = -1.64 \text{ and } z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 68.6}{2.8} = 3.00$$

Next, we sketch a picture and shade the area we wish to find:



Example 11


The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 68.6}{2.8} = -1.64 \text{ and } z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 68.6}{2.8} = 3.00$$

Next, we sketch a picture and shade the area we wish to find:



We can then use technology to compute:

$$P(-1.64 \leq z \leq 3.00) \approx 0.9484 \text{ (rounded)}$$

Example 11


The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of men are normally distributed with a mean of 68.6 in. and a standard deviation of 2.8 in.

Let's find the percentage of men meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 68.6}{2.8} = -1.64 \text{ and } z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 68.6}{2.8} = 3.00$$

Next, we sketch a picture and shade the area we wish to find:



We can then use technology to compute:

$$P(-1.64 \leq z \leq 3.00) \approx 0.9484 \text{ (rounded)}$$

So, we see that about 94.84% of men meet the requirements.

Example 12

The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of women are normally distributed with a mean of 63.7 in. and a standard deviation of 2.9 in.

Let's find the percentage of women meet that requirement.

Example 12

The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of women are normally distributed with a mean of 63.7 in. and a standard deviation of 2.9 in.

Let's find the percentage of women meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 63.7}{2.9} = 0.10$$

Example 12

The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of women are normally distributed with a mean of 63.7 in. and a standard deviation of 2.9 in.

Let's find the percentage of women meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 63.7}{2.9} = 0.10 \quad \text{and} \quad z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 63.7}{2.9} = 4.59$$

Example 12


The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of women are normally distributed with a mean of 63.7 in. and a standard deviation of 2.9 in.

Let's find the percentage of women meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 63.7}{2.9} = 0.10 \quad \text{and} \quad z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 63.7}{2.9} = 4.59$$

Next, we sketch a picture and shade the area we wish to find:



Example 12


The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of women are normally distributed with a mean of 63.7 in. and a standard deviation of 2.9 in.

Let's find the percentage of women meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 63.7}{2.9} = 0.10 \quad \text{and} \quad z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 63.7}{2.9} = 4.59$$

Next, we sketch a picture and shade the area we wish to find:



We can then use technology to compute:

$$P(0.10 \leq z \leq 4.59) \approx 0.4601 \quad (\text{rounded})$$

Example 12


The U.S. Air Force requires that pilots have heights between 64 and 77 in. The heights of women are normally distributed with a mean of 63.7 in. and a standard deviation of 2.9 in.

Let's find the percentage of women meet that requirement.

We start by finding the z-values of the height requirements.

$$z_1 = \frac{x - \mu}{\sigma} = \frac{64 - 63.7}{2.9} = 0.10 \quad \text{and} \quad z_2 = \frac{x - \mu}{\sigma} = \frac{77 - 63.7}{2.9} = 4.59$$

Next, we sketch a picture and shade the area we wish to find:



We can then use technology to compute:

$$P(0.10 \leq z \leq 4.59) \approx 0.4601 \quad (\text{rounded})$$

So, we see that only about 46% of women meet the requirements.

When Finding Values from Known Areas

- Draw a sketch of the graph.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.
- A z score must be negative whenever it is located in the left half of the normal distribution.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.
- A z score must be negative whenever it is located in the left half of the normal distribution.
- Areas are always between 0 and 1, and are never negative.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.
- A z score must be negative whenever it is located in the left half of the normal distribution.
- Areas are always between 0 and 1, and are never negative.

Procedure

- 1 Sketch the normal distribution curve, write the given probability or percentage in the appropriate region of the graph, and identify the x values being sought.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.
- A z score must be negative whenever it is located in the left half of the normal distribution.
- Areas are always between 0 and 1, and are never negative.

Procedure

- 1 Sketch the normal distribution curve, write the given probability or percentage in the appropriate region of the graph, and identify the x values being sought.
- 2 Either use technology or a table to identify the z scores corresponding to that area.

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.
- A z score must be negative whenever it is located in the left half of the normal distribution.
- Areas are always between 0 and 1, and are never negative.

Procedure

- 1 Sketch the normal distribution curve, write the given probability or percentage in the appropriate region of the graph, and identify the x values being sought.
- 2 Either use technology or a table to identify the z scores corresponding to that area.
- 3 Convert to x values: $x = \mu + Z \cdot \sigma$

When Finding Values from Known Areas

- Draw a sketch of the graph.
- Don't confuse z scores and areas.
- Choose the correct side of the graph.
- A z score must be negative whenever it is located in the left half of the normal distribution.
- Areas are always between 0 and 1, and are never negative.

Procedure

- 1 Sketch the normal distribution curve, write the given probability or percentage in the appropriate region of the graph, and identify the x values being sought.
- 2 Either use technology or a table to identify the z scores corresponding to that area.
- 3 Convert to x values: $x = \mu + Z \cdot \sigma$
- 4 Use your sketch to verify that the solution makes sense.

Example 13


When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement.

Example 13

When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?


Example 13

When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?



Example 13


When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?



Using either technology or a table, we find that $z = 1.645$.

Example 13

When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?




Using either technology or a table, we find that $z = 1.645$.

We then need to convert to the x value.

$$X = \mu + Z \cdot \sigma$$

Example 13

When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?




Using either technology or a table, we find that $z = 1.645$. We then need to convert to the x value.

$$x = \mu + z \cdot \sigma = 63.7 + 1.645 \cdot 2.9$$

Example 13

When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?




Using either technology or a table, we find that $z = 1.645$. We then need to convert to the x value.

$$x = \mu + z \cdot \sigma = 63.7 + 1.645 \cdot 2.9 = 68.4705$$

Example 13

When designing equipment, one common criterion is to use a design that accommodates 95% of the population. In Example 12 we saw that only 46% of women satisfy the U.S. Air Force pilot height requirement. What would be the maximum acceptable height of a woman if the requirements were changed to allow the shortest 95% of women?



Using either technology or a table, we find that $z = 1.645$. We then need to convert to the x value.

$$x = \mu + z \cdot \sigma = 63.7 + 1.645 \cdot 2.9 = 68.4705$$

A requirement of a height less than or equal to 68.5 in. would allow 95% of women to be eligible.