

Regression

Colby Community College

Recall

Recall that a linear equation (a straight line) is one that can be written as

$$y = mx + b$$

where m is the slope (“rise over run”) and b is the y-intercept.

Recall

Recall that a linear equation (a straight line) is one that can be written as

$$y = mx + b$$

where m is the slope (“rise over run”) and b is the y-intercept.

Definition

Given a collection of paired sample data, the **regression line** (or **line of best fit**) is the straight line that “best” fits the scatter plot of the data. (We will discuss that “best” means later.)

Definition

The **regression equation** is

$$\hat{y} = b_0 + b_1x$$

algebraically describes the regression line. The regression equation expresses a relationship between x and \hat{y} .

Definition

The **regression equation** is

$$\hat{y} = b_0 + b_1x$$

algebraically describes the regression line. The regression equation expresses a relationship between x and \hat{y} .

Definition

We call x the **explanatory variable**, **predictor variable**, or **independent variable**.

Definition

The **regression equation** is

$$\hat{y} = b_0 + b_1x$$

algebraically describes the regression line. The regression equation expresses a relationship between x and \hat{y} .

Definition

We call x the **explanatory variable**, **predictor variable**, or **independent variable**.

Definition

We call y the **response variable**, or **dependent variable**.

Note

We don't use $y = mx + b$ because the format $y = b_0 + b_1x$ can easily be expanded to include more variables:

$$y = b_0 + b_1x_1 + b_2x_2 + b_3x_3 + \cdots$$

This is used when performing a multiple regression.

Requirements

The requirements for performing a regression are:

- 1 The sample of paired data is a random sample of quantitative data.

Requirements

The requirements for performing a regression are:

- ① The sample of paired data is a random sample of quantitative data.
- ② Visual examination of the scatterplot shows that the points approximate a straight-line pattern.

Requirements

The requirements for performing a regression are:

- ① The sample of paired data is a random sample of quantitative data.
- ② Visual examination of the scatterplot shows that the points approximate a straight-line pattern.
- ③ Outliers can have a strong effect on the regression equation, so remove any outliers if they are known errors.

Slope

The slope of the regression line is

$$b_1 = r \cdot \frac{s_y}{s_x}$$

where r is the linear correlation coefficient, s_y is the standard deviation of the y values, and s_x is the standard deviation of the x values.

Slope

The slope of the regression line is

$$b_1 = r \cdot \frac{s_y}{s_x}$$

where r is the linear correlation coefficient, s_y is the standard deviation of the y values, and s_x is the standard deviation of the x values.

y-intercept

The y-intercept of the regression line is

$$b_0 = \bar{y} - b_1 \bar{x}$$

where \bar{y} is the sample mean of y values and \bar{x} is the sample mean of x values.

Slope

The slope of the regression line is

$$b_1 = r \cdot \frac{s_y}{s_x}$$

where r is the linear correlation coefficient, s_y is the standard deviation of the y values, and s_x is the standard deviation of the x values.

y-intercept

The y-intercept of the regression line is

$$b_0 = \bar{y} - b_1 \bar{x}$$

where \bar{y} is the sample mean of y values and \bar{x} is the sample mean of x values.

Note

Technology will calculate both of these values for you.