

Random Variables

Colby Community College

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

$$\$0 \cdot 25 + \$137 \cdot 55 + (\$137 + \$33) \cdot 25$$

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

$$\$0 \cdot 25 + \$137 \cdot 55 + (\$137 + \$33) \cdot 25 = \$7,535 + \$4,250$$

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

$\$0 \cdot 25 + \$137 \cdot 55 + (\$137 + \$33) \cdot 25 = \$7,535 + \$4,250 = \$11,785$

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

$\$0 \cdot 25 + \$137 \cdot 55 + (\$137 + \$33) \cdot 25 = \$7,535 + \$4,250 = \$11,785$

What is the average revenue per student?

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

$\$0 \cdot 25 + \$137 \cdot 55 + (\$137 + \$33) \cdot 25 = \$7,535 + \$4,250 = \$11,785$

What is the average revenue per student?

$\$11,785 \div 100$ students

Example 1

Two books are assigned for a course:

- A textbook, which costs \$137.
- The accompanying study guide, which costs \$33.

The university bookstore determined:

- 20% of enrolled students do not buy either book.
- 55% of enrolled students only buy the textbook.
- 25% of the enrolled students buy both books.

How many books should be expected to sell if 100 students enrolled?

We expect about 25 students will buy none, 55 will buy just the textbook, and 25 will buy both. A total of $1 \cdot 55 + 2 \cdot 25 = 105$ books.

How much revenue should be expected to be made?

$\$0 \cdot 25 + \$137 \cdot 55 + (\$137 + \$33) \cdot 25 = \$7,535 + \$4,250 = \$11,785$

What is the average revenue per student?

$\$11,785 \div 100 \text{ students} = \$117.85 \text{ per student.}$

Definition

A **random variable** is a random process or variable that has a numerical value, determined by chance, for each outcome.

Definition

A **random variable** is a random process or variable that has a numerical value, determined by chance, for each outcome.

Note

Often capital letters such as X , Y , or Z are used for random variables. Corresponding lower case letters are used for the possible outcomes.

Definition

A **random variable** is a random process or variable that has a numerical value, determined by chance, for each outcome.

Note

Often capital letters such as X , Y , or Z are used for random variables. Corresponding lower case letters are used for the possible outcomes.

Example 2

Consider tossing a coin: We could get either a heads or a tails.

Definition

A **random variable** is a random process or variable that has a numerical value, determined by chance, for each outcome.

Note

Often capital letters such as X , Y , or Z are used for random variables. Corresponding lower case letters are used for the possible outcomes.

Example 2

Consider tossing a coin: We could get either a heads or a tails. If we let X be the number of tails we get in a single flip, then the two possible outcomes are: $x_1 = 0$ and $x_2 = 1$.

Definition

A **random variable** is a random process or variable that has a numerical value, determined by chance, for each outcome.

Note

Often capital letters such as X , Y , or Z are used for random variables. Corresponding lower case letters are used for the possible outcomes.

Example 2

Consider tossing a coin: We could get either a heads or a tails. If we let X be the number of tails we get in a single flip, then the two possible outcomes are: $x_1 = 0$ and $x_2 = 1$.

The probability distribution is:

i	1	2	Total
x_i	0	1	–
$P(X = x_i)$	0.50	0.50	1.00

Example 3

If we let X be the amount a student spends in Example 1, then the probability distribution is:

i	1	2	3	Total
x_i	\$0	\$137	\$170	—
$P(X = x_i)$	0.20	0.55	0.25	1.00

Example 3

If we let X be the amount a student spends in Example 1, then the probability distribution is:

i	1	2	3	Total
x_i	\$0	\$137	\$170	—
$P(X = x_i)$	0.20	0.55	0.25	1.00

Definition

A **discrete random variable** has a collection of values that is finite or countable.

Example 3

If we let X be the amount a student spends in Example 1, then the probability distribution is:

i	1	2	3	Total
x_i	\$0	\$137	\$170	—
$P(X = x_i)$	0.20	0.55	0.25	1.00

Definition

A **discrete random variable** has a collection of values that is finite or countable.

Definition

A **continuous random variable** has infinitely many values, and the collection of values is not countable.

Example 3

If we let X be the amount a student spends in Example 1, then the probability distribution is:

i	1	2	3	Total
x_i	\$0	\$137	\$170	—
$P(X = x_i)$	0.20	0.55	0.25	1.00

Definition

A **discrete random variable** has a collection of values that is finite or countable.

Definition

A **continuous random variable** has infinitely many values, and the collection of values is not countable.

Note

The random variable in Example 3 is a discrete random variable.

Example 4

Let's consider tossing two coins, with the following random variable:

X = number of heads when two coins are tossed

Example 4

Let's consider tossing two coins, with the following random variable:

X = number of heads when two coins are tossed

Is X is a discrete or continuous random variable?

Example 4

Let's consider tossing two coins, with the following random variable:

X = number of heads when two coins are tossed

Is X is a discrete or continuous random variable?

X is discrete since the number of heads can be 0, 1, or 2.

Example 4

Let's consider tossing two coins, with the following random variable:

X = number of heads when two coins are tossed

Is X a discrete or continuous random variable?

X is discrete since the number of heads can be 0, 1, or 2.

The probability distribution is:

i	1	2	3	Total
x_i	0	1	2	—
$P(X = x_i)$	0.25	0.50	0.25	1.00

Example 4

Let's consider tossing two coins, with the following random variable:

X = number of heads when two coins are tossed

Is X a discrete or continuous random variable?

X is discrete since the number of heads can be 0, 1, or 2.

The probability distribution is:

i	1	2	3	Total
x_i	0	1	2	–
$P(X = x_i)$	0.25	0.50	0.25	1.00

Example 5

Hiring managers were asked to identify the biggest mistakes that job applicants make during an interview.

Consider the following table:

i	x_i	$P(X = x_i)$
1	Inappropriate attire	0.50
2	Being late	0.44
3	Lack of Eye Contact	0.33
4	Checking phone or texting	0.30
Total		1.57

Is X and random variable?

Example 5

Hiring managers were asked to identify the biggest mistakes that job applicants make during an interview.

Consider the following table:

i	x_i	$P(X = x_i)$
1	Inappropriate attire	0.50
2	Being late	0.44
3	Lack of Eye Contact	0.33
4	Checking phone or texting	0.30
Total		1.57

Is X a random variable?

- 1 The outcomes are not numerical, they are categorical.

Example 5

Hiring managers were asked to identify the biggest mistakes that job applicants make during an interview.

Consider the following table:

i	x_i	$P(X = x_i)$
1	Inappropriate attire	0.50
2	Being late	0.44
3	Lack of Eye Contact	0.33
4	Checking phone or texting	0.30
Total		1.57

Is X a random variable?

- 1 The outcomes are not numerical, they are categorical.
- 2 $\sum P(X = x_i) = 1.57 \neq 1$

Example 5

Hiring managers were asked to identify the biggest mistakes that job applicants make during an interview.

Consider the following table:

i	x_i	$P(X = x_i)$
1	Inappropriate attire	0.50
2	Being late	0.44
3	Lack of Eye Contact	0.33
4	Checking phone or texting	0.30
Total		1.57

Is X a random variable?

- 1 The outcomes are not numerical, they are categorical.
- 2 $\sum P(X = x_i) = 1.57 \neq 1$

So, we see that X is not a random variable.

Definition


The average outcome of X is called the **expected value** of X .

Definition

The average outcome of X is called the **expected value** of X .

Example 6

In Example 1 the average revenue, \$117.85 per student, is the expected value for the bookstore's revenue.



Expected Value Of A Discrete Random Variable

If X takes outcomes x_1, \dots, x_k with probabilities $P(X = x_1), \dots, P(X = x_k)$, the expected value of X is:

$$\begin{aligned} E(X) &= x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + \dots + x_k \cdot P(X = x_k) \\ &= \sum_{i=1}^k x_i \cdot P(X = x_i) \end{aligned}$$

Expected Value Of A Discrete Random Variable

If X takes outcomes x_1, \dots, x_k with probabilities $P(X = x_1), \dots, P(X = x_k)$, the expected value of X is:

$$\begin{aligned} E(X) &= x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + \dots + x_k \cdot P(X = x_k) \\ &= \sum_{i=1}^k x_i \cdot P(X = x_i) \end{aligned}$$

Note

The Greek letter μ is sometimes used in place of $E(X)$.

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces.

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

If X is the net winnings, then the probability distribution is:

i	1	2	Total
x_i	\$35	-\$1	—
$P(X = x_i)$	$\frac{1}{38}$	$\frac{37}{38}$	1

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

If X is the net winnings, then the probability distribution is:

i	1	2	Total
x_i	\$35	-\$1	—
$P(X = x_i)$	$\frac{1}{38}$	$\frac{37}{38}$	1

The expected value of X is:

$$E(X) = x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + x_3 \cdot P(X = x_3)$$

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

If X is the net winnings, then the probability distribution is:

i	1	2	Total
x_i	\$35	-\$1	—
$P(X = x_i)$	$\frac{1}{38}$	$\frac{37}{38}$	1

The expected value of X is:

$$\begin{aligned} E(X) &= x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + x_3 \cdot P(X = x_3) \\ &= \$35 \cdot \frac{1}{38} + -\$1 \cdot \frac{37}{38} \end{aligned}$$

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

If X is the net winnings, then the probability distribution is:

i	1	2	Total
x_i	\$35	-\$1	—
$P(X = x_i)$	$\frac{1}{38}$	$\frac{37}{38}$	1

The expected value of X is:

$$\begin{aligned} E(X) &= x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + x_3 \cdot P(X = x_3) \\ &= \$35 \cdot \frac{1}{38} + -\$1 \cdot \frac{37}{38} = \$0.9211 - \$0.9737 \end{aligned}$$

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

If X is the net winnings, then the probability distribution is:

i	1	2	Total
x_i	\$35	-\$1	—
$P(X = x_i)$	$\frac{1}{38}$	$\frac{37}{38}$	1

The expected value of X is:

$$\begin{aligned} E(X) &= x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + x_3 \cdot P(X = x_3) \\ &= \$35 \cdot \frac{1}{38} + -\$1 \cdot \frac{37}{38} = \$0.9211 - \$0.9737 \approx -\$0.053 \end{aligned}$$

Example 7

In the casino game Roulette, a wheel with 38 numbered spaces is spun. There are 18 red spaces, 18 black spaces, and 2 green spaces. In one possible bet, the players bet \$1 on a single number. If that number is spun on the wheel, then they receive \$36. Otherwise, they lose their \$1.

If X is the net winnings, then the probability distribution is:

i	1	2	Total
x_i	\$35	-\$1	—
$P(X = x_i)$	$\frac{1}{38}$	$\frac{37}{38}$	1

The expected value of X is:

$$\begin{aligned} E(X) &= x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + x_3 \cdot P(X = x_3) \\ &= \$35 \cdot \frac{1}{38} + -\$1 \cdot \frac{37}{38} = \$0.9211 - \$0.9737 \approx -\$0.053 \end{aligned}$$

On *average*, the player will lose 5.3 cents per bet.

Note

Pick back up here. The following slides need to be edited

Example 8

Suppose an individual has a 0.242% risk of dying during the next year. An insurance company charges \$275 for a life-insurance policy that pays \$100,000 death benefit. What is the expected value for the person buying the insurance?

Example 8

Suppose an individual has a 0.242% risk of dying during the next year. An insurance company charges \$275 for a life-insurance policy that pays \$100,000 death benefit. What is the expected value for the person buying the insurance?

The probabilities and values for the two outcomes are:

Value	Probability
$\$100,000 - \$275 = \$99,725$	0.00242
$-\$275$	$1 - 0.00242 = 0.99758$

Example 8

Suppose an individual has a 0.242% risk of dying during the next year. An insurance company charges \$275 for a life-insurance policy that pays \$100,000 death benefit. What is the expected value for the person buying the insurance?

The probabilities and values for the two outcomes are:

Value	Probability
$\$100,000 - \$275 = \$99,725$	0.00242
$-\$275$	$1 - 0.00242 = 0.99758$

The expected value is $\$99,725 \cdot 0.00242 - \$275 \cdot 0.99758 = -\$33$.

Example 8

Suppose an individual has a 0.242% risk of dying during the next year. An insurance company charges \$275 for a life-insurance policy that pays \$100,000 death benefit. What is the expected value for the person buying the insurance?

The probabilities and values for the two outcomes are:

Value	Probability
$\$100,000 - \$275 = \$99,725$	0.00242
$-\$275$	$1 - 0.00242 = 0.99758$

The expected value is $\$99,725 \cdot 0.00242 - \$275 \cdot 0.99758 = -\$33$.

Note

It makes sense that a insurance policy would have a negative expected value, otherwise the insurance company couldn't stay in business.

The benefit for the consumer is the security that the policy provides.

Example 9

A company estimates that 0.7% of their products will fail after the original warranty period but within 2 years of the purchase, with a replacement cost of \$350. If they offer a 2 year extended warranty for \$48, what is the company's expected value?

Example 9

A company estimates that 0.7% of their products will fail after the original warranty period but within 2 years of the purchase, with a replacement cost of \$350. If they offer a 2 year extended warranty for \$48, what is the company's expected value?

The probabilities and values for the two outcomes are:

Value	Probability
$-\$350 + \$48 = -\$302$	0.007
\$48	$1 - 0.007 = 0.993$

Example 9

A company estimates that 0.7% of their products will fail after the original warranty period but within 2 years of the purchase, with a replacement cost of \$350. If they offer a 2 year extended warranty for \$48, what is the company's expected value?

The probabilities and values for the two outcomes are:

Value	Probability
$-\$350 + \$48 = -\$302$	0.007
\$48	$1 - 0.007 = 0.993$

The expected value is $-\$302 \cdot 0.007 + \$48 \cdot 0.993 = \$45.55$.

The company makes, on average, \$45.55 on each extended warranty.

Example 9

A company estimates that 0.7% of their products will fail after the original warranty period but within 2 years of the purchase, with a replacement cost of \$350. If they offer a 2 year extended warranty for \$48, what is the company's expected value?

The probabilities and values for the two outcomes are:

Value	Probability
$-\$350 + \$48 = -\$302$	0.007
\$48	$1 - 0.007 = 0.993$

The expected value is $-\$302 \cdot 0.007 + \$48 \cdot 0.993 = \$45.55$.

The company makes, on average, \$45.55 on each extended warranty.

Note

The expected value for the consumer may be different. The consumer is likely to pay the more to repair or replace the item out of warranty. (The company pays manufacturing cost, consumer has to pay retail