

Linear Transformations

Department of Mathematics

Salt Lake Community College

(Slides by Adam Wilson)

Linear Transformation

A **Linear Transformation** T on a vector space \mathbb{V} to a vector space \mathbb{W} is a function $T : \mathbb{V} \rightarrow \mathbb{W}$ that preserves *scalar multiplication* and *vector addition*. That is, for all $\vec{u}, \vec{v} \in \mathbb{V}$ and $c \in \mathbb{R}$:

- $T(c\vec{u}) = cT(\vec{u})$
- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$

Linear Transformation

A **Linear Transformation** T on a vector space \mathbb{V} to a vector space \mathbb{W} is a function $T : \mathbb{V} \rightarrow \mathbb{W}$ that preserves *scalar multiplication* and *vector addition*. That is, for all $\vec{u}, \vec{v} \in \mathbb{V}$ and $c \in \mathbb{R}$:

- $T(c\vec{u}) = cT(\vec{u})$
- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$

These conditions are commonly checked together.

$$T(c\vec{u} + d\vec{v}) = cT(\vec{u}) + dT(\vec{v})$$

Linear Transformation

A **Linear Transformation** T on a vector space \mathbb{V} to a vector space \mathbb{W} is a function $T : \mathbb{V} \rightarrow \mathbb{W}$ that preserves *scalar multiplication* and *vector addition*. That is, for all $\vec{u}, \vec{v} \in \mathbb{V}$ and $c \in \mathbb{R}$:

- $T(c\vec{u}) = cT(\vec{u})$
- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$

These conditions are commonly checked together.

$$T(c\vec{u} + d\vec{v}) = cT(\vec{u}) + dT(\vec{v})$$

The vector space \mathbb{V} is called the **domain** for T

The vector space \mathbb{W} is called the **codomain** (or **range** or **target**) for T .

Linear Transformation

A **Linear Transformation** T on a vector space \mathbb{V} to a vector space \mathbb{W} is a function $T : \mathbb{V} \rightarrow \mathbb{W}$ that preserves *scalar multiplication* and *vector addition*. That is, for all $\vec{u}, \vec{v} \in \mathbb{V}$ and $c \in \mathbb{R}$:

- $T(c\vec{u}) = cT(\vec{u})$
- $T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$

These conditions are commonly checked together.

$$T(c\vec{u} + d\vec{v}) = cT(\vec{u}) + dT(\vec{v})$$

The vector space \mathbb{V} is called the **domain** for T

The vector space \mathbb{W} is called the **codomain** (or **range** or **target**) for T .

Image of a Linear Transformation

The **image** of a linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is the set of vectors in \mathbb{W} to which T maps the vectors in \mathbb{V} :

$$\text{Im}(T) = \{\vec{w} \in \mathbb{W} \mid \vec{w} = T(\vec{v}) \text{ for some } \vec{v} \in \mathbb{V}\}$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

$$T(\vec{0}) = \vec{0}$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

$$T(\vec{0}) = \vec{0}$$

Example 2

$$T(\vec{0} + \vec{v}) = T(\vec{0}) + T(\vec{v})$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

$$T(\vec{0}) = \vec{0}$$

Example 2

$$T(\vec{0} + \vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) = T(\vec{0}) + T(\vec{v})$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

$$T(\vec{0}) = \vec{0}$$

Example 2

$$T(\vec{0} + \vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) - T(\vec{v}) = T(\vec{0})$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

$$T(\vec{0}) = \vec{0}$$

Example 2

$$T(\vec{0} + \vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) - T(\vec{v}) = T(\vec{0})$$

$$\vec{0} = T(\vec{v})$$

A linear transformation always associates the zero vector of the domain with the zero vector of the codomain.

Example 1

$$T(0 \cdot \vec{v}) = 0 \cdot T(\vec{v})$$

$$T(\vec{0}) = \vec{0}$$

Example 2

$$T(\vec{0} + \vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) = T(\vec{0}) + T(\vec{v})$$

$$T(\vec{v}) - T(\vec{v}) = T(\vec{0})$$

$$\vec{0} = T(\vec{v})$$

Note

Linear transformations may map nonzero vectors from the domain into the zero vector of the codomain.

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$.
Let's check that this is a linear transformation:

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$.
Let's check that this is a linear transformation:

$$T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle)$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) = T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle)$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle) \\ &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \end{aligned}$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle) \\ &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, cy_1 + dy_2, 0 \rangle \end{aligned}$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle) \\ &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, cy_1 + dy_2, 0 \rangle \\ &= \langle cx_1, cy_1, 0 \rangle + \langle dx_2, dy_2, 0 \rangle \end{aligned}$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle) \\ &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, cy_1 + dy_2, 0 \rangle \\ &= \langle cx_1, cy_1, 0 \rangle + \langle dx_2, dy_2, 0 \rangle \\ &= c \langle x_1, y_1, 0 \rangle + d \langle x_2, y_2, 0 \rangle \end{aligned}$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle) \\ &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, cy_1 + dy_2, 0 \rangle \\ &= \langle cx_1, cy_1, 0 \rangle + \langle dx_2, dy_2, 0 \rangle \\ &= c \langle x_1, y_1, 0 \rangle + d \langle x_2, y_2, 0 \rangle \\ &= cT(\langle x_1, y_1, z_1 \rangle) + dT(\langle x_2, y_2, z_2 \rangle) \end{aligned}$$

Example 3

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, y, 0 \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1, cy_1, cz_1 \rangle + \langle dx_2, dy_2, dz_2 \rangle) \\ &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, cy_1 + dy_2, 0 \rangle \\ &= \langle cx_1, cy_1, 0 \rangle + \langle dx_2, dy_2, 0 \rangle \\ &= c \langle x_1, y_1, 0 \rangle + d \langle x_2, y_2, 0 \rangle \\ &= cT(\langle x_1, y_1, z_1 \rangle) + dT(\langle x_2, y_2, z_2 \rangle) \end{aligned}$$

Therefore, T is a linear transformation.

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

$$T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle)$$

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

$$T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) = T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle)$$

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, 0, 3cx_1 + 3dx_2 \rangle \end{aligned}$$

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by
 $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, 0, 3cx_1 + 3dx_2 \rangle \\ &= c \langle x_1, 0, 3x_1 \rangle + d \langle x_2, 0, 3x_2 \rangle \end{aligned}$$

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by
 $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, 0, 3cx_1 + 3dx_2 \rangle \\ &= c \langle x_1, 0, 3x_1 \rangle + d \langle x_2, 0, 3x_2 \rangle \\ &= cT(\langle x_1, y_1, z_1 \rangle) + dT(\langle x_2, y_2, z_2 \rangle) \end{aligned}$$

Example 4

Consider the mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by
 $T(\langle x, y, z \rangle) = \langle x, 0, 3x \rangle$. Let's check that this is a linear transformation:

$$\begin{aligned} T(c \langle x_1, y_1, z_1 \rangle + d \langle x_2, y_2, z_2 \rangle) &= T(\langle cx_1 + dx_2, cy_1 + dy_2, cz_1 + dz_2 \rangle) \\ &= \langle cx_1 + dx_2, 0, 3cx_1 + 3dx_2 \rangle \\ &= c \langle x_1, 0, 3x_1 \rangle + d \langle x_2, 0, 3x_2 \rangle \\ &= cT(\langle x_1, y_1, z_1 \rangle) + dT(\langle x_2, y_2, z_2 \rangle) \end{aligned}$$

Therefore, T is a linear transformation.

Example 5

Differentiation is a linear transformation. The **derivative operator** $D : \mathcal{C}^1[a, b] \rightarrow \mathcal{C}[a, b]$ is defined by

$$D(f) = f'$$

Example 5

Differentiation is a linear transformation. The **derivative operator** $D : \mathcal{C}^1[a, b] \rightarrow \mathcal{C}[a, b]$ is defined by

$$D(f) = f'$$

We know from calculus that D satisfy both properties:

$$D(cf) = cD(f)$$

$$D(f + g) = D(f) + D(g)$$

Example 5

Differentiation is a linear transformation. The **derivative operator** $D : \mathcal{C}^1[a, b] \rightarrow \mathcal{C}[a, b]$ is defined by

$$D(f) = f'$$

We know from calculus that D satisfy both properties:

$$D(cf) = cD(f)$$

$$D(f + g) = D(f) + D(g)$$

Example 6

Similarly, we can confirm that the **integration operator** $I : \mathcal{C}[a, b] \rightarrow \mathbb{R}$, defined by

$$I(f) = \int_a^b f(t)dt$$

is a linear transformation.

Geometry of Matrix Linear Transformations

If \mathbf{A} is an $m \times n$ matrix and \vec{x} is a column n -vector, then $\mathbf{A}\vec{x}$ can be considered a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, where $T(\vec{x}) = \mathbf{A}\vec{x}$.

In this transformation, the matrix \mathbf{A} allows vectors to be dynamic.

Geometry of Matrix Linear Transformations

If \mathbf{A} is an $m \times n$ matrix and \vec{x} is a column n -vector, then $\mathbf{A}\vec{x}$ can be considered a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, where $T(\vec{x}) = \mathbf{A}\vec{x}$.

In this transformation, the matrix \mathbf{A} allows vectors to be dynamic.

Example 7

The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

defines a **shear** of 1-unit in the x -direction.

Geometry of Matrix Linear Transformations

If \mathbf{A} is an $m \times n$ matrix and \vec{x} is a column n -vector, then $\mathbf{A}\vec{x}$ can be considered a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, where $T(\vec{x}) = \mathbf{A}\vec{x}$.


In this transformation, the matrix \mathbf{A} allows vectors to be dynamic.

Example 7

The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

defines a **shear** of 1-unit in the x -direction.



Geometry of Matrix Linear Transformations

If \mathbf{A} is an $m \times n$ matrix and \vec{x} is a column n -vector, then $\mathbf{A}\vec{x}$ can be considered a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, where $T(\vec{x}) = \mathbf{A}\vec{x}$.


In this transformation, the matrix \mathbf{A} allows vectors to be dynamic.

Example 7

The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

defines a **shear** of 1-unit in the x -direction.



Geometry of Matrix Linear Transformations

If \mathbf{A} is an $m \times n$ matrix and \vec{x} is a column n -vector, then $\mathbf{A}\vec{x}$ can be considered a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, where $T(\vec{x}) = \mathbf{A}\vec{x}$.


In this transformation, the matrix \mathbf{A} allows vectors to be dynamic.

Example 7

The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

defines a **shear** of 1-unit in the x -direction.



Example 8


The *Counterclockwise* rotation about the origin by angle θ is given by:

$$R_{\theta} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

Example 8

The *Counterclockwise* rotation about the origin by angle θ is given by:


$$R_{\theta} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$



Example 8

The *Counterclockwise* rotation about the origin by angle θ is given by:


$$R_{\theta} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$



Example 8

The *Counterclockwise* rotation about the origin by angle θ is given by:



$$R_{\theta} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$



Example 9

Reflection about the x -axis


$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$



Example 9

Reflection about the y-axis



$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$



Example 9

Rotation clockwise about the origin of $\frac{\pi}{4}$



$$\begin{bmatrix} \cos\left(\frac{\pi}{4}\right) & \sin\left(\frac{\pi}{4}\right) \\ -\sin\left(\frac{\pi}{4}\right) & \cos\left(\frac{\pi}{4}\right) \end{bmatrix}$$



Example 9

Rotation counterclockwise about the origin of $\frac{3\pi}{4}$



$$\begin{bmatrix} \cos\left(\frac{3\pi}{4}\right) & -\sin\left(\frac{3\pi}{4}\right) \\ \sin\left(\frac{3\pi}{4}\right) & \cos\left(\frac{3\pi}{4}\right) \end{bmatrix}$$



Example 9

Reflection about the line $y = x$



$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$



Example 9

Reflection about the line $y = 2x$



$$\begin{bmatrix} -\frac{3}{5} & \frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{bmatrix}$$



Example 9

Shear of 2 in the y-direction

$$\begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$



Example 10

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \vec{v}$$

maps

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \quad \text{to} \quad \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

Example 10

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \vec{v}$$

maps

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \quad \text{to} \quad \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

What is the image of T ?

Example 10

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \vec{v}$$

maps

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \quad \text{to} \quad \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

What is the image of T ?

A typical vector in the range is

$$\vec{u} = \mathbf{A}\vec{v} = v_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ 3 \end{bmatrix} + v_3 \begin{bmatrix} 2 \\ 5 \end{bmatrix}$$

Example 10

Consider the transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \vec{v}$$

maps

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \quad \text{to} \quad \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

What is the image of T ?

A typical vector in the range is

$$\vec{u} = \mathbf{A}\vec{v} = v_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ 3 \end{bmatrix} + v_3 \begin{bmatrix} 2 \\ 5 \end{bmatrix}$$

It can be easily verified that $[1, 2]$ and $[1, 3]$ are independent in \mathbb{R}^2 . Which means the image must contain their span, which is exactly \mathbb{R}^2 .

The Standard Matrix for a Linear Transform

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation. The **standard matrix** associated with T is defined by

$$\mathbf{A} = [T(\vec{e}_1) | T(\vec{e}_2) | \cdots | T(\vec{e}_n)]$$

where the columns $T(\vec{e}_j)$ are the images under T of the standard basis vectors $\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n$.

Proof

We can check that this matrix satisfies $T(\vec{v}) = \mathbf{A}\vec{v}$ by

$$T\left(\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}\right) =$$

Proof

We can check that this matrix satisfies $T(\vec{v}) = \mathbf{A}\vec{v}$ by

$$T\left(\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}\right) = T(v_1\vec{e}_1 + v_2\vec{e}_2 + \cdots + v_n\vec{e}_n)$$

Proof

We can check that this matrix satisfies $T(\vec{v}) = \mathbf{A}\vec{v}$ by

$$\begin{aligned} T\left(\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}\right) &= T(v_1\vec{e}_1 + v_2\vec{e}_2 + \cdots + v_n\vec{e}_n) \\ &= v_1T(\vec{e}_1) + v_2T(\vec{e}_2) + \cdots + v_nT(\vec{e}_n) \end{aligned}$$

Proof

We can check that this matrix satisfies $T(\vec{v}) = \mathbf{A}\vec{v}$ by

$$\begin{aligned} T\left(\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}\right) &= T(v_1\vec{e}_1 + v_2\vec{e}_2 + \cdots + v_n\vec{e}_n) \\ &= v_1 T(\vec{e}_1) + v_2 T(\vec{e}_2) + \cdots + v_n T(\vec{e}_n) \\ &= [T(\vec{e}_1) | T(\vec{e}_2) | \cdots | T(\vec{e}_n)] \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \end{aligned}$$

Proof

We can check that this matrix satisfies $T(\vec{v}) = \mathbf{A}\vec{v}$ by

$$\begin{aligned} T\left(\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}\right) &= T(v_1\vec{e}_1 + v_2\vec{e}_2 + \cdots + v_n\vec{e}_n) \\ &= v_1 T(\vec{e}_1) + v_2 T(\vec{e}_2) + \cdots + v_n T(\vec{e}_n) \\ &= [T(\vec{e}_1) | T(\vec{e}_2) | \cdots | T(\vec{e}_n)] \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \\ &= \mathbf{A} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \end{aligned}$$

Example 11

Find the standard matrix that will describe the transformation

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x - y \\ x + y \\ 2x \end{bmatrix}$$

Example 11

Find the standard matrix that will describe the transformation

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x - y \\ x + y \\ 2x \end{bmatrix}$$

We are looking for a matrix **A** that will satisfy

$$\mathbf{A} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x - y \\ x + y \\ 2x \end{bmatrix}$$

Example 11

Find the standard matrix that will describe the transformation

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x - y \\ x + y \\ 2x \end{bmatrix}$$

We are looking for a matrix **A** that will satisfy

$$\mathbf{A} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x - y \\ x + y \\ 2x \end{bmatrix}$$

Thus, for dimensions in the product to match, **A** must be a 3×2 matrix. Which means:

$$\mathbf{A} = \left[T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) \mid T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) \right] = \begin{bmatrix} 1 & -1 \\ 1 & 1 \\ 2 & 0 \end{bmatrix}$$

Example 12

Let $D_2 : \mathbb{P}_3 \rightarrow \mathbb{P}_1$ be the second-derivative operator. So, for a typical cubic polynomial:

$$D_2(ax^3 + bx^2 + cx + d) = 6ax + 2b$$

Example 12

Let $D_2 : \mathbb{P}_3 \rightarrow \mathbb{P}_1$ be the second-derivative operator. So, for a typical cubic polynomial:

$$D_2(ax^3 + bx^2 + cx + d) = 6ax + 2b$$

The standard matrix is made up of

$$\left[D_2 \left(\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) \middle| D_2 \left(\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \right) \middle| D_2 \left(\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right) \middle| D_2 \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right) \right] = \begin{bmatrix} 6 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \end{bmatrix}$$

Example 12

Let $D_2 : \mathbb{P}_3 \rightarrow \mathbb{P}_1$ be the second-derivative operator. So, for a typical cubic polynomial:

$$D_2(ax^3 + bx^2 + cx + d) = 6ax + 2b$$

The standard matrix is made up of



$$\left[D_2 \left(\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) \middle| D_2 \left(\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \right) \middle| D_2 \left(\begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right) \middle| D_2 \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right) \right] = \begin{bmatrix} 6 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \end{bmatrix}$$

Which gives us a matrix that satisfies:

$$\begin{bmatrix} 6 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 6a \\ 2b \end{bmatrix}$$

Example 13

Linear transforms are used extensively in computer graphics, where images or models are just collections of points and line segments. Let us look at a simple example, where we can think of each pixel as a point at its center:



Example 14

We can rotate this image 90° clockwise with the matrix

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

Example 14

We can rotate this image 90° clockwise with the matrix

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

When applying this linear transformation to our image we get

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 2 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 2 & 1 & 2 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

Example 14

We can rotate this image 90° clockwise with the matrix

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

When applying this linear transformation to our image we get

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 2 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 2 & 1 & 2 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

