

# Regression

Colby Community College

## Recall

Recall that a linear equation (a straight line) is one that can be written as

$$y = mx + b$$

where  $m$  is the slope (“rise over run”) and  $b$  is the y-intercept.

## Recall

Recall that a linear equation (a straight line) is one that can be written as

$$y = mx + b$$

where  $m$  is the slope (“rise over run”) and  $b$  is the y-intercept.

## Definition

Given a collection of paired sample data, the **regression line** (or **line of best fit**) is the straight line that “best” fits the scatter plot of the data. (We will discuss that “best” means later.)

## Definition

The **regression equation** is

$$\hat{y} = b_0 + b_1x$$

algebraically describes the regression line. The regression equation expresses a relationship between  $x$  and  $\hat{y}$ .

## Definition

The **regression equation** is

$$\hat{y} = b_0 + b_1x$$

algebraically describes the regression line. The regression equation expresses a relationship between  $x$  and  $\hat{y}$ .

## Definition

We call  $x$  the **explanatory variable**, **predictor variable**, or **independent variable**.

## Definition

The **regression equation** is

$$\hat{y} = b_0 + b_1x$$

algebraically describes the regression line. The regression equation expresses a relationship between  $x$  and  $\hat{y}$ .

## Definition

We call  $x$  the **explanatory variable**, **predictor variable**, or **independent variable**.

## Definition

We call  $y$  the **response variable**, or **dependent variable**.

## Note

We don't use  $y = mx + b$  because the format  $y = b_0 + b_1x$  can easily be expanded to include more variables:

$$y = b_0 + b_1x_1 + b_2x_2 + b_3x_3 + \cdots$$

This is used when performing a multiple regression.

## Requirements

The requirements for performing a regression are:

- 1 The sample of paired data is a random sample of quantitative data.



## Requirements

The requirements for performing a regression are:

- ① The sample of paired data is a random sample of quantitative data.
- ② Visual examination of the scatterplot shows that the points approximate a straight-line pattern.

## Requirements

The requirements for performing a regression are:

- ① The sample of paired data is a random sample of quantitative data.
- ② Visual examination of the scatterplot shows that the points approximate a straight-line pattern.
- ③ Outliers can have a strong effect on the regression equation, so remove any outliers if they are known errors.

## Slope

The slope of the regression line is

$$b_1 = r \cdot \frac{s_y}{s_x}$$

where  $r$  is the linear correlation coefficient,  $s_y$  is the standard deviation of the  $y$  values, and  $s_x$  is the standard deviation of the  $x$  values.

## Slope

The slope of the regression line is

$$b_1 = r \cdot \frac{s_y}{s_x}$$

where  $r$  is the linear correlation coefficient,  $s_y$  is the standard deviation of the  $y$  values, and  $s_x$  is the standard deviation of the  $x$  values.

## y-intercept

The y-intercept of the regression line is

$$b_0 = \bar{y} - b_1 \bar{x}$$

where  $\bar{y}$  is the sample mean of  $y$  values and  $\bar{x}$  is the sample mean of  $x$  values.

## Slope

The slope of the regression line is

$$b_1 = r \cdot \frac{s_y}{s_x}$$

where  $r$  is the linear correlation coefficient,  $s_y$  is the standard deviation of the  $y$  values, and  $s_x$  is the standard deviation of the  $x$  values.

## y-intercept

The y-intercept of the regression line is

$$b_0 = \bar{y} - b_1 \bar{x}$$

where  $\bar{y}$  is the sample mean of  $y$  values and  $\bar{x}$  is the sample mean of  $x$  values.

## Note

Technology will calculate both of these values for you.

## Making Predictions

When making predictions, keep the following in mind:

**Bad Model:** If the regression equation does not appear to be useful for making predictions, don't use the regression equation.

## Making Predictions

When making predictions, keep the following in mind:

**Bad Model:** If the regression equation does not appear to be useful for making predictions, don't use the regression equation.

- The best predicted value of a variable in this case is simply its sample mean.

## Making Predictions

When making predictions, keep the following in mind:

**Bad Model:** If the regression equation does not appear to be useful for making predictions, don't use the regression equation.

- The best predicted value of a variable in this case is simply its sample mean.

**Good Model:** Use the regression equation for predictions only if the graph of the regression line on the scatterplot confirms that the regression line fits the points reasonably well.



## Making Predictions

When making predictions, keep the following in mind:

**Bad Model:** If the regression equation does not appear to be useful for making predictions, don't use the regression equation.

- The best predicted value of a variable in this case is simply its sample mean.

**Good Model:** Use the regression equation for predictions only if the graph of the regression line on the scatterplot confirms that the regression line fits the points reasonably well.

**Correlation:** Use the regression equation for predictions only if the linear correlation coefficient  $r$  indicates that there is a linear correlation between the two variables.

## Making Predictions

When making predictions, keep the following in mind:

**Bad Model:** If the regression equation does not appear to be useful for making predictions, don't use the regression equation.

- The best predicted value of a variable in this case is simply its sample mean.

**Good Model:** Use the regression equation for predictions only if the graph of the regression line on the scatterplot confirms that the regression line fits the points reasonably well.

**Correlation:** Use the regression equation for predictions only if the linear correlation coefficient  $r$  indicates that there is a linear correlation between the two variables.

**Scope:** Use the regression line for predictions only if the data do not go much beyond the scope of available sample data.

## Making Predictions

When making predictions, keep the following in mind:

**Bad Model:** If the regression equation does not appear to be useful for making predictions, don't use the regression equation.

- The best predicted value of a variable in this case is simply its sample mean.

**Good Model:** Use the regression equation for predictions only if the graph of the regression line on the scatterplot confirms that the regression line fits the points reasonably well.

**Correlation:** Use the regression equation for predictions only if the linear correlation coefficient  $r$  indicates that there is a linear correlation between the two variables.

**Scope:** Use the regression line for predictions only if the data do not go much beyond the scope of available sample data.

- Predicting too far beyond the scope of the available sample data is called **extrapolation** and can easily result in bad predictions.


## Example 1

Using the Data Set 16, we can compare a countries chocolate consumption to the number of Nobel laureates.

## Example 1

Using the Data Set 16, we can compare a countries chocolate consumption to the number of Nobel laureates.


Technology give a regression line of  $\hat{y} = -3.37 + 2.49x$  and the scatterplot:



## Example 1

Using the Data Set 16, we can compare a countries chocolate consumption to the number of Nobel laureates.

Technology give a regression line of  $\hat{y} = -3.37 + 2.49x$  and the scatterplot:




This is a good model, so we can approximate the Nobel Laureate rate for a country that consumes 10kg per capita:

$$\hat{y} = -3.37 + 2.49(10)$$

## Example 1

Using the Data Set 16, we can compare a countries chocolate consumption to the number of Nobel laureates.

Technology give a regression line of  $\hat{y} = -3.37 + 2.49x$  and the scatterplot:




This is a good model, so we can approximate the Nobel Laureate rate for a country that consumes 10kg per capita:

$$\hat{y} = -3.37 + 2.49(10) = 21.5$$

## Example 1

Using the Data Set 16, we can compare a countries chocolate consumption to the number of Nobel laureates.

Technology give a regression line of  $\hat{y} = -3.37 + 2.49x$  and the scatterplot:



This is a good model, so we can approximate the Nobel Laureate rate for a country that consumes 10kg per capita:


$$\hat{y} = -3.37 + 2.49(10) = 21.5$$

So, we expect 21.5 Nobel Laureates per 10 million people.




## Example 2

The scatterplot shows sample data recording subject height and IQ score.



## Example 2


The scatterplot shows sample data recording subject height and IQ score.



The correlation is  $r = -0.02731$  and has  $P$ -value 0.909.

## Example 2

The scatterplot shows sample data recording subject height and IQ score.



The correlation is  $r = -0.02731$  and has  $P$ -value 0.909.

This means the regression line is a bad model and should not be used to make predictions.