

Systems of Linear Equations

Adam Wilson

Salt Lake Community College

Linear Systems of Equations

System of Linear Equations

A $m \times n$ **system of linear equations** is a set of m equations in n variables x_1, x_2, \dots, x_n of the form

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \dots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \dots & + & a_{2n}x_n & = & b_2 \\ a_{31}x_1 & + & a_{32}x_2 & + & \dots & + & a_{3n}x_n & = & b_3 \\ \vdots & & \vdots & & & & \vdots & & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \dots & + & a_{mn}x_n & = & b_m \end{array}$$

Linear System of Equations

Notation

There are two primary ways of writing a linear systems using matrices:

Linear System of Equations

Notation

There are two primary ways of writing a linear systems using matrices:

An Augmented Matrix

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ \vdots & \vdots & \ddots & \vdots & \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{array} \right]$$

Linear System of Equations

Notation

There are two primary ways of writing a linear systems using matrices:

An Augmented Matrix

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ \vdots & \vdots & \ddots & \vdots & \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{array} \right]$$

A Matrix Equation (We will look at these in section 3.3)

As the matrix equation $A\vec{x} = \vec{b}$, where:

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}}_{\vec{x}} = \underbrace{\begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}}_{\vec{b}}$$

Row Operations

(for Augmented Matrices)

Elementary Row Operations

- r_i denotes row i *before* the row operation is applied
- R_i denotes row i *after* the row operation is applied

Row Operations

(for Augmented Matrices)

Elementary Row Operations

- r_i denotes row i *before* the row operation is applied
- R_i denotes row i *after* the row operation is applied
- Swap row i and row j :

$$R_i \leftrightarrow R_j \quad (\text{or } R_i = r_j, R_j = r_i)$$

Row Operations

(for Augmented Matrices)

Elementary Row Operations

- r_i denotes row i *before* the row operation is applied
- R_i denotes row i *after* the row operation is applied
- Swap row i and row j :

$$R_i \leftrightarrow R_j \quad (\text{or } R_i = r_j, R_j = r_i)$$

- Multiply row i by a nonzero constant:

$$R_i = c \cdot r_i$$

Row Operations

(for Augmented Matrices)

Elementary Row Operations

- r_i denotes row i *before* the row operation is applied
- R_i denotes row i *after* the row operation is applied
- Swap row i and row j :

$$R_i \leftrightarrow R_j \quad (\text{or } R_i = r_j, R_j = r_i)$$

- Multiply row i by a nonzero constant:

$$R_i = c \cdot r_i$$

- Add row j to row i (leaving row j unchanged):

$$R_i = r_i + r_j$$

Solving Augmented Matrices

Row Echelon Form

Gaussian Elimination

Use row operations until in REF:

$$\left[\begin{array}{ccccc|c} 1 & c_{12} & c_{13} & \cdots & c_{1n} & d_1 \\ 0 & 1 & c_{23} & \cdots & c_{2n} & d_2 \\ 0 & 0 & 1 & \cdots & c_{3n} & d_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & d_m \end{array} \right]$$

Solving Augmented Matrices

Row Echelon Form

Gaussian Elimination

Use row operations until in REF:

$$\left[\begin{array}{cccc|c} 1 & c_{12} & c_{13} & \cdots & c_{1n} & d_1 \\ 0 & 1 & c_{23} & \cdots & c_{2n} & d_2 \\ 0 & 0 & 1 & \cdots & c_{3n} & d_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & d_m \end{array} \right]$$

Then back solve the system:

$$x_1 + c_{12}x_2 + c_{13}x_3 + \cdots + c_{1n}x_n = d_1$$

$$x_2 + c_{23}x_3 + \cdots + c_{2n}x_n = d_2$$

$$\vdots$$

$$x_n = d_m$$

Gaussian Elimination

Example

Example

Consider the system

$$\begin{array}{rrcrcl} x & + & y & + & z & = & 3 \\ 2x & - & 3y & - & z & = & -8 \\ -x & + & 2y & + & 2z & = & 3 \end{array}$$

Gaussian Elimination

Example

Example

Consider the system

$$\begin{array}{rrcrcl} x & + & y & + & z & = & 3 \\ 2x & - & 3y & - & z & = & -8 \\ -x & + & 2y & + & 2z & = & 3 \end{array}$$

We can write this as the augmented matrix:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 2 & -3 & -1 & -8 \\ -1 & 2 & 2 & 3 \end{array} \right]$$

We now want to use row operations to transform this augmented matrix into Row Echelon Form.

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 2 & -3 & -1 & -8 \\ -1 & 2 & 2 & 3 \end{array} \right]$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 2 & -3 & -1 & -8 \\ -1 & 2 & 2 & 3 \end{array} \right] \quad R_2 = r_2 + 2r_3$$

Gaussian Elimination

Example

Example

$$\begin{bmatrix} 1 & 1 & 1 & | & 3 \\ 2 & -3 & -1 & | & -8 \\ -1 & 2 & 2 & | & 3 \end{bmatrix} \quad R_2 = r_2 + 2r_3$$
$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 & | & 3 \\ 0 & 1 & 3 & | & -2 \\ -1 & 2 & 2 & | & 3 \end{bmatrix}$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ -1 & 2 & 2 & 3 \end{array} \right]$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ -1 & 2 & 2 & 3 \end{array} \right] R_3 = r_1 + r_3$$

Gaussian Elimination

Example

Example

$$\begin{bmatrix} 1 & 1 & 1 & | & 3 \\ 0 & 1 & 3 & | & -2 \\ -1 & 2 & 2 & | & 3 \end{bmatrix} \quad R_3 = r_1 + r_3$$
$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 & | & 3 \\ 0 & 1 & 3 & | & -2 \\ 0 & 3 & 3 & | & 6 \end{bmatrix}$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ 0 & 3 & 3 & 6 \end{array} \right]$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ 0 & 3 & 3 & 6 \end{array} \right] R_3 = r_3 - 3r_2$$

Gaussian Elimination

Example

Example

$$\begin{bmatrix} 1 & 1 & 1 & | & 3 \\ 0 & 1 & 3 & | & -2 \\ 0 & 3 & 3 & | & 6 \end{bmatrix} \quad R_3 = r_3 - 3r_2$$
$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 & | & 3 \\ 0 & 1 & 3 & | & -2 \\ 0 & 0 & -6 & | & 12 \end{bmatrix}$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ 0 & 0 & -6 & 12 \end{array} \right]$$

Gaussian Elimination

Example

Example

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ 0 & 0 & -6 & 12 \end{array} \right] R_3 = -\frac{1}{6}r_3$$

Gaussian Elimination

Example

Example

$$\Rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ 0 & 0 & -6 & 12 \end{array} \right] R_3 = -\frac{1}{6}r_3$$
$$\Rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 3 & -2 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

Gaussian Elimination

Example

Example

Now, back solve the system

$$\begin{array}{rclclcl} x & + & y & + & z & = & 3 \\ & & y & + & 3z & = & -2 \\ & & & & z & = & -2 \end{array}$$

Gaussian Elimination

Example

Example

Now, back solve the system

$$\begin{array}{rcrcrcrcrcl} x & + & y & + & z & = & 3 \\ & & y & + & 3z & = & -2 \\ & & & & z & = & -2 \end{array}$$

Start with the third equation: $z = -2$

Gaussian Elimination

Example

Example

Now, back solve the system

$$\begin{array}{rcrcrcrcrcl} x & + & y & + & z & = & 3 \\ & & y & + & 3z & = & -2 \\ & & & & z & = & -2 \end{array}$$

Start with the third equation: $z = -2$

Plug it into the second equation and solve for y :

$$y + 3(-2) = -2 \quad \Rightarrow \quad y = 4$$

Gaussian Elimination

Example

Example

Now, back solve the system

$$\begin{array}{rcrcrcrcrcrl} x & + & y & + & z & = & 3 \\ & & y & + & 3z & = & -2 \\ & & & & z & = & -2 \end{array}$$

Start with the third equation: $z = -2$

Plug it into the second equation and solve for y :

$$y + 3(-2) = -2 \quad \Rightarrow \quad y = 4$$

Plug both into the first equation and solve for x :

$$x + (4) + (-2) = 3 \quad \Rightarrow \quad x = 1$$

Existence and Uniqueness of Solutions

During Gaussian Elimination:

Existence and Uniqueness of Solutions

During Gaussian Elimination:

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid k \neq 0]$$

is encountered, then the system has *no solutions*.

Existence and Uniqueness of Solutions

During Gaussian Elimination:

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid k \neq 0]$$

is encountered, then the system has *no solutions*.

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid 0]$$

is encountered, then the system has *infinitely many solutions*.

Existence and Uniqueness of Solutions

During Gaussian Elimination:

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid k \neq 0]$$

is encountered, then the system has *no solutions*.

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid 0]$$

is encountered, then the system has *infinitely many solutions*.

Some vocabulary:

- If a system has no solutions, it is called **inconsistent**.

Existence and Uniqueness of Solutions

During Gaussian Elimination:

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid k \neq 0]$$

is encountered, then the system has *no solutions*.

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid 0]$$

is encountered, then the system has *infinitely many solutions*.

Some vocabulary:

- If a system has no solutions, it is called **inconsistent**.
- If a system has at least one solution, it is called **consistent**.

Existence and Uniqueness of Solutions

During Gaussian Elimination:

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid k \neq 0]$$

is encountered, then the system has *no solutions*.

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid 0]$$

is encountered, then the system has *infinitely many solutions*.

Some vocabulary:

- If a system has no solutions, it is called **inconsistent**.
- If a system has at least one solution, it is called **consistent**.
 - A system with exactly one solution is called **independent**.

Existence and Uniqueness of Solutions

During Gaussian Elimination:

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid k \neq 0]$$

is encountered, then the system has *no solutions*.

- If a row of the form

$$[0 \quad \cdots \quad 0 \mid 0]$$

is encountered, then the system has *infinitely many solutions*.

Some vocabulary:

- If a system has no solutions, it is called **inconsistent**.
- If a system has at least one solution, it is called **consistent**.
 - A system with exactly one solution is called **independent**.
 - A system with more than one solution is called **dependent**.

Matrix Rank

Reduced Row Echelon Form

An augmented matrix is said to be in **Reduced Row Echelon Form** if:

$$\left[\begin{array}{ccc|c} 1 & \cdots & 0 & k_1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & k_m \end{array} \right]$$

Matrix Rank

Reduced Row Echelon Form

An augmented matrix is said to be in **Reduced Row Echelon Form** if:

$$\left[\begin{array}{ccc|c} 1 & \cdots & 0 & k_1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & k_m \end{array} \right]$$

Rank

The **rank** r of a matrix is equal to how many 1's are in the diagonal of its Reduced Row Echelon Form.

- If r equals the number of variables, there is a unique solution.
- If r is less than the number of variables, the solutions are not unique.