

# The Determinant of a Matrix

Colby Community College

## Determinant of a $2 \times 2$ Matrix

The **determinant of a  $2 \times 2$  matrix** is defined:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

## Determinant of a $2 \times 2$ Matrix

The **determinant of a  $2 \times 2$  matrix** is defined:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

### Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix} = 3 \cdot (-1) - 8 \cdot 5 = -43$$

## Minors of a Matrix

For every element  $a_{ij}$  of a  $n \times n$  matrix  $\mathbf{A}$ , the **minor**  $M_{ij}$  is an  $(n - 1) \times (n - 1)$  matrix obtained by deleting the  $i$ th row and the  $j$ th column of  $\mathbf{A}$ .

## Minors of a Matrix

For every element  $a_{ij}$  of a  $n \times n$  matrix  $\mathbf{A}$ , the **minor**  $M_{ij}$  is an  $(n - 1) \times (n - 1)$  matrix obtained by deleting the  $i$ th row and the  $j$ th column of  $\mathbf{A}$ .

### Example 2

$$\mathbf{A} = \begin{bmatrix} 5 & 4 & -3 \\ 2 & -8 & 1 \\ 9 & 3 & 0 \end{bmatrix} \quad M_{12} =$$

## Minors of a Matrix

For every element  $a_{ij}$  of a  $n \times n$  matrix  $\mathbf{A}$ , the **minor**  $M_{ij}$  is an  $(n - 1) \times (n - 1)$  matrix obtained by deleting the  $i$ th row and the  $j$ th column of  $\mathbf{A}$ .

### Example 2

$$\mathbf{A} = \begin{bmatrix} 5 & 4 & -3 \\ 2 & -8 & 1 \\ 9 & 3 & 0 \end{bmatrix} \qquad M_{12} = \begin{bmatrix} 2 & 1 \\ 9 & 0 \end{bmatrix}$$

## Minors of a Matrix

For every element  $a_{ij}$  of a  $n \times n$  matrix  $\mathbf{A}$ , the **minor**  $M_{ij}$  is an  $(n-1) \times (n-1)$  matrix obtained by deleting the  $i$ th row and the  $j$ th column of  $\mathbf{A}$ .

### Example 2

$$\mathbf{A} = \begin{bmatrix} 5 & 4 & -3 \\ 2 & -8 & 1 \\ 9 & 3 & 0 \end{bmatrix} \quad M_{12} = \begin{bmatrix} 2 & 1 \\ 9 & 0 \end{bmatrix}$$

## Cofactors of a Matrix

For every element  $a_{ij}$  of a  $n \times n$  matrix  $\mathbf{A}$ , the **cofactor** of  $a_{ij}$  is the scalar

$$C_{ij} = (-1)^{(i+j)} |M_{ij}|$$

## Determinants of a $n \times n$ Matrix

For a  $n \times n$  matrix  $\mathbf{A}$ , choose any row or column.



## Determinants of a $n \times n$ Matrix

For a  $n \times n$  matrix  $\mathbf{A}$ , choose any row or column.

**Expansion by the  $i$ th row:**

$$|\mathbf{A}| = \sum_{j=1}^n a_{ij} C_{ij} = \sum_{j=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

## Determinants of a $n \times n$ Matrix

For a  $n \times n$  matrix  $\mathbf{A}$ , choose any row or column.

**Expansion by the  $i$ th row:**

$$|\mathbf{A}| = \sum_{j=1}^n a_{ij} C_{ij} = \sum_{j=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

**Expansion by the  $j$ th column:**

$$|\mathbf{A}| = \sum_{i=1}^n a_{ij} C_{ij} = \sum_{i=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

## Determinants of a $n \times n$ Matrix

For a  $n \times n$  matrix  $\mathbf{A}$ , choose any row or column.

**Expansion by the  $i$ th row:**

$$|\mathbf{A}| = \sum_{j=1}^n a_{ij} C_{ij} = \sum_{j=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

**Expansion by the  $j$ th column:**

$$|\mathbf{A}| = \sum_{i=1}^n a_{ij} C_{ij} = \sum_{i=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

I recommend expanding across the first row.

### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix}$$

### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = 3 \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - 1 \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$

### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + + 3 \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - + 1 \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + -1 \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$

### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + + 3 \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - + 1 \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + - 1 \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$
$$= 3(1 \cdot 2 - 3 \cdot 1) - (2 \cdot 2 - 3 \cdot 0) - (2 \cdot 1 - 1 \cdot 0)$$



### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + + 3 \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - + 1 \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + - 1 \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$
$$= 3(1 \cdot 2 - 3 \cdot 1) - (2 \cdot 2 - 3 \cdot 0) - (2 \cdot 1 - 1 \cdot 0)$$
$$= -3 - 4 - 2$$

### Example 3

Compute the determinant:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{aligned} \begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} &= + + 3 \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - + 1 \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + - 1 \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix} \\ &= 3(1 \cdot 2 - 3 \cdot 1) - (2 \cdot 2 - 3 \cdot 0) - (2 \cdot 1 - 1 \cdot 0) \\ &= -3 - 4 - 2 \\ &= -9 \end{aligned}$$

## Properties of Determinants

- $|\mathbf{A}^T| = |\mathbf{A}|$

## Properties of Determinants

- $|\mathbf{A}^T| = |\mathbf{A}|$
- If  $|\mathbf{A}| \neq 0$ , then  $\mathbf{A}$  is invertable.

## Properties of Determinants

- $|\mathbf{A}^T| = |\mathbf{A}|$
- If  $|\mathbf{A}| \neq 0$ , then  $\mathbf{A}$  is invertable.
- If  $|\mathbf{A}| \neq 0$ , then  $|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|}$

## Properties of Determinants

- $|\mathbf{A}^T| = |\mathbf{A}|$
- If  $|\mathbf{A}| \neq 0$ , then  $\mathbf{A}$  is invertable.
- If  $|\mathbf{A}| \neq 0$ , then  $|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|}$
- If a row (or column) of  $\mathbf{A}$  contains all zeros, then  $|\mathbf{A}| = 0$

## Properties of Determinants

- $|\mathbf{A}^T| = |\mathbf{A}|$
- If  $|\mathbf{A}| \neq 0$ , then  $\mathbf{A}$  is invertable.
- If  $|\mathbf{A}| \neq 0$ , then  $|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|}$
- If a row (or column) of  $\mathbf{A}$  contains all zeros, then  $|\mathbf{A}| = 0$
- If two rows (or two columns) of  $\mathbf{A}$  are equal, then  $|\mathbf{A}| = 0$

## Properties of Determinants

- $|\mathbf{A}^T| = |\mathbf{A}|$
- If  $|\mathbf{A}| \neq 0$ , then  $\mathbf{A}$  is invertable.
- If  $|\mathbf{A}| \neq 0$ , then  $|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|}$
- If a row (or column) of  $\mathbf{A}$  contains all zeros, then  $|\mathbf{A}| = 0$
- If two rows (or two columns) of  $\mathbf{A}$  are equal, then  $|\mathbf{A}| = 0$
- If  $\mathbf{A}$  is an diagonal, upper triangular, or lower triangular matrix, the determinant is the product of the diagonal elements:

$$|\mathbf{A}| = \prod_{i=1}^m a_{ii}$$



## Cramer's Rule

Consider the matrix equation:

$$\mathbf{A}\vec{x} = \vec{b} \quad \text{where} \quad |\mathbf{A}| \neq 0$$


The matrix  $\mathbf{A}_j$  is obtained by replacing the  $j$ th column of  $\mathbf{A}$  with  $\vec{b}$ .

The  $j$ th solution is:

$$x_j = \frac{|\mathbf{A}_j|}{|\mathbf{A}|}$$


## Example 4

The **line of best fit** is the line that gets “closest” to every point.



## Example 4

The **line of best fit** is the line that gets “closest” to every point.



## Least Squares Approximation

A general strategy for finding the line  $y = mx + b$  that best describes a data set is to find  $b$  and  $m$  that minimizes the sums of the squares of the vertical distances between the data points and the line, given by  $F(b, m)$

$$F(b, m) = \sum_{i=1}^n (y_i - (b + mx_i))^2$$

## Least Squares Approximation

A general strategy for finding the line  $y = mx + b$  that best describes a data set is to find  $b$  and  $m$  that minimizes the sums of the squares of the vertical distances between the data points and the line, given by  $F(b, m)$

$$F(b, m) = \sum_{i=1}^n (y_i - (b + mx_i))^2$$

To find such a  $b$  and  $m$ , we need to solve the system:

$$\frac{\partial F}{\partial b} = 0 \quad \text{and} \quad \frac{\partial F}{\partial m} = 0$$

## Least Squares Method


The best-fit straight line for  $n$  data points  $(x_i, y_i)$ ,  $i = 1, 2, \dots, n$ , has y-intercept  $b$  and slope  $m$  as determined by the system

$$\begin{bmatrix} \sum_{i=1}^n 1 & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{bmatrix} \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^n y_i \\ \sum_{i=1}^n x_i y_i \end{bmatrix}$$

## Example 5

Consider the data comparing the high school and college GPA for four students.


$i$	$x_i$	$y_i$
1	1.7	1.1
2	2.3	3.1
3	3.1	2.3
4	4.0	3.8



## Example 5

Consider the data comparing the high school and college GPA for four students.

$i$	$x_i$	$y_i$
1	1.7	1.1
2	2.3	3.1
3	3.1	2.3
4	4.0	3.8



The Least Squares Method system for this dataset is:


$$\begin{bmatrix} 4 & 11.1 \\ 11.1 & 33.79 \end{bmatrix} \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 10.3 \\ 31.33 \end{bmatrix}$$



## Example 5

Consider the data comparing the high school and college GPA for four students.

$i$	$x_i$	$y_i$
1	1.7	1.1
2	2.3	3.1
3	3.1	2.3
4	4.0	3.8




The Least Squares Method system for this dataset is:

$$\begin{bmatrix} 4 & 11.1 \\ 11.1 & 33.79 \end{bmatrix} \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 10.3 \\ 31.33 \end{bmatrix} \Rightarrow \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 0.023 \\ 0.92 \end{bmatrix}$$

## Example 5

Consider the data comparing the high school and college GPA for four students.

$i$	$x_i$	$y_i$
1	1.7	1.1
2	2.3	3.1
3	3.1	2.3
4	4.0	3.8




The Least Squares Method system for this dataset is:

$$\begin{bmatrix} 4 & 11.1 \\ 11.1 & 33.79 \end{bmatrix} \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 10.3 \\ 31.33 \end{bmatrix} \Rightarrow \begin{bmatrix} b \\ m \end{bmatrix} = \begin{bmatrix} 0.023 \\ 0.92 \end{bmatrix}$$

So, the line of best fit is  $y = 0.92x + 0.023$ .


## Example 6

Let us consider some nonlinear approximations.




## Example 6

Let us consider some nonlinear approximations.



## Example 6

Let us consider some nonlinear approximations.



## Example 6

Let us consider some nonlinear approximations.

