

Complements, Conditional Probability, and Bayes's Theorem

Colby Community College

At Least One

When finding the probability of some event occurring “at least once,” we should keep the following in mind:

At Least One

When finding the probability of some event occurring “at least once,” we should keep the following in mind:

- “At least one” has the same meaning as “one or more.”

At Least One

When finding the probability of some event occurring “at least once,” we should keep the following in mind:

- “At least one” has the same meaning as “one or more.”
- The *complement* of getting “at least one” particular event is that you get *no* occurrences of that event.

At Least One

When finding the probability of some event occurring “at least once,” we should keep the following in mind:

- “At least one” has the same meaning as “one or more.”
- The *complement* of getting “at least one” particular event is that you get *no* occurrences of that event.

Example 1

The following are the same event:

- “Not getting at least 1 girl in 10 births.”
- “Getting no girls in 10 births.”
- “Getting 10 boys in 10 births.”

Finding the Probability of “At Least One”

- 1 Let $A =$ “getting at least one event.”

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”

Finding the Probability of “At Least One”

- ① Let A = “getting at least one event.”
- ② Then \bar{A} = “getting none of the event being considered.”
- ③ Find $P(\bar{A})$ = “probability that event A does not occur.”

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”
- 3 Find $P(\bar{A})$ = “probability that event A does not occur.”
- 4 Thus $P(A) = 1 - P(\bar{A})$.

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”
- 3 Find $P(\bar{A})$ = “probability that event A does not occur.”
- 4 Thus $P(A) = 1 - P(\bar{A})$.

Example 2

A study by SquareTrade found that 6% of damaged iPads were damaged by “bags/backpacks.” If 20 damaged iPads are randomly selected, find the probability of getting *at least one* that was damaged in a bag/backpack.

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”
- 3 Find $P(\bar{A})$ = “probability that event A does not occur.”
- 4 Thus $P(A) = 1 - P(\bar{A})$.

Example 2

A study by SquareTrade found that 6% of damaged iPads were damaged by “bags/backpacks.” If 20 damaged iPads are randomly selected, find the probability of getting *at least one* that was damaged in a bag/backpack.

- 1 Let A = “at least 1 iPad was damaged in a bag/backpack.”

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”
- 3 Find $P(\bar{A})$ = “probability that event A does not occur.”
- 4 Thus $P(A) = 1 - P(\bar{A})$.

Example 2

A study by SquareTrade found that 6% of damaged iPads were damaged by “bags/backpacks.” If 20 damaged iPads are randomly selected, find the probability of getting *at least one* that was damaged in a bag/backpack.

- 1 Let A = “at least 1 iPad was damaged in a bag/backpack.”
- 2 Then \bar{A} = “none of the iPads were damaged in a bag/backpack.”

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”
- 3 Find $P(\bar{A})$ = “probability that event A does not occur.”
- 4 Thus $P(A) = 1 - P(\bar{A})$.

Example 2

A study by SquareTrade found that 6% of damaged iPads were damaged by “bags/backpacks.” If 20 damaged iPads are randomly selected, find the probability of getting *at least one* that was damaged in a bag/backpack.

- 1 Let A = “at least 1 iPad was damaged in a bag/backpack.”
- 2 Then \bar{A} = “none of the iPads were damaged in a bag/backpack.”
- 3 Find $P(\bar{A}) = 0.94 \cdot 0.94 \cdot \dots \cdot 0.94 = 0.94^{20} = 0.290$

Finding the Probability of “At Least One”

- 1 Let A = “getting at least one event.”
- 2 Then \bar{A} = “getting none of the event being considered.”
- 3 Find $P(\bar{A})$ = “probability that event A does not occur.”
- 4 Thus $P(A) = 1 - P(\bar{A})$.

Example 2

A study by SquareTrade found that 6% of damaged iPads were damaged by “bags/backpacks.” If 20 damaged iPads are randomly selected, find the probability of getting *at least one* that was damaged in a bag/backpack.

- 1 Let A = “at least 1 iPad was damaged in a bag/backpack.”
- 2 Then \bar{A} = “none of the iPads were damaged in a bag/backpack.”
- 3 Find $P(\bar{A}) = 0.94 \cdot 0.94 \cdot \dots \cdot 0.94 = 0.94^{20} = 0.290$
- 4 Thus $P(A) = 1 - P(\bar{A}) = 1 - 0.290 = 0.710$.

Definition

A **conditional probability** of an event is a probability obtained with the additional information that some other event has already occurred.

Definition

A **conditional probability** of an event is a probability obtained with the additional information that some other event has already occurred.

Intuitive Approach

The conditional probability of B occurring given that A has occurred can be found by assuming that event A has occurred and then calculating the probability that event B will occur.

Definition

A **conditional probability** of an event is a probability obtained with the additional information that some other event has already occurred.

Intuitive Approach

The conditional probability of B occurring given that A has occurred can be found by assuming that event A has occurred and then calculating the probability that event B will occur.

Formal Approach

$$P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$$

Example 3

	Positive Test Result	Negative Test Result
Uses Drugs	45 (True Positive)	5 (False Negative)
Doesn't Use Drugs	25 (False Positive)	480 (True Negative)

Find the following probabilities:

- $P(\text{positive test result} \mid \text{subject uses drugs}) =$

Example 3

	Positive Test Result	Negative Test Result
Uses Drugs	45 (True Positive)	5 (False Negative)
Doesn't Use Drugs	25 (False Positive)	480 (True Negative)

Find the following probabilities:

- $P(\text{positive test result} \mid \text{subject uses drugs}) = \frac{45}{50} = 0.900$

Example 3

	Positive Test Result	Negative Test Result
Uses Drugs	45 (True Positive)	5 (False Negative)
Doesn't Use Drugs	25 (False Positive)	480 (True Negative)

Find the following probabilities:

- $P(\text{positive test result} \mid \text{subject uses drugs}) = \frac{45}{50} = 0.900$
- $P(\text{subject uses drugs} \mid \text{positive test result}) =$

Example 3

	Positive Test Result	Negative Test Result
Uses Drugs	45 (True Positive)	5 (False Negative)
Doesn't Use Drugs	25 (False Positive)	480 (True Negative)

Find the following probabilities:

- $P(\text{positive test result} \mid \text{subject uses drugs}) = \frac{45}{50} = 0.900$
- $P(\text{subject uses drugs} \mid \text{positive test result}) = \frac{45}{70} = 0.643$

Example 3

	Positive Test Result	Negative Test Result
Uses Drugs	45 (True Positive)	5 (False Negative)
Doesn't Use Drugs	25 (False Positive)	480 (True Negative)

Find the following probabilities:

- $P(\text{positive test result} \mid \text{subject uses drugs}) = \frac{45}{50} = 0.900$
- $P(\text{subject uses drugs} \mid \text{positive test result}) = \frac{45}{70} = 0.643$

Note

In general $P(B \mid A) \neq P(A \mid B)$.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Suppose we want to compute $P(C \mid \text{positive test result})$.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Suppose we want to compute $P(C \mid \text{positive test result})$.

We can make the following observations.

- Assume we have 100 subjects. Then, 10 subjects are expected to have cancer. The other 90 do not have cancer.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Suppose we want to compute $P(C \mid \text{positive test result})$.

We can make the following observations.

- Assume we have 100 subjects. Then, 10 subjects are expected to have cancer. The other 90 do not have cancer.
- Among the 90 subjects without cancer, 9 will get a positive result.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Suppose we want to compute $P(C \mid \text{positive test result})$.

We can make the following observations.

- Assume we have 100 subjects. Then, 10 subjects are expected to have cancer. The other 90 do not have cancer.
- Among the 90 subjects without cancer, 9 will get a positive result.
- Among the 90 subjects without cancer, 81 will get a negative result.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Suppose we want to compute $P(C \mid \text{positive test result})$.

We can make the following observations.

- Assume we have 100 subjects. Then, 10 subjects are expected to have cancer. The other 90 do not have cancer.
- Among the 90 subjects without cancer, 9 will get a positive result.
- Among the 90 subjects without cancer, 81 will get a negative result.
- Among the 10 subject with cancer, 8 will get a positive result.

Example 4

Let us assume that cancer has a 1% prevalence, that is 1% of the population has cancer. Let us also assume that C is the event that a randomly selected subject has cancer.

A cancer test has the following performance characteristics:

- $P(C) = 0.01$
- The false positive rate is 10%. That is, $P(\text{positive test result} \mid \bar{C})$
- The true positive rate is 80%. That is, $P(\text{positive test result} \mid C)$

Suppose we want to compute $P(C \mid \text{positive test result})$.

We can make the following observations.

- Assume we have 100 subjects. Then, 10 subjects are expected to have cancer. The other 90 do not have cancer.
- Among the 90 subjects without cancer, 9 will get a positive result.
- Among the 90 subjects without cancer, 81 will get a negative result.
- Among the 10 subjects with cancer, 8 will get a positive result.
- Among the 10 subjects with cancer, 2 will get a negative result.

Example 4

We can summarize the results in the following table.

	Positive Test Result	Negative Test Result	Total
Has Cancer	8 (True Positive)	2 (False Negative)	10
No Cancer	99 (False Positive)	891 (True Negative)	990
Total	107	893	

Example 4

We can summarize the results in the following table.

	Positive Test Result	Negative Test Result	Total
Has Cancer	8 (True Positive)	2 (False Negative)	10
No Cancer	99 (False Positive)	891 (True Negative)	990
Total	107	893	

So, $P(C \mid \text{positive test result}) = 8/107 = 0.0748$.

Example 4

We can summarize the results in the following table.

	Positive Test Result	Negative Test Result	Total
Has Cancer	8 (True Positive)	2 (False Negative)	10
No Cancer	99 (False Positive)	891 (True Negative)	990
Total	107	893	

So, $P(C \mid \text{positive test result}) = 8/107 = 0.0748$.

We can also use the following formula:

$$P(C \mid \text{positive}) = \frac{P(C) \cdot P(\text{positive} \mid C)}{(P(C) \cdot P(\text{positive} \mid C)) + (P(\bar{C}) \cdot P(\text{positive} \mid \bar{C}))}$$

Example 4

We can summarize the results in the following table.

	Positive Test Result	Negative Test Result	Total
Has Cancer	8 (True Positive)	2 (False Negative)	10
No Cancer	99 (False Positive)	891 (True Negative)	990
Total	107	893	

So, $P(C \mid \text{positive test result}) = 8/107 = 0.0748$.

We can also use the following formula:

$$\begin{aligned} P(C \mid \text{positive}) &= \frac{P(C) \cdot P(\text{positive} \mid C)}{(P(C) \cdot P(\text{positive} \mid C)) + (P(\bar{C}) \cdot P(\text{positive} \mid \bar{C}))} \\ &= \frac{0.1 \cdot (8/10)}{(0.01 \cdot 0.80) + (0.99 \cdot 0.10)} \end{aligned}$$

Example 4

We can summarize the results in the following table.

	Positive Test Result	Negative Test Result	Total
Has Cancer	8 (True Positive)	2 (False Negative)	10
No Cancer	99 (False Positive)	891 (True Negative)	990
Total	107	893	

So, $P(C \mid \text{positive test result}) = 8/107 = 0.0748$.

We can also use the following formula:

$$\begin{aligned}P(C \mid \text{positive}) &= \frac{P(C) \cdot P(\text{positive} \mid C)}{(P(C) \cdot P(\text{positive} \mid C)) + (P(\bar{C}) \cdot P(\text{positive} \mid \bar{C}))} \\&= \frac{0.1 \cdot (8/10)}{(0.01 \cdot 0.80) + (0.99 \cdot 0.10)} \\&= 0.0748\end{aligned}$$

Bayes' Theorem

$$P(A | B) = \frac{P(A) P(B | A)}{P(A)} = \frac{P(A) \cdot P(B | A)}{(P(A) \cdot P(B | A)) + (P(\bar{A}) \cdot P(B | \bar{A}))}$$

Bayes' Theorem

$$P(A | B) = \frac{P(A) P(B | A)}{P(A)} = \frac{P(A) \cdot P(B | A)}{(P(A) \cdot P(B | A)) + (P(\bar{A}) \cdot P(B | \bar{A}))}$$

Note

Bayes' Theorem is useful for sequential events, where new information is obtained for a subsequent event, and the new information is used to revise the probability of the original event.

Bayes' Theorem

$$P(A | B) = \frac{P(A) P(B | A)}{P(A)} = \frac{P(A) \cdot P(B | A)}{(P(A) \cdot P(B | A)) + (P(\bar{A}) \cdot P(B | \bar{A}))}$$

Note

Bayes' Theorem is useful for sequential events, where new information is obtained for a subsequent event, and the new information is used to revise the probability of the original event.

Definition

A **prior probability** is an initial probability value originally obtained before any additional information is obtained.

Bayes' Theorem

$$P(A | B) = \frac{P(A) P(B | A)}{P(A)} = \frac{P(A) \cdot P(B | A)}{(P(A) \cdot P(B | A)) + (P(\bar{A}) \cdot P(B | \bar{A}))}$$

Note

Bayes' Theorem is useful for sequential events, where new information is obtained for a subsequent event, and the new information is used to revise the probability of the original event.

Definition

A **prior probability** is an initial probability value originally obtained before any additional information is obtained.

Definition

A **posterior probability** is a probability value that has been revised by using additional information that is later obtained.