

The Determinant of a Matrix

Department of Mathematics

Salt Lake Community College

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix}$$

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix} = 3 \cdot (-1) - 8 \cdot 5$$

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix} = 3 \cdot (-1) - 8 \cdot 5 = -43$$

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix} = 3 \cdot (-1) - 8 \cdot 5 = -43$$

Example 2

$$\begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix}$$

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix} = 3 \cdot (-1) - 8 \cdot 5 = 43$$

Example 2

$$\begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 3 \cdot (1) - (-2) \cdot 6$$

Determinant of a 2×2 Matrix

The **determinant of a 2×2 matrix** is defined to be:

$$|\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Example 1

$$\begin{vmatrix} 3 & 8 \\ 5 & -1 \end{vmatrix} = 3 \cdot (-1) - 8 \cdot 5 = 43$$

Example 2

$$\begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 3 \cdot (1) - (-2) \cdot 6 = 15$$

Minors of a Matrix

For every element a_{ij} of a $n \times n$ matrix \mathbf{A} , the **minor** M_{ij} is an $(n - 1) \times (n - 1)$ matrix obtained by deleting the i th row and the j th column of \mathbf{A} .

Minors of a Matrix

For every element a_{ij} of a $n \times n$ matrix \mathbf{A} , the **minor** M_{ij} is an $(n - 1) \times (n - 1)$ matrix obtained by deleting the i th row and the j th column of \mathbf{A} .

Example 3

$$\mathbf{A} = \begin{bmatrix} 5 & 4 & -3 \\ 2 & -8 & 1 \\ 9 & 3 & 0 \end{bmatrix} \quad M_{12} =$$

Minors of a Matrix

For every element a_{ij} of a $n \times n$ matrix \mathbf{A} , the **minor** M_{ij} is an $(n - 1) \times (n - 1)$ matrix obtained by deleting the i th row and the j th column of \mathbf{A} .

Example 3

$$\mathbf{A} = \begin{bmatrix} 5 & 4 & -3 \\ 2 & -8 & 1 \\ 9 & 3 & 0 \end{bmatrix} \qquad M_{12} = \begin{bmatrix} 2 & 1 \\ 9 & 0 \end{bmatrix}$$

Minors of a Matrix

For every element a_{ij} of a $n \times n$ matrix \mathbf{A} , the **minor** M_{ij} is an $(n-1) \times (n-1)$ matrix obtained by deleting the i th row and the j th column of \mathbf{A} .

Example 3

$$\mathbf{A} = \begin{bmatrix} 5 & 4 & -3 \\ 2 & -8 & 1 \\ 9 & 3 & 0 \end{bmatrix} \quad M_{12} = \begin{bmatrix} 2 & 1 \\ 9 & 0 \end{bmatrix}$$

Cofactors of a Matrix

For every element a_{ij} of a $n \times n$ matrix \mathbf{A} , the **cofactor** of a_{ij} is the scalar

$$C_{ij} = (-1)^{(i+j)} |M_{ij}|$$

Determinants of a $n \times n$ Matrix

For a $n \times n$ matrix **A**, choose any row or column.

Determinants of a $n \times n$ Matrix

For a $n \times n$ matrix \mathbf{A} , choose any row or column.

Expansion by the i th row:

$$|\mathbf{A}| = \sum_{j=1}^n a_{ij} C_{ij} = \sum_{j=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

Determinants of a $n \times n$ Matrix

For a $n \times n$ matrix \mathbf{A} , choose any row or column.

Expansion by the i th row:

$$|\mathbf{A}| = \sum_{j=1}^n a_{ij} C_{ij} = \sum_{j=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

Expansion by the j th column:

$$|\mathbf{A}| = \sum_{i=1}^n a_{ij} C_{ij} = \sum_{i=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

Determinants of a $n \times n$ Matrix

For a $n \times n$ matrix \mathbf{A} , choose any row or column.

Expansion by the i th row:

$$|\mathbf{A}| = \sum_{j=1}^n a_{ij} C_{ij} = \sum_{j=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

Expansion by the j th column:

$$|\mathbf{A}| = \sum_{i=1}^n a_{ij} C_{ij} = \sum_{i=1}^n a_{ij} (-1)^{(i+j)} |\mathbf{M}_{ij}|$$

I recommend you always expand across the first row.

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + (+3) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix}$$

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + (+3) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - (+1) \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix}$$

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + (+3) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - (+1) \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + (+3) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - (+1) \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$
$$= 3(1 \cdot 2 - 3 \cdot 1) - (2 \cdot 2 - 3 \cdot 0) - (2 \cdot 1 - 1 \cdot 0)$$

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} = + (+3) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - (+1) \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix}$$
$$= 3(1 \cdot 2 - 3 \cdot 1) - (2 \cdot 2 - 3 \cdot 0) - (2 \cdot 1 - 1 \cdot 0)$$
$$= -3 - 4 - 2$$

Example 4

Compute the determinant

$$\begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{aligned} \begin{vmatrix} 3 & 1 & -1 \\ 2 & 1 & 3 \\ 0 & 1 & 2 \end{vmatrix} &= + (+3) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} - (+1) \begin{vmatrix} 2 & 3 \\ 0 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix} \\ &= 3(1 \cdot 2 - 3 \cdot 1) - (2 \cdot 2 - 3 \cdot 0) - (2 \cdot 1 - 1 \cdot 0) \\ &= -3 - 4 - 2 \\ &= -9 \end{aligned}$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix} = + (+3) \begin{vmatrix} 6 & 2 \\ -2 & 3 \end{vmatrix}$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix} = + (+3) \begin{vmatrix} 6 & 2 \\ -2 & 3 \end{vmatrix} - (+0) \begin{vmatrix} 4 & 2 \\ 8 & 3 \end{vmatrix}$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix} = + (+3) \begin{vmatrix} 6 & 2 \\ -2 & 3 \end{vmatrix} - (+0) \begin{vmatrix} 4 & 2 \\ 8 & 3 \end{vmatrix} + (-1) \begin{vmatrix} 4 & 6 \\ 8 & -2 \end{vmatrix}$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix} = + (+3) \begin{vmatrix} 6 & 2 \\ -2 & 3 \end{vmatrix} - (+0) \begin{vmatrix} 4 & 2 \\ 8 & 3 \end{vmatrix} + (-1) \begin{vmatrix} 4 & 6 \\ 8 & -2 \end{vmatrix}$$
$$= 3(6 \cdot 3 - 2 \cdot (-2)) + 0(4 \cdot 3 - 2 \cdot 8) - (4 \cdot (-2) - 6 \cdot 8)$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix} = + (+3) \begin{vmatrix} 6 & 2 \\ -2 & 3 \end{vmatrix} - (+0) \begin{vmatrix} 4 & 2 \\ 8 & 3 \end{vmatrix} + (-1) \begin{vmatrix} 4 & 6 \\ 8 & -2 \end{vmatrix} \\ = 3(6 \cdot 3 - 2 \cdot (-2)) + 0(4 \cdot 3 - 2 \cdot 8) - (4 \cdot (-2) - 6 \cdot 8) \\ = 66 + 0 + 56$$

Example 5

Compute the determinant

$$\begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix}$$

Expanding across the first row gives:

$$\begin{aligned} \begin{vmatrix} 3 & 0 & -1 \\ 4 & 6 & 2 \\ 8 & -2 & 3 \end{vmatrix} &= + (+3) \begin{vmatrix} 6 & 2 \\ -2 & 3 \end{vmatrix} - (+0) \begin{vmatrix} 4 & 2 \\ 8 & 3 \end{vmatrix} + (-1) \begin{vmatrix} 4 & 6 \\ 8 & -2 \end{vmatrix} \\ &= 3(6 \cdot 3 - 2 \cdot (-2)) + 0(4 \cdot 3 - 2 \cdot 8) - (4 \cdot (-2) - 6 \cdot 8) \\ &= 66 + 0 + 56 \\ &= 122 \end{aligned}$$

Properties of Determinants

- If any row (or any column) of \mathbf{A} contains all zeros, then $|\mathbf{A}| = 0$

Properties of Determinants

- If any row (or any column) of \mathbf{A} contains all zeros, then $|\mathbf{A}| = 0$
- If any two rows (or any two columns) of \mathbf{A} are equal, then $|\mathbf{A}| = 0$

Properties of Determinants

- If any row (or any column) of \mathbf{A} contains all zeros, then $|\mathbf{A}| = 0$
- If any two rows (or any two columns) of \mathbf{A} are equal, then $|\mathbf{A}| = 0$
- If \mathbf{A} is an diagonal, upper triangular, or lower triangular matrix, the determinant is the product of the diagonal elements.

Properties of Determinants

- If any row (or any column) of \mathbf{A} contains all zeros, then $|\mathbf{A}| = 0$
- If any two rows (or any two columns) of \mathbf{A} are equal, then $|\mathbf{A}| = 0$
- If \mathbf{A} is an diagonal, upper triangular, or lower triangular matrix, the determinant is the product of the diagonal elements.
- The value of $|\mathbf{A}|$ changes sign if any two rows (or columns) are interchanged.

Properties of Determinants

- If any row (or any column) of \mathbf{A} contains all zeros, then $|\mathbf{A}| = 0$
- If any two rows (or any two columns) of \mathbf{A} are equal, then $|\mathbf{A}| = 0$
- If \mathbf{A} is an diagonal, upper triangular, or lower triangular matrix, the determinant is the product of the diagonal elements.
- The value of $|\mathbf{A}|$ changes sign if any two rows (or columns) are interchanged.
- If any row (or any column) of $|\mathbf{A}|$ is multiplied by a nonzero number k , the value of $|\mathbf{A}|$ is also changed by a factor of k .

Properties of Determinants

- If any row (or any column) of \mathbf{A} contains all zeros, then $|\mathbf{A}| = 0$
- If any two rows (or any two columns) of \mathbf{A} are equal, then $|\mathbf{A}| = 0$
- If \mathbf{A} is an diagonal, upper triangular, or lower triangular matrix, the determinant is the product of the diagonal elements.
- The value of $|\mathbf{A}|$ changes sign if any two rows (or columns) are interchanged.
- If any row (or any column) of $|\mathbf{A}|$ is multiplied by a nonzero number k , the value of $|\mathbf{A}|$ is also changed by a factor of k .
- If the entries of any row (or any column) of $|\mathbf{A}|$ are multiplied by a nonzero number k and the result is added to another row, the value of $|\mathbf{A}|$ remains unchanged.

Cramer's Rule

Consider the system:

$$\begin{array}{rclcl} ax & + & by & = & s \\ cx & + & dy & = & t \end{array}$$

Cramer's Rule

Consider the system:

$$\begin{array}{rclcl} ax & + & by & = & s \\ cx & + & dy & = & t \end{array}$$

We then have the following three determinants:

$$|\mathbf{D}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} \quad |\mathbf{D}_x| = \begin{vmatrix} s & b \\ t & d \end{vmatrix} \quad |\mathbf{D}_y| = \begin{vmatrix} a & s \\ c & t \end{vmatrix}$$

Cramer's Rule

Consider the system:

$$\begin{array}{rclcl} ax & + & by & = & s \\ cx & + & dy & = & t \end{array}$$

We then have the following three determinants:

$$|\mathbf{D}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} \quad |\mathbf{D}_x| = \begin{vmatrix} s & b \\ t & d \end{vmatrix} \quad |\mathbf{D}_y| = \begin{vmatrix} a & s \\ c & t \end{vmatrix}$$

The solutions is:

$$x = \frac{|D_x|}{|D|} = \frac{\begin{vmatrix} s & b \\ t & d \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}} \qquad y = \frac{|D_y|}{|D|} = \frac{\begin{vmatrix} a & s \\ c & t \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}}$$

Cramer's Rule

Consider the system:

$$\begin{array}{rclcl} ax & + & by & = & s \\ cx & + & dy & = & t \end{array}$$

We then have the following three determinants:

$$|D| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} \quad |D_x| = \begin{vmatrix} s & b \\ t & d \end{vmatrix} \quad |D_y| = \begin{vmatrix} a & s \\ c & t \end{vmatrix}$$

The solutions is:

$$x = \frac{|D_x|}{|D|} = \frac{\begin{vmatrix} s & b \\ t & d \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}} \qquad y = \frac{|D_y|}{|D|} = \frac{\begin{vmatrix} a & s \\ c & t \end{vmatrix}}{\begin{vmatrix} a & b \\ c & d \end{vmatrix}}$$

Note

This method can be extended to any size system of equations.

Example 6

Consider the system

$$\begin{array}{rclcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Example 6

Consider the system

$$\begin{array}{rcrcrcrcrcrl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix}$$

$$|D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix}$$

$$|D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix}$$

Example 6

Consider the system

$$\begin{array}{rclcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \qquad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} \qquad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix}$$

Example 6

Consider the system

$$\begin{array}{rclcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \quad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \quad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix}$$

Example 6

Consider the system

$$\begin{array}{rcrcrcrcrcrl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \qquad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \qquad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

Example 6

Consider the system

$$\begin{array}{rclcrcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \qquad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \qquad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

We can now find x

$$x = \frac{|D_x|}{|D|}$$

Example 6

Consider the system

$$\begin{array}{rclcrcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \quad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \quad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

We can now find x

$$x = \frac{|D_x|}{|D|} = \frac{-1}{-1}$$

Example 6

Consider the system

$$\begin{array}{rclcrcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \quad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \quad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

We can now find x

$$x = \frac{|D_x|}{|D|} = \frac{-1}{-1} = 1$$

Example 6

Consider the system

$$\begin{array}{rclcrcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \quad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \quad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

We can now find x and y

$$x = \frac{|D_x|}{|D|} = \frac{-1}{-1} = 1 \quad y = \frac{|D_y|}{|D|}$$

Example 6

Consider the system

$$\begin{array}{rclcrcl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \quad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \quad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

We can now find x and y

$$x = \frac{|D_x|}{|D|} = \frac{-1}{-1} = 1 \quad y = \frac{|D_y|}{|D|} = \frac{-2}{-1}$$

Example 6

Consider the system

$$\begin{array}{rcrcrcrcrcrl} x & + & 2y & = & 5 \\ 2x & + & 3y & = & 8 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \quad |D_x| = \begin{vmatrix} 5 & 2 \\ 8 & 3 \end{vmatrix} = -1 \quad |D_y| = \begin{vmatrix} 1 & 5 \\ 2 & 8 \end{vmatrix} = -2$$

We can now find x and y

$$x = \frac{|D_x|}{|D|} = \frac{-1}{-1} = 1 \quad y = \frac{|D_y|}{|D|} = \frac{-2}{-1} = 2$$

Example 7

Consider the system

$$\begin{array}{rclcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Example 7

Consider the system

$$\begin{array}{rcrcrcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix}$$

$$|D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix}$$

$$|D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

We can now find x

$$x = \frac{|D_x|}{|D|}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

We can now find x

$$x = \frac{|D_x|}{|D|} = \frac{30}{15}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

We can now find x

$$x = \frac{|D_x|}{|D|} = \frac{30}{15} = 2$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

We can now find x and y

$$x = \frac{|D_x|}{|D|} = \frac{30}{15} = 2 \quad y = \frac{|D_y|}{|D|}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

We can now find x and y

$$x = \frac{|D_x|}{|D|} = \frac{30}{15} = 2 \quad y = \frac{|D_y|}{|D|} = \frac{15}{15}$$

Example 7

Consider the system

$$\begin{array}{rclcrcl} 3x & - & 2y & = & 4 \\ 6x & + & y & = & 13 \end{array}$$

Let us solve this system using Cramer's Rule.

This means we need to calculate the following determinants.

$$|D| = \begin{vmatrix} 3 & -2 \\ 6 & 1 \end{vmatrix} = 15 \quad |D_x| = \begin{vmatrix} 4 & -2 \\ 13 & 1 \end{vmatrix} = 30 \quad |D_y| = \begin{vmatrix} 3 & 4 \\ 6 & 13 \end{vmatrix} = 15$$

We can now find x and y

$$x = \frac{|D_x|}{|D|} = \frac{30}{15} = 2 \quad y = \frac{|D_y|}{|D|} = \frac{15}{15} = 1$$