


Laplace Transforms

Adam Wilson


Salt Lake Community College

In this chapter we shall study the Laplace transformation, which will allow us to reframe some hard problems into equivalent easier problems.

In this chapter we shall study the Laplace transformation, which will allow us to reframe some hard problems into equivalent easier problems.



In this chapter we shall study the Laplace transformation, which will allow us to reframe some hard problems into equivalent easier problems.



Laplace Transform

The **Laplace Transform** $\mathcal{L}\{f(t)\}$ of a suitable function $f(t)$ defined on $[0, \infty)$ is the function $F(s)$ given by

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt = \lim_{b \rightarrow \infty} \int_0^b e^{-st} f(t) dt$$

where s may be complex.

Laplace Transform

The **Laplace Transform** $\mathcal{L}\{f(t)\}$ of a suitable function $f(t)$ defined on $[0, \infty)$ is the function $F(s)$ given by

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt = \lim_{b \rightarrow \infty} \int_0^b e^{-st} f(t) dt$$

where s may be complex.

Linearity of the Laplace Transform

If $F(s) = \mathcal{L}\{f(t)\}$ and $G(s) = \mathcal{L}\{g(t)\}$, then by the properties of integrals

$$\mathcal{L}\{af(t) + bg(t)\} = aF(s) + bG(s) \quad \text{for } a, b \in \mathbb{C}$$

Laplace Transform

The **Laplace Transform** $\mathcal{L}\{f(t)\}$ of a suitable function $f(t)$ defined on $[0, \infty)$ is the function $F(s)$ given by

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt = \lim_{b \rightarrow \infty} \int_0^b e^{-st} f(t) dt$$

where s may be complex.

Linearity of the Laplace Transform

If $F(s) = \mathcal{L}\{f(t)\}$ and $G(s) = \mathcal{L}\{g(t)\}$, then by the properties of integrals

$$\mathcal{L}\{af(t) + bg(t)\} = aF(s) + bG(s) \quad \text{for } a, b \in \mathbb{C}$$

Existence Theorem for Laplace Transform

If $f(t)$ is piecewise continuous on $[0, \infty)$ and of exponential order α , then the Laplace transform $F(s) = \mathcal{L}\{f(t)\}$ exists for $s > \alpha$.

Example

Let us consider the constant function $f(t) = 1$.

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\}$$

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{1\}$$

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{1\} = \int_0^{\infty} e^{-st} \cdot 1 dt$$

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{1\} = \int_0^{\infty} e^{-st} \cdot 1 dt = \lim_{b \rightarrow \infty} \left[-\frac{e^{-st}}{s} \right]_0^b$$

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{1\} = \int_0^{\infty} e^{-st} \cdot 1 dt = \lim_{b \rightarrow \infty} \left[-\frac{e^{-st}}{s} \right]_0^b = \lim_{b \rightarrow \infty} \left[-\frac{e^{-sb}}{s} + \frac{1}{s} \right]$$

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{1\} = \int_0^{\infty} e^{-st} \cdot 1 dt = \lim_{b \rightarrow \infty} \left[-\frac{e^{-st}}{s} \right]_0^b = \lim_{b \rightarrow \infty} \left[-\frac{e^{-sb}}{s} + \frac{1}{s} \right]$$

If $s > 0$, then $e^{-sb} = \frac{1}{e^{sb}} \rightarrow 0$ as $b \rightarrow \infty$.

Example

Let us consider the constant function $f(t) = 1$.

For $t \geq 0$, we can calculate



$$\mathcal{L}\{f(t)\} = \mathcal{L}\{1\} = \int_0^{\infty} e^{-st} \cdot 1 dt = \lim_{b \rightarrow \infty} \left[-\frac{e^{-st}}{s} \right]_0^b = \lim_{b \rightarrow \infty} \left[-\frac{e^{-sb}}{s} + \frac{1}{s} \right]$$

If $s > 0$, then $e^{-sb} = \frac{1}{e^{sb}} \rightarrow 0$ as $b \rightarrow \infty$.

$$\mathcal{L}\{1\} = F(s) = \frac{1}{s} \quad s > 0$$

Example

Let us consider the constant function $f(t) = 1$.



Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\}$$

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{e^{at}\}$$

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{e^{at}\} = \int_0^{\infty} e^{-st} \cdot e^{at} dt$$

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{e^{at}\} = \int_0^{\infty} e^{-st} \cdot e^{at} dt = \lim_{b \rightarrow \infty} \int_0^b e^{-(s-a)t} dt$$

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate

$$\begin{aligned}\mathcal{L}\{f(t)\} &= \mathcal{L}\{e^{at}\} = \int_0^{\infty} e^{-st} \cdot e^{at} dt = \lim_{b \rightarrow \infty} \int_0^b e^{-(s-a)t} dt \\ &= \lim_{b \rightarrow \infty} \left[-\frac{e^{-(s-a)b}}{s-a} + \frac{1}{s-a} \right]\end{aligned}$$

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate

$$\begin{aligned}\mathcal{L}\{f(t)\} &= \mathcal{L}\{e^{at}\} = \int_0^{\infty} e^{-st} \cdot e^{at} dt = \lim_{b \rightarrow \infty} \int_0^b e^{-(s-a)t} dt \\ &= \lim_{b \rightarrow \infty} \left[-\frac{e^{-(s-a)b}}{s-a} + \frac{1}{s-a} \right]\end{aligned}$$

If $s > 0$ and $s > a$, then $e^{-(s-a)b} = \frac{1}{e^{(s-a)b}} \rightarrow 0$ as $b \rightarrow \infty$.

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.

For $t \geq 0$, we can calculate



$$\begin{aligned}\mathcal{L}\{f(t)\} &= \mathcal{L}\{e^{at}\} = \int_0^{\infty} e^{-st} \cdot e^{at} dt = \lim_{b \rightarrow \infty} \int_0^b e^{-(s-a)t} dt \\ &= \lim_{b \rightarrow \infty} \left[-\frac{e^{-(s-a)b}}{s-a} + \frac{1}{s-a} \right]\end{aligned}$$

If $s > 0$ and $s > a$, then $e^{-(s-a)b} = \frac{1}{e^{(s-a)b}} \rightarrow 0$ as $b \rightarrow \infty$.

$$\mathcal{L}\{e^{at}\} = F(s) = \frac{1}{s-a} \quad s > a$$

Example

Let us consider the constant function $f(t) = e^{at}$, $a \in \mathbb{R}$.



Example

Let us calculate the Laplace transform for $f(t) = 5 - 3e^{-2t}$

Example

Let us calculate the Laplace transform for $f(t) = 5 - 3e^{-2t}$

$$\mathcal{L}\{f(t)\}$$

Example

Let us calculate the Laplace transform for $f(t) = 5 - 3e^{-2t}$

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{5 \cdot 1 - 3e^{-2t}\}$$

Example

Let us calculate the Laplace transform for $f(t) = 5 - 3e^{-2t}$

$$\begin{aligned}\mathcal{L}\{f(t)\} &= \mathcal{L}\{5 \cdot 1 - 3e^{-2t}\} \\ &= 5\mathcal{L}\{1\} - 3\mathcal{L}\{e^{-2t}\}\end{aligned}$$

Example

Let us calculate the Laplace transform for $f(t) = 5 - 3e^{-2t}$

$$\begin{aligned}\mathcal{L}\{f(t)\} &= \mathcal{L}\{5 \cdot 1 - 3e^{-2t}\} \\ &= 5\mathcal{L}\{1\} - 3\mathcal{L}\{e^{-2t}\} \\ &= 5\left(\frac{1}{s}\right) - 3\left(\frac{1}{s+2}\right)\end{aligned}$$

Example

Let us calculate the Laplace transform for $f(t) = 5 - 3e^{-2t}$

$$\begin{aligned}\mathcal{L}\{f(t)\} &= \mathcal{L}\{5 \cdot 1 - 3e^{-2t}\} \\ &= 5\mathcal{L}\{1\} - 3\mathcal{L}\{e^{-2t}\} \\ &= 5\left(\frac{1}{s}\right) - 3\left(\frac{1}{s+2}\right) \\ &= \frac{2s+10}{s(s+2)}\end{aligned}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\} = \mathcal{L}\{\cos(kt)\} + i\mathcal{L}\{\sin(kt)\}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\} = \mathcal{L}\{\cos(kt)\} + i\mathcal{L}\{\sin(kt)\}$$

But, we know how to calculate e^{ikt} :

$$\mathcal{L}\{e^{ikt}\} = \frac{1}{s - ik}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\} = \mathcal{L}\{\cos(kt)\} + i\mathcal{L}\{\sin(kt)\}$$

But, we know how to calculate e^{ikt} :

$$\mathcal{L}\{e^{ikt}\} = \frac{1}{s - ik} \cdot \frac{s + ik}{s + ik}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\} = \mathcal{L}\{\cos(kt)\} + i\mathcal{L}\{\sin(kt)\}$$

But, we know how to calculate e^{ikt} :

$$\begin{aligned}\mathcal{L}\{e^{ikt}\} &= \frac{1}{s - ik} \cdot \frac{s + ik}{s + ik} \\ &= \frac{s + ik}{s^2 + k^2}\end{aligned}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\} = \mathcal{L}\{\cos(kt)\} + i\mathcal{L}\{\sin(kt)\}$$

But, we know how to calculate e^{ikt} :

$$\begin{aligned}\mathcal{L}\{e^{ikt}\} &= \frac{1}{s - ik} \cdot \frac{s + ik}{s + ik} \\ &= \frac{s + ik}{s^2 + k^2} \\ &= \frac{s}{s^2 + k^2} + i \frac{k}{s^2 + k^2}\end{aligned}$$

Example

Instead of directly calculating the Laplace transform for \sin and \cos , let us use Euler's formula to make the job easier.

$$\mathcal{L}\{e^{ikt}\} = \mathcal{L}\{\cos(kt) + i \sin(kt)\} = \mathcal{L}\{\cos(kt)\} + i\mathcal{L}\{\sin(kt)\}$$

But, we know how to calculate e^{ikt} :

$$\begin{aligned}\mathcal{L}\{e^{ikt}\} &= \frac{1}{s - ik} \cdot \frac{s + ik}{s + ik} \\ &= \frac{s + ik}{s^2 + k^2} \\ &= \frac{s}{s^2 + k^2} + i \frac{k}{s^2 + k^2}\end{aligned}$$

So, if we equate the real and imaginary parts, we get:

$$\mathcal{L}\{\cos(kt)\} = \frac{s}{s^2 + k^2} \quad \text{and} \quad \mathcal{L}\{\sin(kt)\} = \frac{k}{s^2 + k^2}$$

Inverse Laplace Transform

A function $f(t)$ whose transform is $F(s)$ is called the **inverse Laplace transform** of F , and we write

$$f(t) = \mathcal{L}^{-1}\{F(s)\}$$

Example

Let us calculate the inverse Laplace transform for

$$F(s) = \frac{2s - 14}{(s + 1)(s - 3)}$$

Example

Let us calculate the inverse Laplace transform for

$$F(s) = \frac{2s - 14}{(s + 1)(s - 3)}$$

the Partial Fractions Decomposition of $F(s)$ is

$$F(s) = \frac{4}{s + 1} - 2\frac{2}{s - 3} = 4 \cdot \frac{1}{s + 1} - 2 \cdot \frac{1}{s - 3}$$

Example

Let us calculate the inverse Laplace transform for

$$F(s) = \frac{2s - 14}{(s + 1)(s - 3)}$$

the Partial Fractions Decomposition of $F(s)$ is

$$F(s) = \frac{4}{s + 1} - 2\frac{2}{s - 3} = 4 \cdot \frac{1}{s + 1} - 2 \cdot \frac{1}{s - 3}$$

So, we can use linearity to find $f(t)$.

$$F(s) = 4\mathcal{L}\{e^{-t}\} - 2\mathcal{L}\{3t\}$$

Example

Let us calculate the inverse Laplace transform for

$$F(s) = \frac{2s - 14}{(s + 1)(s - 3)}$$

the Partial Fractions Decomposition of $F(s)$ is

$$F(s) = \frac{4}{s + 1} - 2\frac{2}{s - 3} = 4 \cdot \frac{1}{s + 1} - 2 \cdot \frac{1}{s - 3}$$

So, we can use linearity to find $f(t)$.

$$F(s) = 4\mathcal{L}\{e^{-t}\} - 2\mathcal{L}\{3t\} = \mathcal{L}\{4e^{-t} - 2e^{3t}\}$$

Example

Let us calculate the inverse Laplace transform for

$$F(s) = \frac{2s - 14}{(s + 1)(s - 3)}$$

the Partial Fractions Decomposition of $F(s)$ is

$$F(s) = \frac{4}{s + 1} - 2\frac{2}{s - 3} = 4 \cdot \frac{1}{s + 1} - 2 \cdot \frac{1}{s - 3}$$

So, we can use linearity to find $f(t)$.

$$F(s) = 4\mathcal{L}\{e^{-t}\} - 2\mathcal{L}\{3t\} = \mathcal{L}\{4e^{-t} - 2e^{3t}\}$$

Thus,

$$f(t) = 4e^{-t} - 2e^{3t}$$

Some Laplace Transforms

$f(t)$	$\mathcal{L}\{f(t)\}$	
1	$\frac{1}{s}$	$s > 0$
t^n	$\frac{n!}{s^{n+1}}$	$s > 0, n \in \mathbb{N}^+$
e^{at}	$\frac{1}{s-a}$	$s > a$
$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$	$s > a, n \in \mathbb{N}^+$
$\sin(bt)$	$\frac{b}{s^2 + b^2}$	$s > 0$
$\cos(bt)$	$\frac{s}{s^2 + b^2}$	$s > 0$

Some More Laplace Transforms

$f(t)$	$\mathcal{L}\{f(t)\}$	
$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2 + b^2}$	$s > a$
$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2 + b^2}$	$s > a$
$\sinh(bt)$	$\frac{b}{s^2 - b^2}$	$s > b $
$\cosh(bt)$	$\frac{s}{s^2 - b^2}$	$s > b $

Example

Let us find the inverse Laplace transform for

$$F(s) = \frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)}$$

Example

Let us find the inverse Laplace transform for

$$F(s) = \frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)}$$

We can again use Partial Fraction Decomposition.

$$\frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)} = \frac{2}{s - 1} - \frac{1}{s^2 + 5}$$

Example

Let us find the inverse Laplace transform for

$$F(s) = \frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)}$$

We can again use Partial Fraction Decomposition.

$$\frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)} = \frac{2}{s - 1} - \frac{1}{s^2 + 5}$$

Now, if we let $b = \sqrt{5}$, then F becomes

$$F(s) = 2 \underbrace{\left(\frac{1}{s - 1} \right)}_{\mathcal{L}\{e^t\}} - \frac{1}{\sqrt{5}} \underbrace{\left(\frac{\sqrt{5}}{s^2 + (\sqrt{5})^2} \right)}_{\mathcal{L}\{\sin(\sqrt{5}t)\}}$$

Example

Let us find the inverse Laplace transform for

$$F(s) = \frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)}$$

We can again use Partial Fraction Decomposition.

$$\frac{2s^2 - s + 11}{(s - 1)(s^2 + 5)} = \frac{2}{s - 1} - \frac{1}{s^2 + 5}$$

Now, if we let $b = \sqrt{5}$, then F becomes

$$F(s) = 2 \underbrace{\left(\frac{1}{s - 1} \right)}_{\mathcal{L}\{e^t\}} - \frac{1}{\sqrt{5}} \underbrace{\left(\frac{\sqrt{5}}{s^2 + (\sqrt{5})^2} \right)}_{\mathcal{L}\{\sin(\sqrt{5}t)\}}$$

$$\text{Thus, } f(t) = \mathcal{L}^{-1}\{F(s)\} = 2e^t - \frac{1}{\sqrt{5}} \sin(\sqrt{5}t).$$

Example

Consider

$$F(s) = \frac{s + 1}{s^2 + 4s + 13}$$

Example

Consider

$$F(s) = \frac{s + 1}{s^2 + 4s + 13}$$

To find $\mathcal{L}\{F(s)\}$, we will need to rearrange things a bit.

$$\frac{s + 1}{s^2 + 4s + 13} = \frac{(s + 2) - 1}{(s^2 + 4s + 4) + (9)}$$

Example

Consider

$$F(s) = \frac{s + 1}{s^2 + 4s + 13}$$

To find $\mathcal{L}\{F(s)\}$, we will need to rearrange things a bit.

$$\begin{aligned}\frac{s + 1}{s^2 + 4s + 13} &= \frac{(s + 2) - 1}{(s^2 + 4s + 4) + (9)} \\ &= \frac{(s + 2) - 1}{(s + 2)^2 + 3^2}\end{aligned}$$

Example

Consider

$$F(s) = \frac{s + 1}{s^2 + 4s + 13}$$

To find $\mathcal{L}\{F(s)\}$, we will need to rearrange things a bit.

$$\begin{aligned}\frac{s + 1}{s^2 + 4s + 13} &= \frac{(s + 2) - 1}{(s^2 + 4s + 4) + (9)} \\ &= \frac{(s + 2) - 1}{(s + 2)^2 + 3^2} \\ &= \frac{s + 2}{(s + 2)^2 + 3^2} - \frac{1}{(s + 2)^2 + 3^2}\end{aligned}$$

Example

Consider

$$F(s) = \frac{s + 1}{s^2 + 4s + 13}$$

To find $\mathcal{L}\{F(s)\}$, we will need to rearrange things a bit.

$$\begin{aligned}\frac{s + 1}{s^2 + 4s + 13} &= \frac{(s + 2) - 1}{(s^2 + 4s + 4) + (9)} \\ &= \frac{(s + 2) - 1}{(s + 2)^2 + 3^2} \\ &= \frac{s + 2}{(s + 2)^2 + 3^2} - \frac{1}{(s + 2)^2 + 3^2} \\ &= \frac{s + 2}{(s + 2)^2 + 3^2} - \frac{1}{3} \frac{3}{(s + 2)^2 + 3^2}\end{aligned}$$

Example

Consider

$$F(s) = \frac{s+1}{s^2+4s+13}$$

To find $\mathcal{L}\{F(s)\}$, we will need to rearrange things a bit.

$$\begin{aligned}\frac{s+1}{s^2+4s+13} &= \frac{(s+2)-1}{(s^2+4s+4)+(9)} \\ &= \frac{(s+2)-1}{(s+2)^2+3^2} \\ &= \frac{s+2}{(s+2)^2+3^2} - \frac{1}{(s+2)^2+3^2} \\ &= \frac{s+2}{(s+2)^2+3^2} - \frac{1}{3} \frac{3}{(s+2)^2+3^2} \\ &= \mathcal{L}\{e^{-2t} \cos(3t)\} - \frac{1}{3} \mathcal{L}\{s^{-2t} \sin(3t)\}\end{aligned}$$

Example

Consider

$$F(s) = \frac{s+1}{s^2+4s+13}$$

To find $\mathcal{L}\{F(s)\}$, we will need to rearrange things a bit.

$$\begin{aligned}\frac{s+1}{s^2+4s+13} &= \frac{(s+2)-1}{(s^2+4s+4)+(9)} \\ &= \frac{(s+2)-1}{(s+2)^2+3^2} \\ &= \frac{s+2}{(s+2)^2+3^2} - \frac{1}{(s+2)^2+3^2} \\ &= \frac{s+2}{(s+2)^2+3^2} - \frac{1}{3} \frac{3}{(s+2)^2+3^2} \\ &= \mathcal{L}\{e^{-2t} \cos(3t)\} - \frac{1}{3} \mathcal{L}\{s^{-2t} \sin(3t)\}\end{aligned}$$

Thus, by linearity we have $f(t) = e^{-2t} \cos(3t) - \frac{1}{3} s^{-2t} \sin(3t)$.