

Linear Equations

Department of Mathematics

Salt Lake Community College

(Slides by Adam Wilson)

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15$$

$$4x - 2y + 3\sqrt{z} = 12$$

$$2x - 3y + 4z + 3 = w$$

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15 \quad \text{No}$$

$$4x - 2y + 3\sqrt{z} = 12$$

$$2x - 3y + 4z + 3 = w$$

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15 \quad \text{No}$$

$$4x - 2y + 3\sqrt{z} = 12 \quad \text{No}$$

$$2x - 3y + 4z + 3 = w$$

Definition

An equation $F(x_1, x_2, \dots, x_n) = C$ is **linear** if it is of the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = C$$

where a_1, a_2, \dots, a_n and C are constants.

If $C = 0$, the equation is said to be **homogeneous**.

Example 1

Which of the following are linear equations?

$$4x - 3e^x = 15 \quad \text{No}$$

$$4x - 2y + 3\sqrt{z} = 12 \quad \text{No}$$

$$2x - 3y + 4z + 3 = w \quad \text{Yes}$$

Definition

A differential equation $F(y, y', y'', \dots, y^{(n)}) = f(t)$ is **linear** if it is of the form

$$a_n(t) \frac{d^n y}{dt^n} + a_{n-1}(t) \frac{d^{n-1} y}{dt^{n-1}} + \cdots + a_1(t) \frac{dy}{dt} + a_0(t)y = f(t)$$

where all functions of t are assumed to be defined over some common interval I .

If $f(t) = 0$ over the interval I , the differential equation is said to be **homogeneous**.

Definition

A differential equation $F(y, y', y'', \dots, y^{(n)}) = f(t)$ is **linear** if it is of the form

$$a_n(t) \frac{d^n y}{dt^n} + a_{n-1}(t) \frac{d^{n-1} y}{dt^{n-1}} + \cdots + a_1(t) \frac{dy}{dt} + a_0(t)y = f(t)$$

where all functions of t are assumed to be defined over some common interval I .

If $f(t) = 0$ over the interval I , the differential equation is said to be **homogeneous**.

First and Second Order Notation

It is common to write first-order differential equations as

$$y' + p(t)y = f(t)$$

and second-order differential equations as

$$y'' + p(t)y' + q(t)y = f(t)$$

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
-----------------------	-------	---------	--------------	--------------

$$y' + ty = 1$$

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes		

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
-----------------------	-------	---------	--------------	--------------

$$y' + ty = 1$$

1

Yes

No

Variable

$$y'' + yy' + y = t$$

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes		

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$				

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3			

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3	Yes		

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3	Yes	No	

Example 2

Let us classify the following differential equations.

Differential Equation	Order	Linear?	Homogeneous?	Coefficients
$y' + ty = 1$	1	Yes	No	Variable
$y'' + yy' + y = t$	2	No	—	—
$y'' + ty' + y^2 = 0$	2	No	—	—
$y'' + 3y' + 2y = 0$	2	Yes	Yes	Constant
$y'' + y = \sin(y)$	2	No	—	—
$y''' + 3y' + y = \sin(t)$	3	Yes	No	Variable

Notation

We will use a **vector** notation to represent a whole set of variables:

Linear Algebraic Equations:

$$\vec{x} = [x_1, x_2, \dots, x_n]$$

Linear Differential Equations:

$$\vec{y} = [y^{(n)}, y^{(n-1)}, \dots, y', y]$$

Notation

We will use a **vector** notation to represent a whole set of variables:

Linear Algebraic Equations:

$$\vec{x} = [x_1, x_2, \dots, x_n]$$

Linear Differential Equations:

$$\vec{y} = [y^{(n)}, y^{(n-1)}, \dots, y', y]$$

Definition

A **linear operator** L is an entire operation performed on a set of variables.

Linear Algebraic Equations:

$$L(\vec{x}) = a_1x_1 + a_2x_2 + \dots + a_nx_n$$

Linear Differential Equations:

$$L(\vec{y}) = a_n(t) \frac{d^n y}{dt^n} + a_{n-1}(t) \frac{d^{n-1} y}{dt^{n-1}} + \dots + a_1(t) \frac{dy}{dt} + a_0(t)y$$

Example 3

What is the linear operator for the following linear differential equations?

$$y' + ty = 1$$

Example 3

What is the linear operator for the following linear differential equations?

$$y' + ty = 1 \quad \rightarrow \quad L(\vec{y}) = y' + ty$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{l} y' + ty = 1 \\ y'' + 2y = 3y' + t \end{array} \quad \rightarrow \quad L(\vec{y}) = y' + ty$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{l} y' + ty = 1 \\ y'' - 3y' + 2y = t \end{array} \quad \rightarrow \quad L(\vec{y}) = y' + ty$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \end{array}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \\ y^{(4)} + 3y = \sin(t) & \end{array}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \\ y^{(4)} + 3y = \sin(t) & \rightarrow L(\vec{y}) = y^{(4)} + 3y \end{array}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{array}{ll} y' + ty = 1 & \rightarrow L(\vec{y}) = y' + ty \\ y'' - 3y' + 2y = t & \rightarrow L(\vec{y}) = y'' - 3y' + 2y \\ y^{(4)} + 3y = \sin(t) & \rightarrow L(\vec{y}) = y^{(4)} + 3y \end{array}$$

Linear Operator Properties

$$\begin{aligned} L(k\vec{u}) &= kL(\vec{u}), \quad k \in \mathbb{R} \\ L(\vec{u} + \vec{w}) &= L(\vec{u}) + L(\vec{w}) \end{aligned}$$

Example 3

What is the linear operator for the following linear differential equations?

$$\begin{aligned}y' + ty &= 1 & \rightarrow & L(\vec{y}) = y' + ty \\y'' - 3y' + 2y &= t & \rightarrow & L(\vec{y}) = y'' - 3y' + 2y \\y^{(4)} + 3y &= \sin(t) & \rightarrow & L(\vec{y}) = y^{(4)} + 3y\end{aligned}$$

Linear Operator Properties

$$\begin{aligned}L(k\vec{u}) &= kL(\vec{u}), \quad k \in \mathbb{R} \\L(\vec{u} + \vec{w}) &= L(\vec{u}) + L(\vec{w})\end{aligned}$$

Proof

The properties can be proved directly for algebraic operators.

For differential operators, the proof follows from the derivative properties:

- $(kf)' = kf'$
- $(f + g)' = f' + g'$

Superposition Principle for Linear Homogeneous Equations

Let \vec{u}_1 and \vec{u}_2 be any solutions of the *homogeneous linear* equation

$$L(\vec{u}) = 0$$

- The sum $\vec{u} = \vec{u}_1 + \vec{u}_2$ is also a solution.
- For any constant k , $\vec{u} = k\vec{u}_1$ is also a solution.

Superposition Principle for Linear Homogeneous Equations

Let \vec{u}_1 and \vec{u}_2 be any solutions of the *homogeneous linear* equation

$$L(\vec{u}) = 0$$

- The sum $\vec{u} = \vec{u}_1 + \vec{u}_2$ is also a solution.
- For any constant k , $\vec{u} = k\vec{u}_1$ is also a solution.

Proof

The proof of the Superposition Principle follows directly from the properties of linear operators from the previous slides.

$$L(\vec{u}) = L(\vec{u}_1 + \vec{u}_2) = L(\vec{u}_1) + L(\vec{u}_2) = 0 + 0 = 0$$

$$L(\vec{u}) = L(k\vec{u}_1) = kL(\vec{u}_1) = k \cdot 0 = 0$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

$$y'' - 4y = (8e^{2t} + 12e^{-2t}) - 4(2e^{2t} + 3e^{-2t})$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

$$\begin{aligned} y'' - 4y &= (8e^{2t} + 12e^{-2t}) - 4(2e^{2t} + 3e^{-2t}) \\ &= 8e^{2t} + 12e^{-2t} - 8e^{2t} - 12e^{-2t} \end{aligned}$$

Example 4

The point $(1, 3)$ is on the line $y = 3x$. So is the point $(2, 6) = (2 \cdot 1, 2 \cdot 3)$. Additionally, the point $(3, 9) = (1 + 2, 3 + 6) = (1, 3) + (2, 6)$ is on the line.

Example 5

The differential equation

$$y'' - 4y = 0$$

has the solutions $y = e^{2t}$ and $y = e^{-2t}$.

By superposition, $y = 2e^{2t} + 3e^{-2t}$ must also be a solution.

This is easily verified:

$$y' = 4e^{2t} - 6e^{-2t}$$

$$y'' = 8e^{2t} + 12e^{-2t}$$

$$\begin{aligned} y'' - 4y &= (8e^{2t} + 12e^{-2t}) - 4(2e^{2t} + 3e^{-2t}) \\ &= 8e^{2t} + 12e^{-2t} - 8e^{2t} - 12e^{-2t} = 0 \end{aligned}$$

Nonhomogeneous Principle

Let \vec{u}_p be any solution (called a particular solution) to *linear nonhomogeneous* equation

$$L(\vec{u}) = C \quad (\text{algebraic})$$

or

$$L(\vec{u}) = f(t) \quad (\text{differential})$$

Then,

$$\vec{u} = \vec{u}_h + \vec{u}_p$$

is also a solution, here \vec{u}_h is a solution to the **associated homogeneous** equation

$$L(\vec{u}) = 0$$

Furthermore, *every solution of the nonhomogeneous equation must be of the form $\vec{u} = \vec{u}_h + \vec{u}_p$.*

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$L(\vec{u}_q - \vec{u}_p) =$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$L(\vec{u}_q - \vec{u}_p) = L(\vec{u}_q) + L(-\vec{u}_p)$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$\begin{aligned} L(\vec{u}_q - \vec{u}_p) &= L(\vec{u}_q) + L(-\vec{u}_p) \\ &= L(\vec{u}_q) - L(\vec{u}_p) \end{aligned}$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$\begin{aligned} L(\vec{u}_q - \vec{u}_p) &= L(\vec{u}_q) + L(-\vec{u}_p) \\ &= L(\vec{u}_q) - L(\vec{u}_p) \\ &= f(t) - f(t) = 0 \end{aligned}$$

Proof

It is easy to show that $\vec{u} = \vec{u}_h + \vec{u}_p$ is a solution.

$$L(\vec{u}) = L(\vec{u}_h + \vec{u}_p) = L(\vec{u}_h) + L(\vec{u}_p) = 0 + f(t) = f(t)$$

To show that every solution has to be of this form, suppose that \vec{u}_q is any solution. Note that $\vec{u}_q = \vec{u}_p + (\vec{u}_q - \vec{u}_p)$.

We can then show that $\vec{u}_q - \vec{u}_p$ is also a solution to $L(\vec{u}) = 0$:

$$\begin{aligned} L(\vec{u}_q - \vec{u}_p) &= L(\vec{u}_q) + L(-\vec{u}_p) \\ &= L(\vec{u}_q) - L(\vec{u}_p) \\ &= f(t) - f(t) = 0 \end{aligned}$$

Process for Solving Nonhomogeneous Linear Equations

Step 1: Find all solutions \vec{u}_h of $L(\vec{u}) = 0$.

Step 2: Find any solution \vec{u}_p of $L(\vec{u}) = f$.

Step 3: Add $\vec{u}_h + \vec{u}_p = \vec{u}$ to find all solutions of $L(\vec{u}) = f$.

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1:

Step 2:

Step 3:

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1: Solve the associated homogeneous equation $y' - y = 0$, or $y' = y$. (Note: first-order homogeneous linear differential equations are always separable.)

$$y_h = ce^t, \quad \text{for any } c \in \mathbb{R}$$

Step 2:

Step 3:

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1: Solve the associated homogeneous equation $y' - y = 0$, or $y' = y$. (Note: first-order homogeneous linear differential equations are always separable.)

$$y_h = ce^t, \quad \text{for any } c \in \mathbb{R}$$

Step 2: We can verify by differentiation and substitution that $y_p = -t - 1$ is a particular solution.

Step 3:

Example 6

Consider

$$y' - y = t$$

To solve using superposition we need to complete three steps.

Step 1: Solve the associated homogeneous equation $y' - y = 0$, or $y' = y$. (Note: first-order homogeneous linear differential equations are always separable.)

$$y_h = ce^t, \quad \text{for any } c \in \mathbb{R}$$

Step 2: We can verify by differentiation and substitution that $y_p = -t - 1$ is a particular solution.

Step 3: Superposition tells us that


$$y = y_h + y_p = ce^t - t - 1$$

is a solution for any $c \in \mathbb{R}$.

Example 6


Consider

$$y' - y = t$$




$\{y_h\}$

+



y_p

=



$\{y_h\} + y_p$

Euler-Lagrange Two-Stage Method

We have seen that the general solution for the linear first-order differential equation

$$y' + p(t)y = f(t)$$

has the form $y = y_h + y_p$.

Euler-Lagrange Two-Stage Method

We have seen that the general solution for the linear first-order differential equation

$$y' + p(t)y = f(t)$$

has the form $y = y_h + y_p$.

We solved the corresponding homogenous equation using separation of variables, getting a one-parameter family

$$y_h = ce^{-\int p(t)dt}$$

where $c \in \mathbb{R}$.

Euler-Lagrange Two-Stage Method

We have seen that the general solution for the linear first-order differential equation

$$y' + p(t)y = f(t)$$

has the form $y = y_h + y_p$.

We solved the corresponding homogenous equation using separation of variables, getting a one-parameter family

$$y_h = ce^{-\int p(t)dt}$$

where $c \in \mathbb{R}$.

The second step is to find a particular solution, which we will accomplish using **variation of parameters**, which was developed by French mathematician Joseph Louis Lagrange.

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

and change the constant c to a function $v(t)$ and try a solution of the form

$$y_p(t) = v(t)e^{-\int p(t)dt}$$

where the unknown function $v(t)$ is called the **varying parameter**.

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

and change the constant c to a function $v(t)$ and try a solution of the form

$$y_p(t) = v(t)e^{-\int p(t)dt}$$

where the unknown function $v(t)$ is called the **varying parameter**.

Our goal is to find $v(t)$, to do so we need to substitute y_p into the DE.

$$\underbrace{\left(v'(t)e^{-\int p(t)dt} - p(t)v(t)e^{-\int p(t)dt} \right)}_{y_p'} + \underbrace{p(t)v(t)e^{-\int p(t)dt}}_{p(t)y_p} = f(t)$$

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

and change the constant c to a function $v(t)$ and try a solution of the form

$$y_p(t) = v(t)e^{-\int p(t)dt}$$

where the unknown function $v(t)$ is called the **varying parameter**.

Our goal is to find $v(t)$, to do so we need to substitute y_p into the DE.

$$v'(t)e^{-\int p(t)dt} = f(t)$$

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

and change the constant c to a function $v(t)$ and try a solution of the form

$$y_p(t) = v(t)e^{-\int p(t)dt}$$

where the unknown function $v(t)$ is called the **varying parameter**.

Our goal is to find $v(t)$, to do so we need to substitute y_p into the DE.

$$v'(t) = f(t)e^{\int p(t)dt}$$

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

and change the constant c to a function $v(t)$ and try a solution of the form

$$y_p(t) = v(t)e^{-\int p(t)dt}$$

where the unknown function $v(t)$ is called the **varying parameter**.

Our goal is to find $v(t)$, to do so we need to substitute y_p into the DE.

$$v(t) = \int f(t)e^{\int p(t)dt} dt$$

Variation of Parameters

The idea of variation of parameters is to start with

$$y_h(t) = ce^{-\int p(t)dt}$$

and change the constant c to a function $v(t)$ and try a solution of the form

$$y_p(t) = v(t)e^{-\int p(t)dt}$$

where the unknown function $v(t)$ is called the **varying parameter**.

Our goal is to find $v(t)$, to do so we need to substitute y_p into the DE.

$$v(t) = \int f(t)e^{\int p(t)dt} dt$$

Now that we have $v(t)$, we have determined a particular solution.

$$y_p(t) = v(t)e^{-\int p(t)dt} = e^{-\int p(t)dt} \int f(t)e^{\int p(t)dt} dt$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Let us assume for the moment that $y \neq 0$ and use separation of variables.

$$\frac{dy}{y} = -\frac{dt}{t+1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Let us assume for the moment that $y \neq 0$ and use separation of variables.

$$\ln |y| = -\ln(t+1) + c$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Let us assume for the moment that $y \neq 0$ and use separation of variables.

$$|y| = e^{\ln(t+1)^{-1} + c}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Let us assume for the moment that $y \neq 0$ and use separation of variables.

$$|y| = e^c (t+1)^{-1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Let us assume for the moment that $y \neq 0$ and use separation of variables.

$$y_h = \pm \frac{e^c}{t+1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 1. We start by solving the associated homogeneous equation.

$$y' + \left(\frac{1}{t+1} \right) y = 0$$

Let us assume for the moment that $y \neq 0$ and use separation of variables.

$$y_h = \frac{k}{t+1}$$

where $k = \pm e^c$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 2. Next, using variation of parameters, we try

$$y_p = \frac{v(t)}{t+1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 2. Next, using variation of parameters, we try

$$y_p = \frac{v(t)}{t+1}$$

which gives:

$$v'(t)e^{-\int p(t)dt} = f(t)$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 2. Next, using variation of parameters, we try

$$y_p = \frac{v(t)}{t+1}$$

which gives:

$$\frac{v'(t)}{t+1} = 2$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 2. Next, using variation of parameters, we try

$$y_p = \frac{v(t)}{t+1}$$

which gives:

$$v'(t) = 2t + 2$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 2. Next, using variation of parameters, we try

$$y_p = \frac{v(t)}{t+1}$$

which gives:

$$v(t) = t^2 + 2t + c$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 2. Next, using variation of parameters, we try

$$y_p = \frac{v(t)}{t+1}$$

which gives:

$$v(t) = t^2 + 2t + c$$

But, we only need a single $v(t)$, so we can let $c = 0$, giving

$$y_p = \frac{t^2 + 2t}{t+1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 3. Thus, the general solution is:

$$y(t) = y_h + y_p = \frac{k}{t+1} + \frac{t^2 + 2t}{t+1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 3. Thus, the general solution is:

$$y(t) = y_h + y_p = \frac{k}{t+1} + \frac{t^2 + 2t}{t+1}$$

Step 4. Substituting the initial condition into the general solution gives:

$$0 = y(0) = \frac{k}{(0)+1} + \frac{(0)^2 + 2(0)}{(0)+1}$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 3. Thus, the general solution is:

$$y(t) = y_h + y_p = \frac{k}{t+1} + \frac{t^2 + 2t}{t+1}$$

Step 4. Substituting the initial condition into the general solution gives:

$$0 = y(0) = \frac{k}{(0)+1} + \frac{(0)^2 + 2(0)}{(0)+1} \Rightarrow k = 0$$

Example 7

Consider the IVP

$$y' + \left(\frac{1}{t+1} \right) y = 2, \quad y(0) = 0, \quad t \geq 0$$

Step 3. Thus, the general solution is:

$$y(t) = y_h + y_p = \frac{k}{t+1} + \frac{t^2 + 2t}{t+1}$$

Step 4. Substituting the initial condition into the general solution gives:

$$0 = y(0) = \frac{k}{(0)+1} + \frac{(0)^2 + 2(0)}{(0)+1} \Rightarrow k = 0$$

Which means that the solution to the IVP is

$$y(t) = \frac{t^2 + 2t}{t+1}$$

Euler-Lagrange Method for Solving Linear First-Order DEs

To solve a linear differential equation

$$y' + p(t)y = f(t)$$

where p and f are continuous on a domain I , use the following steps.

Step 1. Solve the corresponding homogenous equation $y' + p(t)y = 0$ to obtain the one-parameter family.

$$y_h = ce^{-\int p(t)dt}$$

Step 2. Solve

$$v'(t)e^{-\int p(t)dt} = f(t)$$

for $v(t)$ to obtain a particular solution $y_p = v(t)e^{-\int p(t)dt}$.

Step 3. Combine the results of Step 1 and Step 2 to form the general solution

$$y(t) = y_h + y_p$$

Step 4. If you are solving an IVP, only after Step 3 can you plug in the initial condition.

Note

Variation of Parameters is a very powerful method, and you will see it again in a proper differential equations course. But, for first-order (and *only* first-order) equations we have a second method, called the **Integrating Factor Method** which may also be used.

Note

Variation of Parameters is a very powerful method, and you will see it again in a proper differential equations course. But, for first-order (and *only* first-order) equations we have a second method, called the **Integrating Factor Method** which may also be used.

For the differential equation

$$y' + p(t)y = f(t)$$

we will break this new method down into two cases:

- $p(t)$ is constant.
- $p(t)$ is variable.

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

Let us start with the differential equation.

$$y' + ay = f(t)$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

We first multiply both sides of the equation by e^{at} .

$$e^{at} (y' + ay) = e^{at} f(t)$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

We then apply Euler's observation to the left-hand side.

$$\frac{d}{dt} (e^{at} y) = e^{at} f(t)$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

Next we integrate both sides.

$$e^{at} y = \int e^{at} f(t) dt + c$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

Solving for y gives:

$$y(t) = e^{-at} \int e^{at} f(t) dt + ce^{-at}$$

Integrating Factor Method (Constant Coefficient)

Let us look at the first-order linear differential equation

$$y' + ay = f(t), \quad a \in \mathbb{R}$$

This method uses a simple observation made by Euler:

$$e^{at} (y' + ay) = \frac{d}{dt} (e^{at} y)$$

Solving for y gives:

$$y(t) = e^{-at} \int e^{at} f(t) dt + ce^{-at}$$

Note

This is the same answer we got from Variation of Parameters, though achieved through a different route. We have obtained both y_h and y_p at the same time.

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We seek a function $\mu(t)$ that satisfies Euler's observation, i.e.

$$\mu(t) \cdot (y' + p(t)y) = \frac{d}{dt} (\mu(t) \cdot y)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

Let us carry out the differentiation on the right-hand side

$$\mu(t)y' + p(t)\mu(t)y = \mu'(t)y + \mu(t)y'$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

If we assume $y(t) \neq 0$, this simplifies to

$$\mu'(t) = p(t)\mu(t)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\frac{\mu'(t)}{\mu(t)} = p(t)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\ln |\mu(t)| = \int p(t) dt$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

We now know the integrating factor, and perform the same steps as before.

$$y' + p(t)y = f(t)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

Multiply both sides by the integrating factor.

$$\mu(t) \cdot (y' + p(t)y) = \mu(t) \cdot f(t)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

Apply the property $\mu(t) \cdot (y' + p(t)y) = (\mu(t) \cdot y)'$ to the left-hand side.

$$(\mu(t)y)' = \mu(t)f(t)$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

Integrate both sides.

$$\mu(t)y(t) = \int \mu(t)f(t)dt + c$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

Assuming $\mu(t) \neq 0$, we can solve for y .

$$y(t) = \frac{1}{\mu(t)} \int \mu(t)f(t)dt + \frac{c}{\mu(t)}$$

Integrating Factor Method (Variable Coefficient)

Now let us look at the more general first-order differential equation

$$y' + p(t)y = f(t)$$

We can find a solution $\mu(t) > 0$ by Separation of Variables.

$$\mu(t) = e^{\int p(t)dt}$$

Assuming $\mu(t) \neq 0$, we can solve for y .

$$y(t) = \frac{1}{\mu(t)} \int \mu(t)f(t)dt + \frac{c}{\mu(t)}$$

Note

We have again found y_h and y_p at the same time.

Integrating Factor Method for First-Order Linear DEs

To solve the linear first-order DE, where p and f are continuous on a domain I .

$$y' + p(t)y = f(t)$$

Step 1. Find the integrating factor $\mu(t) = e^{\int p(t)dt}$, where $\int p(t)dt$ represents *any* anti-derivative of $p(t)$.

Step 2. Multiply both sides of the DE by $\mu(t)$, which always simplifies to:

$$\left(e^{\int p(t)dt} y(t) \right)' = e^{\int p(t)dt} f(t)$$

Step 3. Find the anti-derivative to get:

$$e^{\int p(t)dt} y(t) = \int e^{\int p(t)dt} f(t) dt + c$$

Step 4. Solve algebraically for y .

$$y = e^{-\int p(t)dt} \int e^{\int p(t)dt} f(t) dt + ce^{-\int p(t)dt}$$

Step 5. For IVPs, substitute the initial conditions in to find c .

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 1. Find the integrating factor:

$$\mu(t) = e^{\int p(t)dt}$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 1. Find the integrating factor:

$$\mu(t) = e^{\int (-1)dt}$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 1. Find the integrating factor:

$$\mu(t) = e^{\int (-1)dt} = e^{-t}$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 1. Find the integrating factor:

$$\mu(t) = e^{\int (-1)dt} = e^{-t}$$

Step 2. Multiply both sides of the DE by $\mu(t)$:

$$e^{-t} (y' - y) = e^{-t}$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 1. Find the integrating factor:

$$\mu(t) = e^{\int (-1)dt} = e^{-t}$$

Step 2. Multiply both sides of the DE by $\mu(t)$:

$$e^{-t} (y' - y) = e^{-t}$$

Which reduces to:

$$(e^{-t}y)' = te^{-t}$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt = e^{-t}(-t - 1) + c$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt = e^{-t}(-t - 1) + c$$

Step 4. Solve for y :

$$y(t) = e^t (e^{-t}) (-t - 1) + ce^t$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt = e^{-t}(-t - 1) + c$$

Step 4. Solve for y :

$$y(t) = e^t (e^{-t}) (-t - 1) + ce^t = -t - 1 + ce^t$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt = e^{-t}(-t - 1) + c$$

Step 4. Solve for y :

$$y(t) = e^t (e^{-t}) (-t - 1) + ce^t = -t - 1 + ce^t$$

Step 5. Plug in the initial conditions to find the solution to the IVP:

$$1 = y(0) = -0 - 1 + ce^0$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt = e^{-t}(-t - 1) + c$$

Step 4. Solve for y :

$$y(t) = e^t (e^{-t}) (-t - 1) + ce^t = -t - 1 + ce^t$$

Step 5. Plug in the initial conditions to find the solution to the IVP:

$$1 = y(0) = -0 - 1 + ce^0 \Rightarrow c = 2$$

Example 8

Consider the IVP

$$y' - y = t, \quad y(0) = 1$$

Let us solve this DE using the Integrating Factor method.

Step 3. Find the antiderivative:

$$e^{-t}y = \int te^{-t}dt = e^{-t}(-t - 1) + c$$

Step 4. Solve for y :

$$y(t) = e^t (e^{-t}) (-t - 1) + ce^t = -t - 1 + ce^t$$

Step 5. Plug in the initial conditions to find the solution to the IVP:

$$1 = y(0) = -0 - 1 + ce^0 \Rightarrow c = 2$$

Thus, the solution to the IVP is $y(t) = -t - 1 + 2e^t$