

# Properties of Linear Transformations

Adam Wilson


Salt Lake Community College

## Injectivity


A function  $f : \mathbb{X} \rightarrow \mathbb{Y}$  is **one-to-one**, or **injective**, provided that  $f(u) = f(v)$  implies  $u = v$ .

## Injectivity

A function  $f : \mathbb{X} \rightarrow \mathbb{Y}$  is **one-to-one**, or **injective**, provided that  $f(u) = f(v)$  implies  $u = v$ .



(a)  $f(x) = x^3$  is injective




(b)  $g(x) = x^2$  is not injective

## Surjectivity


The set of output values of a function  $f : \mathbb{X} \rightarrow \mathbb{Y}$  is a subset of the codomain  $\mathbb{Y}$  and is called the **image** of the function. If the image is all of  $\mathbb{Y}$ , the function  $f$  is said to map **onto**  $\mathbb{Y}$ , or to be **surjective**.

## Surjectivity

The set of output values of a function  $f : \mathbb{X} \rightarrow \mathbb{Y}$  is a subset of the codomain  $\mathbb{Y}$  and is called the **image** of the function. If the image is all of  $\mathbb{Y}$ , the function  $f$  is said to map **onto**  $\mathbb{Y}$ , or to be **surjective**.



(a)  $f(x) = -x$  is surjective



(b)  $g(x) = |x|$  is not surjective

## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix}$$

## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of  $T$ ?



## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of  $T$ ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of  $T$ ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \text{ are independent.}$$

So, **Col**  $\mathbf{A}$ , the column space of  $\mathbf{A}$ , is the image of  $T$ .

## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of  $T$ ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \text{ are independent.}$$

So, **Col**  $\mathbf{A}$ , the column space of  $\mathbf{A}$ , is the image of  $T$ .

We can derive the equation of this plane by looking at the system

$$u_1 = v_1 + v_2, \quad u_2 = v_1 - v_2, \quad u_3 = 2v_1 + v_2$$

## Example

The linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \Rightarrow \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 \\ v_1 - v_2 \\ 2v_1 + v_2 \end{bmatrix}$$

What is the image of  $T$ ?

$$\vec{u} = v_1 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + v_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \text{ and } \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \text{ are independent.}$$

So, **Col A**, the column space of  $\mathbf{A}$ , is the image of  $T$ .

We can derive the equation of this plane by looking at the system

$$u_1 = v_1 + v_2, \quad u_2 = v_1 - v_2, \quad u_3 = 2v_1 + v_2$$

Eliminating  $v_1$  and  $v_2$  gives us

$$3u_1 + u_2 - 2u_3 = 0$$

## Image Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with image  $\mathbf{Im}(T)$ . Then:

## Image Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with image  $\mathbf{Im}(T)$ . Then:

- $\mathbf{Im}(T)$  is a subspace of  $\mathbb{W}$ .

## Image Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with image  $\mathbf{Im}(T)$ . Then:

- $\mathbf{Im}(T)$  is a subspace of  $\mathbb{W}$ .
- $T$  is surjective if and only if  $\mathbf{Im}(T) = \mathbb{W}$ .

## Image Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with image  $\mathbf{Im}(T)$ . Then:

- $\mathbf{Im}(T)$  is a subspace of  $\mathbb{W}$ .
- $T$  is surjective if and only if  $\mathbf{Im}(T) = \mathbb{W}$ .

## Rank of a Linear Transformation

The dimension of the image of a linear transformation  $T$  is called its **rank**

$$\mathbf{rank}(T) = \mathbf{dim}(\mathbf{Im}(T))$$



## Example

For  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

## Example

For  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

## Example

For  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Which means

$$\text{Im}(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \end{bmatrix} \right\}$$

which is a subset of  $\mathbb{R}^2$ .

## Example

For  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Which means

$$\text{Im}(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \end{bmatrix} \right\}$$

which is a subset of  $\mathbb{R}^2$ . To determine the dimension we need to determine how many pivot columns  $\mathbf{A}$  has:

$$\begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

## Example

For  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \vec{w}$$

We can write

$$\vec{w} = v_1 \begin{bmatrix} 2 \\ -1 \end{bmatrix} + v_2 \begin{bmatrix} -4 \\ 2 \end{bmatrix} + v_3 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + v_4 \begin{bmatrix} 6 \\ -3 \end{bmatrix}$$

Which means

$$\text{Im}(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \begin{bmatrix} 6 \\ -3 \end{bmatrix} \right\}$$

which is a subset of  $\mathbb{R}^2$ . To determine the dimension we need to determine how many pivot columns  $\mathbf{A}$  has:

$$\begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 0 & 3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Thus,  $\text{rank}(T) = \text{dim}(\text{Im}(T)) = \text{dim}(\text{Col } \mathbf{A}) = 2$ .

## Rank of a Matrix Multiplication Operator

For any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T(\vec{x}) = \mathbf{A}\vec{x}$ , where  $\mathbf{A} \in \mathbb{M}_{mn}$  and  $\vec{x} \in \mathbb{V}$ , the image of  $T$  is the column space of  $A$ . (That's is,  $\mathbf{Im}(T) = \mathbf{Col} \mathbf{A}$ .)

The pivot columns of  $A$  form a basis for  $\mathbf{Im}(T)$ .

Consequently,

$$\begin{aligned}\mathbf{rank}(T) &= \mathbf{dim}(\mathbf{Im}(T)) \\ &= \mathbf{dim}(\mathbf{Col} \mathbf{A}) \\ &= \text{The number of pivot columns in } \mathbf{A}.\end{aligned}$$

## Rank of a Matrix Multiplication Operator

For any linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T(\vec{x}) = \mathbf{A}\vec{x}$ , where  $\mathbf{A} \in \mathbb{M}_{mn}$  and  $\vec{x} \in \mathbb{V}$ , the image of  $T$  is the column space of  $A$ . (That's is,  $\mathbf{Im}(T) = \mathbf{Col} \mathbf{A}$ .)

The pivot columns of  $A$  form a basis for  $\mathbf{Im}(T)$ .

Consequently,

$$\begin{aligned}\mathbf{rank}(T) &= \mathbf{dim}(\mathbf{Im}(T)) \\ &= \mathbf{dim}(\mathbf{Col} \mathbf{A}) \\ &= \text{The number of pivot columns in } \mathbf{A}.\end{aligned}$$

## Reminder

The basis of  $\mathbf{Col} \mathbf{A}$  must come from  $\mathbf{A}$ , *not* from the RREF of  $\mathbf{A}$ .

Recall that a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$  must map the zero vector of  $\mathbb{V}$  to the zero vector of  $\mathbb{W}$ .



Recall that a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$  must map the zero vector of  $\mathbb{V}$  to the zero vector of  $\mathbb{W}$ .

## Kernel of a Linear Transformation

The **kernel** (or **nullspace**) of a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$ , denoted  $\ker(T)$ , is the set of vectors in  $\mathbb{V}$  that are mapped by  $T$  to the zero vector of  $\mathbb{W}$ .

$$\ker(T) = \left\{ \vec{v} \in \mathbb{V} \mid T(\vec{v}) = \vec{0} \right\}$$

Recall that a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$  must map the zero vector of  $\mathbb{V}$  to the zero vector of  $\mathbb{W}$ .

## Kernel of a Linear Transformation

The **kernel** (or **nullspace**) of a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$ , denoted  $\ker(T)$ , is the set of vectors in  $\mathbb{V}$  that are mapped by  $T$  to the zero vector of  $\mathbb{W}$ .

$$\ker(T) = \left\{ \vec{v} \in \mathbb{V} \mid T(\vec{v}) = \vec{0} \right\}$$

### Example

Consider the projection  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by

$$T(x, y, z) = (x, y, 0)$$

What is the kernel of  $T$ ?

Recall that a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$  must map the zero vector of  $\mathbb{V}$  to the zero vector of  $\mathbb{W}$ .

## Kernel of a Linear Transformation

The **kernel** (or **nullspace**) of a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$ , denoted  $\ker(T)$ , is the set of vectors in  $\mathbb{V}$  that are mapped by  $T$  to the zero vector of  $\mathbb{W}$ .

$$\ker(T) = \left\{ \vec{v} \in \mathbb{V} \mid T(\vec{v}) = \vec{0} \right\}$$

### Example

Consider the projection  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by

$$T(x, y, z) = (x, y, 0)$$

What is the kernel of  $T$ ?

$$\ker(T) = \{(0, 0, z) \mid z \in \mathbb{R}\}$$

## Example

Consider the transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

## Example

Consider the transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to  $\vec{0}$ , we have to solve the system:

$$\begin{array}{rcrcrcrcrcrcl} v_1 & + & v_2 & + & 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + & 5v_3 & = & 0 \end{array}$$

## Example

Consider the transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to  $\vec{0}$ , we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to  $\vec{0}$ , we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to  $\vec{0}$ , we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

So, we see that  $v_3$  is a free variable and if  $v_3 = s$  is a parameter, we have  $v_1 = -s$ ,  $v_2 = -s$ , and  $v_3 = s$ .



## Example

Consider the transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{v}) = \mathbf{A}\vec{v} = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} v_1 + v_2 + 2v_3 \\ 2v_1 + 3v_2 + 5v_3 \end{bmatrix}$$

To find the vectors that are mapped to  $\vec{0}$ , we have to solve the system:

$$\begin{array}{rrcr} v_1 & + & v_2 & + 2v_3 & = & 0 \\ 2v_1 & + & 3v_2 & + 5v_3 & = & 0 \end{array} \rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 2 & 0 \\ 2 & 3 & 5 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

So, we see that  $v_3$  is a free variable and if  $v_3 = s$  is a parameter, we have  $v_1 = -s$ ,  $v_2 = -s$ , and  $v_3 = s$ .

Thus, the kernel is any scalar multiple of  $\langle -1, -1, 1 \rangle$ :

$$\ker(T) = \text{span} \left\{ \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix} \right\}$$

### Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right]$$

Thus,

$$\ker(T) = \{\vec{0}\}$$

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$



## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & | & 0 \\ 1 & 1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

## Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix}$$

What is the kernel of  $T$ ?

To find the kernel we need to solve  $A\vec{v} = \vec{0}$ :

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cc|c} 1 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{array} \right]$$

In this case we have  $v_1 = -\frac{1}{2}v_2$  and so, if we let our parameter be  $v_2 = s$  we have

$$\ker(T) = \left\{ s \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right\}$$

### Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

### Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

It should be clear that this transformation maps all vectors  $\vec{v} \in \mathbb{R}^2$  to  $\vec{0}$ .

### Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

It should be clear that this transformation maps all vectors  $\vec{v} \in \mathbb{R}^2$  to  $\vec{0}$ .

Thus,

$$\ker(T) = \mathbb{R}^2$$

### Example

Consider the transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by the matrix

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

It should be clear that this transformation maps all vectors  $\vec{v} \in \mathbb{R}^2$  to  $\vec{0}$ .

Thus,

$$\ker(T) = \mathbb{R}^2$$

These examples seem to suggest that the kernel of the linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$  is a subspace of  $\mathbb{W}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\mathbf{ker}(T)$ .

Then,

- 1  $\mathbf{ker}(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\mathbf{ker}(T) = \{\vec{0}\}$ .



## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- 1  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify 1, so let us prove 2.

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- 1  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify 1, so let us prove 2.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- ①  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- ②  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify ①, so let us prove ②.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

Then, for  $\vec{w} \in \ker(T)$ , we have  $T(\vec{w}) = \vec{0}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- 1  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify 1, so let us prove 2.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

Then, for  $\vec{w} \in \ker(T)$ , we have  $T(\vec{w}) = \vec{0}$ .

Which means  $T(\vec{w}) = T(\vec{0})$  and hence  $\vec{w} = \vec{0}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- ①  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- ②  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify ①, so let us prove ②.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

Then, for  $\vec{w} \in \ker(T)$ , we have  $T(\vec{w}) = \vec{0}$ .

Which means  $T(\vec{w}) = T(\vec{0})$  and hence  $\vec{w} = \vec{0}$ . Thus,  $\ker(T) = \{\vec{0}\}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\mathbf{ker}(T)$ .

Then,

- 1  $\mathbf{ker}(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\mathbf{ker}(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify 1, so let us prove 2.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

Then, for  $\vec{w} \in \mathbf{ker}(T)$ , we have  $T(\vec{w}) = \vec{0}$ .

Which means  $T(\vec{w}) = T(\vec{0})$  and hence  $\vec{w} = \vec{0}$ . Thus,  $\mathbf{ker}(T) = \{\vec{0}\}$ .

Now, suppose that we know  $\mathbf{ker}(T) = \{\vec{0}\}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- 1  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify 1, so let us prove 2.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

Then, for  $\vec{w} \in \ker(T)$ , we have  $T(\vec{w}) = \vec{0}$ .

Which means  $T(\vec{w}) = T(\vec{0})$  and hence  $\vec{w} = \vec{0}$ . Thus,  $\ker(T) = \{\vec{0}\}$ .

Now, suppose that we know  $\ker(T) = \{\vec{0}\}$ .

So  $T(\vec{u}) = T(\vec{v})$  implies  $T(\vec{u}) - T(\vec{v}) = T(\vec{u} - \vec{v}) = T(\vec{0}) = \vec{0}$ .

## Kernel Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from vector space  $\mathbb{V}$  to vector space  $\mathbb{W}$  with kernel  $\ker(T)$ .

Then,

- 1  $\ker(T)$  is a subspace of  $\mathbb{V}$ .
- 2  $T$  is injective if and only if  $\ker(T) = \{\vec{0}\}$ .

## Proof

It is straightforward to verify 1, so let us prove 2.

Suppose  $T$  is injective, which means that  $T(\vec{u}) = T(\vec{v})$  implies  $\vec{u} = \vec{v}$ .

Then, for  $\vec{w} \in \ker(T)$ , we have  $T(\vec{w}) = \vec{0}$ .

Which means  $T(\vec{w}) = T(\vec{0})$  and hence  $\vec{w} = \vec{0}$ . Thus,  $\ker(T) = \{\vec{0}\}$ .

Now, suppose that we know  $\ker(T) = \{\vec{0}\}$ .

So  $T(\vec{u}) = T(\vec{v})$  implies  $T(\vec{u}) - T(\vec{v}) = T(\vec{u} - \vec{v}) = T(\vec{0}) = \vec{0}$ .

Thus,  $\vec{u} - \vec{v}$  is in the kernel, which means  $\vec{u} - \vec{v} = \vec{0}$  and thus  $T$  is injective.



## Example

Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$ , where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

## Example

Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$ , where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[ \begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$ , where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[ \begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

## Example

Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$ , where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[ \begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

So, we see that  $v_1 = 2v_2 - 3v_4$  and  $v_3 = 0$ . If we let  $v_2 = r$  and  $v_4 = s$ ,

$$\vec{v} = \begin{bmatrix} 2r - 3s \\ r + 0s \\ 0r + 0s \\ 0r + s \end{bmatrix} = r \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

## Example

Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$ , where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[ \begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

So, we see that  $v_1 = 2v_2 - 3v_4$  and  $v_3 = 0$ . If we let  $v_2 = r$  and  $v_4 = s$ ,

$$\vec{v} = \begin{bmatrix} 2r - 3s \\ r + 0s \\ 0r + 0s \\ 0r + s \end{bmatrix} = r \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \rightarrow \ker(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

The dimension of the kernel of  $T$  is 2.

## Example

Consider the transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  defined by  $T(\vec{v}) = \mathbf{A}\vec{v}$ , where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 & 3 & 6 \\ -1 & 2 & -2 & -3 \end{bmatrix}$$

We can determine the kernel by solving the system:

$$\left[ \begin{array}{cccc|c} 2 & -4 & 3 & 6 & 0 \\ -1 & 2 & -2 & -3 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{array} \right]$$

So, we see that  $v_1 = 2v_2 - 3v_4$  and  $v_3 = 0$ . If we let  $v_2 = r$  and  $v_4 = s$ ,

$$\vec{v} = \begin{bmatrix} 2r - 3s \\ r + 0s \\ 0r + 0s \\ 0r + s \end{bmatrix} = r \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \rightarrow \ker(T) = \text{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

The dimension of the kernel of  $T$  is 2.

(Remember that the dimension of the image of  $T$  was 2.)

## Dimension Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from a finite vector space  $\mathbb{V}$ .

Then

$$\mathbf{dim}(\ker(T)) + \mathbf{dim}(\mathbf{Im}(T)) = \mathbf{dim}(\mathbb{V})$$

## Dimension Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from a finite vector space  $\mathbb{V}$ .

Then

$$\mathbf{dim}(\ker(T)) + \mathbf{dim}(\mathbf{Im}(T)) = \mathbf{dim}(\mathbb{V})$$

## Example

From earlier examples:



## Dimension Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from a finite vector space  $\mathbb{V}$ .

Then

$$\mathbf{dim}(\ker(T)) + \mathbf{dim}(\mathbf{Im}(T)) = \mathbf{dim}(\mathbb{V})$$

## Example

From earlier examples:

- $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by  $T(x, y, z) = (x, y, 0)$  had

$$\mathbf{dim}(\ker(T)) + \mathbf{dim}(\mathbf{Im}(T)) = 1 + 2 = 3 = \mathbf{dim}(\mathbb{R}^3)$$

## Dimension Theorem

Let  $T : \mathbb{V} \rightarrow \mathbb{W}$  be a linear transformation from a finite vector space  $\mathbb{V}$ .

Then

$$\mathbf{\dim(\ker(T)) + \dim(\text{Im}(T)) = \dim(\mathbb{V})}$$

## Example

From earlier examples:

- $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by  $T(x, y, z) = (x, y, 0)$  had

$$\mathbf{\dim(\ker(T)) + \dim(\text{Im}(T)) = 1 + 2 = 3 = \dim(\mathbb{R}^3)}$$

- $D_2 : \mathbb{P}_3 \rightarrow \mathbb{P}_1$  had

$$\mathbf{\ker(D_2) = \{cx + d\} \quad \text{and} \quad \text{Im}(D_2) = \{6ax + 2b\}}$$

$$\mathbf{\dim(\ker(D_2)) + \dim(\text{Im}(D_2)) = 2 + 2 = 4 = \dim(\mathbb{P}_3)}$$