

Eigenvalues and Eigenvectors

Department of Mathematics

Salt Lake Community College

(Slides by Adam Wilson)

Example 1

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$


Example 1

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

We can see how T maps a few vectors:

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \longrightarrow$$




Example 1

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

We can see how T maps a few vectors:

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ -2 \end{bmatrix} \quad \longrightarrow$$




Example 1

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

We can see how T maps a few vectors:

$$T\left(\begin{bmatrix} -1 \\ -1 \end{bmatrix}\right) = \begin{bmatrix} -3 \\ 0 \end{bmatrix} \quad \longrightarrow$$




Example 1

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

But something interesting happens for some special vectors.

$$T\left(\begin{bmatrix} 2 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 2 \end{bmatrix} = 2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \longrightarrow$$




Example 1

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(\vec{u}) = \mathbf{A}\vec{u}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}$$

But something interesting happens for some special vectors.

$$T\left(\begin{bmatrix} 1 \\ -2 \end{bmatrix}\right) = \begin{bmatrix} -3 \\ 6 \end{bmatrix} = -3 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \rightarrow$$



Eigenvalues and Eigenvectors

Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation from vector space \mathbb{V} into itself. A scalar λ is a **eigenvalue** of T if there is a *nonzero* vector $\vec{v} \in \mathbb{V}$ such that

$$T(\vec{v}) = \lambda \vec{v}$$

Such a nonzero vector \vec{v} is called an **eigenvector** of T corresponding to λ .

Eigenvalues and Eigenvectors

Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation from vector space \mathbb{V} into itself. A scalar λ is a **eigenvalue** of T if there is a *nonzero* vector $\vec{v} \in \mathbb{V}$ such that

$$T(\vec{v}) = \lambda \vec{v}$$

Such a nonzero vector \vec{v} is called an **eigenvector** of T corresponding to λ .

Note

Eigenvalues are also called **proper values** or **characteristic values**.

Eigenvalues and Eigenvectors

Let $T : \mathbb{V} \rightarrow \mathbb{V}$ be a linear transformation from vector space \mathbb{V} into itself. A scalar λ is a **eigenvalue** of T if there is a *nonzero* vector $\vec{v} \in \mathbb{V}$ such that

$$T(\vec{v}) = \lambda \vec{v}$$

Such a nonzero vector \vec{v} is called an **eigenvector** of T corresponding to λ .

Note

Eigenvalues are also called **proper values** or **characteristic values**.

Note

If the linear transformation T is represented by an $n \times n$ matrix \mathbf{A} , where $\mathbb{V} = \mathbb{R}^n$ and $T(\vec{v}) = \mathbf{A}\vec{v}$, then λ and \vec{v} are characterized by the equation

$$\mathbf{A}\vec{v} = \lambda \vec{v}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

While this equation always has the trivial solution $\vec{v} = \vec{0}$, we are looking for any non-zero solutions. Therefore, we are looking for when

$$|\mathbf{A} - \lambda\mathbf{I}_n| = 0$$

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

While this equation always has the trivial solution $\vec{v} = \vec{0}$, we are looking for any non-zero solutions. Therefore, we are looking for when

$$|\mathbf{A} - \lambda\mathbf{I}_n| = 0$$

This is called the **characteristic equation** of matrix A .

Computing Eigenvalues and Eigenvectors

If \mathbf{A} is a $n \times n$ matrix, and \mathbf{I}_n is the $n \times n$ identity matrix, then

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

$$\mathbf{A}\vec{v} = \lambda\mathbf{I}_n\vec{v}$$

$$(\mathbf{A} - \lambda\mathbf{I}_n)\vec{v} = \vec{0}$$

While this equation always has the trivial solution $\vec{v} = \vec{0}$, we are looking for any non-zero solutions. Therefore, we are looking for when

$$|\mathbf{A} - \lambda\mathbf{I}_n| = 0$$

This is called the **characteristic equation** of matrix \mathbf{A} .

The polynomial in λ denoted by

$$p(\lambda) = |\mathbf{A} - \lambda\mathbf{I}_n|$$

is called the **characteristic polynomial** of \mathbf{A} .

Summary of Steps for Finding Eigenvalues and Eigenvectors

- 1 Write the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}_n| = 0$.
- 2 Solve the characteristic equation for λ .
- 3 For each eigenvalue λ_i , find the corresponding eigenvector \vec{v}_i by solving the system of equations

$$(\mathbf{A} - \lambda_i \mathbf{I}_n) \vec{v}_i = \vec{0}$$

Summary of Steps for Finding Eigenvalues and Eigenvectors

- 1 Write the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}_n| = 0$.
- 2 Solve the characteristic equation for λ .
- 3 For each eigenvalue λ_i , find the corresponding eigenvector \vec{v}_i by solving the system of equations

$$(\mathbf{A} - \lambda_i \mathbf{I}_n) \vec{v}_i = \vec{0}$$

Note

Eigenvectors are *not* unique. An eigenvector is just a direction, any nonzero multiple of \vec{v}_i works just as well.

Summary of Steps for Finding Eigenvalues and Eigenvectors

- 1 Write the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}_n| = 0$.
- 2 Solve the characteristic equation for λ .
- 3 For each eigenvalue λ_i , find the corresponding eigenvector \vec{v}_i by solving the system of equations

$$(\mathbf{A} - \lambda_i \mathbf{I}_n) \vec{v}_i = \vec{0}$$

Note

Eigenvectors are *not* unique. An eigenvector is just a direction, any nonzero multiple of \vec{v}_i works just as well.

Note

For large matrices these steps become cumbersome, so computer algebra systems are often employed.

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0$$
$$\rightarrow (1-\lambda)(-2-\lambda) - 4 = 0$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0$$
$$\rightarrow (1-\lambda)(-2-\lambda) - 4 = 0 \rightarrow \lambda^2 + \lambda - 6 = 0$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \end{aligned}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 \\ 2 & -4 \end{bmatrix} \begin{array}{c} 0 \\ 0 \end{array}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \left[\begin{array}{cc|c} -1 & 2 & 0 \\ 2 & -4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 1 & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 4 & 2 & | & 0 \\ 2 & 1 & | & 0 \end{bmatrix}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 4 & 2 & | & 0 \\ 2 & 1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Example 2

In our first example we saw two eigenvectors, let us verify these using the characteristic equation.

$$\begin{aligned} \left| \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| &= 0 \rightarrow \begin{vmatrix} 1-\lambda & 2 \\ 2 & -2-\lambda \end{vmatrix} = 0 \\ \rightarrow (1-\lambda)(-2-\lambda) - 4 &= 0 \rightarrow \lambda^2 + \lambda - 6 = 0 \\ \rightarrow (\lambda - 2)(\lambda + 3) &= 0 \rightarrow \lambda_1 = 2 \quad \text{and} \quad \lambda_2 = -3 \end{aligned}$$

To find the eigenvector for λ_1 we need to solve:

$$\begin{bmatrix} 1-(2) & 2 \\ 2 & -2-(2) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -1 & 2 & | & 0 \\ 2 & -4 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve:

$$\begin{bmatrix} 1-(-3) & 2 \\ 2 & -2-(-3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 4 & 2 & | & 0 \\ 2 & 1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_2 = s \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \left[\begin{array}{cc|c} -2 & 1 & 0 \\ 4 & -2 & 0 \end{array} \right]$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \quad \rightarrow \quad \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \quad \rightarrow \quad (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 2 & 1 & | & 0 \\ 4 & 2 & | & 0 \end{bmatrix}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 2 & 1 & | & 0 \\ 4 & 2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Example 3

Let us find the eigenvalues and eigenvectors for the matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}_2| = 0 \rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{vmatrix} = 0 \rightarrow (1 - \lambda)^2 - 4 = 0$$

Which has solutions $\lambda_1 = 3$ and $\lambda_2 = -1$.

To find the eigenvector for λ_1 we need to solve

$$\begin{bmatrix} 1 - (3) & 1 \\ 4 & 1 - (3) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} -2 & 1 & | & 0 \\ 4 & -2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -\frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_1 = s \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

To find the eigenvector for λ_2 we need to solve

$$\begin{bmatrix} 1 - (-1) & 1 \\ 4 & 1 - (-1) \end{bmatrix} \vec{v} = \vec{0} \rightarrow \begin{bmatrix} 2 & 1 & | & 0 \\ 4 & 2 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \rightarrow \vec{v}_2 = s \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Which simplifies to:

$$\lambda^3 - 2\lambda^2 - \lambda + 2 = 0$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Which simplifies to:

$$\begin{aligned} \lambda^3 - 2\lambda^2 - \lambda + 2 &= 0 \\ (\lambda - 2)(\lambda - 1)(\lambda + 1) &= 0 \end{aligned}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

The characteristic equation is:

$$|\mathbf{A} - \lambda \mathbf{I}_2| = \begin{vmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{vmatrix} = 0$$

Which simplifies to:

$$\begin{aligned} \lambda^3 - 2\lambda^2 - \lambda + 2 &= 0 \\ (\lambda - 2)(\lambda - 1)(\lambda + 1) &= 0 \end{aligned}$$

So, the eigenvalues are $\lambda_1 = 2$, $\lambda_2 = 1$, and $\lambda_3 = -1$.

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\begin{bmatrix} 1 - (2) & 1 & -2 \\ -1 & 2 - (2) & 1 \\ 0 & 1 & -1 - (2) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\begin{bmatrix} -1 & 1 & -2 \\ -1 & 0 & 1 \\ 0 & 1 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\left[\begin{array}{ccc|c} -1 & 1 & -2 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 1 & -3 & 0 \end{array} \right]$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 2$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = v_3$ and $v_2 = 3v_3$. Replacing v_3 with parameter s gives

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - (1) & 1 & -2 \\ -1 & 2 - (1) & 1 \\ 0 & 1 & -1 - (1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\begin{bmatrix} 0 & 1 & -2 \\ -1 & 1 & 1 \\ 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & -2 & 0 \\ -1 & 1 & 1 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right]$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = 3v_3$ and $v_2 = 2v_3$. Replacing v_3 with parameter s gives

$$\vec{v}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & -2 \\ -1 & 2 - \lambda & 1 \\ 0 & 1 & -1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\begin{bmatrix} 1 - (-1) & 1 & -2 \\ -1 & 2 - (-1) & 1 \\ 0 & 1 & -1 - (-1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\begin{bmatrix} 2 & 1 & -2 \\ -1 & 3 & 1 \\ 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 2 & 1 & -2 & 0 \\ -1 & 3 & 1 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right]$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 4

Let us find the eigenvalues and eigenvectors for

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

To find the eigenvector for $\lambda_3 = -1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = v_3$ and $v_2 = 0$. Replacing v_3 with parameter s gives

$$\vec{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Special Cases

Triangular Matrices: The eigenvalues of an upper (or lower) triangular matrix appear on the main diagonal.

Special Cases

Triangular Matrices: The eigenvalues of an upper (or lower) triangular matrix appear on the main diagonal.

2×2 Matrices: For

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The eigenvalues are the solutions to

$$\lambda^2 - (\text{tr } \mathbf{A})\lambda + |\mathbf{A}| = 0$$

Special Cases

Triangular Matrices: The eigenvalues of an upper (or lower) triangular matrix appear on the main diagonal.

2×2 Matrices: For

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

The eigenvalues are the solutions to

$$\lambda^2 - (\text{tr } \mathbf{A})\lambda + |\mathbf{A}| = 0$$

Trace

The **trace** of a matrix, **tr \mathbf{A}** , is the sum of all elements in the diagonal.

Eigenspace Theorem for Linear Transformations

For each eigenvalue λ of a linear transformations $T : \mathbb{V} \rightarrow \mathbb{V}$, the **eigenspace**, defined by

$$\mathbb{E}_\lambda = \{\vec{v} \in \mathbb{V} \mid T(\vec{v}) = \lambda\vec{v}\}$$

is a subspace of \mathbb{V} .


Eigenspace Theorem for Linear Transformations

For each eigenvalue λ of a linear transformations $T : \mathbb{V} \rightarrow \mathbb{V}$, the **eigenspace**, defined by

$$\mathbb{E}_\lambda = \{\vec{v} \in \mathbb{V} \mid T(\vec{v}) = \lambda \vec{v}\}$$

is a subspace of \mathbb{V} .

Example 5



Example 6

For the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

we had the following eigenvectors:

$$\begin{aligned} \lambda_1 = 2 \quad \vec{v}_1 &= \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \\ \lambda_2 = 1 \quad \vec{v}_2 &= \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \\ \lambda_3 = -1 \quad \vec{v}_3 &= \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

Example 6

For the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$$

we had the following eigenvectors:

$$\begin{array}{lll} \lambda_1 = 2 & \vec{v}_1 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} & \mathbb{E}_{\lambda_1} = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \right\} \\ \lambda_2 = 1 & \vec{v}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} & \mathbb{E}_{\lambda_2} = \text{span} \left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} \right\} \\ \lambda_3 = -1 & \vec{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} & \mathbb{E}_{\lambda_3} = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\} \end{array}$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\vec{v}_2 = c\vec{v}_1 \quad \text{where } c \neq 0$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\vec{v}_2 = c\vec{v}_1 \quad \text{where } c \neq 0$$

$$\lambda_2 \vec{v}_2 = c\lambda_2 \vec{v}_1$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

But, we could also have multiplied by \mathbf{A}

$$\mathbf{A}\vec{v}_2 = c\mathbf{A}\vec{v}_1$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

But, we could also have multiplied by \mathbf{A}

$$\begin{aligned}\mathbf{A}\vec{v}_2 &= c\mathbf{A}\vec{v}_1 \\ \lambda_2 \vec{v}_2 &= c\lambda_1 \vec{v}_1\end{aligned}$$

Distinct Eigenvalue Theorem

Let \mathbf{A} be an $n \times n$ matrix. If $\lambda_1, \lambda_2, \dots, \lambda_p$ are distinct eigenvalues with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p$, then $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_p\}$ is a set of linearly independent vectors.

Proof (sketch)

If we have two eigenvalues with $\lambda_1 \neq \lambda_2$, then if the associated eigenvectors \vec{v}_1 and \vec{v}_2 were linearly dependent, we have

$$\begin{aligned}\vec{v}_2 &= c\vec{v}_1 \quad \text{where } c \neq 0 \\ \lambda_2 \vec{v}_2 &= c\lambda_2 \vec{v}_1\end{aligned}$$

But, we could also have multiplied by \mathbf{A}

$$\begin{aligned}\mathbf{A}\vec{v}_2 &= c\mathbf{A}\vec{v}_1 \\ \lambda_2 \vec{v}_2 &= c\lambda_1 \vec{v}_1\end{aligned}$$

Which would imply that $\lambda_1 = \lambda_2$,

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| &= 0 \\ \lambda(\lambda + 3)^2 &= 0 \end{aligned}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| &= 0 \\ \lambda(\lambda + 3)^2 &= 0 \end{aligned}$$

So, the eigenvalues are $\lambda_1 = 0$, $\lambda_2 = -3$. (Note that -3 is a repeated root.)

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\begin{bmatrix} -2 - \lambda & 1 & 1 \\ 1 & -2 - \lambda & 1 \\ 1 & 1 & -2 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\begin{bmatrix} -2 - (0) & 1 & 1 \\ 1 & -2 - (0) & 1 \\ 1 & 1 & -2 - (0) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\left[\begin{array}{ccc|c} -2 & 1 & 1 & 0 \\ 1 & -2 & 1 & 0 \\ 1 & 1 & -2 & 0 \end{array} \right]$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_1 = 0$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = v_3$ and $v_2 = v_3$. Replacing v_3 with parameter s gives

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\begin{bmatrix} -2 - \lambda & 1 & 1 \\ 1 & -2 - \lambda & 1 \\ 1 & 1 & -2 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\begin{bmatrix} -2 - (-3) & 1 & 1 \\ 1 & -2 - (-3) & 1 \\ 1 & 1 & -2 - (-3) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \end{array} \right]$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

To find the eigenvector for $\lambda_2 = -3$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_1 = -v_2 - v_3$. This means we need two parameters, $v_2 = r$ and $v_3 = s$. Which means we have two linearly independent eigenvectors.

$$\vec{v}_2 = \begin{bmatrix} -r - s \\ 1 \\ 1 \end{bmatrix} = r \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + s \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

This means the eigenspace is

$$\mathbb{E}_{\lambda_2} = \mathbf{span} \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

which is a two-dimensional subspace of \mathbb{R}^3 .

Example 7

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{bmatrix}$$

This means the eigenspace is

$$\mathbb{E}_{\lambda_2} = \text{span} \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

which is a two-dimensional subspace of \mathbb{R}^3 .

Any linear combination of these two vectors is also an eigenvector, which means that the eigenspace is a plane.

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

Since this is an upper triangular matrix, we know that the eigenvalue is $\lambda = 1$, with multiplicity of 3.

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - \lambda & 1 & 1 \\ 0 & 1 - \lambda & 1 \\ 0 & 0 & 1 - \lambda \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\begin{bmatrix} 1 - (1) & 1 & 1 \\ 0 & 1 - (1) & 1 \\ 0 & 0 & 1 - (1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_2 + v_3 = 0$ and $v_3 = 0$. Replacing v_1 with parameter s gives

$$\vec{v} = \begin{bmatrix} s \\ 0 \\ 0 \end{bmatrix} = s \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Example 8

Consider the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

To find the eigenvector for $\lambda = 1$ we need to solve the system:

$$\left[\begin{array}{ccc|c} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

So, we have $v_2 + v_3 = 0$ and $v_3 = 0$. Replacing v_1 with parameter s gives

$$\vec{v} = \begin{bmatrix} s \\ 0 \\ 0 \end{bmatrix} = s \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Which means the eigenspace has dimension 1.

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0$$

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0$$

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

The only solutions to this equation is $\lambda = \pm i$.

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

The only solutions to this equation is $\lambda = \pm i$.

The situation here is like that with quadratic equations: there are some things which just cannot be done in a real-valued world.

Example 9

Consider

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$$

Which has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0 \quad \rightarrow \quad \lambda^2 + 1 = 0$$

The only solutions to this equation is $\lambda = \pm i$.

The situation here is like that with quadratic equations: there are some things which just cannot be done in a real-valued world.

We can continue in the same way and find that the eigenvectors are

$$\vec{\mathbf{v}}_1 = \begin{bmatrix} -i \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{\mathbf{v}}_2 = \begin{bmatrix} -1 \\ -i \end{bmatrix}$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad \rightarrow \quad \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \end{aligned}$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \end{aligned}$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \\ &\rightarrow \cos(\theta) - \lambda = \pm i \sin(\theta) \end{aligned}$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is

$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \\ &\rightarrow \cos(\theta) - \lambda = \pm i \sin(\theta) \\ &\rightarrow \lambda = \cos(\theta) \pm i \sin(\theta) \end{aligned}$$

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

The characteristic equation is


$$\begin{aligned} |\mathbf{A} - \lambda \mathbf{I}| = 0 &\rightarrow \begin{vmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{vmatrix} = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &\rightarrow (\cos(\theta) - \lambda)^2 = \sin^2(\theta) \\ &\rightarrow \cos(\theta) - \lambda = \pm i \sin(\theta) \\ &\rightarrow \lambda = \cos(\theta) \pm i \sin(\theta) \end{aligned}$$

Which means these eigenvalues rotate a vector, instead of scaling it.

Example 10

Consider the rotation transformation

$$\mathbf{A} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$



Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.
- \mathbf{A} has a zero eigenvalue if and only if $|\mathbf{A}| = 0$

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{v} = \vec{0}$ has a nontrivial solution.
- \mathbf{A} has a zero eigenvalue if and only if $|\mathbf{A}| = 0$
- \mathbf{A} and \mathbf{A}^T have the same characteristic polynomials and the same eigenvalues.

Eigenvalue Properties

Let \mathbf{A} be an $n \times n$ matrix

- λ is an eigenvalue of \mathbf{A} if and only if $|\mathbf{A} - \lambda \mathbf{I}| = 0$.
- λ is an eigenvalue of \mathbf{A} if and only if $(\mathbf{A} - \lambda \mathbf{I})\vec{\mathbf{v}} = \vec{\mathbf{0}}$ has a nontrivial solution.
- \mathbf{A} has a zero eigenvalue if and only if $|\mathbf{A}| = 0$
- \mathbf{A} and \mathbf{A}^T have the same characteristic polynomials and the same eigenvalues.
- If λ is an eigenvalue of an invertible matrix \mathbf{A} , then $\frac{1}{\lambda}$ is an eigenvalue of \mathbf{A}^{-1} .

Note

Characteristic roots of a linear homogeneous DEs are eigenvalues.

Note

Characteristic roots of a linear homogeneous DEs are eigenvalues.

Properties of Linear Homogeneous DEs with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

Note

Characteristic roots of a linear homogeneous DEs are eigenvalues.

Properties of Linear Homogeneous DEs with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.

Note

Characteristic roots of a linear homogeneous DEs are eigenvalues.

Properties of Linear Homogeneous DEs with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.
- The solution set is also a vector space of vector functions.

Note

Characteristic roots of a linear homogeneous DEs are eigenvalues.

Properties of Linear Homogeneous DEs with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.
- The solution set is also a vector space of vector functions.
- The eigenspace for each eigenvalue is a one-dimensional line in the direction of a vector in \mathbb{R}^n .

Note

Characteristic roots of a linear homogeneous DEs are eigenvalues.

Properties of Linear Homogeneous DEs with Distinct Eigenvalues

For the DE $\vec{x}' = \mathbf{A}\vec{x}$ with distinct eigenvalues, the following hold:

- The domain of the linear transformation is a vector space of vector functions.
- The solution set is also a vector space of vector functions.
- The eigenspace for each eigenvalue is a one-dimensional line in the direction of a vector in \mathbb{R}^n .

Note

We will explore the connection between eigenvalues and solutions to differential equations next chapter.