

S1211Q Introduction to Statistics



Lecture 4

Wei Wang

July 6, 2012

More Probability Properties

- Consider an experiment whose sample space is S . For each event A (B) in S , we assume that a number $P(A)$ is defined and satisfies the following rules:
 - $0 \leq P(A) \leq 1$.
 - $P(S)=1$.
 - $P(A^c)=1-P(A)$.
 - If A and B are disjoint, then $P(A \cup B)=P(A)+P(B)$.
 - For any two events A and B , $P(A \cup B)=P(A)+P(B)-P(A \cap B)$.



Example

Ex. A store accepts either VISA or Mastercard. 50% of the stores customers have VISA, 30% have Mastercard and 10% have both. What is the probability that a customer has a credit card the store accepts?


A = customers has VISA

B = customers has Mastercard

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= 0.5 + 0.3 - 0.1 = 0.7 \end{aligned}$$

Example cont.

What is the probability that a customer has either a VISA or MC, but not both?



$$\begin{aligned} P(\text{A or B but not both}) &= P(A) + P(B) - 2P(A \cap B) \\ &= 0.5 + 0.3 - 0.2 = 0.6 \end{aligned}$$


Example Cont.

What is the probability that a customer has a VISA but no MC?

$$\begin{aligned}P(\text{A but not both}) &= P(A) - P(A \cap B) \\ &= 0.5 - 0.1 = 0.4\end{aligned}$$

What is the probability that a customer has a MC but no VISA?


$$\begin{aligned}P(\text{B but not both}) &= P(B) - P(A \cap B) \\ &= 0.3 - 0.1 = 0.2\end{aligned}$$



Three Events

- For any three events A, B and C,

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$



- ▶ Recall that for discrete variables and all outcomes are equally likely to occur, the probability of event A is given by

$$P(A) = \frac{\text{Number of outcomes in } A}{\text{Number of outcome in Sample Space } S}$$

- ▶ Counting Techniques are essential to efficiently calculate the numerator. Specifically, we will talk about **Permutations** and **Combinations**.

Product Rule

- ▶ If the first object of an ordered pair can be selected in n_1 ways, and for each of these n_1 ways the second objects can be selected in n_2 ways, then the number of pairs is $n_1 n_2$
- ▶ This rule applies when we have multiple stages.
- ▶ Here the key is that the stages are independent of each other.

Permutations

- ▶ How many different ordered arrangements of the letters a , b and c are possible?

Permutations

- ▶ How many different ordered arrangements of the letters a , b and c are possible?
- ▶ By direct enumeration we see that there are 6, namely, abc , acb , bac , bca , cab and cba . Each arrangement is known as a *permutation*.

Permutations

- ▶ How many different ordered arrangements of the letters a , b and c are possible?
- ▶ By direct enumeration we see that there are 6, namely, abc , acb , bac , bca , cab and cba . Each arrangement is known as a *permutation*.
- ▶ Suppose now that we have n objects. Reasoning similar to that we have just used for the 3 letters then shows that there are

$$n(n - 1)(n - 2) \dots 3 \cdot 2 \cdot 1 = n!$$

different permutations of the n objects.

Example 1

- ▶ How many different batting orders are possible for a baseball team consisting of 9 players?

Example 1

- ▶ How many different batting orders are possible for a baseball team consisting of 9 players?
- ▶ **Solution:** There are $9! = 362880$ possible batting orders.

Example 2

- ▶ Ms. Davis has 10 books that she is going to put on her bookshelf. Of these, 4 are mathematics books, 3 are physics books, 2 are literature books and 1 is a language book. Ms. Mortimer wants to arrange her books so that all the books dealing with the same subject are together on her shelf. How many different arrangements are possible?

Example 2

- ▶ Ms. Davis has 10 books that she is going to put on her bookshelf. Of these, 4 are mathematics books, 3 are physics books, 2 are literature books and 1 is a language book. Ms. Mortimer wants to arrange her books so that all the books dealing with the same subject are together on her shelf. How many different arrangements are possible?
- ▶ **Solution:** There are $4!3!2!1!$ arrangements such that the mathematics books are first in line, then the physics books, then the history books, and then the language books. Similarly, for each possible ordering of the subjects, there are $4!3!2!1!$ possible arrangements. Hence, as there are $4!$ possible ordering of the subjects, the desired answer is $4!4!3!2!1! = 6912$.

Formal Definition of Permutations

- ▶ An ordered subset is called a permutation.

Formal Definition of Permutations

- ▶ An ordered subset is called a permutation.
- ▶ The number of permutations of size k that can be formed from n objects is denoted by $P_{k,n}$

Formal Definition of Permutations

- ▶ An ordered subset is called a permutation.
- ▶ The number of permutations of size k that can be formed from n objects is denoted by $P_{k,n}$
- ▶ $P_{k,n} = \frac{n!}{(n-k)!}$

Formal Definition of Permutations

- ▶ An ordered subset is called a permutation.
- ▶ The number of permutations of size k that can be formed from n objects is denoted by $P_{k,n}$
- ▶ $P_{k,n} = \frac{n!}{(n-k)!}$
- ▶ In particular, $P_{n,n} = n!$ and $P_{1,n} = n$

Combinations

- ▶ Combinations are similar to Permutations, except that order is irrelevant here.

Combinations

- ▶ Combinations are similar to Permutations, except that order is irrelevant here.
- ▶ A typical combination question is:
How many different groups of 3 could be selected from the 5 items *A*, *B*, *C*, *D* and *E*?

Combinations

- ▶ Combinations are similar to Permutations, except that order is irrelevant here.
- ▶ A typical combination question is:
How many different groups of 3 could be selected from the 5 items A, B, C, D and E ?
- ▶ There are thus $P_{3,5} = 5 \times 4 \times 3$ ways of selecting a group of 3 when the order in which the items are selected is relevant.

Combinations

- ▶ Combinations are similar to Permutations, except that order is irrelevant here.
- ▶ A typical combination question is:
How many different groups of 3 could be selected from the 5 items A, B, C, D and E ?
- ▶ There are thus $P_{3,5} = 5 \times 4 \times 3$ ways of selecting a group of 3 when the order in which the items are selected is relevant.
- ▶ But every group is counted $P_{3,3} = 3 \times 2 \times 1$ times. So there are

$$\frac{P_{3,5}}{3!} = 10$$

groups.

Formal Definition of Combinations

- ▶ An unordered subset is called a combination.

Formal Definition of Combinations

- ▶ An unordered subset is called a combination.
- ▶ The number of combinations of size k that can be formed from n objects is denoted by $C_{k,n}$ or $\binom{n}{k}$.

Formal Definition of Combinations

- ▶ An unordered subset is called a combination.
- ▶ The number of combinations of size k that can be formed from n objects is denoted by $C_{k,n}$ or $\binom{n}{k}$.
- ▶ $\binom{n}{k} = P_{k,n}/k!$

Formal Definition of Combinations

- ▶ An unordered subset is called a combination.
- ▶ The number of combinations of size k that can be formed from n objects is denoted by $C_{k,n}$ or $\binom{n}{k}$.
- ▶ $\binom{n}{k} = P_{k,n}/k!$
- ▶ In particular, $\binom{n}{n} = 1$ and $\binom{n}{1} = n$.

Example 1

- ▶ A committee of 3 is to be formed from a group of 20 people. How many different committees are possible?

Example 1

- ▶ A committee of 3 is to be formed from a group of 20 people. How many different committees are possible?
- ▶ **Solution:** There are $\binom{20}{3} = \frac{20 \times 19 \times 18}{3 \times 2 \times 1} = 1140$ possible committees.

Example 2

- ▶ From a group of 5 women and 7 men, how many different committees consisting of 2 women and 3 men can be formed?

Example 2

- ▶ From a group of 5 women and 7 men, how many different committees consisting of 2 women and 3 men can be formed?
- ▶ **Solution:** As there $\binom{5}{2}$ possible groups of 2 women and $\binom{7}{3}$ possible groups of 3 men, it follows that there are $\binom{5}{2} \binom{7}{3} = 350$ possible committees consisting of 2 women and 3 men.

Example 3, tricky

- ▶ In the last example, what if 2 of the men are feuding and refuse to serve on the committee together?

Example 3, tricky

- ▶ In the last example, what if 2 of the men are feuding and refuse to serve on the committee together?
- ▶ **Solution:** Now suppose that 2 of them refuse to serve together, because a total of $\binom{2}{2}\binom{5}{1}$ out of the $\binom{7}{3} = 35$ possible groups of 3 men contain both of the feuding men, it follows that there are $35 - 5 = 30$ groups that do not contain both of the feuding men. Because there are still $\binom{5}{2}$ ways to choose 2 women, there are $30 \cdot 10 = 300$ possible committees in this case.