

W1211 Introduction to Statistics

Lecture 13

Wei Wang

Oct 17, 2012

Independent rv's

- Recall the definition of independence of two random events A and B.

$$P(A \cap B) = P(A) P(B)$$

- We say two random variables X and Y are **independent** if and only if

$$P(X=x, Y=y) = P(X=x) P(Y=y), \text{ for any } x \text{ and } y.$$

- More specifically, two random variables X and Y are said to be independent if for every pair x and y values,

$$p(x, y) = p_X(x) p_Y(y), \text{ when } X \text{ and } Y \text{ are discrete;}$$

or

$$f(x, y) = f_X(x) f_Y(y), \text{ when } X \text{ and } Y \text{ are continuous.}$$

Ex. The second case of the previous example.

Multiple Random Variables

- If X_1, X_2, \dots, X_n are all discrete random variables, the joint pmf of the variables is the function

$$p(x_1, x_2, \dots, x_n) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

If the variables are continuous, the joint pdf of X_1, X_2, \dots, X_n is the function

$f(x_1, x_2, \dots, x_n)$ such that for any n intervals $[a_1, b_1], \dots, [a_n, b_n]$,

$$P(a_1 \leq X_1 \leq b_1, \dots, a_n \leq X_n \leq b_n) = \int_{a_1}^{b_1} \cdots \int_{a_n}^{b_n} f(x_1, \dots, x_n) dx_1 \dots dx_n$$

- What should be the regularity conditions for $p(x_1, x_2, \dots, x_n)$ and $f(x_1, x_2, \dots, x_n)$?
- How do get the marginal distributions of X_1, X_2, \dots by using $p(x_1, x_2, \dots, x_n)$ and $f(x_1, x_2, \dots, x_n)$?

Independence

- Proposition:

The random variables X_1, X_2, \dots, X_n , are said to be independent if for every subset $X_{i_1}, X_{i_2}, \dots, X_{i_k}$, of the variables (each pair, each triple, and so on), the joint pmf or pdf of the subset is equal to the product of the marginal pmf's or pdf's.

- $$p(x_1, x_2, \dots, x_n) = \prod_{i=1}^n p_{X_i}(x_i)$$

- $$f(x_1, x_2, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i)$$

Conditional dist.

- Using the marginal distributions, one can calculate the conditional distribution of one rv given the other.
- Let X and Y be two conditional rv's with joint pdf $f(x, y)$ and marginal X pdf $f_X(x)$. Then for any X value x for which $f_X(x) > 0$, the **conditional probability density function** of Y given that $X=x$ is

$$f_{Y|X}(y|x) = \frac{f(x, y)}{f_X(x)} \quad -\infty < y < \infty.$$

- If X and Y are discrete, replace pdf's by pmf's in this definition gives the **conditional probability mass function** of Y when $X=x$.

Expectation of Functions

- Recall how we compute $E[h(X)]$. A similar result also holds for a function $h(X, Y)$ of two jointly distributed rv's.
- Let X and Y be jointly distributed rv's with pmf $p(x, y)$, if they are discrete; or pdf $f(x, y)$, if they are continuous. The expected value of a function $h(X, Y)$, denoted by $E[h(X, Y)]$ is given by

$$E[h(X, Y)] = \begin{cases} \sum_x \sum_y h(x, y) \cdot p(x, y) & \text{if } X \text{ and } Y \text{ are discrete} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x, y) \cdot f(x, y) dx dy & \text{if } X \text{ and } Y \text{ are continuous} \end{cases}$$

- This result can also be extended to multiple (>2) rv case.

Expectation of Linear Function of Multiple RV's

- ▶ Linearity is well preserved in expectation.

$$E(a \cdot X + b \cdot Y + c) = a \cdot E(X) + b \cdot E(Y) + c$$

Expectation of Product of Multiple RV's

- ▶ Unlike the linear case, expectation of product in general doesn't equal to the product of expectations

$$E(XY) \neq E(X)E(Y)$$

Expectation of Product of Multiple RV's

- ▶ Unlike the linear case, expectation of product in general doesn't equal to the product of expectations

$$E(XY) \neq E(X)E(Y)$$

- ▶ But if X and Y are independent, then

$$\begin{aligned} E(XY) &= \int \int xyf(x, y)dx dy = \int \int xyf_X(x)f_Y(y)dx dy \\ &= \int xf_X(x)dx \int yf_Y(y)dy = E(X)E(Y) \end{aligned}$$

Expectation of Product of Multiple RV's

- ▶ Unlike the linear case, expectation of product in general doesn't equal to the product of expectations

$$E(XY) \neq E(X)E(Y)$$

- ▶ But if X and Y are independent, then

$$\begin{aligned} E(XY) &= \int \int xyf(x, y)dxdy = \int \int xyf_X(x)f_Y(y)dxdy \\ &= \int xf_X(x)dx \int yf_Y(y)dy = E(X)E(Y) \end{aligned}$$

- ▶ And for independent RV's, in general

$$E[g(X)h(Y)] = E[g(X)]E[h(Y)]$$

Covariance

- When two random variables X and Y are not independent, it is often of interest to assess how strongly they are related to one another.
- A popular measurement to characterize the dependence of two rv's is called **correlation**. To calculate correlation of two rv's, we'll have calculate the **covariance** of the two rv's.
- The **covariance** between two rv's X and Y is

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - \mu_X)(Y - \mu_Y)] \\ &= \begin{cases} \sum_x \sum_y (x - \mu_X)(y - \mu_Y) \cdot p(x, y) & X, Y \text{ discrete} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - \mu_X)(y - \mu_Y) \cdot f(x, y) dx dy & X, Y \text{ continuous} \end{cases}\end{aligned}$$

Short cut

- Proposition:

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$$

- What happens if we set $Y=X$?

Covariance and Variance

- ▶ As we can see, variance is a special case of covariance, where $X = Y$.

Covariance and Variance

- ▶ As we can see, variance is a special case of covariance, where $X = Y$.
- ▶ Variance of linear function of multiple RV's is given by

$$\text{Var}(aX + bY) = a^2 \text{Var}(X) + b^2 \text{Var}(Y) + 2ab \cdot \text{Cov}(X, Y)$$

Example

Ex. Suppose the joint distribution of X and Y are

$$f(x, y) = \begin{cases} 24xy & 0 \leq x \leq 1, 0 \leq y \leq 1, x + y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

What is the covariance of X and Y?

$$f_X(x) = \int_y f(x, y) dy = \int_0^{1-x} 24xy dy = 12x(1-x)^2$$

$$f_Y(y) = 12y(1-y)^2$$

$$E(X) = \int_0^1 x \cdot 12x(1-x)^2 dx = \frac{2}{5} = E(Y)$$

$$E(XY) = \int \int_{x,y} xy f(x, y) dx dy = \int_0^1 \int_0^{1-y} 24x^2 y^2 dx dy = \frac{2}{15}$$

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = \frac{2}{15} - \left(\frac{2}{5}\right)^2 = -\frac{2}{75}$$

Correlation

- The **correlation coefficient** of X and Y , denoted by $\text{Corr}(X, Y)$ or $\rho_{X,Y}$ is defined by

$$\rho_{X,Y} = \frac{\text{Cov}(X, Y)}{\sigma_X \cdot \sigma_Y}$$

- Because of Cauchy-Schwarz inequality, we have

$$\text{Cov}^2(X, Y) \leq \text{Var}(X)\text{Var}(Y) \implies |\rho_{X,Y}| \leq 1$$

- The correlation coefficient $\rho_{X,Y}$ is **NOT** a completely general measure of the strength of a relationship. $\rho_{X,Y}$ is actually a measure of the degree of **linear** relationship between X and Y .