

W1211 Introduction to Statistics

Lecture 19

Wei Wang

Nov 12, 2012

Point Estimation

- ▶ A point estimate of a parameter θ is a single number that can be regarded as a sensible value for θ . A **point estimate** is obtained by selecting a suitable statistic and computing its value from the given sample data. The selected statistic is called **the point estimator** of θ , denoted by $\hat{\theta}$.

Point Estimation

- ▶ A point estimate of a parameter θ is a single number that can be regarded as a sensible value for θ . A **point estimate** is obtained by selecting a suitable statistic and computing its value from the given sample data. The selected statistic is called **the point estimator** of θ , denoted by $\hat{\theta}$.
- ▶ For example, if we flip a biased coin n times and get X heads, and we want to estimate the probability of getting heads p , it is intuitive to use sample proportion(mean) X/n as the estimator. If we observe $n = 100$ and $X = 73$, then our estimate of p is 0.73.

Measure of Good Estimators

- ▶ First of all estimator is a random variable. In the biased coin example, we use the sample proportion X/n as our estimator of p . Can you say something about the distribution of \hat{p} ?

Measure of Good Estimators

- ▶ First of all estimator is a random variable. In the biased coin example, we use the sample proportion X/n as our estimator of p . Can you say something about the distribution of \hat{p} ?
- ▶ From the knowledge of sampling distribution, we know $E(\hat{p}) = p$ and $Var(\hat{p}) = p(1 - p)/n$. And if n is large, \hat{p} approximately follows a normal distribution.

Measure of Good Estimators

- ▶ First of all estimator is a random variable. In the biased coin example, we use the sample proportion X/n as our estimator of p . Can you say something about the distribution of \hat{p} ?
- ▶ From the knowledge of sampling distribution, we know $E(\hat{p}) = p$ and $Var(\hat{p}) = p(1 - p)/n$. And if n is large, \hat{p} approximately follows a normal distribution.
- ▶ In principle, there are many estimators for a given parameters. Two properties of estimators are desired.
 - ▶ **Unbiasedness** This is about how faithful the estimator is. A point estimator $\hat{\theta}$ is an unbiased estimator of θ if $E(\hat{\theta}) = \theta$ for every possible value of θ . If $\hat{\theta}$ is not unbiased, then $E(\hat{\theta}) - \theta$ is called the bias of $\hat{\theta}$.

Measure of Good Estimators

- ▶ First of all estimator is a random variable. In the biased coin example, we use the sample proportion X/n as our estimator of p . Can you say something about the distribution of \hat{p} ?
- ▶ From the knowledge of sampling distribution, we know $E(\hat{p}) = p$ and $Var(\hat{p}) = p(1 - p)/n$. And if n is large, \hat{p} approximately follows a normal distribution.
- ▶ In principle, there are many estimators for a given parameters. Two properties of estimators are desired.
 - ▶ **Unbiasedness** This is about how faithful the estimator is. A point estimator $\hat{\theta}$ is an unbiased estimator of θ if $E(\hat{\theta}) = \theta$ for every possible value of θ . If $\hat{\theta}$ is not unbiased, then $E(\hat{\theta}) - \theta$ is called the bias of $\hat{\theta}$.
 - ▶ **Small Variance** This is about how stable the estimator is. Unbiased estimators are faithful in the long run, but might have large fluctuation when sample size is small.

Principles of Selecting Estimators

- ▶ First, choose the estimators that are unbiased.
- ▶ Then, among the unbiased estimators, choose the one with the smallest variance.

Two General Results about Unbiasedness

- ▶ Let $X_1, X_2, X_3, \dots, X_n$ be a random sample from a distribution with mean μ and variance σ^2 , and if we use sample mean \bar{X} as the estimator of population mean μ , then $\hat{\mu} = \bar{X}$ is an unbiased estimator of μ

Two General Results about Unbiasedness

- ▶ Let $X_1, X_2, X_3, \dots, X_n$ be a random sample from a distribution with mean μ and variance σ^2 , and if we use sample mean \bar{X} as the estimator of population mean μ , then $\hat{\mu} = \bar{X}$ is an unbiased estimator of μ
- ▶ Let $X_1, X_2, X_3, \dots, X_n$ be a random sample from a distribution with mean μ and variance σ^2 , and if we use sample variance S^2 as the estimator of population variance σ^2 , then $\hat{\sigma}^2$ is an unbiased estimator of σ^2 , i.e.,

$$E\left(\frac{\sum (X_i - \bar{X})^2}{n - 1}\right) = \sigma^2$$

MVUE

- For unbiased estimators, what are their MSE's?

$$E(\hat{\theta} - \theta)^2 = E(\hat{\theta} - E(\hat{\theta}))^2 = \text{Var}(\hat{\theta})$$

- Among all estimators of θ that are unbiased, choose the one that has minimum variance. The resulting $\hat{\theta}$ is called the **minimum variance unbiased estimator (MVUE)** of θ .
- One needs more knowledge to actually identify if some estimator is really MVUE. But in a special case, we have the following theorem.

Let X_1, X_2, \dots, X_n be an i.i.d. sequence of random samples from a **normal distribution** with mean μ and σ . Then the estimator $\hat{\mu} = \bar{X}$ is the **MVUE** for μ .

The Standard Error

- When reporting a point estimator, one also reports the **standard error** associated with it.
- The **standard error** of an estimator $\hat{\theta}$ is its standard deviation $\sigma_{\hat{\theta}} = \sqrt{\text{Var}(\hat{\theta})}$. If the standard error itself involves unknown parameters whose values can be estimated, substitution of these estimates into $\sigma_{\hat{\theta}}$ yields the **estimated standard error** of the estimator, which we denote as $\hat{\sigma}_{\hat{\theta}}$.
- The associated standard error gives us an idea of how good/accurate the estimators are.

Estimator, Its Standard Error and Estimated Standard Error

- ▶ Now we are trying to estimate the probability of getting heads of a biased coin, so each flip X_i is a Bernoulli RV with parameter p , the estimator of parameter p is the sample mean/proportion

$$\hat{p} = \frac{\sum_{i=1}^n X_i}{n}$$

Estimator, Its Standard Error and Estimated Standard Error

- ▶ Now we are trying to estimate the probability of getting heads of a biased coin, so each flip X_i is a Bernoulli RV with parameter p , the estimator of parameter p is the sample mean/proportion

$$\hat{p} = \frac{\sum_{i=1}^n X_i}{n}$$

- ▶ If we flipped 100 times and observed 75 heads, then our estimate of p is

$$\hat{p} = \frac{75}{100} = 0.75$$

Estimator, Its Standard Error and Estimated Standard Error

- ▶ Now we are trying to estimate the probability of getting heads of a biased coin, so each flip X_i is a Bernoulli RV with parameter p , the estimator of parameter p is the sample mean/proportion

$$\hat{p} = \frac{\sum_{i=1}^n X_i}{n}$$

- ▶ If we flipped 100 times and observed 75 heads, then our estimate of p is

$$\hat{p} = \frac{75}{100} = 0.75$$

- ▶ Also, we need to report how good our estimator is through its Standard Error. This is also related to the Interval Estimation.

Estimator, Its Standard Error and Estimated Standard Error

- ▶ The standard error is $Var(\hat{p}) = \frac{p(1-p)}{n}$, but we cannot report it since we don't know what p is.
- ▶ So we can only report the estimated standard error of the estimator \hat{p}

$$\widehat{Var(\hat{p})} = \frac{\hat{p}(1 - \hat{p})}{n}$$

Another Example

- ▶ Now we have X_1, X_2, \dots, X_N IID with mean μ and variance σ^2 , what's the estimator of μ ?

Another Example

- ▶ Now we have X_1, X_2, \dots, X_N IID with mean μ and variance σ^2 , what's the estimator of μ ?

▶

$$\hat{\mu} = \bar{X}$$

Another Example

- ▶ Now we have X_1, X_2, \dots, X_N IID with mean μ and variance σ^2 , what's the estimator of μ ?

▶

$$\hat{\mu} = \bar{X}$$

- ▶ The standard error of $\hat{\mu}$ is $\sqrt{\text{Var}(\hat{\mu})} = \frac{\sigma}{\sqrt{n}}$. Can we report $\frac{\sigma}{\sqrt{n}}$?

Another Example

- ▶ Now we have X_1, X_2, \dots, X_N IID with mean μ and variance σ^2 , what's the estimator of μ ?

▶

$$\hat{\mu} = \bar{X}$$

- ▶ The standard error of $\hat{\mu}$ is $\sqrt{\text{Var}(\hat{\mu})} = \frac{\sigma}{\sqrt{n}}$. Can we report $\frac{\sigma}{\sqrt{n}}$?
- ▶ It really depends on whether or not we know σ . If we know it, then we can report $\frac{\sigma}{\sqrt{n}}$; otherwise, we can only report $\frac{\hat{\sigma}}{\sqrt{n}}$.