

P-Value

- To report the result of a hypothesis-testing analysis is to simply say whether the null hypothesis was rejected at a specified level of significance. This type of statement is somewhat inadequate because **it says nothing about whether the conclusion was a very close call or quite clear cut.**
- **P-value** is a quantity that conveys much information about the strength of evidence against H_0 and allows an individual decision maker to draw a conclusion at any specified level α .
- The **P-value** (*observed significance level*) is the probability, under the null hypothesis, that **the test statistic is more *extreme* than the observed statistic.**

What P-Values are not

- ▶ The P-value is not the probability that H_0 is true.
- ▶ The P-value is not Type I Error α .
- ▶ The P-value is not the significance level.
- ▶ The P-value is not Type II Error β

Comparison Between P-value and Type I Error α

- ▶ P-value = $P(\text{Test Statistic is more extreme than observed Test Statistic Value under Null Hypothesis})$
- ▶ Type I Error = $P(\text{Test Statistic falls into Rejection Region under Null Hypothesis})$

Remarks

- ▶ **The smaller the P-value, the more evidence there is in the sample data against the null hypothesis and for the alternative hypothesis.**
- ▶ P-values can be seen as a more flexible procedure of Hypothesis Testing. The practical advantage is that it is easier to switch to a test of different significance level
- ▶ The decision rule based on P-values

Decision rule based on the *P*-value


Select a significance level α (as before, the desired type I error probability).
Then

reject H_0 if $P\text{-value} \leq \alpha$
do not reject H_0 if $P\text{-value} > \alpha$

- ▶ The P-value is the smallest significance level α at which the null hypothesis can be rejected.

P-values and Tails

- Like Rejection Region, P-values are also related to the type of test we are concerning, upper-tailed, lower-tailed or two-tailed.



Two sample tests

- A new drug is claimed to significantly reduce the blood pressure for high blood pressure patients. What kind of tests can we use to verify the claim?
- A new drug is claimed to perform much better in terms of reducing blood pressure than an old drug. What kind of tests can we use to verify the claim?

Things to cover

- As in the one sample testing problem, we will cover the following cases:
 1. Two **normal** populations with **known** variance.
 2. Two populations with **unknown** distribution and **large sample** size.
 3. Two **normal** populations with **unknown** variance.
 4. Two population **proportions** with **large sample** size.
 5. Tests about variances. (NOT required.)
- Basic assumptions for comparing population means:
 1. X_1, X_2, \dots, X_m is a random sample (i.i.d.) from a population with mean μ_1 and variance σ_1^2 .
 2. Y_1, Y_2, \dots, Y_n is a random sample (i.i.d.) from a population with mean μ_2 and variance σ_2^2 .
 3. The X and Y samples are independent of one another.

Test statistics

- Since we are comparing the population means, a natural test statistic to use would be the difference of two sample means. Because of independence we have,

$$\begin{aligned}E(\bar{X} - \bar{Y}) &= \mu_1 - \mu_2 \\Var(\bar{X} - \bar{Y}) &= \frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}\end{aligned}$$

Case I: normal, known variance

$$H_0 : \mu_1 - \mu_2 = \Delta_0$$

$$\text{Test statistic: } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} \sim N(0,1)$$

vs Alternative Hypothesis:

$$H_a : \mu_1 - \mu_2 > \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} > Z_\alpha$$

$$H_a : \mu_1 - \mu_2 < \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} < -Z_\alpha$$

$$H_a : \mu_1 - \mu_2 \neq \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} < -Z_{\alpha/2} \text{ or } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n}}} > Z_{\alpha/2}$$

Questions

- How to compute P-value for case I?
- How to compute type II errors for case I?
- In a balanced design, derive the sample size calculation formula (for alternative “>”):

$$m = n = \frac{(\sigma_1^2 + \sigma_2^2)(Z_\alpha + Z_\beta)^2}{(\Delta' - \Delta_0)^2}$$

Case II: large sample

$$H_0 : \mu_1 - \mu_2 = \Delta_0$$

$$\text{Test statistic: } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} \sim \text{AN}(0,1)$$

vs Alternative Hypothesis:

$$H_a : \mu_1 - \mu_2 > \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} > Z_\alpha$$

$$H_a : \mu_1 - \mu_2 < \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} < -Z_\alpha$$

$$H_a : \mu_1 - \mu_2 \neq \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} < -Z_{\alpha/2} \text{ or } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} > Z_{\alpha/2}$$

Questions

- How to construct confidence interval for $\mu_1 - \mu_2$ in case II?

Case III: normal, unknown variance

$$H_0 : \mu_1 - \mu_2 = \Delta_0$$

Test statistic: $\frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} \sim t_\nu$, ν is the df of the t-distribution and it's approximately estimated

by the sampled data: $\nu = \frac{\left(\frac{s_1^2}{m} + \frac{s_2^2}{n}\right)^2}{\frac{(s_1^2 / m)^2}{m-1} + \frac{(s_2^2 / n)^2}{n-1}}$, and round ν down to the nearest integer.

Case III cont.

vs Alternative Hypothesis:

$$H_a : \mu_1 - \mu_2 > \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} > t_{\alpha, \nu}$$

$$H_a : \mu_1 - \mu_2 < \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} < -t_{\alpha, \nu}$$

$$H_a : \mu_1 - \mu_2 \neq \Delta_0, \text{ reject if } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} < -t_{\alpha/2, \nu} \text{ or } \frac{\bar{X} - \bar{Y} - \Delta_0}{\sqrt{\frac{s_1^2}{m} + \frac{s_2^2}{n}}} > t_{\alpha/2, \nu}$$

Questions

- How to compute P-values of the test?
- How to construct confidence interval for $\mu_1 - \mu_2$ in case III?
- What if we know that $\sigma_1^2 = \sigma_2^2$?

The *pooled estimator* of $\sigma^2 = \sigma_1^2 = \sigma_2^2$ is given by

$$S_p^2 = \frac{m-1}{m+n-2} \cdot S_1^2 + \frac{n-1}{m+n-2} \cdot S_2^2$$

Case IV

$$H_0 : p_1 - p_2 = 0$$

Test statistic: $\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{m} + \frac{1}{n}\right)}}$, $\hat{p} = \frac{m}{m+n}\hat{p}_1 + \frac{n}{m+n}\hat{p}_2$ (the *weighted* average of \hat{p}_1

and \hat{p}_2)

Case IV cont.

vs Alternative Hypothesis:

$$H_a : p_1 - p_2 > 0, \text{ reject if } \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{m} + \frac{1}{n}\right)}} > Z_{\alpha}$$

$$H_a : p_1 - p_2 < 0, \text{ reject if } \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{m} + \frac{1}{n}\right)}} < -Z_{\alpha}$$

$$H_a : p_1 - p_2 \neq 0, \text{ reject if } \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{m} + \frac{1}{n}\right)}} > Z_{\alpha/2} \text{ or } \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p})\left(\frac{1}{m} + \frac{1}{n}\right)}} < -Z_{\alpha/2}$$

Paired t-test

- As in the previous example, the data is paired, the two scores (before and after) recorded for each individual are **dependent**, but the between individuals the pairs are **independent**.
- Thus in order to test $H_0: \mu_1 - \mu_2 = 0$, one has to look at the difference of each pair. The problem eventually becomes a **one sample t-test problem**.