

(Chapter 38 P 316 - 319.)

Let  $n \geq 0$  be an integer. We introduce the following notation:

$$n! = (1)(2)(3) \cdots (n-1)(n).$$

Read:  $n$  factorial

Example:

$$0! = 1$$

$$1! = 1 = 1$$

$$2! = (1)(2) = 2$$

$$3! = (1)(2)(3) = 6$$

$$4! = (1)(2)(3)(4) = 24.$$

$$5! = (1)(2)(3)(4)(5) = 120.$$

In this lecture, we want to prove the following theorem.

Theorem (38.2 Binomial Theorem) The binomial coefficients in the expansion

$$(A+B)^n = \binom{n}{0} A^n + \binom{n}{1} A^{n-1} B + \cdots + \binom{n}{k} A^{n-k} B^k + \cdots + \binom{n}{n} B^n$$

are given by the formula:

$$\binom{n}{k} \stackrel{\textcircled{1}}{=} \frac{n(n-1) \cdots (n-k+1)}{k!} \stackrel{\textcircled{2}}{=} \frac{n!}{k!(n-k)!}$$

Example: (1)  $\binom{4}{2} = \frac{4!}{(2!)(2!)} = \frac{24}{(2)(2)} = 6$

(2)  $\binom{5}{2} = \frac{5!}{(2!)(3!)} = \frac{120}{(2)(6)} = 10$

(3)  $\binom{n}{0} = \frac{n!}{(0!)(n!)} = \frac{n!}{(1)(n!)} = \frac{n!}{n!} = 1$

(4)  $\binom{n}{n} = \frac{n!}{(n!)(0!)} = \frac{n!}{(n!)(1)} = \frac{n!}{n!} = 1.$

Proof of theorem:

We first prove ②:  $\frac{n(n-1)\cdots(n-k+1)}{k!} = \frac{n!}{k!(n-k)!}$

$$\frac{n(n-1)\cdots(n-k+1)}{k!} = \frac{n(n-1)\cdots(n-k+1)(n-k)(n-k-1)\cdots(2)(1)}{k! \cdot (n-k)(n-k-1)\cdots(2)(1)}$$

$$= \frac{n!}{k!(n-k)!} \quad \checkmark$$

Proof of ①: before the proof, we can look at an example:

we want to calculate  $\binom{4}{2}$

$\binom{4}{2}$  = coefficient of  $A^2B^2$  in  $(A+B)^4$ .

$$(A+B)^4 = (A+B)(A+B)(A+B)(A+B).$$

We want to look at  $A^2B^2$ . This means:

we can consider each parenthesis as a box  
(we have 4 boxes in total)

A, B      A, B      A, B      A, B

we want to pick up 2 "A" and 2 "B" in 4 boxes.

(Indeed, we only need to pick up 2 "A" since the left ones are "B")

To pick up one A, we have 4 choices  
(since we have 4 boxes.)

After pick up one A, we want to pick one A again,  
we have 3 choices.

(since we have 3 boxes left)

Therefore, we have  $(4)(3) = 12$  choices.

However, in this process, we overcounted:

Step I Pick up A in 1st box

Step II Pick up A in 3rd box

Step I' : Pick up A in 3rd box

Step II' : Pick up A in 1st box.

They are different when we are picking up A

but they give the same results for where we picked up A.

Indeed, in our process, the "order" of picking up A does not matter. That's why we overcounted.

Therefore, we need to divide the total ways by the order.

Since we are picking up 2 A, the number of order is  $2! = 2$ .

This implies: 
$$\binom{4}{2} = \frac{(4)(3)}{2!}$$

The proof of general case is similar:

$$\binom{n}{k} = \text{coefficient of } A^{n-k} B^k \text{ in } (A+B)^n$$



n boxes.

Pick up k "A"

Total number is  $= n(n-1)(n-2) \cdots (n-k+1)$ .

We need to divide by the order. Since we are picking up  $k$  "A", the number of order is  $k!$

Therefore 
$$\binom{n}{k} = \frac{n(n-1)(n-2) \cdots (n-k+1)}{k!} = \frac{n!}{k!(n-k)!} \quad \square.$$

Remark: The explicit formula also gives an proof for 
$$\binom{n}{k} = \binom{n}{n-k}.$$