

## Remerciements

*Ret e voe din dibab 'tre div vuhez,  
En-dro d'ar familh, en-dro d'an enorioù.  
Met n'eus ket a enor hep familh,  
Setu perak emañ ar gerioù-mañ.*

*Papa, Maman, me 'zo aet kuit  
War hent ar studioù, pell diouzh an ti,  
Pell diouzh ho karantez, pell diouzh ar c'hi,  
Hag ar c'hazh sioul tal an tan.*

*Kavet em eus ur garantez,  
Kollet em eus ur mignon,  
Gouelañ a ran noz-ha-deiz,  
Ne vo ken a «ken ar c'hentañ».*

*Soñjal a ran eus ma zud-kozh,  
D'an amzer a dremen,  
D'ar glav ha d'an amzer vat,  
O c'hortoz an deiz, o tremen hep bezañ.*

# Contents

|                                                                            |    |
|----------------------------------------------------------------------------|----|
| 1. Introduction (4-5 pages)                                                | 3  |
| 1.1. Biological and pharmacological context                                | 3  |
| 1.2. Lipid membranes as physicochemical systems                            | 3  |
| 1.3. Ligand–membrane coupling                                              | 5  |
| 1.4. Objectives                                                            | 6  |
| 2. Methods (3-4 pages)                                                     | 7  |
| 2.1. All-atom molecular dynamics                                           | 7  |
| 2.1.1. Membrane compositions, system preparation                           | 7  |
| 2.1.2. Force field and simulation parameters                               | 8  |
| 2.1.3. Umbrella sampling and PMF calculations                              | 9  |
| 2.1.4. Observables                                                         | 10 |
| 2.2. Coarse-grained simulations                                            | 11 |
| 2.3. Computational performances                                            | 11 |
| 3. Atomistic results: lipid dependent ligand-membrane coupling (5-6 pages) | 11 |
| 3.1. Insertion profiles, orientational behaviors                           | 11 |
| 3.2. Free energy profiles (PMF)                                            | 15 |
| 3.3. Structural and mechanical response                                    | 15 |
| 4. Extension to lipidomic context: Coarse-grained results (4-5 pages)      | 19 |
| 4.1. Multiscale consistency                                                | 19 |
| 4.2. Realistic membrane compositions                                       | 19 |
| 4.3. Lateral organisation and partitioning                                 | 20 |
| 5. Discussion (4-5 pages)                                                  | 20 |
| 5.1. Lipid composition as a modulator of ligand-membrane coupling          | 20 |
| 5.2. Implications for lipidomic alterations                                | 20 |
| 5.3. Methodological strengths and limitations                              | 20 |
| 5.4. Perspectives                                                          | 20 |
| 6. Conclusion (3 pages)                                                    | 20 |
| Bibliography                                                               | 21 |
| A. Glossary                                                                | 23 |

## **1. Introduction (4-5 pages)**

### **1.1. Biological and pharmacological context**

Cellular communication relies on molecular signals that must be detected and interpreted at the cell surface. In many cases, this process involves membrane receptors embedded in the plasma membrane, which translate extracellular or membrane-associated signals into intracellular responses. The membrane is therefore not only a physical boundary separating the cell from its environment, but also the cellular compartment in which many signaling events are initiated.


Traditional pharmacological frameworks implicitly assume that neurotransmitters diffuse through aqueous media before encountering their receptor [1]. This picture, while adequate for hydrophilic signaling molecules, becomes incomplete in the case of the endocannabinoid system (ECS) — a central regulator of neural activity, and a major pharmacological target in neurological and metabolic disorders. Endocannabinoids are lipid-derived molecules synthesized on demand from the hydrolysis of membrane phospholipids [2]. This biochemical origin confers lipophilic properties, meaning that these molecules preferentially partition into hydrophobic environments rather than the surrounding aqueous phase.

Among the main molecular targets of the ECS, the cannabinoid receptor type 1 (CB1) is one of the most abundantly expressed G protein-coupled receptors (GPCRs) in the mammalian central nervous system. Structural and pharmacological evidence suggests that access to its orthosteric binding pocket may occur laterally from within the bilayer rather than directly from the extracellular bulk [3], [4]. This pathway is relevant not only for endogenous cannabinoids such as anandamide and 2-arachidonoylglycerol (2-AG), but also for exogenous ligands targeting CB1 [5]. These include phytocannabinoids such as  $\Delta^9$ -tetrahydrocannabinol (THC), as well as synthetic compounds developed for pharmacological purposes, among which CP55,940 and WIN55.212-2 are potent agonists, and rimonabant (SR141716A) a well-known inverse agonist.

The presynaptic membrane environment, which constitutes an essential component of endocannabinoid signaling, is enriched in sterols, phospholipids, and sphingolipids [6], [7]. Lipidomic analyses have revealed significant compositional alterations under pathological conditions, including depletion of  $\omega$ -3 and  $\omega$ -6 polyunsaturated fatty acids (PUFAs) in patients experiencing a first psychotic episode [8], [9]. From a biophysical standpoint, such compositional shifts modify membrane properties including acyl chain order, interfacial hydration, bilayer thickness and bending rigidity — parameters that directly influence ligand partitioning and mobility within the bilayer. Understanding how membrane composition modulates these ligand-membrane interactions therefore appears essential for describing the early physicochemical steps preceding CB1 engagement.

### **1.2. Lipid membranes as physicochemical systems**

Lipid membranes are not homogeneous solvents but instead complex physicochemical systems, exhibiting rich phase behavior. This complexity is mainly brought by the diversity of their molecular constituents: headgroups, acyl chain saturation, and sterol content.



**Figure 1** — Schematic representation of phospholipid amphiphilicity and bilayer organization. Hydrophilic headgroups orient toward the aqueous phase, whereas hydrophobic acyl chains assemble into the membrane core, leading to the spontaneous formation of lipid bilayers. Due to its amphiphilic character and high lipophilicity, CP55,940 is expected to preferentially partition into this heterogeneous membrane environment rather than remain in the surrounding aqueous phase. Adapted from ... insérer ref depuis [ce lien](#).

The nomenclature of phospholipids reflects this structural organization. Species are typically designated by a four-letter code in which the first two letters indicate the acyl chains, while the last two specify the headgroup. For instance, POPC is for *1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine*, while SDPS denotes *1-stearoyl-2-docosahexaenoyl-sn-glycero-3-phospho-L-serine*.


Headgroups vary in size, charge distribution and hydrogen-bonding capability, thereby defining the interfacial character of the membrane [10]. These variations impose distinct geometric constraints at the molecular level, which propagate to the collective organization of the bilayer. *Phosphatidylcholines* (PC headgroup) exhibit an approximately cylindrical geometry favoring lamellar structures, whereas smaller headgroups such as *phosphatidylethanolamine* (PE) introduce packing asymmetry and curvature stress.

Beneath this interfacial layer lies the hydrophobic core, mainly determined by the composition and chemical structure of acyl chains, also known as fatty acids [11]. Chain length and degree of unsaturation constitute the two primary levers to modulate the core. For example, increasing unsaturation enhances conformational flexibility, *i.e.* *cis* double bonds introduce kinks that disrupt tight packing, hence reduce chain order and increase membrane fluidity.

Cell membranes also contain other essential lipids such as sterols and sphingomyelins. Among sterols, cholesterol plays a role in regulating membrane organization, fluidity and structural stability. Through their collective interactions, these different lipid species give rise to distinct membrane phase states. The phase diagram shown in Figure 2 corresponds to a model membrane composed of DPPC and cholesterol. The horizontal axis represents the cholesterol mole fraction, *i.e.* the proportion of cholesterol molecules relative to the total number of lipid molecules in the membrane mixture, while the vertical axis corresponds to temperature. Depending on these two parameters, the membrane may adopt different physical states, including the gel phase ( $L_\beta$ ), the liquid-disordered phase ( $L_\alpha$ , also commonly denoted  $L_d$ ), and the liquid-ordered phase ( $L_o$ ).

The gel phase ( $L_\beta$ ) corresponds to a highly ordered state in which lipid acyl chains are tightly packed, with reduced lateral mobility, resulting in a relatively rigid membrane. In contrast, the liquid-disordered phase ( $L_\alpha$  or  $L_d$ ) is characterized by lower chain ordering, enhanced confor-

mational flexibility and increased lateral diffusion. The liquid-ordered phase ( $L_o$ ), typically associated with cholesterol-rich membranes, combines features of both regimes by maintaining relatively ordered acyl chains while preserving substantial lipid mobility.



**Figure 2** — Phase diagram of DPPC-cholesterol mixture as a function of temperature.

Adapted from [12]. © Elsevier (2010), reproduced with permission.<sup>1</sup>

Broad coexistence regions also appear in the diagram, such as  $L_\alpha + L_o$  or  $L_\beta + L_o$ , indicating that different phases may coexist within the same membrane system depending on cholesterol concentration and temperature. Such coexistence reflects the heterogeneous nature of multi-component membranes, in which local lipid organization and dynamics strongly depend on composition and thermodynamic conditions. Although phase equilibria are temperature-dependent, physiological conditions typically constrain this parameter within a narrow range.

In multicomponent systems, compositional heterogeneity may emerge at larger length scales. Lipid-lipid interactions between different species can drive partial lateral demixing [ref heberle, ref florentin allemand], leading to the formation of distinct domains within the same bilayer. Often discussed in the context of lipid rafts, this mesoscale structure reflects the balance between entropic mixing and favorable intermolecular interactions, and results in regions exhibiting various mechanical properties.

### 1.3. Ligand-membrane coupling

While lipophilicity is often invoked to rationalize membrane affinity, a single partition coefficient ( $\log P$ ) cannot fully describe the complexity of ligand insertion into lipid bilayers, even though it is true that most CB1 ligands exhibit relatively high  $\log P$  values [ref si rien littérature, faire un tableau nous meme]. One reason is that the experimental determination of  $\log P$  typically

<sup>1</sup>License number n°6241360511955

relies on octanol–water partitioning [ref method], a simplified system that lacks the headgroup region — which defines a chemically distinct interfacial environment.

Another limitation is that logP constitutes a purely thermodynamic descriptor and therefore provides little insight into the microscopic mechanism of insertion. It does not inform on whether a ligand preferentially enters through hydrophobic regions, whether it should reorient upon insertion, or how it organizes once embedded within the bilayer. Generally, it is advised to compute potential of means force (PMFs) [ref] — that is, free-energy profiles evaluated along a chosen reaction coordinate — if one seeks better descriptions. PMFs provide access to the energetic cost associated with membrane entry, identify possible interfacial minima and reveal insertion barriers that could not be inferred from a global partition coefficient alone.

From this perspective, the membrane can be treated as an energy landscape experienced by the ligand. While such an approach is necessary to quantify insertion and identify potential barriers, it remains a reduced description of the interaction. The question arises: what is the physical nature of this interaction? A first level of consideration is steric. Much like inclusions in liquid crystals, an inserted ligand can be expected to behave as a local defect, perturbing lipid packing and altering the orientational order of neighboring acyl chains. Additional mechanical effects may also emerge; the ligand may locally displace lipids upon insertion, generating curvature stress or elastic deformation of the surrounding [ref].

Chemical and electrostatic contributions must also be considered. The membrane interface is characterized by a certain distribution of charges and dipoles, and screening effects are not necessarily uniform across the interfacial region. From a physical standpoint, the bilayer constitutes an electromagnetic environment, within which local interactions may further modulate ligand behavior.

Within this framework, CP55,940 emerges as a particularly relevant model ligand for CB1. Beyond its established pharmacological profile [ref] as a potent CB1 agonist, it presents physicochemical characteristics that make it especially suitable for investigating ligand–membrane coupling. It exhibits a relatively high partition coefficient, consistent with strong membrane affinity. Its molecular architecture combines an aromatic core and a hydrophobic aliphatic moiety, favoring insertion toward the bilayer interior, with additional polar functional groups capable of interacting with interfacial regions. Such amphiphilic balance makes it structurally compatible with prolonged residence within a bilayer while maintaining the ability to sample different insertion depths and orientations. It is therefore reasonable to treat CP55,940 as the molecular probe it has effectively become within the CB1 literature.

## **1.4. Objectives**

The present work seeks to characterize how ligand–membrane coupling evolves as a function of lipid composition, in the specific context of CB1 ligands.

We aim, on the one hand, to quantify how ligand insertion perturbs the local structural organization of the bilayer. On the other hand, we examine how the membrane itself constrains and modulates ligand behavior, in a setting where the bilayer may constitute a privileged access pathway to CB1. This study is therefore deliberately positioned upstream of explicit receptor simulations.

To address these questions, we perform molecular dynamics simulations at different level of resolutions. First, relying on all-atom simulations, we aim at characterizing ligand insertion and local ordering at high structural details – being sensitive to fine structural rearrangements. These simulations are conducted in compositionally simple membranes, within well-identified phase states, in order to isolate elementary physicochemical mechanisms.

We then extend the analysis to a coarse-grained level and to larger systems and longer timescales. This allows us to explore compositional effects at a broader scale and to assess whether local trends identified at the atomistic level persist in more complex membrane environments.

In particular, coarse-grained models make it possible to move beyond idealized binary mixtures and incorporate lipid compositions inspired by experimental lipidomics data. This is especially relevant for investigating variations in PUFAs content, whose imbalance has been reported in pathological contexts.

Accordingly, our main analytical tools consist of established membrane descriptors: membrane thickness, acyl chain order parameters, density profiles, PMFs, hydration profiles ... These indicators provide complementary structural and thermodynamic information and will be introduced and discussed systematically throughout the study. In line with this approach, we rely on a single ligand – CP55,940 – used as a probe in order to isolate membranotropic effects from ligand-specific variability.

Fundamentally, this work is guided by the idea that membrane composition must be regarded as an essential variable rather than a passive background. If the coupling is sensitive to lipid organization, then any variations in acyl chain saturation, sterol content, or PUFA balance may alter drug behavior in ways that cannot be captured by protein affinity alone. The broader perspective of this research is to eventually clarify to what extent the membrane itself participates in shaping pharmacological outcomes ; however, our current study remains focused on the mechanics of coupling.

## **2. Methods (3-4 pages)**

### **2.1. All-atom molecular dynamics**

#### **2.1.1. Membrane compositions, system preparation**

All-atom membrane systems were, for fluid systems, constructed using the CHARMM-GUI membrane builder [ref à mettre]. Gel-like membranes were built by hand from a Slipid pre-equilibrated patch [ref à slipid]. Symmetric bilayers composed of 100 lipids per leaflet were generated under periodic boundary conditions, resulting in lamellar multilayer systems separated by aqueous slabs.

Two reference phospholipids were primarily considered in order to explore distinct membrane physical states. DOPC bilayers were used to represent a fluid liquid-disordered phase at physiological temperature, while DPPC membranes were simulated at 25°C, a temperature well below the main phase transition temperature of DPPC ( $\sim 41^\circ\text{C}$ ). This ensures that the membrane remains in a stable gel-like phase during the simulations and avoids the strong fluctuations associated with the transition regime [refs]. Working with these two systems allows us to

contrast ligand behavior in environments characterized by markedly different packing and chain ordering properties.

In addition to these reference membranes, compositional perturbations were introduced by partially substituting the host lipids with polyunsaturated species. In practice, 10 % of the lipids were replaced by either SAPC ( $\omega$ -6) or SDPC ( $\omega$ -3), leading to six distinct membrane compositions in total. These substitutions introduce highly unsaturated acyl chains into otherwise well-defined bilayers and allow us to probe the influence of polyunsaturated lipids on ligand-membrane interactions.

Due to the planar geometry of the bilayer and the use of periodic boundary conditions, the simulated systems naturally adopt a stacked lamellar configuration. To avoid interactions between periodic images of adjacent membranes, a sufficiently large aqueous slab was introduced, resulting in a hydration level of approximately 45 water molecules per lipid [ref]. Sodium and chloride ions were added to reproduce an isotonic salt concentration (0.15 M NaCl) representative of physiological conditions.

System preparation followed the standard equilibration procedure provided by CHARMM-GUI. After energy minimization, a restrained NVT equilibration was performed for 500 ps, followed by a restrained NPT equilibration of 1 ns using Berendsen pressure coupling. The production equilibration runs were then performed for 100–300 ns, depending on membrane composition, to allow the bilayers to reach equilibrium before further analysis. This was especially important for gel-phase membranes, which exhibit slower lipid dynamics and therefore require longer equilibration times [ref].

The ligand CP55,940 was introduced at concentrations reaching up to 10 mol% relative to the lipid content, after equilibration of membranes. To ease spontaneous insertion while avoiding artificial aggregation in the aqueous phase, ligand molecules were initially positioned close to the membrane surface. A pulling restraint along the membrane normal was applied during the early stages of equilibration in order to keep the ligands near the bilayer interface. After ligand insertion, an additional system relaxation step was performed. The systems were first subjected to energy minimization in order to remove possible steric clashes introduced during ligand placement. A short equilibration stage was then carried out before starting the production simulations.

### **2.1.2. Force field and simulation parameters**

All-atoms molecular dynamics simulations were performed using GROMACS 2018.2. Interactions were described using the CHARMM36m force field, which provides a well-established parametrization for lipid membranes and is widely used for phospholipid bilayer simulations [13].

Alternative membrane force fields exist, including so-called Berger model for lipids. However, CHARMM36m is known to correctly treat polar functional groups such as hydroxyl (-OH) moieties, which are relevant for describing ligand-membrane interactions involving hydrogen bonding at the membrane interface.

Ligand parameters for CP55,940 were generated using CGenFF, through the ligand reader & modeler module of CHARMM-GUI [14]. Water molecules were represented using the TIP3P



model, consistent with the CHARMM parametrization. This model is commonly employed in membrane simulations and remains computationally efficient compared to more complex water models due to its lower number of degrees of freedom.

The equations of motion were integrated using a 2 fs timestep, with all bonds involving hydrogen atoms constrained using the LINCS algorithm. Long-range electrostatic interactions were treated using the particle mesh Ewald (PME) method, with a real-space cutoff of 12 Å. The same cutoff was applied to short-range van der Waals interactions.

Temperature was controlled using a Nosé–Hoover thermostat, while pressure was maintained using a Parrinello–Rahman barostat with semi-isotropic pressure coupling, allowing independent fluctuations in the membrane plane and along the bilayer normal. The pressure coupling time constant was set to 5 ps.

### 2.1.3. Umbrella sampling and PMF calculations

**NOTES** Vérifier les valeurs des constantes dans le dernier paragraphe.

To characterize ligand insertion beyond the simple partition coefficient, we considered potentials of mean force along a chosen reaction coordinate. In the present case, this coordinate was defined as the projection of the ligand center of mass along the membrane normal (z-axis), which provides a natural description of the insertion process.

From statistical mechanics, the PMF corresponds to a free-energy profile obtained by projecting the phase space distribution onto this reduced reaction coordinate. If  $x$  denotes a microscopic configuration of the system, and if  $U(x)$  is its associated potential energy, the canonical probability distribution is given by the Boltzmann law,

$$P(x) = \frac{1}{Z} e^{-\beta U(x)}$$

where  $\beta = 1/k_B T$  and  $Z$  is the partition function. Then, the probability associated with a given value of the reaction coordinate  $\xi$  is obtained by integrating out all remaining degrees of freedom, yielding,

$$P(\xi) = \frac{1}{Z} \int dx e^{-\beta U(x)} \delta(\xi - \xi(x))$$

with  $\delta$  the Dirac distribution. Finally, the corresponding free-energy profile is defined, up to an additive constant, as

$$F(\xi) = -k_B T \ln P(\xi) + \text{constant}$$

Broadly speaking, this quantity can be interpreted as an effective free-energy obtained after averaging over all microscopic configurations compatible with a given position of the ligand along the membrane normal.

In that sense, the free-energy profile can be reconstructed by varying the reaction coordinate of the ligand and, at each fixed position, estimating the associated probability distribution while allowing the ligand to explore the remaining degrees of freedom. Repeating this procedure over a series of overlapping windows gives access to the full profile. This procedure is known as

umbrella sampling, and the ligand is maintained around a given value of the reaction coordinate by means of a harmonic biasing potential.

In practice, initial configurations for these windows were generated using *steered molecular dynamics* (SMD). After equilibration of the membrane–ligand system, the ligand was gradually pulled along the membrane normal, starting from the aqueous phase toward the bilayer center.

Then, 35 configurations were extracted along this trajectory and then used as starting points for the umbrella sampling simulations, of which each was running for 100ns. The distributions obtained in each window were subsequently combined using the weighted histogram analysis method (WHAM) in order to reconstruct the unbiased PMF, since the harmonic potential introduces a bias.

In the present work, PMF calculations were restricted to DOPC membranes (64 lipids per leaflet) using GROMACS 2026.1. This choice was motivated by their fluid nature, which ensures sufficient molecular mobility and facilitates convergence of the free–energy profile along the chosen reaction coordinate. During steered molecular dynamics, the ligand was pulled along the membrane normal using a harmonic biasing potential, with a force constant of approximately  $400\text{kJ mol}^{-1}\text{nm}^{-2}$  and a constant pulling rate of  $1 \times 10^{-4}\text{nm ps}^{-1}$ .

#### **2.1.4. Observables**

We characterized the structural and thermodynamic properties of the membrane using a set of standard observables, commonly employed in the field, probing the global state of the membrane and its local response to ligand insertion.

The first observable is the so-called area per lipid (APL), defined as the average surface occupied by a lipid in the membrane plane. APL is particularly sensitive to membrane composition and phase state, and provides a direct measure of lipid packing in a given leaflet. It also serves as a practical indicator of equilibration, as reference values for pure DOPC and DPPC membranes are well documented in the literature [ref]. The area per lipid was computed using a Voronoi tessellation of the membrane plane [ref dans lipophilic]. Each lipid was assigned an effective area corresponding to its Voronoi cell, constructed from the lateral positions of lipid headgroups. The APL was then obtained by averaging over all lipids within a leaflet.

The membrane thickness was measured as the average distance between the phosphate groups of the two opposing leaflets. Variations in thickness may reflect both compositional effects and asymmetry induced by local perturbations. Density profiles were evaluated by the use of histogram of atomic positions along a given axis, providing the average spatial distribution of molecular species and allowing the ligand position within the bilayer to be identified.

Hydration profiles were computed by counting the number of water molecules in the vicinity of the polar headgroup, using a distance-based cutoff, typically of the order of the order of  $3.5\text{ \AA}$  — usually used for studying structuring waters within GPCR. Hydration provides a measure of solvent penetration at the membrane interface and is particularly relevant for characterizing this region, where polar interactions and hydrogen bonding play a significant role.

The orientational order of lipid acyl chains was also quantified, using the deuterium order parameter. This quantity is a direct measure of chain alignements with respect to the membrane normal and is defined as,

$$S_{CH} = \frac{1}{2} \langle 3 \cos^2 \theta - 1 \rangle$$

where  $\theta$  is the angle between a given C-H bond vector and the normal, and where  $\langle \cdot \rangle$  denotes a statistical average over time and lipids of the same species. The order parameter was computed for each carbon atom along the acyl chains using gorder [ref], resulting in a position-dependent profile that reflects the variation of chain ordering from the headgroup region to the bilayer core. It is often understood that a higher value corresponds to a more rigid (more ordered) system, whereas values closer to zero indicate a more fluid system. However, this interpretation should be treated with caution, since the parameter only reflects an angular measurement.

Beyond these observables, molecular dynamics simulations can still be viewed as *in silico* experiments<sup>2</sup> and the trajectories provides additional qualitative insight into ligand behavior, such as insertion pathways or fine details of molecular arrangements.

## 2.2. Coarse-grained simulations

- Préciser la version de gromacs
- Martini
- Stratégie de mapping
- S'inspirer de données de lipidomiques
- Modulation d'omega 6 et de médicament
- Observables, validation

## 2.3. Computational performances

- Jean-Zay, Mesocentre
- Taille des systemes
- ns/day performance
- GPU/CPU usage
- coût total en temps de calcul
- Comment être reproductible? les replica


# 3. Atomistic results: lipid dependent ligand-membrane coupling (5-6 pages)

## 3.1. Insertion profiles, orientational behaviors

In order to characterize the location of CP55,940 within the bilayer, density profiles were computed along the membrane normal for the different molecular groups — composing the ligand, and the lipids.

---

<sup>2</sup>Unfortunately, “simulation” has become increasingly misused to mean nothing more than “calculation.” – William L. Jorgensen [ref]




**Figure 3** — Density profiles along the membrane normal for CP55,940 moieties and representative lipid groups in (a) DOPC, (b) DPPC, (c) DPPC+SAPC (9:1), (d) DPPC+SDPC (9:1) membranes. The cyclohexanol, phenol and alkyl-chain profiles correspond to the ligand, and corresponding densities are multiplied by 10 for better visualization.

As shown in Figure 3, in pure DOPC layer (a), the ligand displays a preferential location within the interfacial region of the membrane, as its aromatic moieties remain predominantly localized near the glycerol backbone of the lipids. The aliphatic chain penetrates a bit deeper into the membrane – it tends to align with the surrounding chains, instead of completely folding on itself. This is expected for an amphiphilic insertion mode. We also observed a symmetric behavior upon enrichment with SAPC or SDPC lipids, suggesting that ligand localization is well preserved in these more disordered membranes.

A different behavior is reported in DPPC-containing systems (b-d). In contrast with the simple interfacial distribution in DOPC, the ligand density profiles exhibit multiple maxima extending deeper into the membrane. In particular, all the moieties display a bimodal distribution, with one population remaining close to glycerol and another penetrating significantly further into the core. This may be due to the coexistence of several metastable insertion states, possibly arising from slow relaxation dynamics imposed by more the ordered environment, leading to incomplete sampling on molecular dynamics timescales and contributing to the observed bimodal distribution.


Figure 4 illustrates the orientational distributions of CP55,940 aromatic core relative to the membrane normal. Fluid DOPC membrane (b) exhibits a relatively broad distribution centered around intermediate tilt angles (35–55°). In this configuration, the aromatic core remains close to the polar heads, allowing the hydroxyl groups to act as interfacial anchors in a hydrated region of the membrane.



**Figure 4** — Orientational distributions of the aromatic core of CP55,940 relative to the membrane normal in (b) DOPC, (c) DPPC and (d) DPPC+SDPC (9:1) membranes. (a) Schematic representation of the tilt-angle definition.

Compared with DOPC, DPPC membranes (c-d) display less defined distributions, without a clearly preferred tilt angle, suggesting a more heterogeneous orientational landscape with multiple nearly equivalent configurations. This broader distribution may reflect a higher packing frustration and a more constrained interfacial environment, where the ligand orientation is less collectively organized and more sensitive to local membrane structure.


This interpretation is supported by Figure 5, where we report orientational distributions of the alkyl chain of the ligand. The DOPC membrane (b) shows an almost isotropic distribution, with no preferential tilt angle over a wide angular range. This behavior suggests a highly fluid and permissive environment, where the ligand retains orientational freedom and can explore multiple configurations. In contrast, the DPPC system (c) display a marked low-angle preference (20°), indicative of a more tightly packed surrounding that constrains the alkyl chain. An intermediate behavior is observed for DPPC+SDPC (d), supporting the idea that lipid disorder progressively restores flexibility and reduce confinement effects.



**Figure 5** — Orientational distributions of the alkyl chain of CP55,940 relative to the membrane normal in (b) DOPC, (c) DPPC and (d) DPPC+SDPC (9:1) membranes.  
(a) Schematic representation of the tilt-angle definition.

Taken together, these observations suggest that the insertion behavior of CP55,940 results from a balance between chemical affinity for the membrane interface and steric constraints imposed by the lipid matrix. In fluid condition, the ligand appears to be primarily guided by favorable chemical interactions, as the hydroxyl groups remain anchored near the hydrated interfacial region while the rest of the molecule retains conformational freedom. This picture is consistent with previous experimental NMR results obtained for CP55,940 in POPC membranes, a fluid phosphatidylcholine bilayer close to DOPC, where the ligand was also reported to adopt an interfacial insertion mode [ref]. It contrasts with ordered membranes, where DPPC imposes stronger packing constraints, leading to broader insertion distributions and less well-defined orientational preferences. In these systems, local membrane organization likely competes with the intrinsic amphiphilic character of the ligand in determining its insertion mode. The introduction of unsaturations progressively relaxes these constraints, partially restoring the configurational flexibility and interfacial behavior.

### 3.2. Free energy profiles (PMF)



**Figure 6** — Potential of mean force computed for a CP55,940 ligand approaching a DOPC membrane in all-atoms. An indicative illustration is set in the background to visualize the leaflet.


Having established from insertion profiles that CP55,940 adopts preferentially an interfacial insertion mode in fluid DOPC membranes, we next quantified the energetic cost associated to it. The free-energy profile shown in Figure 6 was reconstructed by taking the aqueous phase as the reference state.

Starting from bulk water, the PMF rapidly decreases as the ligand approaches the interface, indicating a thermodynamic driving force for membrane insertion, confirming that the membrane constitutes a favorable environment for the ligand. Interestingly, a small energy barrier is observed close to the water-lipid interface ; it likely corresponds to the energetic cost associated with crossing the phosphate headgroup region and locally perturbing lipid packing in order to access the hydrophobic core of the bilayer.

Once this barrier is crossed, the free-energy decreases toward a pronounced minimum located inside the membrane, in good agreement with the density profiles and orientational analyses. Finally, the PMF rises again when approaching the bilayer center, usually depleted in lipids. In practice, this free-energy increase should act as a barrier against spontaneous flip-flop events on accessible simulation timescales.

### 3.3. Structural and mechanical response

We next examined how the membrane itself responds to ligand incorporation. To this end, we analyzed the area per lipid, the deuterium order parameter and the local hydration environment of the glycerol, which together should provide complementary information on membrane packing, ordering and interfacial organization. In mixed membranes, order parameter analyses were restricted to the host lipids (DOPC or DPPC). Indeed, the relatively small number of SAPC or SDPC molecules in the simulated patches did not allow statistically robust profiles to be extracted for the polyunsaturated species themselves.




**Figure 7** — Area per lipid (APL) measured for DOPC-based and DPPC-based membranes in the absence (control) and presence of CP55,940. For mixed systems, the reported values correspond to the average area per host lipid within the membrane plane.

The first important thing to note is that the control systems reproduce the expected physical trends associated with membrane composition and phase state. As shown in Figure 7 and Figure 9, DPPC-based membranes display smaller APL values together with substantially higher order parameters than DOPC systems (Figure 8), reflecting the tightly packed and more ordered nature of the gel phase. DPPC membranes enriched with SAPC or SDPC display though slightly higher order parameters than pure DPPC. At first sight, this may appear counterintuitive, since the introduction of unsaturated chains is generally associated with increased disorder. However, this observation should be interpreted carefully. In the gel phase, DPPC acyl chains are strongly tilted with respect to the membrane normal, while remaining very straight, as suggested by the flat-looking curve. Still, since the order parameter only probes the angle formed by C–H bonds relative to the normal, such collective tilt can artificially reduce the measured  $S_{CH}$  despite the membrane remaining rigid. The introduction of polyunsaturated lipids likely shifts the membrane away from the gel organization toward a more fluid but still ordered state, possibly closer to a liquid-ordered regime. In this configuration, the average chain tilt relative to the membrane normal is reduced, which may contribute to the larger apparent order parameters observed here. We also noted that the profiles become more rounded and less flat, suggesting that part of the extreme packing characteristic of the gel phase is progressively lost, together with some restored fluidity. Conversely, DOPC bilayers exhibit larger APL values and lower order parameters, in agreement with their liquid-disordered nature at physiological temperature. The addition of unsaturations further shifts the systems toward more disordered configurations.




Overall, these observations are consistent with the expected hierarchy between the different membrane compositions and therefore support the physical coherence of the simulated systems.



**Figure 8** — Deuterium order parameter profiles ( $S_{CH}$ ) computed for DOPC acyl chains in pure DOPC, DOPC+SAPC (9:1) and DOPC+SDPC (9:1) membranes, in the absence (control) and presence of CP55,940. The upper and lower panels correspond respectively to the sn-1 and sn-2 chains of DOPC. Order parameters were computed only for the host DOPC lipids.

In this background, the effect of CP55,940 appears to be dependent on the initial membrane organization. In all systems, ligand insertion induces an increase in area per lipid, indicating a global lateral expansion — depacking — of the bilayer. This effect remains moderate in fluid DOPC membranes but becomes more pronounced in DPPC systems. From a physical perspective, this behavior is compatible with the ligand acting as a local inclusion within the membrane, perturbing the packing and generating additional free volume around the insertion region. This expansion is not systematically accompanied by a decrease in chain ordering. In DOPC membranes, the deuterium parameter profiles remain almost unchanged upon ligand insertion, suggesting that the fluid bilayer can accommodate the perturbation without major structural reorganization. By contrast, DPPC-based membranes display a noticeable increase in chain ordering in the presence of the ligand, which we interpret as a trace of packing rearrangement around the ligand, where the membrane reorganizes in order to compensate for the perturbation introduced by insertion.


Hydration profiles further support the observed differences between the various membrane compositions. As represented in Figure 10, DOPC-based membranes exhibit systematically larger hydration levels around the phosphate group than DPPC systems. The introduction of polyunsaturated lipids increases hydration in both membrane families, an effect that is particularly pronounced in DPPC systems. This behavior is coherent with the increase in area per lipid and the partial loss of the highly packed gel organization discussed previously. The insertion of unsaturated chains likely introduces local packing defects and additional free volume, thereby facilitating water penetration toward the interfacial region.



**Figure 9** — Deuterium order parameter profiles ( $S_{CH}$ ) computed for DPPC acyl chains in pure DPPC, DPPC+SAPC (9:1) and DPPC+SDPC (9:1) membranes, for both control systems and membranes containing CP55,940. The upper and lower panels correspond respectively to the sn-1 and sn-2 chains of DPPC. Order parameters were evaluated only on the host DPPC lipids.

The presence of CP55,940 also shifts the distributions toward larger hydration values. This effect remains moderate in fluid DOPC membranes but becomes much more visible in DPPC systems. Altogether, these observations support the idea that both polyunsaturated lipids and ligand insertion contribute to relaxing the compact organization of ordered membranes and promote a more hydrated interfacial environment.

For clarity, all hydration distributions were normalized by the number of host lipids (DOPC or DPPC), allowing direct comparison between the different membrane compositions despite the presence of mixed systems.



**Figure 10** — Hydration distributions computed around the phosphate moieties of the host lipids for (a) DOPC and DOPC+SAPC (9:1) membranes, (b) DOPC and DOPC+SDPC (9:1) membranes, (c) DPPC and DPPC+SAPC (9:1) membranes, and (d) DPPC and DPPC+SDPC (9:1) membranes, in the absence and presence of CP55,940. The distributions are normalized by the number of host lipids (DOPC or DPPC).

## 4. Extension to lipidomic context: Coarse-grained results (4-5 pages)

À voir où on met le multiscale consistency. En premier j'oserais dire.

### 4.1. Multiscale consistency

- Comparison between atomistics and CG results
- Conserved vs scale-dependent features
- PMF.

### 4.2. Realistic membrane compositions

- w6 modulation
- cholesterol content
- headgroups

### **4.3. Lateral organisation and partitioning**

- Ligand spatial distribution
- Phase preference
- Demixing, rafts, collective membrane behavior

## **5. Discussion (4-5 pages)**

### **5.1. Lipid composition as a modulator of ligand-membrane coupling**

### **5.2. Implications for lipidomic alterations**

- w6 depletion & potential pharmacological consequences
- Positioning within broader scientific landscape

### **5.3. Methodological strengths and limitations**

- Absence of receptor
- Sampling limitations
- Force field consideration

### **5.4. Perspectives**

- Inclusion of CB1 in future work
- Multi ligand comparison
- Ongoing collaboration with CBMN

## **6. Conclusion (3 pages)**

## Bibliography

- [1] J. Ritter, R. J. Flower, G. Henderson, Y. K. Loke, D. J. MacEwan, and H. P. Rang, *Rang and Dale's pharmacology*, Ninth edition. Edinburgh: Elsevier, 2020.
- [2] L. Venance, R. Maldonado, and O. Manzoni, “Le système endocannabinoïde central,” *médecine/sciences*, vol. 20, no. 1, pp. 45–53, Jan. 2004, doi: [10.1051/medsci/200420145](https://doi.org/10.1051/medsci/200420145).
- [3] D. P. Hurst *et al.*, “A Lipid Pathway for Ligand Binding Is Necessary for a Cannabinoid G Protein-coupled Receptor,” *Journal of Biological Chemistry*, vol. 285, no. 23, pp. 17954–17964, Jun. 2010, doi: [10.1074/jbc.M109.041590](https://doi.org/10.1074/jbc.M109.041590).
- [4] J. Jakowiecki, U. Orzeł, S. Chawananon, P. Misztal, and S. Filipek, “The Hydrophobic Ligands Entry and Exit from the GPCR Binding Site-SMD and SuMD Simulations,” *Molecules*, vol. 25, no. 8, p. 1930, Apr. 2020, doi: [10.3390/molecules25081930](https://doi.org/10.3390/molecules25081930).
- [5] G. Thakur, S. Nikas, and A. Makriyannis, “CB1 Cannabinoid Receptor Ligands,” *Mini-Reviews in Medicinal Chemistry*, vol. 5, no. 7, pp. 631–640, Jul. 2005, doi: [10.2174/1389557054368772](https://doi.org/10.2174/1389557054368772).
- [6] R. Ahrends, S. R. Ellis, S. H. Verhelst, and M. R. Kreutz, “Synaptoneurolipidomics: lipidomics in the study of synaptic function,” *Trends in Biochemical Sciences*, vol. 50, no. 2, pp. 156–170, Feb. 2025, doi: [10.1016/j.tibs.2024.12.004](https://doi.org/10.1016/j.tibs.2024.12.004).
- [7] E. Lauwers, R. Goodchild, and P. Verstreken, “Membrane Lipids in Presynaptic Function and Disease,” *Neuron*, vol. 90, no. 1, pp. 11–25, Apr. 2016, doi: [10.1016/j.neuron.2016.02.033](https://doi.org/10.1016/j.neuron.2016.02.033).
- [8] A. T. P. Le *et al.*, “Analysis of polyunsaturated fatty acids in antipsychotic-free individuals with at-risk mental state and patients with first-episode schizophrenia,” *Frontiers in Psychiatry*, vol. 14, p. 1188452, Jul. 2023, doi: [10.3389/fpsy.2023.1188452](https://doi.org/10.3389/fpsy.2023.1188452).
- [9] A. Frajerman *et al.*, “Lipides membranaires dans la schizophrénie et la psychose débutive : de potentiels biomarqueurs et pistes thérapeutiques ?,” *L'Encéphale*, vol. 46, no. 3, pp. 209–216, Jun. 2020, doi: [10.1016/j.encep.2019.11.009](https://doi.org/10.1016/j.encep.2019.11.009).
- [10] K.-H. Kim and B. C. Yoo, “Decoding Membrane Lipids: Analytical Barriers and Technological Advances in Modern Lipidomics,” *International Journal of Molecular Sciences*, vol. 27, no. 3, p. 1472, Feb. 2026, doi: [10.3390/ijms27031472](https://doi.org/10.3390/ijms27031472).
- [11] O. Ali and A. Szabó, “Review of Eukaryote Cellular Membrane Lipid Composition, with Special Attention to the Fatty Acids,” *International Journal of Molecular Sciences*, vol. 24, no. 21, p. 15693, Oct. 2023, doi: [10.3390/ijms242115693](https://doi.org/10.3390/ijms242115693).
- [12] D. Marsh, “Liquid-ordered phases induced by cholesterol: A compendium of binary phase diagrams,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1798, no. 3, pp. 688–699, Mar. 2010, doi: [10.1016/j.bbamem.2009.12.027](https://doi.org/10.1016/j.bbamem.2009.12.027).
- [13] Y. Yu and J. B. Klauda, “Update of the CHARMM36 United Atom Chain Model for Hydrocarbons and Phospholipids,” *The Journal of Physical Chemistry B*, vol. 124, no. 31, pp. 6797–6812, Aug. 2020, doi: [10.1021/acs.jpcc.0c04795](https://doi.org/10.1021/acs.jpcc.0c04795).

- [14] S. Kim, J. Lee, S. Jo, C. L. Brooks, H. S. Lee, and W. Im, “CHARMM-GUI ligand reader and modeler for CHARMM force field generation of small molecules: CHARMM-GUI Ligand Reader and Modeler for CHARMM Force Field Generation of Small Molecules,” *Journal of Computational Chemistry*, vol. 38, no. 21, pp. 1879–1886, Jun. 2017, doi: [10.1002/jcc.24829](https://doi.org/10.1002/jcc.24829).

## **A. Glossary**

- ECS
- CB1
- GPCR
- PUFA
- affinity
- potency
- efficacy
- lipid
- omega 3, omega 6
- scaffold