

Reinforcement **Learning** for Business, Economics, and Social Sciences


Unit 2-1: Markov Processes

Davud Rostam-Afschar (Uni Mannheim)

How to predict transitions?

Markov Chains

Unrolling the Problem



Goal: Learn to choose actions that maximize rewards

Unrolling the Problem

- ▶ Modeling environment dynamics
- ▶ Unrolling the control loop leads to a sequence of states, actions and rewards:

$$s_0, a_0, r_0, s_1, a_1, r_1, s_2, a_2, r_2, \dots$$

- ▶ This sequence forms a stochastic process (due to some uncertainty in the dynamics of the process)

Common Properties

- ▶ Processes are rarely arbitrary
- ▶ They often exhibit some structure
 - ▶ Laws of the process do not change
 - ▶ Short history sufficient to predict future

Common Properties

- ▶ Processes are rarely arbitrary
- ▶ They often exhibit some structure
 - ▶ Laws of the process do not change
 - ▶ Short history sufficient to predict future
- ▶ **Example:** weather prediction
 - ▶ Same model can be used everyday to predict weather
 - ▶ Weather measurements of past few days sufficient to predict weather


Common Properties

- ▶ Processes are rarely arbitrary
- ▶ They often exhibit some structure
 - ▶ Laws of the process do not change
 - ▶ Short history sufficient to predict future
- ▶ **Example:** weather prediction
 - ▶ Same model can be used everyday to predict weather
 - ▶ Weather measurements of past few days sufficient to predict weather
- ▶ **Example:** text prediction
 - ▶ Same model can be used in every conversation to predict next utterance
 - ▶ letter sequences of past texts sufficient to predict new sentences

Markovian and Stationary Processes

Stochastic Process

- ▶ Consider the sequence of states only
- ▶ Definition
 - ▶ Set of States: S
 - ▶ Stochastic dynamics: $\mathbb{P}(s_t | s_{t-1}, \dots, s_0)$



Stochastic Process

- ▶ Problem:
 - ▶ Infinitely large conditional distributions
- ▶ Solutions:
 - ▶ **Stationary process:**
Dynamics do not change over time
 - ▶ **Markov assumption:**
Current state depends only on a finite history of past states
 - ▶ Russell and Norvig (2016, Section 15.1)


K-Order Markov Process

- ▶ Assumption: last k states sufficient

K-Order Markov Process

- ▶ Assumption: last k states sufficient
- ▶ First-order Markov Process

- ▶ $\mathbb{P}(s_t | s_{t-1}, \dots, s_0) = \mathbb{P}(s_t | s_{t-1})$




K-Order Markov Process

- ▶ Assumption: last k states sufficient


- ▶ First-order Markov Process

- ▶ $\mathbb{P}(s_t | s_{t-1}, \dots, s_0) = \mathbb{P}(s_t | s_{t-1})$



- ▶ Second-order Markov Process

- ▶ $\mathbb{P}(s_t | s_{t-1}, \dots, s_0) = \mathbb{P}(s_t | s_{t-1}, s_{t-2})$



Markov Process

- ▶ Commonly, a Markov Process refers to a
 - ▶ First-order process

$$\mathbb{P}(s_t \mid s_{t-1}, s_{t-2}, \dots, s_0) = \mathbb{P}(s_t \mid s_{t-1}) \forall t$$

- ▶ Stationary process

$$\mathbb{P}(s_t \mid s_{t-1}) = \mathbb{P}(s_{t'} \mid s_{t'-1}) \forall t'$$


- ▶ **Advantage:**
can specify the entire process with a single concise conditional distribution

$$\mathbb{P}(s' \mid s)$$

Examples



Examples

- ▶ Marrying decision of young women
 - ▶ **States:** relationship history
 - ▶ **Dynamics:** age



Examples


- ▶ Marrying decision of young women
 - ▶ **States:** relationship history
 - ▶ **Dynamics:** age
- ▶ Robotic control
 - ▶ **States:** $\langle x, y, z, \theta \rangle$
coordinates of joints
 - ▶ **Dynamics:** constant motion



Examples


► Marrying decision of young women

- **States:** relationship history
- **Dynamics:** age



► Robotic control

- **States:** $\langle x, y, z, \theta \rangle$
coordinates of joints
- **Dynamics:** constant motion



► Inventory management

- **States:** inventory level
- **Dynamics:** constant (stochastic) demand



Inference in Markov Processes

- ▶ Common task is prediction: $\mathbb{P}(s_{t+k} \mid s_t)$
- ▶ Computation:

$$\mathbb{P}(s_{t+k} \mid s_t) = \sum s_{t+k} \dots s_{t+k-1} \prod_{i=1}^k \mathbb{P}(s_{t+i} \mid s_{t+i-1})$$

- ▶ Discrete states (matrix operations):
 - ▶ Let T be a $|S| \times |S|$ matrix representing $\mathbb{P}(s_{t+k} \mid s_t)$
 - ▶ Then $\mathbb{P}(s_{t+k} \mid s_t) = T^k$
 - ▶ Complexity: $\mathcal{O}(k|S|^3)$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

		never married	married
$T =$	never married	0.5	0.5
	married	0	1

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

	never married	married
never married	0.5	0.5
married	0	1

 $\rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:

$$\begin{array}{c} \text{Year } k \\ \hline p_{t+k} = p_t T^k \end{array}$$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:

$$\begin{array}{c|c} \text{Year } k & p_{t+k} = p_t T^k \\ \hline 1 & [0.250000 \ 0.750000] \end{array}$$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:

Year k	$p_{t+k} = p_t T^k$
1	$[0.250000 \ 0.750000]$
2	$[0.125000 \ 0.875000]$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:

Year k	$p_{t+k} = p_t T^k$
1	$[0.250000 \ 0.750000]$
2	$[0.125000 \ 0.875000]$
3	$[0.062500 \ 0.937500]$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:

Year k	$p_{t+k} = p_t T^k$
1	$[0.250000 \ 0.750000]$
2	$[0.125000 \ 0.875000]$
3	$[0.062500 \ 0.937500]$
4	$[0.031250 \ 0.968750]$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \ 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:

Year k	$p_{t+k} = p_t T^k$
1	$[0.250000 \ 0.750000]$
2	$[0.125000 \ 0.875000]$
3	$[0.062500 \ 0.937500]$
4	$[0.031250 \ 0.968750]$
5	$[0.015625 \ 0.984375]$

Example: Marrying as 2-State Markov Process

Setup: Initial distribution $p_t = [0.5_{\text{never married}} \quad 0.5_{\text{married}}]$

$$T = \begin{array}{c|cc} & \text{never married} & \text{married} \\ \hline \text{never married} & 0.5 & 0.5 \\ \text{married} & 0 & 1 \end{array} \rightarrow T^2 = \begin{pmatrix} 0.25 & 0.75 \\ 0 & 1 \end{pmatrix}, \dots$$

Predicted Distributions:


Year k	$p_{t+k} = p_t T^k$
1	$[0.250000 \quad 0.750000]$
2	$[0.125000 \quad 0.875000]$
3	$[0.062500 \quad 0.937500]$
4	$[0.031250 \quad 0.968750]$
5	$[0.015625 \quad 0.984375]$

Long Run:

$$\pi = \lim_{k \rightarrow \infty} p_{t+k} = [0 \quad 1] \quad (\text{everyone eventually marries})$$

How Quickly Get Young Women Married?

Probability of Young Women born 1950 of Being Never Married
US National Longitudinal Survey (1968-1988)



```
xtsteadystate nev_mar if birth_yr ==50, tw 3dists ini ss pred twowayopt(.)
```

Non-Markovian and/or Non-Stationary Processes

- ▶ What if the process is not Markovian and/or not stationary?
- ▶ Solution: add new state components until dynamics are Markovian and stationary

Non-Markovian and/or Non-Stationary Processes

- ▶ What if the process is not Markovian and/or not stationary?
- ▶ Solution: add new state components until dynamics are Markovian and stationary
 - ▶ Marrying: probability of marrying may depend on: How long a woman has been single, her past relationship history, norms in 1970 vs. 1980
 - ▶ Add time since last relationship, number of prior marriages, cohort, ...

Non-Markovian and/or Non-Stationary Processes

- ▶ What if the process is not Markovian and/or not stationary?
- ▶ Solution: add new state components until dynamics are Markovian and stationary
 - ▶ Marrying: probability of marrying may depend on: How long a woman has been single, her past relationship history, norms in 1970 vs. 1980
 - ▶ Add time since last relationship, number of prior marriages, cohort, ...
 - ▶ Where do we stop?

Markovian Stationary Process

- ▶ **Problem:** adding components to the state description to force a process to be Markovian and stationary may significantly increase computational complexity
- ▶ **Solution:** try to find the smallest state description that is self-sufficient (i.e., Markovian and stationary)

Decision Making

- ▶ Predictions by themselves are useless
- ▶ They are only useful when they will influence future decisions
- ▶ Hence the ultimate task is decision making
- ▶ How can we influence the process to visit desirable states?
 - ▶ Model: Markov Decision Process

References I

RUSSELL, S. J., AND P. NORVIG (2016): *Artificial intelligence: a modern approach*. Pearson.

Takeaways

How Can we Use Markov Processes to Predict Future States?

- ▶ Model sequences of states with probabilistic transitions
 - ▶ First-order Markov and stationarity assumptions simplify prediction
 - ▶ Adding state components can restore Markovian/stationary properties—at a computational cost
 - ▶ Prediction relies on transition matrices
 - ▶ Real goal: use predictions for decision-making
- Markov Decision Processes

Appendix

Prediction and Steady State via Eigendecomposition

Objective: Predict future state distributions $\mathbb{P}(s_{t+k} \mid s_t)$ and compute the steady-state distribution using eigendecomposition

Inputs:

- ▶ Initial distribution: p_t
- ▶ Transition matrix: T where $T_{ij} = \mathbb{P}(s_{t+1} = j \mid s_t = i)$
- ▶ Horizon: k (number of steps ahead)

Procedure:

1. Eigendecompose: $T = U\Lambda U^{-1}$
2. Compute predicted distribution:

$$p_{t+k} = T^k p_t = U\Lambda^k U^{-1} p_t$$

3. Steady state distribution:

$$\pi = \lim_{k \rightarrow \infty} p_{t+k}$$